

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Faculty Publications: Department of Entomology

Entomology, Department of

2016

Predominantly Left-Dominant Mandibular Chirality In Coleoptera

Mathew L. Brust

Chadron State College, mbrust@csc.edu

W. Wyatt Hoback

University of Nebraska at Kearney, whoback@okstate.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

Brust, Mathew L. and Hoback, W. Wyatt, "Predominantly Left-Dominant Mandibular Chirality In Coleoptera" (2016). *Faculty Publications: Department of Entomology*. 476.

<https://digitalcommons.unl.edu/entomologyfacpub/476>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

PREDOMINANTLY LEFT-DOMINANT MANDIBULAR CHIRALITY IN COLEOPTERA

MATHEW L. BRUST
Department of Biology
Chadron State College, 1000 Main St.
Chadron, NE 69337, U.S.A.
mbrust@csc.edu

AND

W. WYATT HOBACK
Department of Entomology and Plant Pathology
Oklahoma State University, 127 Noble Research Center
Stillwater, OK 74078, U.S.A.

ABSTRACT

Patterns of mandible chirality have been virtually unexplored in beetles, apart from a single study in 2010. Here we present the mandible chirality trends found in 58 beetle species representing the families Carabidae (52 species), Cerambycidae (three species), and Silphidae (three species) that display overlapping mandibles. Mandible chirality was not random in the groups examined, all of which exhibited a dominant trend towards left-superiority. The degree to which each taxonomic group displayed the left-superior trend differed, with tiger beetles showing the greatest dominance (99%) and longhorned beetles the weakest (71%).

Key Words: morphology, mouthparts, mandible, “handedness”, dentition, Carabidae, Cerambycidae, Silphidae

Beetles display enormous diversity in feeding habits, but all possess chewing mouthparts (Triplehorn and Johnson 2005) adapted to different feeding methods. For example, some beetles feed by clipping bits of vegetation. In such cases, mandibles that meet directly are common. Other beetles, particularly predaceous species, often have mandibles that overlap rather than meet directly. In addition to feeding, many male beetles utilize their mandibles to grasp females during mating (Triplehorn and Johnson 2005). Thus, at least two major factors likely affect selection pressure in regard to mandible alignment in beetles.

Despite selection for different mandible orientation, very little research on this exists for insects in general. Crowson (1981) mentions that in most beetles the tip of one mandible overlaps the other, but does not present information about which crosses which. Richardson (2010) showed a strong chirality trend in the tiger beetles *Omus audouini* Reiche and *Omus dejeani* Reiche, both of which usually exhibited left-superiority, *i.e.*, the left mandible crossing over the right mandible. We are not aware of any other published study on this subject in beetles. The benefit of overlapping mandibles in predaceous beetles is unclear. However, the mandibles in such species also tend to be slightly curved inward at the tips, have prominent dentition, or both. The mechanism of mandible cross-

ing also has not been examined in detail, nor has the ability of individuals to control which mandible crosses over the other. Nevertheless, overlap of mandibles may help to minimize wear (Richardson 2010) and improve the maceration efficiency. Mandible overlap may also be important in mating of species such as tiger beetles where males guard mates by grasping the female metathoracic copulatory sulcus with the mandible (Freitag 1974). Sexual dimorphism of tiger beetle mandibles has been documented (*e.g.*, Kritsky and Simon 1995), but mandible overlap was not examined.

Museum specimens allow an opportunity to examine mandible overlap. Preliminary examination of hundreds of specimens within the tiger beetle genus *Cicindela* Linnaeus (*sensu lato*) showed a pattern consistent with the findings of Richardson (2010). Thus, we hypothesized that mandibular chirality in predaceous beetles is not random and is consistent within taxonomic groups.

MATERIAL AND METHODS

Data were gathered from the insect collection at Chadron State College as well as the personal tiger beetle collection of the first author. In larger species and those with large mandibles (*i.e.*, most tiger beetles), mandibular chirality, defined as crossing of the tips of the mandibles, was determined

Table 1. List of taxa examined, including the number of specimens showing left-superiority in mandible chirality (Left) or right-superiority (Right), the total number of each species examined (*n*), and the proportion of each species' cohort showing left-superiority in mandible chirality.

Carabidae: Cicindelinae	Left	Right	<i>n</i>	Proportion left-dominant
<i>Amblycheila cylindriciformis</i> Say	22	0	22	1.0000
<i>Omus californicus</i> Eschscholtz	35	0	35	1.0000
<i>Omus audouinii</i> Reiche	6	0	6	1.0000
<i>Omus submetallicus</i> G. Horn	2	0	2	1.0000
<i>Omus dejeani</i> Reiche	9	0	9	1.0000
<i>Tetracha virginica</i> (Linnaeus)	51	0	51	1.0000
<i>Tetracha carolina</i> (Linnaeus)	34	0	34	1.0000
<i>Chaetodera regalis</i> (Dejean)	12	0	12	1.0000
<i>Cicindela sylvatica</i> Linnaeus	14	0	14	1.0000
<i>Cicindela granulata</i> Gebler	7	0	7	1.0000
<i>Cicindela soluta</i> Dejean	16	0	16	1.0000
<i>Sophiodela chinensis</i> DeGeer	7	0	7	1.0000
<i>Cicindela nebraskana</i> Casey, <i>C. longilabris</i> Say, and hybrids	330	0	330	1.0000
<i>Cicindela repanda</i> Dejean	683	1	684	0.9985
<i>Cicindela duodecimguttata</i> Dejean	170	0	170	1.0000
<i>Cicindela oregona</i> Dejean	111	0	111	1.0000
<i>Cicindela hirticollis</i> Say	357	0	357	1.0000
<i>Cicindela limbata</i> Say	266	0	266	1.0000
<i>Cicindela bellissima</i> Leng	9	0	9	1.0000
<i>Cicindela formosa</i> Say	1,061	1	1,062	0.9991
<i>Cicindela purpurea</i> Olivier	741	0	741	1.0000
<i>Cicindela splendida</i> Hentz	490	1	491	0.9980
<i>Cicindela denverensis</i> Casey	379	0	379	1.0000
<i>Cicindela limbalis</i> Klug	561	0	561	1.0000
<i>Cicindela decemnotata</i> Say	45	0	45	1.0000
<i>Cicindela sexguttata</i> Fabricius	596	0	596	1.0000
<i>Cicindela denikei</i> Brown	107	0	107	1.0000
<i>Cicindela patruela</i> Dejean	183	0	183	1.0000
<i>Cicindela pulchra</i> Say	168	0	168	1.0000
<i>Cicindela fulgida</i> Say	343	0	343	1.0000
<i>Cicindela parowana</i> Wickham	9	0	9	1.0000
<i>Cicindela scutellaris</i> Say	934	1	935	0.9989
<i>Cicindela tranquebarica</i> (Herbst)	608	0	608	1.0000
<i>Cicindela ancocisconensis</i> (T. W. Harris)	11	0	11	1.0000
<i>Cicindela lengi</i> (Horn)	264	0	264	1.0000
<i>Cicindelidia willistoni</i> (LeConte)	82	0	82	1.0000
<i>Cicindelidia schauppii</i> (G. H. Horn)	9	0	9	1.0000
<i>Cicindelidia punctulata</i> (Olivier)	780	1	781	0.9987
<i>Cicindelidia obsoleta</i> (Say)	24	0	24	1.0000
<i>Cicindelidia ocellata</i> (Klug)	23	0	23	1.0000
<i>Cicindelidia sedecimpunctata</i> (Klug)	15	0	15	1.0000
<i>Habrosclimorpha circumpecta</i> (LaFerté-Sénéctère)	148	0	148	1.0000
<i>Cylindera terricola</i> (Say)	116	0	116	1.0000
<i>Ellipsoptera waplery</i> (LeConte)	10	0	10	1.0000
<i>Ellipsoptera cuprascens</i> (LeConte)	66	0	66	1.0000
<i>Dromochorus pruina</i> (Casey)	3	0	3	1.0000
Carabidae: non-Cicindelinae				
<i>Pasimachus elongatus</i> LeConte	31	0	31	1.0000
<i>Calosoma luxatum</i> Say	7	1	8	0.8750
<i>Calosoma callidum</i> (Fabricius)	32	12	44	0.7273
<i>Harpalus pennsylvanicus</i> (DeGeer)	5	0	5	1.0000
<i>Chlaenius</i> sp. (purple)	13	3	16	0.8125
<i>Chlaenius</i> sp. (green)	4	0	4	1.0000

Continued on next page

Table 1. Continued.

	Left	Right	<i>n</i>	Proportion left-dominant
Silphidae				
<i>Nicrophorus marginatus</i> Fabricius	169	11	180	0.9389
<i>Nicrophorus carolinus</i> (Linnaeus)	229	18	247	0.9271
<i>Nicrophorus obscurus</i> Kirby	5	0	5	1.0000
Cerambycidae				
<i>Tetraopes femoratus</i> LeConte	16	4	20	0.8000
<i>Tetraopes tetraphthalmus</i> (Forster)	37	24	61	0.6066
<i>Tetraopes annulatus</i> LeConte	24	3	27	0.8889

through naked eye observation, otherwise chirality was determined using a dissecting scope. A total of 10,575 specimens representing 58 species in the beetle families Carabidae (52 species, 46 being tiger beetles (subfamily Cicindelinae)), Silphidae (three species of *Nicrophorus* Fabricius), and Cerambycidae (three species of *Tetraopes* Dalman) were examined in this study. For each specimen, the chirality was observed and recorded based on which mandible was superior, thus, each specimen was recorded as either right-superior or left-superior. Specimens with mandibles open were not included.

RESULTS

Mandibular chirality was found to be non-random in all beetle groups examined, with all species and groups exhibiting a strong trend toward left-superior chirality (Table 1). However, the dominance of left-superiority differed among beetle groups. For example, all but five of the 9,922 tiger beetle specimens examined displayed left-superiority; the five specimens not displaying left-superiority each represented a single individual of different species. Thus, 41 of 46 tiger beetle species exhibited 100% left-superiority. In contrast, left-superiority in non-tiger beetle Carabidae averaged 90% (range = 73–100%) (Table 2). Left-superiority in *Nicrophorus* spp. (Silphidae) averaged 96% (range = 93–100%), while in *Tetraopes* spp. (Cerambycidae) it averaged only 77% (range = 61–89%).

DISCUSSION

The trends observed in mandibular chirality in this study suggest that 1) mandibular chirality in beetles with overlapping mandibles is not random and 2) left-superior chirality is dominant but to differing degrees among beetle groups. In tiger beetles, the left-superior state was observed in more than 99% of specimens examined, including more

than 9,900 specimens representing 46 species in 11 genera. Left-superiority was also dominant in non-tiger beetle Carabidae (average 90%), Silphidae (average 96%), and Cerambycidae (77%). These three beetle families are distantly related to each other and represent two different suborders, suggesting that left-dominant mandibular chirality may be common throughout the order. Differences in left-superiority could relate to sex, though it was untested in this study and would be expected to result in close to 50:50 ratios. None of the right-superior specimens were noticeably teneral.

The results of our study suggest that mandibular chirality in beetles with overlapping mandibles may be a fixed genetic character that is not controlled by the individual beetle. Additional questions that warrant further investigation include: 1) whether the left-superior trait is a conserved trait that has only evolved once in beetles or has evolved independently more than once; 2) which genes are involved in mandibular chirality and whether they influence other physical traits, most significantly mandibular asymmetry; 3) what other physical modifications are influenced by mandibular chirality, for example, whether one mandible is positioned higher than the other or the head also exhibits asymmetry; 4) if the degree of mandibular dentition or curvature is correlated with degree of mandibular chirality; and 5) what is the interaction in selection pressures on mandible

Table 2. Proportion of specimens per broad taxonomic group showing left-superiority in mandible chirality. *n* = number of specimens examined within each group (*n*).

Group	Proportion left-dominant	<i>n</i>
Carabidae: Cicindelinae	0.9994	9,922
Carabidae: non-Cicindelinae	0.8518	108
Silphidae	0.9329	432
Cerambycidae	0.7130	108

shape for feeding in both sexes versus mate grasping by males.

ACKNOWLEDGMENTS

We thank Chadron State College for their support and Oklahoma State University Department of Entomology and Plant Pathology, and the Oklahoma Agricultural Experiment Station for providing resources. Drs. Bruce Noden and Rick Grantham provided helpful comments on an earlier version of this manuscript.

REFERENCES CITED

- Crowson, R. A. 1981.** The Biology of the Coleoptera. Academic Press, New York, NY.
- Freitag, R. 1974.** Selection for a non-genitalic mating structure in female tiger beetles of the genus *Cicindela* (Coleoptera: Cicindelidae). The Canadian Entomologist 106: 561–568.
- Kritsky, G., and S. Simon. 1995.** Mandibular sexual dimorphism in *Cicindela* Linnaeus (Coleoptera: Cicindelidae). The Coleopterists Bulletin 49(2): 143–148.
- Richardson, R. K. 2010.** Mandibular chirality in tiger beetles. The Coleopterists Bulletin 64(4): 386–387.
- Triplehorn, C. A., and N. F. Johnson. 2005.** Borror and DeLong's Introduction to the Study of Insects, 7th Edition. Thompson Brooks/Cole, Belmont, CA.

(Received 17 July 2015; accepted 19 January 2016. Publication date 18 March 2016.)