

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

U.S. National Park Service Publications and Papers

National Park Service

2010

Aquatic Invertebrate Monitoring at Agate Fossil Beds National Monument

Lusha Tronstad
University of Wyoming

Follow this and additional works at: <http://digitalcommons.unl.edu/natlpark>

Tronstad, Lusha, "Aquatic Invertebrate Monitoring at Agate Fossil Beds National Monument" (2010). *U.S. National Park Service Publications and Papers*. 165.
<http://digitalcommons.unl.edu/natlpark/165>

This Article is brought to you for free and open access by the National Park Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in U.S. National Park Service Publications and Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Aquatic Invertebrate Monitoring at Agate Fossil Beds National Monument

2010 Annual Report

Natural Resource Technical Report NPS/NGPN/NRTR—2012/654



ON THE COVER

Niobrara River, Agate Fossil Beds National Monument

Photograph by: Lusha Tronstad, Wyoming Natural Diversity Database

Aquatic Invertebrate Monitoring at Agate Fossil Beds National Monument

2010 Annual Report

Natural Resource Technical Report NPS/NGPN/NRTR—2012/654

Lusha Tronstad, Ph.D.
Wyoming Natural Diversity Database
1000 East University Ave., Department 3381
University of Wyoming
Laramie, Wyoming 82071

December 2012
U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Technical Report Series is used to disseminate results of scientific studies in the physical, biological, and social sciences for both the advancement of science and the achievement of the National Park Service mission. The series provides contributors with a forum for displaying comprehensive data that are often deleted from journals because of page limitations.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the Northern Great Plains Inventory & Monitoring Network website (<http://science.nature.nps.gov/im/units/NGPN>), the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/nrpm/>), and the WYNDD website (<http://www.uwyo.edu/wyndd/reports-and-publications/>).

Please cite this publication as:

Tronstad, L. 2012. Aquatic invertebrate monitoring at Agate Fossil Beds National Monument: 2010 annual report. Natural Resource Technical Report NPS/NGPN/NRTR—2012/654. National Park Service, Fort Collins, Colorado.

Contents

| | Page |
|-----------------------------|------|
| Figures..... | iv |
| Tables..... | v |
| Acknowledgments..... | vi |
| Introduction..... | 1 |
| Study Area | 3 |
| Methods..... | 5 |
| General Measurements | 5 |
| Hester-Dendy Samples | 5 |
| Hess Samples..... | 5 |
| Core Samples | 6 |
| Invertebrate Analysis | 6 |
| Results..... | 7 |
| Discussion..... | 13 |
| Literature Cited | 15 |

Figures

Page

Figure 1. a. Debris dam caused by the Hester-Dendy samplers. b. The debris dam likely increases variability in samples, such as the sampler shown that was pushed out of the water by the debris dam. Photos by Marcia Wilson. 2

Figure 2. We sampled 3 sites along the Niobrara River at Agate Fossil Beds National Monument. The black line is the Monument boundary and the transparent white areas are private land within the Monument boundary. The inset map shows the location of Agate Fossil Beds National Monument (star)..... 3

Figure 3. Photos of a. Agate Springs Ranch, b. Agate Middle, and c. Agate East. Location information located in Table 1..... 4

Figure 4. Invertebrate bioassessment metrics for 3 sites along the Niobrara River collected with Hester-Dendy multiplate sampler (a, b), Hess sampler (c, d), and corer (e, f). Higher values for taxa diversity, taxa evenness, number of EPT taxa/number of taxa, taxa richness, and EPT richness indicate better ecosystem quality, while lower values of HBI indicate higher ecosystem quality. Error bars are standard errors..... 9

Figure 5. Invertebrate bioassessment metrics for 3 sites along the Niobrara River, Agate Springs Ranch (a, b), Agate Middle (c, d), and Agate East (e, f), collected with core, Hess sampler, and Hester-Dendy multiplate sampler. Higher values for taxa diversity, taxa evenness, number of EPT taxa/number of taxa, taxa richness, and EPT richness indicate better ecosystem quality, while lower values of HBI indicate higher ecosystem quality. Error bars are standard errors. 10

Figure 6. Cluster analysis of the taxonomic composition of invertebrates, as sampled by different devices. Hester-Dendy (green), Hess (red), and core (blue) samplers collected similar taxa. 12

Tables

| | Page |
|--|------|
| Table 1. Location (NAD83) and elevation of each site along the Niobrara River. Water quality data collected on 15 July 2010 when the Hester-Dendy samplers were deployed..... | 7 |
| Table 2. Water quality, air temperature, and stream width collected on 16-17 August 2010 when all the invertebrate samples were collected..... | 8 |
| Table 3. Stream depth (cm) behind each Hester-Dendy sampler. Sampler 1 was on the south side of the Niobrara River and sampler 7 was on the north side of the river. Parallel depth is the actual water depth. Vertical displacement is an index of stream discharge, where larger numbers indicate higher flows. See methods for more details..... | 8 |

Acknowledgments

I thank Bryan Tronstad, Marcia Wilson, Robert Manasek, James Hill, and Gina Hadan who helped with logistics and field work.

Introduction

Aquatic invertebrates are excellent indicators of ecosystem quality, and these animals have been used to monitor ecosystems since the 1870s (Cairns and Pratt 1993). Growth, survival, and reproduction of aquatic invertebrates are sensitive to declines in environmental quality allowing analysis of assemblage structure to monitor lakes, streams, and wetlands (Johnson et al. 1993). Aquatic invertebrates are ideal indicator species, because they are ubiquitous, abundant, relatively long-lived, diverse, and typically sedentary (Rosenberg and Resh 1993b). In contrast, water samples can be directly analyzed for suspected pollutants, but such samples are snapshots of water quality only and can easily miss discrete discharges of pollutants or other undesirable conditions. In addition, analyzing water samples can be costly, making aquatic invertebrates a cost-effective alternative. Aquatic invertebrates integrate ecosystem quality throughout their lives, and much research has focused on how pollution alters assemblages (e.g., Rosenberg and Resh 1993a).

Choosing the best method of collecting aquatic invertebrates can be difficult, especially because there are many different devices available (Merritt et al. 2008). The ecosystem type (e.g., stream, lake or wetland), substrate type (e.g., gravel, clay, bedrock), vegetation (e.g., rooted macrophytes, floating macrophytes, periphyton), type of aquatic invertebrates targeted (e.g., benthic, hyporheic, neustic, emerging adults), and study question should all be considered when deciding on what sampler to use. Whether to collect data quantitatively (statistically rigorous information; e.g., Hess sampler), semi-quantitatively (sampling area defined; e.g., kick net), or qualitatively (estimate presence and absence of taxa; e.g., dip net; Merritt et al. 2008) depends upon the research question. All sampling devices have advantages and disadvantages; therefore, choosing a sampler that minimizes sampling bias in a given ecosystem is vital.

The National Park Service has been monitoring fish at Agate Fossil Beds National Monument since 1979 (Stasiak 1990). Over that time, ecosystem quality declined. One explanation for the decline is the invasion of yellow flag iris (*Iris pseudacorus*) (Bowles 2010; Stasiak et al. in prep). Yellow flag iris probably slows water velocity and increases organic matter in the stream leading to large daily swings in dissolved oxygen concentrations. Another explanation for the decline in ecosystem quality is the introduction of invasive northern pike (*Esox lucius*) in the Niobrara River (Stasiak et al. in prep). Pike are piscivores and likely reduced the fish assemblage from 11 species to 3 species between 1989 and 2011. Stasiak et al. (in prep) speculated that pike currently feed on crayfish, because other fish are scarce in the river. Introducing pike may have caused a trophic cascade that changed the abundance and assemblage of invertebrates in the Niobrara River.

Hester-Dendy multi-plate samplers have been used to monitor aquatic invertebrates in the Niobrara River since 1989; however, the National Park Service would like to consider other sampling methods. In the Niobrara River, the Hester-Dendy plates imitate rooted macrophyte leaves and aquatic invertebrates colonize the plates for 30 days before being collected. These samplers can be effective, but have limitations. In the Niobrara River, large debris dams form upstream of the samplers, bias the invertebrates collected and introduce greater variability (Figure 1). Therefore, my objective was to compare Hester-Dendy, Hess, and core samples collected at 3 sites along the Niobrara River at Agate Fossil Beds National Monument. My specific questions were: 1) How does the number of individuals collected from each sampler

compare? 2) How do the bioassessment metrics compare among samplers? and 3) Do these samplers collect similar taxa?




Figure 1. a. Debris dam caused by the Hester-Dendy samplers. b. The debris dam likely increases variability in samples, such as the sampler shown that was pushed out of the water by the debris dam. Photos by Marcia Wilson.

Study Area

The headwaters of the Niobrara River are located around Lusk, Wyoming, and flow eastward into Nebraska and eventually to the Missouri River near Niobrara, Nebraska. The Niobrara River Basin covers 32,600 km², of which the majority is grassland in northern Nebraska (Galat et al. 2005). Over 95% of the land within the basin is used for cattle and crop production. The Niobrara River flows through Agate Fossil Beds National Monument in western Nebraska about 23 km from the Wyoming border. At this point the Niobrara River is a low order stream flowing through grassland. Agate Fossil Beds National Monument includes about 1,100 ha in a valley bottom and about 18 km of river that flows through the 6 km wide monument (Figure 2). The riparian vegetation in the Monument is dominated by cattails (*Typha* sp.) and the invasive yellow flag iris. The substrate in the river consists of fine particles (e.g., sand). Pike, white suckers (*Catostomus commersonii*), and green sunfish (*Lepomis cyanellus*) currently inhabit the river within the Monument (Stasiak et al. in prep).




Figure 2. We sampled three sites along the Niobrara River at Agate Fossil Beds National Monument. The black line is the Monument boundary and the transparent white areas are private land within the Monument boundary. The inset map shows the location of Agate Fossil Beds National Monument (star).

I sampled 3 sites along the Niobrara River (Figure 2). The most upstream site (Agate Springs Ranch) is located near the west Monument boundary. Agate Springs Ranch has an overstory of plains cottonwood (*Populus deltoides*), and cattails are more abundant than iris (Figure 3a). The middle site, Agate Middle, is shallower and lacks an overstory (Figure 3b). Both iris and cattails are abundant here. Finally, Agate East, the site located before the Niobrara River flows out of the Monument, is the deepest site (Figure 3c). The vegetation is dominated by iris with a few willow (*Salix* spp.).




Figure 3. Photos of a. Agate Springs Ranch, b. Agate Middle, and c. Agate East. Location information located in Table 1.

Methods

General Measurements

To assess the general habitat characteristics of the Niobrara River I measured several features including water quality, turbidity, sediment composition, and depth. I measured dissolved oxygen (percent saturation and mg/L), pH, water temperature, specific conductivity, and oxidation-reduction potential using a Yellow Springs Instruments Professional Plus. The probe was calibrated on site before use. I measured turbidity by estimating the depth at which a Secchi disk disappeared from sight. The composition of sediment was estimated by sampling sediment across the width of the stream and selecting the percent class for clay, silt, sand, gravel, cobble, boulder, bedrock, and hardpan/shale on a scale of 0 -7 (0 = none, 1 = trace, 2 = 1-5%, 3 = 5-25%, 4 = 25-50%, 5 = 50-75%, 6 = 75-95% and 7 = 95-100%; Peterson et al. 1999). Clay was defined as fine particles forming a ribbon after removing water, whereas silt did not form a ribbon. Sand was particles 0.06-2 mm in diameter, gravel was 2-64 mm in diameter, cobble was 64-256 mm in diameter, boulder was 256-4000 mm in diameter, bedrock was >4000 mm in diameter, and hardpan/shale was firm, consolidated fine substrate. I measured the depth of the water with a meter stick (3.2 mm width) parallel and perpendicular to flow across the width of the stream. By subtracting the two measurements, I calculated an estimate of water velocity (vertical displacement); the greater the vertical displacement of the water, the higher the water velocity. Finally, I recorded the location and elevation of each site using a global positioning system (GPS; Garmin eTrex Vista HCx).

Hester-Dendy Samples

I deployed seven Hester-Dendy multi-plate samplers (76 mm by 76 mm, 9 plates, Wildlife Supply Company) at each site on 15 July 2010. Five Hester-Dendy samplers were used for analysis and two extra samplers were deployed in case any were lost. I strung a rope across the stream between two permanent posts and tied seven loops to separate the Hester-Dendy samplers. From each loop, I tied another rope with the Hester-Dendy samplers hanging about 0.3 meters below to allow for a drop in water level. I retrieved the samplers on 16 and 17 August 2010 by approaching the site from downstream and placing a dip net (212 μ m mesh) under each sampler. Hester-Dendy samplers were immediately placed in a container with 80% ethanol, and any organisms in the dip net were removed and placed in the same container. After returning to the laboratory, I dismantled the Hester-Dendy samplers to remove invertebrates that colonized the plates, rinsed samples using a 212 μ m sieve, and preserved samples in 80% ethanol. Replicates two through six were typically used for analysis.

Hess Samples

To sample invertebrates that live in the emergent vegetation that is abundant along the Niobrara River, I collected three Hess samples (500 μ m mesh, 860 cm² sampling area, Wildlife Supply Company) from each site on 16 and 17 August 2010. I placed the Hess sampler over cattails and yellow flag iris to collect invertebrates that lived on the vegetation and in the surrounding sediment. The vegetation and sediment were vigorously agitated using our hands and a brush, and the samples were preserved in 80% ethanol.

Core Samples

To collect invertebrates that live in the sediment of the Niobrara River, I took three core samples (Hand corer, 16.6 cm² sampling area, Wildlife Supply Company) from each site. Sediment was sieved through 212 µm mesh to remove fine particles and the samples were preserved in 80% ethanol.

Invertebrate Analysis

Invertebrates were sorted from the debris and identified to genus (Insecta, Turbellaria, Isopoda, and Amphipoda), family (Decapoda, Pelecypoda (Bivalvia), Gastropoda), class (Annelida, Acarina) or phylum (Nematoda) with two exceptions (order: Collembola, family: Diptera) according to Peterson et al. (1999). If invertebrates were extremely numerous (>500 individuals) in any sample, I subsampled as follows. First, I removed the large and rare invertebrates, then I subsampled the remaining invertebrates. Invertebrates were identified under a dissecting microscope using Merritt et al. (2008) for insects, and Thorp and Covich (2010) and Smith (2001) for non-insect invertebrates.

Several bioassessment metrics have been calculated since 1989 to estimate stream quality based on the invertebrates collected: Hilsenhoff Biotic Index (HBI), Ephemeroptera, Plecoptera, and Trichoptera (EPT) richness, number of EPT taxa divided by the total number of taxa collected, taxa diversity (Shannon index), taxa richness, and taxa evenness (Bowles 2010). To distinguish among sites, I used ANOVA to compare abundance (per sampler) and bioassessment metrics for each sampler (DataDesk6.1). To evaluate sampling devices, I used ANOVA to compare abundance (per sampler) and each of these six bioassessment metrics among sampler types. Differences among samplers and sites were distinguished using Bonferroni multiple comparison tests. Differences were significant when $p < 0.017$ ($0.05/3$; where I had three samplers and three sites). Finally, I used cluster analysis to estimate the degree to which the taxonomic composition of samples varied by sampler type (PC-ORD version 6).

Results

Water quality was similar between deploying and retrieving the Hester-Dendy samplers. In general, Agate Springs Ranch had the highest dissolved oxygen concentrations and Agate East had the lowest concentrations (Tables 1 and 2). All dissolved oxygen concentrations were collected mid-morning, except Agate Springs Ranch and Agate Middle on 16 August 2010, which were collected in the afternoon. pH was highest at Agate Springs Ranch on both dates. Specific conductivity increased from Agate Springs Ranch to Agate East. Reducing conditions (<200 mV; oxidation-reduction potential) appeared to dominate in the Niobrara River. At all sites, I could view the Secchi disk at the bottom of the stream. Stream width was consistent among sites; however, the Niobrara River was widest at Agate Springs Ranch and narrowest at Agate Middle. The water reached into the wetland vegetation at both Agate Middle and Agate East when I collected samples. Agate East was the deepest site and Agate Middle was the shallowest (Table 3). Stream discharge was highest at Agate Middle and lowest at Agate East. Overall, the substrate in the Niobrara River at each site was dominated by sand. At Agate Springs Ranch, I estimated that 1-5% was silt and 95-100% was sand. At Agate Middle, 25-50% was sand, 50-75% was gravel, and there was a trace of silt. Agate East was composed of 5-25% silt and 75-95% sand.

Table 1. Location (NAD83) and elevation of each site along the Niobrara River. Water quality data collected on 15 July 2010 when the Hester-Dendy samplers were deployed.

| Site description | Agate Springs Ranch | Agate Middle | Agate East |
|-------------------------------|---------------------|--------------|------------|
| Easting | 599323 | 602143 | 604495 |
| Northing | 4697497 | 4696844 | 4697913 |
| Elevation (m) | 1354 | 1350 | 1343 |
| DO (% saturation) | 90 | 67 | 57 |
| DO (mg/L) | 6.8 | 5.2 | 4.5 |
| pH | 8.18 | 7.78 | 7.6 |
| Water temperature (°C) | 21.2 | 19.9 | 19 |
| Specific conductivity (µS/cm) | 417.1 | 451.2 | 455.6 |
| ORP (mV) | 182.1 | 171.6 | 219.8 |

Table 2. Water quality, air temperature, and stream width collected on 16-17 August 2010 when all the invertebrate samples were collected.

| Site description | Agate Springs Ranch | Agate Middle | Agate East |
|-------------------------------|---------------------|--------------|------------|
| DO (% saturation) | 98 | 93 | 81 |
| DO (mg/L) | 7.6 | 7.3 | 6.7 |
| pH | 8.15 | 7.88 | 7.99 |
| Water temperature (°C) | 19.2 | 19.1 | 16.9 |
| Specific conductivity (µS/cm) | 419.5 | 423.7 | 428.5 |
| ORP (mV) | 188.7 | 45 | 107.6 |
| Secchi disk depth (cm) | >64 | >60 | >97 |
| Air temperature (°C) | 27.5 | 19 | 28 |
| Stream width (m) | | | |
| Open water | 2.18 | 1.9 | 2.1 |
| In vegetation, south | 0 | 0.36 | 0.23 |
| In vegetation, north | 0 | 0.74 | 0.33 |

Table 3. Stream depth (cm) behind each Hester-Dendy sampler. Sampler 1 was on the south side of the Niobrara River and sampler 7 was on the north side of the river. Parallel depth is the actual water depth. Vertical displacement is an index of stream discharge, where larger numbers indicate higher flows. See methods for more details.

| Sampler # | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Mean |
|----------------------------|----|----|----|-----|-----|-----|----|------|
| Agate Springs Ranch | | | | | | | | |
| Parallel depth | 53 | 60 | 64 | 63 | 62 | 59 | 54 | 59 |
| Perpendicular depth | 53 | 62 | 65 | 70 | 72 | 59 | 54 | 62 |
| Verticle displacement | 0 | 2 | 1 | 7 | 10 | 0 | 0 | 3 |
| Agate Middle | | | | | | | | |
| Parallel depth | 56 | 58 | 60 | 59 | 56 | 52 | 40 | 54 |
| Perpendicular depth | 56 | 76 | 82 | 77 | 70 | 55 | 41 | 65 |
| Verticle displacement | 0 | 18 | 22 | 18 | 14 | 3 | 1 | 11 |
| Agate East | | | | | | | | |
| Parallel depth | 77 | 89 | 93 | 97 | 96 | 99 | 96 | 92 |
| Perpendicular depth | 77 | 91 | 93 | 100 | 103 | 101 | 97 | 95 |
| Verticle displacement | 0 | 2 | 0 | 3 | 7 | 2 | 1 | 2 |

I identified 37 invertebrate taxa using Hester-Dendy multi-plate samplers in the Niobrara River, from four phylum (Nematoda, Annelida, Mollusca, and Arthropoda). Overall, Diptera, Ephemeroptera, Crustacea, and Mollusca were the most numerous groups collected in order of decreasing abundance. Hester-Dendy samplers from Agate East (1,987 ind/sampler) contained the most invertebrates and Agate Springs Ranch (420 ind/sampler) had the fewest ($F = 5.6$, $df = 2$, $p = 0.019$). Taxa diversity was lowest at Agate Springs Ranch (Bonferroni: $p < 0.005$), and

similar between Agate Middle and Agate East (Figure 4a; $F = 11.9$, $df = 2$, $p = 0.0014$). Agate East had the highest taxa richness (Figure 4b; $F = 12.57$, $df = 2$, $p = 0.0011$; Bonferroni: $p < 0.004$), and Agate Middle had higher taxa evenness than Agate Springs Ranch ($F = 12.3$, $df = 2$, $p = 0.0025$; Bonferroni: $p = 0.002$). I collected three EPT taxa at both Agate Springs Ranch and Agate East, and three EPT taxa at Agate Middle. Agate Springs Ranch contained the highest proportion of EPT taxa. The average tolerance value for invertebrates was higher at Agate East and Agate Middle, and much lower at Agate Springs Ranch ($F = 105.6$, $df = 2$, $p < 0.0001$; Bonferroni: $p < 0.0001$).




Figure 4. Invertebrate bioassessment metrics for three sites along the Niobrara River collected with Hester-Dendy multi-plate sampler (a, b), Hess sampler (c, d), and corer (e, f). Higher values for taxa diversity, taxa evenness, number of EPT taxa/number of taxa, taxa richness, and EPT richness indicate better ecosystem quality, while lower values of HBI indicate higher ecosystem quality. Error bars are standard errors.

I collected 34 taxa of invertebrates from four phylum using a Hess sampler in the Niobrara River. Overall, Diptera, Mollusca, Odonata, and Ephemeroptera were the most numerous invertebrates in decreasing order of abundance. Agate East had the highest density of invertebrates (2,710 ind/sampler) and Agate Middle had the lowest density (152 ind/sampler; $F = 6.4$, $df = 2$, $p = 0.033$; Bonferroni: $p > 0.05$). Taxa diversity was similar among sites (Figure 4c; $F = 1.4$, $df = 2$, $p = 0.31$). I collected the same number of taxa at Agate Springs Ranch and




Figure 5. Invertebrate bioassessment metrics for three sites along the Niobrara River, Agate Springs Ranch (a, b), Agate Middle (c, d), and Agate East (e, f), collected with core, Hess sampler, and Hester-Dendy multi-plate sampler. Higher values for taxa diversity, taxa evenness, number of EPT taxa/number of taxa, taxa richness, and EPT richness indicate better ecosystem quality, while lower values of HBI indicate higher ecosystem quality. Error bars are standard errors.

Agate East, and half the number of taxa at Agate Middle (Figure 4d; $F = 22.3$, $df = 2$, $p = 0.0017$; Bonferroni: $p < 0.004$). Taxa evenness was highest at Agate Middle and lowest at Agate East, but values were not statistically significant among sites ($F = 4.5$, $df = 2$, $p = 0.064$). I collected three EPT taxa at Agate Springs Ranch and Agate East, and two EPT taxa at Agate Middle. The proportion of EPT taxa ($F = 0.047$, $df = 2$, $p = 0.95$) and mean tolerance values ($F = 4$, $df = 2$, $p = 0.079$) were similar among sites.

I collected 14 taxa using a sediment corer in the Niobrara River. Overall, Diptera, Crustacea, and Mollusca were the most numerous groups in decreasing order of abundance. Agate East had the highest density of invertebrates (28 ind/sample) and Agate Springs Ranch had the lowest abundance (5 ind/sample). Taxa diversity and evenness were similar among sites (Figure 4e). Taxa richness, EPT richness, and the proportion of EPT taxa were highest at Agate East (Figure 4f), but not statistically different among sites ($p > 0.05$). Finally, mean tolerance values for taxa were similar among sites ($F = 3.5$, $df = 2$, $p = 0.10$).

In general, Hester-Dendy and Hess samplers collected similar samples, but core samples typically differed (Figure 5). The total number of individuals collected per sample was similar between Hester-Dendy and Hess samples ($F = 5.19$, $df = 2$, $p = 0.012$, Bonferroni: $p = 0.999$), but core samples collected fewer individuals than Hester-Dendy samples (Bonferroni: $p = 0.017$). All samplers collected a similar number of Mollusca, Crustacea, and Coleoptera ($p > 0.05$), but Hester-Dendy samplers collected more Ephemeroptera (mean values; Hester-Dendy = 146, Hess = 17, and core = 0.1, Bonferroni < 0.017) and Diptera (mean values; Hester-Dendy = 182, Hess = 50, and core = 6.4, Bonferroni < 0.017) per sample than Hess or core samples ($F = 9.38$ and 14.26 , $df = 2$, $p = 0.008$ and < 0.0001 , respectively). The assemblage of taxa did not vary by the sampler, as cluster analysis did not group samples by sampler type, with the exception of core samples (Figure 6). However, Hester-Dendy and Hess samplers were intermixed throughout the tree.

Bioassessment metrics were similar between Hester-Dendy and Hess samples, but values differed for core samples. Taxa diversity was similar among samplers, but taxa richness, taxa evenness, EPT richness, HBI, and EPT taxa/total taxa differed among samplers. Core samples produced different values for taxa richness ($F = 43.57$, $df = 2$, $p < 0.0001$), taxa evenness ($F = 8.2$, $df = 2$, $p = 0.0016$), EPT richness ($F = 58.56$, $df = 2$, $p < 0.0001$), and EPT taxa/total taxa ($F = 72.7$, $df = 2$, $p < 0.0001$) compared to Hester-Dendy and Hess samplers (Bonferroni: $p < 0.017$). However, HBI values differed between Hester-Dendy (mean = 5.5) and Hess (mean = 7.8) samplers ($F = 12.28$, $df = 2$, $p = 0.0001$, Bonferroni: $p < 0.017$). The lower values for Hester-Dendy samplers may be from a large number of *Paraleptophebia* (tolerance value = 1.2) collected with Hester-Dendy samplers at Agate Springs Ranch site.




Figure 6. Cluster analysis of the taxonomic composition of invertebrates, as sampled by different devices. Hester-Dendy (green), Hess (red), and core (blue) samplers collected similar taxa.

Discussion

Prairie streams can be difficult to sample for aquatic invertebrates. Prairie streams often have fine substrates, yet most quantitative aquatic invertebrate samplers are designed for gravel or cobble bottom streams (e.g., Surber sampler). Therefore, alternative sampling methods are necessary. Core samplers have the advantage of directly targeting fine substrates; however, cores typically sample a small area and the samples can be rather time intensive to sort. Additionally, fewer invertebrates may live in the sediment, depending on conditions. Others have used semi-quantitative box samplers in prairie wetlands that consist of placing a 4-sided rectangle in the wetland and collecting invertebrates within the area using a dip net (Meyer and Whiles 2008). However, box samplers do not work well in deep habitats, streams, or areas with dense vegetation. Therefore, an appropriate sampling method often depends upon the conditions, habitat, and the question.

Another option for collecting aquatic invertebrates is using Hester-Dendy plates, which is an artificial substrate sampler. Hester-Dendy samplers provide solid substrate in habitats that may lack such areas. Additionally, these samplers may mimic snags or macrophytes which can be difficult to sample. Invertebrate density is typically calculated based on the surface area of the plates; however, surface area in natural habitats (e.g., surface area of macrophyte leaves) is seldom calculated and would be extremely labor intensive. Therefore, density or biomass of aquatic invertebrates collected with Hester-Dendy samplers can only be compared to aquatic invertebrates from other ecosystems that were collected using the same method. In the Niobrara River, macrophytes in the wetland area provide abundant substrate for aquatic invertebrates. The wetland area differs from the main channel in that the wetland area can be large (0.4 km wide in places) and water velocity is significantly slower. Hester-Dendy samplers were placed in the main channel of the stream where water velocities were higher causing large debris dams to form. Debris dams were periodically cleared from the Hester-Dendy samplers; however, I have observed debris dams that were >0.3 m (>1 ft) deep and >1.8 m (>6 ft) in length when I retrieved the samplers (Figure 1). Because of these large debris dams, I collected species that normally would not be collected with a Hester-Dendy sampler, such as large crayfish (Decapods). Also, debris dams may cause higher variability in the samples because either more (including debris) or fewer invertebrates (removing debris may displace individuals) may be collected depending on how the samplers are gathered. I have also observed Hester-Dendy samplers pushed out of the water by debris dams (Figure 1b).

Hess samplers collect natural densities of aquatic invertebrates that are comparable to other quantitative methods among ecosystems (per m² of stream bottom). Hess samplers are traditionally used similarly to Surber samplers, but they have the advantage of surrounding the sampling area. I chose to use a Hess sampler to collect aquatic invertebrates in the Niobrara River, because I could sample the macrophytes and sediments to estimate natural densities. I slipped the Hess sampler over the macrophytes at the edge of the main channel, and scoured the vegetation and sediment. Therefore, I sampled invertebrates that lived in both habitats (vegetation and sediment) and that had multiple habits (e.g., crawlers, clingers, etc.). Hess samplers have shortfalls too; for example, Hess samplers cannot be used in deep water. However, modifications can be made to minimize bias.

In general, I collected similar aquatic invertebrates using Hester-Dendy and Hess samplers, but core samples contained less diversity. Hester-Dendy samplers contained a few taxa that I did not collect in Hess samples. For example, I found low densities of Hirudinea (leeches), *Heptagenia* (mayfly), Anisoptera (dragonflies), and *Polycentropus* (caddisfly) in Hester-Dendy samples, but these taxa did not appear in Hess samples. Based on the habits of these taxa, I would expect Hess samples to collect them. However, these are rarer taxa and I collected fewer Hess samples than Hester-Dendy samples, which may explain why I collected greater diversity in the latter. Additionally, I collected ostracods in Hester-Dendy samples. Ostracods are small crustaceans that probably flowed through the 500 μm mesh of the Hess sampler, but were retained in the 212 μm mesh of the sieve for the Hester-Dendy samplers. Conversely, I collected Stratiomyidae (true fly) only in the Hess samples. Stratiomyidae appeared to be rare, which explains why I did not collect them in other samplers. Finally, I collected far less diversity in the core samples, but I only collected *Corbicula* using the core sampler at the East site.

Using core and Hess samplers together, the overall density of invertebrates in the Niobrara could be calculated given the available habitat in each area. In August, the stream channel was between 63 and 100% of the habitat available to aquatic insects, while only 0 to 37% of water was in the wetland vegetation. Therefore, invertebrates that can live in the sediments (collected with a core sampler) had the greatest area to inhabit. A much larger area of the riparian habitat is flooded earlier in the year before water is withdrawn for irrigation. Floodplains can provide large areas for invertebrates to colonize and produce higher biomass compared to other habitats within the ecosystem (Benke 2001).

I recommend using a Hess sampler to collect aquatic invertebrates in the Niobrara River. Collecting invertebrates with a Hess sampler compared to a Hester-Dendy sampler will reduce the number of visits to the sites along the Niobrara River from potentially two (deploying, three visits to clear debris dams, and collecting) to only one (collecting). By collecting invertebrates on natural substrate there may be less variability in the samples, because of the difficulties using Hester-Dendy samplers in the Niobrara River. In general, Hester-Dendy and Hess samplers collected a similar number and assemblage of aquatic invertebrates in 2010. I do not recommend collecting core samples, because these samples contained significantly lower diversity and I collected the same taxa in Hess samples with the exception of rare *Corbicula*. Hess samples should be collected in early August, when water levels are slightly higher and more water extends into the riparian habitat, which will expedite sampling.

Literature Cited

- Benke, A. C. 2001. Importance of flood regime to invertebrate habitat in an unregulated river-floodplain ecosystem. *Journal of the North American Benthological Society* 20:225-240.
- Bowles, D. E. 2010. Aquatic invertebrate monitoring at Agate Fossil Beds National Monument, 1989-2009 trend report. Heartland Inventory and Monitoring Network, Springfield, Missouri.
- Cairns, J., and J. R. Pratt. 1993. A history of biological monitoring using benthic macroinvertebrates. Pages 10-27 *in* D. M. Rosenberg and V. H. Resh, editors. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York, NY.
- Galat, D. L., C. R. Berry, E. J. Peters, and R. G. White. 2005. Missouri River Basin. Pages 427-480 *in* A. C. Benke and C. E. Cushing, editors. *Rivers of North America*. Elsevier, New York, NY.
- Johnson, R. K., T. Wiederholm, and D. M. Rosenberg. 1993. Freshwater biomonitoring using individual organisms, populations, and species assemblages of benthic macroinvertebrates. Pages 40-158 *in* D. M. Rosenberg and V. H. Resh, editors. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York, NY.
- Merritt, R. W., K. W. Cummins, and M. B. Berg, editors. 2008. *An Introduction to the Aquatic Insects of North America*, 4th edition. Kendall Hunt Publishing, Dubuque, IA.
- Meyer, C. K., and M. R. Whiles. 2008. Macroinvertebrate communities in restored and natural Platte River slough wetlands. *Journal of the North American Benthological Society* 27:626-639.
- Peterson, J. T., W. M. Rizzo, E. D. Schneider, and G. D. Willson. 1999. Macroinvertebrate biomonitoring protocol for four prairie streams. United States Geological Survey and United States Forest Service.
- Rosenberg, D. M., and V. H. Resh, editors. 1993a. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York.
- Rosenberg, D. M., and V. H. Resh. 1993b. Introduction to freshwater biomonitoring and benthic macroinvertebrates. Pages 1-9 *in* D. M. Rosenberg and V. H. Resh, editors. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York, NY.
- Smith, D. G. 2001. *Pennak's Freshwater Invertebrates of the United States*, 4th edition. John Wiley and Sons, Inc., New York.
- Stasiak, R. H. 1990. Fishes of the Niobrara River at Agate Fossil Beds National Monument 1979 and 1989. Unpublished Report.
- Stasiak, R. H., G. R. Cunningham, S. Flash, A. Wagner, and A. Barela. In prep. Fishes of the Niobrara River at Agate Fossil Beds National Monument: 2011 survey. University of Nebraska at Omaha, Omaha.

Thorp, J. H., and A. P. Covich, editors. 2010. Ecology and Classification of North American Freshwater Invertebrates, 3rd edition. Elsevier, New York.

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 165/119284, December 2012

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525
www.nature.nps.gov

EXPERIENCE YOUR AMERICA™