

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Sociology Department, Faculty Publications

Sociology, Department of

10-20-2016

Infertility and fertility intentions, desires, and outcomes among US women

Karina M. Shreffler

Oklahoma State University, karina.shreffler@okstate.edu

Stacy Tiemeyer

Oklahoma State University, stacy.tiemeyer@okstate.edu

Cassandra Dorius

Iowa State University, cdorius@iastate.edu

Tiffany Spierling


Oklahoma State University, tiffany.spierling@okstate.edu

Arthur L. Greil

Alfred University, fgreil@alfred.edu

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/sociologyfacpub>

 Part of the [Family, Life Course, and Society Commons](#), and the [Social Psychology and Interaction Commons](#)

Shreffler, Karina M.; Tiemeyer, Stacy; Dorius, Cassandra; Spierling, Tiffany; Greil, Arthur L.; and McQuillan, Julia, "Infertility and fertility intentions, desires, and outcomes among US women" (2016). *Sociology Department, Faculty Publications*. 587.
<http://digitalcommons.unl.edu/sociologyfacpub/587>

This Article is brought to you for free and open access by the Sociology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Sociology Department, Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Karina M. Shreffler, Stacy Tiemeyer, Cassandra Dorius, Tiffany Spierling, Arthur L. Greil, and Julia McQuillan



DEMOGRAPHIC RESEARCH

A peer-reviewed, open-access journal of population sciences

DEMOGRAPHIC RESEARCH

VOLUME 35, ARTICLE 39, PAGES 1149–1168

PUBLISHED 20 OCTOBER 2016

<http://www.demographic-research.org/Volumes/Vol35/39/>

DOI: 10.4054/DemRes.2016.35.39

Research Article

Infertility and fertility intentions, desires, and outcomes among US women

Karina M. Shreffler

Stacy Tiemeyer

Cassandra Dorius

Tiffany Spierling

Arthur L. Greil

Julia McQuillan

©2016 Karina M. Shreffler et al.

This open-access work is published under the terms of the Creative Commons Attribution NonCommercial License 2.0 Germany, which permits use, reproduction & distribution in any medium for non-commercial purposes, provided the original author(s) and source are given credit. See <http://creativecommons.org/licenses/by-nc/2.0/de/>

Contents

1	Introduction	1150
1.1	Connecting the infertility experience and identity with demographic research on fertility	1151
1.2	What does it mean to “intend to give birth” in the context of infertility?	1152
2	Data and methods	1153
2.1	Sample	1153
2.2	Measures	1154
2.3	Analytic strategy	1155
3	Results	1156
4	Conclusions	1161
	References	1165

Infertility and fertility intentions, desires, and outcomes among US women

Karina M. Shreffler¹

Stacy Tiemeyer²

Cassandra Dorius³

Tiffany Spierling⁴

Arthur L. Greil⁵

Julia McQuillan⁶

Abstract

BACKGROUND AND OBJECTIVE

Little is known about how the experience of infertility or identification as someone with infertility shapes women's fertility intentions, desires, or birth outcomes. The purpose of this paper is to help fill this gap in knowledge for fertility-intentions research.

METHODS

Using data from the National Survey of Fertility Barriers (NSFB), we use linear and logistic regression methods to assess how infertility and parity statuses are associated with fertility intentions and desires, as well as how statuses at one point in time predict birth three years later.

RESULTS

We find that infertility is associated with lower fertility intentions. Women who have experienced infertility and/or identify as a person with infertility, however, express greater desires to have a baby and a higher ideal number of children. Women who meet the medical criteria for infertility are less likely than fecund women to give birth, despite greater desires.

¹ Oklahoma State University, USA. E-Mail: karina.shreffler@okstate.edu.

² Oklahoma State University, USA.

³ Iowa State University, USA.

⁴ Oklahoma State University, USA.

⁵ Alfred University, USA.

⁶ University of Nebraska-Lincoln, USA.

CONCLUSION

These findings have important theoretical implications for our understanding of the meaning of fertility intentions for those who think their ability to achieve their intentions is uncertain, as well as for empirical research on fertility.

1. Introduction

Demographers interested in fertility trends and projections have long considered the role of infertility, particularly in analyses of developing countries (see Rutstein and Shah 2004). When surveyed about infertility, the lay public also understands the term ‘infertility’ to mean a permanent inability to give birth (Maill 1994). Yet the medical definition of infertility – 12 months or more of unprotected, heterosexual intercourse without conception (American Society for Reproductive Medicine [ASRM] 2008), sometimes referred to as subfecundity – does not indicate permanent involuntary childlessness. The proportion of women in the US who experience infertility using the ASRM definition is 7% to 15.5% in a given year depending upon specific measurement procedures (Thoma et al. 2013). The percentage is substantially higher (51.8%) for women who have met the ASRM criteria at some point throughout their reproductive lifespan (Greil et al. 2011a). Little is known about how women who experience infertility (e.g., meet the medical criteria) or identify as having a fertility problem (i.e., with or without meeting the medical criteria) think about their fertility intentions and desires compared to women without infertility experiences. Further, it is not known how infertility affects birth outcomes in countries such as the United States, where over 10% of all women trying to become pregnant seek medical treatment for infertility (Simonsen, Baksh, and Stanford 2012). Fears of infertility and decisions about how to respond to failures to conceive may influence fertility intentions, desires, and outcomes. Yet scant research has examined these relationships. We therefore use the National Survey of Fertility Barriers (NSFB) to explore the associations between infertility and fertility intentions, fertility desires, and the ideal number of children among US women ages 25–45 by fertility status (not infertile, medically infertile, self-identifying as infertile without meeting the medical definition of infertility, and self-identifying as infertile with experience of infertility) and motherhood status (childless and having one or more children). We further examine how infertility and motherhood predict the likelihood of a birth in the three years after the initial interview.

1.1 Connecting the infertility experience and identity with demographic research on fertility

Infertility is an interesting phenomenon for fertility researchers and theorists who study fertility intentions. Though seemingly straightforward, infertility does not necessarily have clear implications for fertility considerations. Demographers have long realized the importance of infertility at the societal level for measures of fertility trends. For example, Davis and Blake (1956) identified intermediate fertility variables including sterility, which included noncontraceptive sterility, or what we think of today as infertility or subfecundity. Calculations predicting fertility trends measure infertility in different ways: Stover (1998) suggested including a measure of sterility in fertility calculations that includes women in a union who are not menopausal, postpartum amenorrheic, or pregnant but have not given birth within the past five years despite not using contraception, as well as women who declare themselves to be infertile. Mascarenhas and colleagues (2012) have applied this approach to Demographic Health Survey data from 53 countries, for example, in an effort to determine the global prevalence of infertility.

Thus, whereas the demographic approach often infers infertility from questions on couple status, contraception use, and fertility desires (Mascarenhas et al. 2012), the medical approach utilizes questions focused on meeting the medical criteria for infertility (e.g., “Have you ever had a period of at least 12 months of regular, unprotected sexual intercourse without becoming pregnant?”) and perceptions of a fertility problem (e.g., “Do you see yourself as someone who has had trouble trying to conceive?”). The medical approach, therefore, can identify individuals or couples who may benefit from medical assistance to achieve conception, whereas the demographic approach is typically seeking to identify women who are not likely to become pregnant. In the demographic- and social-science literatures, we are perhaps most familiar with infertile women who are involuntarily childless. This is a meaningful group for both demographic and social scientists, as this group has important implications for fertility-rate calculations (Bongaarts and Potter 2013) and because infertility is particularly distressing for women who are childless and have strong fertility desires (Schwerdtfeger and Shreffler 2009). Yet the majority of women who experience infertility do not remain childless (Schwerdtfeger and Shreffler 2009). Further, while we might expect a great deal of overlap between experiencing a period of infertility (e.g., not getting pregnant after at least a year of regular, unprotected heterosexual intercourse) and realizing that there might be a fertility problem, there are surprising discrepancies. A substantial proportion of women (35%) who meet the medical criteria do not view themselves as being infertile (White et al. 2006). Surprisingly, there are also women who do not appear to meet the medical criteria for infertility but believe that they have a fertility problem or expect to have a problem when they start trying to

conceive. A study of unmarried adults aged 18–29, for example, found that 19% of women believed that they were “very likely” to be infertile, and having this belief was associated with contraceptive behaviors (Polis and Zabin 2012). Infertility, therefore, could affect fertility in two directions. Women who meet the medical criteria for infertility are not getting pregnant, at least temporarily. Women who identify as infertile but do not meet the medical criteria may actually be more likely to give birth due to their beliefs that contraception to prevent birth is unnecessary. Moreover, if women who are infertile do not think that they will become pregnant, these births could be classified as ‘unintended’ due to the current measurement of fertility intentions..

1.2 What does it mean to “intend to give birth” in the context of infertility?

Theoretical conceptualizations of fertility intentions suggest that they are shaped by the desire for a particular outcome, a belief that taking an action will result in that desired outcome, and a commitment to perform the action (Malle, Moses, and Baldwin 2003). Ajzen and Klobas (2013) suggest that one factor affecting fertility intentions is the degree of perceived control that individuals have over their fertility. Women who are infertile may not believe that taking an action (e.g., stopping contraception, predicting ovulation) will result in pregnancy. Yet the experience of infertility alone may not be enough to influence fertility intentions and behaviors. It may require conscious processing of this experience, resulting in identification of a fertility problem. Thus, there needs to be a consideration of both the experience of infertility and the perception of infertility. Not all women who meet the medical criteria for infertility are aware that they have a problem (Greil and McQuillan 2010). Indeed, it is the perception of infertility that matters more for psychological outcomes than merely meeting the medical criteria for infertility (Greil et al. 2011b).

Therefore, women who have given thought to their fertility experiences and have come to realize that they have trouble becoming pregnant may be particularly susceptible to influences of infertility on their fertility intentions. Fertility-intentions questions seemingly assume the ability to carry out intentions; for example, the 2006–2010 National Survey of Family Growth (NSFG) asked respondents, “Looking to the future, do you intend to have (a/another) child at some time?” followed by, “How sure are you that you (will/will not) have a baby?” Often, these questions are coded into one measure ranging from ‘very sure, do not intend’ to ‘very sure, intend’ (e.g., Thomson 1997), which is then used to predict future fertility (e.g., Schoen et al. 1999) or intendedness of a birth (e.g., Maximova and Quesnel-Vallée 2009).

Yet, infertility calls into question the meaning of fertility intentions when the outcomes are uncertain, especially if respondents are asked to report intentions within a

particular time frame. McQuillan et al. (2015) suggest a stronger emphasis on identity is needed in studies of fertility intentions in addition to the more structural (i.e., life course) and cultural (i.e., values) approaches often used. Though they focused on motherhood identity, we suggest that identifying as infertile may be another salient identity in the study of fertility intentions. Because perception of infertility and meeting the medical criteria of infertility are often not synonymous constructs, measuring them separately allows for more-refined investigation of associations between infertility and fertility intentions, desires, ideals, and outcomes. For example, the uncertainty of the outcome following perceived infertility raises interesting questions about fertility intentions. What do intentions mean for women who may strongly want to give birth or avoid giving birth but are uncertain about their ability to be able to conceive? Do they report strong intentions (e.g., very sure that they intend or do not intend to give birth), or do they downgrade or downplay their intentions even if they might want to have a baby, or alternatively, do they downplay intentions if they plan to avoid pregnancy because it seems unnecessary to have strong intentions not to have a baby while infertile? Do infertile women reduce their intentions, and possibly their preferred number of children, to meet their expectations of lowered fertility, perhaps because they are trying to be realistic about what they believe is possible? On the other hand, does concern over infertility increase their desire to have a baby and/or preferred family size, resulting in inflated preferences? And finally, are infertile women able to have the number of children that they want to have, or does meeting the medical criteria for infertility predict a lower likelihood of a birth?

2. Data and methods

2.1 Sample

Our data comes from the National Survey of Fertility Barriers (NSFB), a random digit-dialing telephone survey of 4,712 women of childbearing ages (25–45) and a subset of their husbands and partners. The study was designed to assess social and health factors related to reproductive choices and fertility for US women. The first wave was collected from 2004 to 2006, and the second three years later. The data is nationally representative, with an oversample of Black and Hispanic women and women with fertility problems. Analyses for this study are weighted to account for the oversamples. Our sample for fertility intentions and desires analyses is restricted to 2,978 women who are not surgically sterilized or in a heterosexual marriage or cohabiting relationship with a man who has been surgically sterilized. Our sample for the analysis examining

birth odds by infertility and parity status includes 1,547 women from the first sample who participated in the wave-two interview.

2.2 Measures

Dependent variables. In our first set of multivariate analyses utilizing wave-one data only, we examine three dependent variables targeting fertility intentions, wants, and ideals. In a subsequent analysis, we utilize these variables as predictor variables of whether or not a birth occurred between survey waves. The first dependent variable, *fertility intentions*, is based on two questions that are combined to create an ordinal measure of fertility intentions. Respondents were asked, “Do you intend to have a baby?” and “Of course, sometimes things do not work out exactly as we intend them to, or something makes us change our minds. In your case, how sure are you that you will/will not have a child?” Responses were coded so that low scores indicate, “Very sure, do not intend,” (–2) and high scores indicate, “Very sure, do intend” (+2). Women who said they “don’t know” their intentions, who said they cannot have children, or who said they would let God or nature decide are coded 0 (the center of the scale). These questions are similar to those used in the National Survey of Families and Households; we recoded the response categories so that a positive score indicates intending and a negative score indicates not intending to have a baby. Our second dependent variable, *want a baby*, is measured by a question asking, “In the future, would you like to have a(nother) baby?” Responses are coded from 1 (definitely no) to 4 (definitely yes). *Ideal number of children* was measured by asking, “How many children would you consider to be ideal for you?” and coded from 0 to 5, with 5 including preferences of five or more children. *Birth between waves* is an indicator variable where 1 means that the respondent gave birth between waves.

Infertility and parity groups. We categorized women as *medically infertile* if they had ever had a period of 12 months or more during which they had unprotected heterosexual intercourse and did not get pregnant. We considered women to be *self-identifying as infertile* if they answered yes to either, “Do you think of yourself as someone who has, has had, or might have trouble getting pregnant?” or “Do you think of yourself as someone who has or has had fertility problems?” Three distinct groups were created: women who were medically infertile but did not self-identify as infertile, women who did not meet the medical criteria for infertility but did identify as infertile, and women who both met medical criteria and self-identified as infertile. A fourth group, of non-infertile women (answering no to both the self-identification and medically infertile questions), was also included in the study. These four groups were

further split by parity, comparing women who were childless to women who had at least one biological child.

Sociodemographic control variables. *Age* is a continuous variable and ranges from 25 to 45 in our sample. *Married* was measured by a question about marital status, with 1 indicating that the respondent was married at the time of the interview. *Cohabiting* was coded by either a voluntary response to the marital status question or a yes to a follow-up question, “Are you currently living with a romantic partner?” Single women who are neither married nor in a cohabiting relationship are the reference category. *Education* (in years) is a continuous variable, ranging from 2 to 22 in our sample. *Race/ethnicity* is included as a dummy variable for Black, Hispanic, and “Other race,” with White respondents as the reference category.

2.3 Analytic strategy

We first provide an overview of the sample, comparing descriptive statistics (means/percentages) of all study variables by infertility and parity status. To examine differences within infertility and parity groups, we employed Bonferroni post-hoc tests. The differences in distributions of our variables by parity motivated our decision to run separate multivariate analyses for women with young children and women with 1 or more children.

Our first multivariate analysis uses stereotype logistic regression (SLM) to predict fertility intentions and desires. Both fertility intentions and fertility desires are measured on an ordinal scale, but ordinal logistic regression assumes that the distance between each category is the same (proportional odds) (Long and Freese 2006). We tested the proportional-odds assumption using a likelihood-ratio test and a Brant test. Our sensitivity analyses suggested that coefficients for both fertility intentions and fertility desires were significantly different, which violates the proportional-odds assumption. SLM does not require proportional odds and is therefore a more appropriate method for variables that have unequal distances between response options.

Next, we conducted an ordinary least squares (OLS) regression analysis to examine the predictors of one’s ideal number of children. Because ‘ideal number of children’ is a count variable ranging from 0 to 5, we also performed a sensitivity analysis using a Poisson modeling approach; the results were robust across modeling strategies. We present findings from the more widely used OLS models because they are more intuitive to interpret. Finally, we conducted a logistic regression analysis to estimate associations between infertility statuses and intentions by parity and the odds of giving birth in the three years between waves of the study.

3. Results

Results indicate that infertility experience and identity shape fertility intentions and desires in interesting ways. The means and percentages of fertility intentions, desires for a baby, and ideal number of children are provided in Table 1 by infertility- and parity-group statuses. Findings reveal that childless women at the extremes of the infertility continuum have the highest overall fertility intentions, including women who are not infertile and those who are both medically infertile and self-identifying as infertile; all other groups had negative scores on the intentions variable, indicating that the average response was a no to the question on intentions to have a child. The ‘desire to have a baby’ and ‘ideal number of children’ variables reveal a different pattern, however. Childless women in every group had higher scores for wanting to have a baby than women with children. Women without children who both identify as infertile and meet medical criteria for infertility, however, had the highest mean score ($M=3.36$), indicating their responses to wanting a baby were between “definitely yes” and “probably yes.” Further, when asked how many children would be ideal for them, women who already had at least one child reported more children as ideal across every category of infertility. Among childless women, however, those who experienced infertility and identified as infertile reported the highest ideal number of children ($M=2.35$). At the bivariate level, therefore, it appears that women at the extreme end of infertility – those who are both medically infertile and identifying as infertile – have higher fertility intentions, are more likely to report wanting to have a baby, and report a higher ideal number of children than women in other groups. Women who already had at least one child at the first wave and were not infertile were significantly more likely to have given birth by wave two compared to women who only met the medical criteria and women with children who met the medical criteria and identified as infertile.

Stereotype regression analyses, presented in Table 2, indicate similar patterns. Regardless of parity, women who have experienced a fertility problem and identify as having a fertility problem report higher levels of fertility intentions and desire to have a baby. Fertility status was not associated with significantly higher levels of fertility intentions for childless women. Among women with children, those who self-identified only and those who identified and experienced infertility had higher odds of being very sure that they intend to have more children versus being very sure that they will not have more children relative to women without fertility problems ($OR=2.13$; $OR=1.64$). Among childless women, the odds of definitely wanting a child versus definitely not increase significantly by an odds ratio of 7.62. Among mothers, identifying a fertility problem (with [$OR=4.12$] and without [$OR=2.34$] meeting medical criteria for infertility) the odds of definitely wanting another child are also significantly higher as compared to mothers without fertility problems.

Table 1: Means of dependent and sociodemographic variables by fertility and parity status group for US women 25–45 years of age, N=2,978

	Childless				1+Child					
	Not infertile	Medically infertile	Self-ID infertile	Infertile & ID	Post hoc ²	Not infertile	Medically infertile	Self-ID infertile	Infertile & ID	Post hoc ²
Intentions and desires										
Fertility intentions	.25	-.02	-.02	.10		-.52	-.63	-.41	-.39	
Want to have a baby	2.93	2.93	2.81	3.36	IS>N,M,S	2.26	2.22	2.48	2.66	IS>N,M
Ideal number of children	2.01	2.04	1.89	2.35	IS>N	2.82	2.97	2.79	2.91	
Birth between waves ¹	.23	.09	.28	.26		.33	.16	.31	.13	IS<N; M<N
Sociodemographic variables										
Age (yrs)	32.44	33.80	34.16	35.09	IS>N	33.77	33.95	35.31	35.58	IS>N; IS>M
Married (%)	.31	.38	.35	.54	IS>N	.70	.59	.66	.74	IS<M; M<N
Cohabiting (%)	.15	.18	.09	.15		.09	.12	.13	.09	
Education (yrs)	15.81	14.88	15.31	14.21	IS<N	13.65	13.41	13.85	13.73	
Black (%)	.12	.14	.08	.19		.11	.23	.14	.19	IS>N; M>N
Hispanic (%)	.08	.13	.11	.12		.26	.25	.19	.19	
Other race/ethnicity (%)	.11	.17	.16	.11		.07	.06	.12	.06	
N	608	122	79	200		948	573	80	368	

Source: Wave 1 & Wave 2 National Survey of Fertility Barriers, N=2,978.

¹ Birth between waves reflect a smaller subset of respondents re-interviewed at Wave 2, N=1,547.

² N= Not infertile, M= Medically infertile only, S= Self-ID infertile only, IS= Medically infertile & Self-ID
Means are presented unless otherwise noted.

Table 2: Stereotype logistic regression analyses of the association among fertility-status indicators, sociodemographic characteristics, with fertility intentions and wants by parity status, N=2,978

	Fertility intentions ^a			Want to have a baby ^b							
	Childless			1+Child		Childless		1+Child			
	OR	se	z	OR	se	OR	se	OR	se	z	
Infertility status											
Not Infertile (Ref)											
Medically infertile only	.67	(.30)	(-.89)	.76	(.18)	(-1.15)	1.20	(.47)	.98	(.20)	(-.11)
Self-ID only	.77	(.45)	(-.44)	2.13*	(.80)	(2.01)	1.00	(.46)	2.34*	(.93)	(2.14)
Infertile & Self-ID	1.37	(.56)	(.77)	1.64*	(.44)	(1.84)	7.62***	(3.44)	4.12***	(.96)	(6.07)
Sociodemographic variables											
Age (Centered)	.69***	(.02)	(-12.58)	.78***	(.02)	(-8.28)	.80***	(.02)	(-10.43)	(.02)	(-8.66)
Unmarried (Ref)											
Married	2.99**	(1.04)	(3.16)	1.24	(.34)	(.80)	1.68*	(.50)	.86	(.21)	(-.59)
Cohabiting	1.71	(.75)	(1.22)	1.49	(.53)	(1.12)	1.14	(.45)	.75	(.24)	(-.88)
Education yrs (Centered)	1.13*	(.06)	(2.06)	1.15*	(.07)	(2.40)	1.03	(.05)	1.09*	(.04)	(2.04)
White (Ref)											
Black	2.04*	(.74)	(1.97)	.79	(.28)	(-.65)	1.16	(.36)	.77	(.25)	(-.80)
Hispanic	1.04	(.41)	(.10)	.83	(.29)	(-.53)	1.46	(.58)	1.23	(.31)	(.80)
Other race/ethnicity	2.77	(1.89)	(1.50)	1.10	(.54)	(-.19)	2.75	(1.78)	1.11	(.62)	(-.18)
N	1009			1969			1009		1969		
chi2	182.59***			99.55***			136.80		122.43		

Source: Wave 1 National Survey of Fertility Barriers.

Exponentiated coefficients; Standard errors in parentheses

^a. Fertility Intentions Comparison of Extremes: Very Sure Intend vs. Very Sure Not Intending

^b. Want to Have Baby Comparison of Extremes: Definitely Yes vs. Definitely No

* $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$, **** $p < 0.001$

In Table 3 we present the linear regression analyses of the association between fertility status, sociodemographic variables, and the ideal number of children by parity status. Among childless women, those who were both medically infertile and self-identified as infertile reported wanting more children than their fertile counterparts, supporting the notion that infertility may be driving the higher preferences for childbearing among infertile women. The pattern is not the same among women who already had children (parity of one or more). Among mothers, those who were medically infertile wanted significantly more children than fertile women, net of other factors.

Table 3: OLS regression analyses of the association between fertility, sociodemographic variables, and ideal number of children by parity status, N=2,978

	Ideal number of children			
	Childless		1+Child	
	b	se	b	se
Infertility status				
Not infertile (Ref)				
Medically infertile only	.06	(.13)	.17 [†]	(.07)
Self-ID only	-.05	(.16)	.01	(.13)
Infertile & Self-ID	.41 ^{***}	(.12)	.10	(.08)
Sociodemographic variables				
Age (Centered)	-.05 ^{***}	(.01)	.00	(.01)
Unmarried (Ref)				
Married	.08	(.09)	.17 [†]	(.08)
Cohabiting	-.09	(.12)	.01	(.11)
Education yrs (Centered)	-.02	(.02)	-.05 ^{***}	(.01)
White (Ref)				
Black	.16	(.11)	-.12	(.10)
Hispanic	.10	(.13)	.16 [*]	(.08)
Other race/ethnicity	.13	(.16)	-.32 ^{**}	(.11)
Intercept	1.84 ^{***}	(.08)	2.64 ^{***}	(.08)
<i>N</i>	1009		1969	
<i>R</i> ²	.10		.05	

Source: Wave 1 National Survey of Fertility Barriers.

^{*} $p < 0.10$, [†] $p < 0.05$, ^{**} $p < 0.01$, ^{***} $p < 0.001$

Are fertility and parity statuses associated with the odds of a birth between waves? To answer this question, on Table 4, we present odds ratios from multivariate logistic regression models exploring whether fertility status, intentions, and desires influenced the likelihood of giving birth between waves, controlling for a rich array of sociodemographic characteristics. We ran separate sets of models for childless women and women with children. Our first model examines only fertility status. Among childless women, those who were medically infertile were 66% less likely to give birth between waves, relative to fertile women (OR=.34). And among those who were already mothers, medically infertile women were about 60% less likely to give birth between waves, relative to fertile women (OR=.38). Likewise, mothers who were both medically infertile and self-identified as infertile were significantly less likely than fertile women to have children between waves (OR=.31). Next, we added fertility intentions and desires in model two. Lastly, our third model includes all variables. Fertility intentions and desires were positively associated with the likelihood of having a child between waves. Taking into account fertility intentions and desires largely reduced the difference between fertile and medically infertile mothers. Conversely, mothers who experienced infertility and identified as infertile were significantly less likely to have children, even when we take intentions and desires into account. Among childless women, those who met the medical criteria only remained significantly less likely to give birth between waves compared to fertile childless women. Our findings suggest that birth outcomes differ by parity. Among childless women, those who met the medical criteria only were significantly less likely to give birth between waves. Interestingly, we find that among mothers, women who meet medical criteria for infertility and self-identify as infertile were least likely to meet their expectations, despite having the highest preferences for having children. As expected, women who meet the medical criteria for infertility have significantly fewer children on average than women who do not meet criteria for infertility.

Table 4: Logistic regression analyses of the association between infertility, fertility, sociodemographic variables, and birth between waves by parity status, N=1,547

	Childless						1+ Child					
	m1		m2		m3		m1		m2		m3	
	OR	se	OR	se	OR	se	OR	se	OR	se	OR	se
Infertility and fertility variables												
Not infertile (Ref)												
Medically infertile only	.34 [†]	(.16)	.29 [†]	(.16)	.15 ^{***}	(.09)	.38 ^{***}	(.10)	.56 ⁺	(.18)	.53 ⁺	(.19)
Self-ID only	1.29	(.52)	1.29	(.51)	1.66	(.69)	.92	(.36)	1.20	(.47)	1.57	(.67)
Infertile & Self-ID	1.18	(.35)	1.64	(.64)	1.14	(.45)	.31 ^{***}	(.08)	.39 ^{**}	(.12)	.44 ^{**}	(.13)
Fertility intentions W1 (Centered)			2.28 ^{***}	(.38)	1.71 ^{**}	(.32)			1.71 ^{**}	(.30)	1.47 [†]	(.24)
Want a baby W1 (Centered)			1.63 [†]	(.36)	1.92 ^{**}	(.49)			1.35 ⁺	(.24)	1.37 ⁺	(.25)
Ideal number of children (Centered)			1.23	(.17)	1.19	(.19)			1.19	(.13)	1.16	(.14)
Sociodemographic variables												
Age					.90 ^{***}	(.03)					.91 ^{***}	(.02)
Unmarried (Ref)												
Married					8.32 ^{***}	(3.00)					4.75 ^{***}	(2.05)
Cohabiting					3.23 ^{**}	(1.42)					1.35	(.83)
Education years (Centered)					.92	(.06)					.99	(.05)
White (Ref)												
Black					1.06	(.47)					.86	(.29)
Hispanic					2.32 ⁺	(1.07)					2.40 [†]	(.84)
Other Race					4.48 ^{**}	(2.36)					.77	(.46)
Intercept	.30 ^{***}	(.04)	.13 ^{***}	(.03)	.03 ^{***}	(.01)	.49 ^{***}	(.07)	.34 ^{***}	(.06)	.07 ^{***}	(.03)
N	630		630		630		917		917		917	
pseudo R ²	.02		.24		.36		.04		.18		.26	

Source: Wave 1 and Wave 2 National Survey of Fertility Barriers.

Exponentiated coefficients; Standard errors in parentheses.

Birth between waves reflect a smaller subset of respondents re-interviewed at Wave 2, N=1,547.

[†] $p < 0.10$, ⁺ $p < 0.05$, ^{**} $p < 0.01$, ^{***} $p < 0.001$

4. Conclusions

Despite a growing body of literature on infertility/subfecundity in demographic and social-science fields, little is known about the associations between infertility and fertility intentions, desires, and birth outcomes. Further, extant research typically fails

to differentiate between those who meet the medical criteria but do not realize they are infertile, those who perceive themselves to have a fertility problem although they do not meet the medical criteria for infertility, and those who both meet the medical criteria and identify as infertile. A few exceptions are Greil and colleagues' work on the "hidden infertile" (Greil et al. 2009) and the link between perceiving a fertility problem and distress about infertility (Greil et al. 2011b).

The findings presented here highlight the importance of both the experience of infertility and identification as a person with fertility problems for fertility intentions, desires, and birth outcomes. Supporting theories of intentions that suggest that an outcome needs to be certain to formulate an intention (Ajzen and Klobas 2013; Malle, Moses, and Baldwin 2003), we found that fertility intentions were highest for women who had never experienced infertility and were childless at the time of the first interview. Taken at face value, this might suggest that women who meet the medical criteria or perceive that they are infertile are not trying to get pregnant. Yet when we explored fertility desires further, we found that women who both meet the medical criteria and identify as infertile report significantly greater desires to have a baby. Among childless women, they also have a much higher ideal number of children than other women in the sample. Of course, simply wanting a birth may not predict one, especially for women who have met the medical criteria for infertility in the past. Indeed, our findings indicate that, regardless of parity, women who met the medical criteria for infertility but did not identify as having a fertility problem at the first survey wave were less likely to give birth by the second wave. Women who identified as infertile but did not meet the medical criteria at the first wave were actually more likely to give births between waves than women who were not infertile, though the difference was not significant.

These findings suggest that fertility intentions may not be the best predictors of births for women who have experienced infertility and/or identify as someone with fertility problems. The uncertainty of whether or not they will be able to give birth may prevent them from reporting a strong intention to have a baby. Bachrach and Morgan (2013) have called for greater theory development "at the intersection of cognitive science, social science, and social demography" (page 480) to better understand fertility intentions and their realization. In particular, they argue that fertility researchers need improved understanding of how mental and social phenomena are related to intentions (Bachrach and Morgan 2013). This study provides an example of when intentions may not be as meaningful or predictive for fertility outcomes. Because of the substantial minority of women who experience infertility each year, our findings have important implications for theory and research regarding fertility intentions.

Our findings also highlight the importance of distinguishing between experiencing infertility and identifying as infertile; moreover, these classifications of infertility

matter in different ways for fertility intentions vs. outcomes. Not surprisingly, identity matters for fertility intentions; women who view themselves as having a fertility problem, especially when they also have met the medical criteria for infertility, are reporting lower fertility intentions but greater desires to have a baby and higher family size preferences. This is problematic because women who believe that they have fertility problems may be assigned lower values on fertility intentions measures despite wanting (and perhaps trying) to have a baby, and any births may be misidentified as unplanned or unwanted when they do give birth. Interestingly, it appears that women who fail to realize a fertility problem when, in fact, they meet the criteria for infertility, are least likely to give birth. This finding has important implications as well; this may be one reason why fertility outcomes do not always match intentions. There are also important medical implications, as women who do not identify as having fertility problems may be less likely to seek treatment that could help them achieve a desired birth (White et al. 2006).

Of course, the results of this study should be interpreted with caution in light of its limitations. First, respondents' infertility statuses relied on retrospective reports of periods of 12 months or more during which respondents had regular, unprotected heterosexual intercourse with no pregnancy. To our knowledge, there have been no studies examining the validity of retrospective reports of infertility episodes. Integrating medical records with survey data would provide a more accurate classification of infertility episodes. Given our significant findings for infertility experience, however, we are confident that, if anything, our findings provide conservative estimates. Second, although the study participants were of reproductive age and not surgically sterilized, the timing of the infertility episode(s) varied; in some cases, the infertility episodes occurred more than 10 years before the survey interview. Again, we suspect that, if anything, this might make our findings more conservative than a study that includes only women currently experiencing infertility. Due to the much smaller percentage of women who experience infertility in a given year as compared to women who ever experience infertility, it is difficult to assess these relationships with such small samples. Another limitation related to this issue of small cell counts when creating four groups of infertility statuses is that we were only able to utilize two groups of parity statuses (women with and without children).

Future research should explore infertility-group membership and possible selectivity. It is unclear, for example, why many women who have not met the medical criteria for infertility perceive themselves as infertile. It is also unclear why some women meet the medical criteria for infertility but do not recognize that they might have a fertility problem. Future prospective data-collection efforts are encouraged to include questions that allow for the classification of women into infertility categories, including identity and experience, and follow women over time to more definitively

illustrate the ways in which infertility affects fertility. Future research is also encouraged to investigate the reverse relationship between fertility intentions, wants, and ideals and perception of a fertility problem. The meaning of the period of infertility may matter for outcomes; not all women who are not contracepting are actively “trying” to get pregnant (Johnson et al. 2014). Whereas some women may experience a relatively long period of unprotected sex without conception and not consider it a problem, those who are hoping or trying for a pregnancy may self-identify a problem before they meet the medical criteria for infertility

Despite limitations, our findings suggest that women who meet the medical criteria for infertility and/or identify as having a fertility problem are a special case for fertility-intentions and -outcomes research and theory. This study highlights the need for more in-depth probing of fertility plans, desires, and contraceptive behaviors when women believe that they may not be able to conceive. Simply asking these women if they intend to give birth may be an inadequate question.

References

- Ajzen, I. and Klobas, J. (2013). Fertility intentions: An approach based on the theory of planned behavior. *Demographic Research* 29(8): 203–232. doi:[10.4054/DemRes.2013.29.8](https://doi.org/10.4054/DemRes.2013.29.8).
- American Society for Reproductive Medicine (ASRM). (2008). Definitions of infertility and recurrent pregnancy loss. *Fertility and Sterility* 90(5) supplement: S60. doi:[10.1016/j.fertnstert.2008.08.065](https://doi.org/10.1016/j.fertnstert.2008.08.065).
- Bachrach, C.A. and Morgan, S.P. (2014). A cognitive–social model of fertility intentions. *Population and Development Review* 39(3): 459–485. doi:[10.1111/j.1728-4457.2013.00612.x](https://doi.org/10.1111/j.1728-4457.2013.00612.x).
- Bongaarts, J. and Potter, R.E. (2013). *Fertility, biology, and behavior: An analysis of the proximate determinants*. New York: Academic Press.
- Davis, K. and Blake, J. (1956). Social structure and fertility: An analytic framework. *Economic development and cultural change* 4(2): 211–235.
- Greil, A.L. and McQuillan, J. (2010). ‘Trying times’: Medicalization, intent, and ambiguity in the definition of infertility. *Medical Anthropology Quarterly* 24(2): 137–156. doi:[10.1111/j.1548-1387.2010.01094.x](https://doi.org/10.1111/j.1548-1387.2010.01094.x).
- Greil, A.L., McQuillan, J., Johnson, K., Blevins-Slauson, K., and Shreffler, K.M. (2009). The hidden infertile: Infertile women without pregnancy intent in the United States. *Fertility and Sterility* 93(6): 2080–2083. doi:[10.1016/j.fertnstert.2009.08.024](https://doi.org/10.1016/j.fertnstert.2009.08.024).
- Greil, A.L., McQuillan, J., Lowry, M., and Shreffler, K.M. (2011a). Infertility treatment and fertility specific distress: A longitudinal analysis of a population-based sample of U.S. women. *Social Science and Medicine* 73(1): 87–94. doi:[10.1016/j.socscimed.2011.04.023](https://doi.org/10.1016/j.socscimed.2011.04.023).
- Greil, A.L., Shreffler, K.M., Schmidt, L., and McQuillan, J. (2011b). Variation in distress among women with infertility: Evidence from a population-based sample. *Human Reproduction* 26(8): 2101–2112. doi:[10.1093/humrep/der148](https://doi.org/10.1093/humrep/der148).
- Johnson, K.M., McQuillan, J., Greil, A.L., and Shreffler, K.M. (2014). Towards a more inclusive framework for understanding fertility barriers. In: Nash, M. (ed.). *Reframing reproduction: Sociological perspectives on gender, sexuality, and reproduction in late modernity*. New York: Palgrave Macmillan: 23–38. doi:[10.1057/9781137267139_2](https://doi.org/10.1057/9781137267139_2).

- Long, S.J. and Freese, J. (2006). *Regression models for categorical dependent variables using Stata*. 2nd ed. College Station: Stata Press.
- Mail, C.E. (1994). Community constructs on involuntary childlessness: Sympathy, stigma, and social support. *Canadian Review of Sociology and Anthropology* 31(4): 392–421. doi:[10.1111/j.1755-618X.1994.tb00828.x](https://doi.org/10.1111/j.1755-618X.1994.tb00828.x).
- Malle, B.F., Moses, L.J., and Baldwin, D.A. (2003). *Intentions and intentionality: Foundations of social cognition*. Cambridge: MIT Press.
- Mascarenhas, M.N., Cheung, H., Mathers, C.D., and Stevens, G.A. (2012). Measuring infertility in populations: Constructing a standard definition for use with demographic and reproductive health surveys. *Population Health Metrics* 10(17): 1–11. doi:[10.1186/1478-7954-10-17](https://doi.org/10.1186/1478-7954-10-17).
- Maximova, K. and Quesnel-Vallée, A. (2009). Mental health consequences of unintended childlessness and unplanned births: Gender differences and life course dynamics. *Social Science & Medicine* 68(5): 850–857. doi:[10.1016/j.socscimed.2008.11.012](https://doi.org/10.1016/j.socscimed.2008.11.012).
- McQuillan, J., Greil, A.L., Shreffler, K.M., and Bedrous, A.V. (2015). The importance of motherhood and fertility intentions among women in the United States. *Sociological Perspectives* 58(1): 20–35. doi:[10.1177/0731121414534393](https://doi.org/10.1177/0731121414534393).
- Polis, C.B. and Zabin, L.S. (2012). Missed conceptions or misconceptions? Perceived infertility among unmarried young adults in the United States. *Perspectives on Sexual and Reproductive Health* 44(1): 30–38. doi:[10.1363/4403012](https://doi.org/10.1363/4403012).
- Rutstein, S.O. and Shah, I.H. (2004). *Infecundity, infertility, and childlessness in developing countries*. DHS Comparative Reports No. 9. Calverton: ORC Macro and the World Health Organization.
- Schoen, R., Astone, N.M., Kim, Y.T., Nathanson, C.A., and Fields, J.M. (1999). Do fertility intentions affect fertility behavior? *Journal of Marriage and Family* 61(3): 790–799. doi:[10.2307/353578](https://doi.org/10.2307/353578).
- Schwerdtfeger, K.L. and Shreffler, K.M. (2009). Trauma of pregnancy loss and infertility among mothers and involuntarily childless women in the United States. *Journal of Loss and Trauma* 14(3): 211–227. doi:[10.1080/15325020802537468](https://doi.org/10.1080/15325020802537468).
- Simonsen, S.E., Baksh, L., and Stanford, J.B. (2012). Infertility treatment in a population-based sample: 2004–2005. *Maternal and Child Health Journal* 16(4): 877–886. doi:[10.1007/s10995-011-0809-6](https://doi.org/10.1007/s10995-011-0809-6).

- Stover, J. (1998). Revising the proximate determinants of fertility framework: What have we learned in the past 20 years? *Studies in Family Planning* 29(3): 255–267. doi:10.2307/172272.
- Thoma, M.E., McLain, A.C., Louis, J.F., King, R.B., Trumble, A.C., Sundaram, R., and Buck Louis, G.M. (2013). The prevalence of infertility in the United States as estimated by the current duration approach and a traditional constructed approach. *Fertility and Sterility* 99(5): 1324–1331. doi:10.1016/j.fertnstert.2012.11.037.
- Thomson, E. (1997). Couple childbearing desires, intentions, and births. *Demography* 34(3): 343–354. doi:10.2307/3038288.
- White, L., McQuillan, J., Greil, A.L., and Johnson, D.L. (2006). Infertility: Testing a helpseeking model. *Social Science & Medicine* 62(4): 1031–1041. doi:10.1016/j.socscimed.2005.11.012.

