


1948

## Ground-Water Levels in the Lower Platte River Valley, Nebraska

H. A. Waite

*University of Nebraska-Lincoln*

Follow this and additional works at: <http://digitalcommons.unl.edu/conservationsurvey>

 Part of the [Geology Commons](#), [Geomorphology Commons](#), [Hydrology Commons](#), [Paleontology Commons](#), [Sedimentology Commons](#), [Soil Science Commons](#), and the [Stratigraphy Commons](#)

---

Waite, H. A., "Ground-Water Levels in the Lower Platte River Valley, Nebraska" (1948). *Conservation and Survey Division*. 638.  
<http://digitalcommons.unl.edu/conservationsurvey/638>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Conservation and Survey Division by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# *Nebraska Water Survey Paper*

Number 3

May 1948

**GROUND-WATER LEVELS  
IN THE  
LOWER PLATTE RIVER VALLEY,  
NEBRASKA**

**H. A. WAITE**



**UNIVERSITY OF NEBRASKA  
CONSERVATION AND SURVEY DIVISION**

# NEBRASKA WATER SURVEY PAPER

NUMBER 3

MAY 1948

## GROUND-WATER LEVELS IN THE LOWER PLATTE RIVER VALLEY, NEBRASKA

BY

H. A. WAITE

Library Copy



PUBLISHED BY THE UNIVERSITY OF NEBRASKA  
CONSERVATION AND SURVEY DIVISION, LINCOLN

# THE UNIVERSITY OF NEBRASKA

R. G. GUSTAVSON, *Chancellor*

## Board of Regents

F. W. JOHNSON, <i>President</i> , Lexington	STANLEY D. LONG, Grand Island
C. Y. THOMPSON, <i>V.Pres.</i> , West Point	J. L. WELSH, Omaha
ROBERT W. DEVOE, Lincoln	GEORGE LIGGETT, Utica
JOHN SELLECK, <i>Corporation Secretary</i>	

## CONSERVATION AND SURVEY DIVISION

G. E. Condra, *Chief and State Geologist*

E. C. Reed, *Associate Chief and Associate State Geologist*

As defined by law, the mission of the Conservation and Survey Division of the University is to survey and describe the geology, topography, natural resources, industries, and conservation problems of Nebraska and serve as a factual Information Bureau relating to the conservation and development of the state's resources.

Reports of the Division are published by its departments: Geological Survey, Water Survey, Soil Survey, Resource and Industry Survey, Biological Survey, and Conservation Problem Survey. The Information Bureau service is carried on by correspondence, field examination, hearings, conferences, lectures, special reports, and by use of departmental publications.

### WATER SURVEY PUBLICATIONS

This publication is the third of a series of Water Survey Papers designed to include special reports and articles of less comprehensive nature than Water Survey Bulletins. More formal reports of cooperative investigations in Nebraska are published as Water Supply Papers by the U. S. Geological Survey in cooperation with the Nebraska Water Survey of the Conservation and Survey Division.

This paper was prepared for presentation at the fifty-fifth annual convention of the Nebraska State Irrigation Association held at North Platte, Nebraska, December 4-6, 1947. Immediately after the paper had been presented, Mr. Lloyd C. Thomas, president of the Association, stated that he believed the paper contained information that would be of interest to practically all of the members of the Association and asked Dr. Condra if the paper could be published by his Division, so that copies could be distributed to the members of the Association and others as soon as possible. Dr. Condra replied in the affirmative provided that the members so desired, and provided that approval of the Director of the U. S. Geological Survey could be obtained for publication of the paper. A vote was taken and President Thomas announced that it was unanimous. The approval of the Director of the Geological Survey has been obtained and the paper is herewith published by the Conservation and Survey Division of the University of Nebraska.

# Ground-water Levels in the Lower Platte River Valley, Nebraska<sup>1</sup>

BY HERBERT A. WAITE<sup>2</sup>

Ground-water levels in the Lower Platte River Valley, Nebraska, are of interest because ground water sustains many activities that are of concern to the State, the community, the businessman, and the farmer. Many individuals have a vital stake in the ground-water resources of the Lower Platte River Valley in Nebraska; water from wells plays no small part in the economy of that region. The public water supplies and the rural domestic supplies are obtained from wells. Industry and irrigation, too, depend heavily upon ground water as a basic factor in their profitable operations.

The systematic collection of water-level records is a basic procedure in even the simplest investigation of ground-water resources. Practically all interpretations of changes in storage in ground-water reservoirs are based on these critical ground-water data. Periodic water-level observations are instrumental in supplying a wealth of water facts concerning ground-water withdrawals, the rises and declines of the water table, spacing of wells, possible recharge to underground reservoirs, and other related data. When extended over sufficiently long periods of time, water-level records furnish valuable information in regard to seasonal fluctuations and long-term trends of the water table, the value of which increases as time goes by and demands on ground-water sources become progressively greater.

The first comprehensive studies of the behavior of the water table in the Platte River Valley were begun in 1930 as one phase of the cooperative ground-water program started in that year by the United States Geological Survey and the Conservation and Survey Division of the University of Nebraska. Periodic measurements of the depth to the water table were made at about 100 wells in the segment of the Platte Valley between Gothenburg

<sup>1</sup> Published with the permission of the Director of the Geological Survey, U. S. Department of the Interior.

<sup>2</sup> District Geologist of U. S. Geological Survey in charge of cooperative State-Federal ground-water investigations in Nebraska.

and Grand Island.<sup>3</sup> The wells were selected so that fluctuations of the water table could be observed in all parts of the area. During the early part of the investigation measurements were made in only 60 wells, but other wells were added later as it became desirable to study the fluctuation of the water table in more detail.

As a result of increased interest in ground-water levels, a State-wide program of water-level measurements was begun in 1934, and about 60 of the wells in the Platte Valley on which measurements were begun in 1930 and 1931 were included in this program. In 1945 the scope of the water-level observation program in the Lower Platte River Valley was greatly expanded in connection with a comprehensive program of ground-water investigations in the Missouri River Basin in Nebraska begun by the Ground Water Division, United States Geological Survey, in cooperation with the United States Bureau of Reclamation and with the continued cooperation of the Conservation and Survey Division of the University of Nebraska.

The area considered in this paper is confined to the Platte River Valley bottom lands and terraces from the range line two miles west of North Platte to the range line one mile east of Fremont, and aggregates approximately 2,500 square miles. Periodic measurements of the depth to water in about 260 observation wells are now being made. The locations of all observation wells in the Lower Platte River Valley are shown in figure 1. The Central Nebraska Public Power and Irrigation District and the Loup River Public Power District have installed observation wells in the vicinities of their respective projects and many of these wells are situated in the valley. A total of 38 of these wells has been incorporated in the present water-level observation program. In order to achieve more nearly uniform coverage in the valley, 74 privately owned wells were added to the observation-well program and 66 small-diameter wells, equipped with screened drive points, were installed by the Geological Survey in selected localities. Water-level measurements since 1930 and prior to 1945 in all observation wells in the Platte Valley have been published in

<sup>3</sup> Lugn, A. L., and Wenzel, L. K., Geology and ground-water resources of south-central Nebraska, with special reference to the Platte River Valley between Chapman and Gothenburg; U. S. Geol. Survey Water-Supply Paper 779, pp. 105-108, 1938.

annual reports of the Geological Survey.<sup>4</sup> Subsequent measurements (since January 1, 1945) will be published in similar reports of this series.

The net changes in water level in the lower Platte River Valley for the periods December 1930 to December 1939, December 1939 to December 1946, and for the over-all period December 1930 to December 1946 are shown in three maps which have been prepared for this purpose. (Figs. 2, 3, and 4, respectively.)

The amount of rise or decline in water levels during the respective periods is shown by distinctive patterns. Figure 2 indicates that a net decline of the water table was predominant over much of the area during the period December 1930 to December 1939. By contrast, figure 3 shows that a general net rise in water levels occurred in much of the area during the period December 1939 to December 1946. However, the amount of general rise, as indicated on figure 3, was not sufficient to balance the general decline indicated in all areas of figure 2. This is borne out by figure 4, which covers the over-all period December 1930 to December 1946 and shows net changes in water level for the combined periods of the first two maps.

The amount of the net decline shown in the vicinity of Cozad and Lexington for the period December 1939 to December 1946 is sufficiently small that it is entirely possible that it could be eliminated in one or two seasons of more favorable precipitation. However, it is possible that this decline may be a forerunner of a long-term water-level trend. The decline of the water table in western Hall County, on the other hand, has been progressive, even through the period December 1939 to December 1946 when water levels in the remainder of the Lower Platte River Valley were rising. During the period from 1930 until the present, the net declines in water levels in observation wells in this area ranged up to a maximum of about 7 feet. On the basis of this known water-level history, it would seem that any additional expansion of pump irrigation in this rather critical area should proceed cautiously lest the area be faced eventually with problems of local overdevelopment and attendant increasing pumping lifts.

<sup>4</sup>U. S. Geol. Survey Water-Supply Papers 777 (1935); 817 (1936); 840 (1937); 845 (1938); 886 (1939); 908 (1940); 938 (1941); 946 (1942); 988 (1943); and 1018 (1944).

The net changes in water level are least in the vicinity of the river and are greatest in areas that are farther from the river, where conditions are less favorable for recharge. The Platte River apparently tends to stabilize the water table beneath the lands lying immediately adjacent to it. Fluctuations of stage in the river probably affect the ground-water levels at greater distances from the river in the Platte Valley than in most other valleys, because the water table for several miles adjacent to the river is in approximate equilibrium with the water surface of the river.

The map covering the period December 1939 to December 1946 (fig. 3) shows that the water levels in observation wells in some parts of this area, notably south of the river between North Platte and Lexington, have risen as much as 17 feet. This amounts to a rise of about 12 to 15 feet above the normal rise for the region during this period and suggests that the water table has been artificially raised owing to seepage losses from canals and reservoirs in the vicinity.

The Lower Platte River Valley has been arbitrarily subdivided into six segments for purposes of comparison of changes in ground-water storage. (See fig. 1). The "East of Merrick County" segment includes the valley area between the range line 1 mile east of Fremont and the range line 12 miles west of Columbus. The "Merrick County" segment extends from the range line 12 miles west of Columbus to the east line of Hall County. The "Hall County" segment is restricted to the valley portion of that county and to a very small portion of extreme northwestern Adams County. The "Buffalo County" area includes all of the valley portion of that county and the valley portions of Kearney and Phelps Counties lying directly south of Buffalo County. Likewise, the "Dawson County" segment includes all of the valley portion of that county and the small valley portions of Phelps and Gosper Counties lying directly south of Dawson County. The "Lincoln County" area is restricted to the valley portion of Lincoln County up to the range line 2 miles west of North Platte.

Graphs showing changes in ground-water storage in each of the six segments of the Lower Platte River Valley are presented in figure 5, together with the monthly precipitation at the following stations of the U. S. Weather Bureau: Schuyler, Central City, Grand Island, Kearney, Lexington, and North Platte.






Figure 5. Graphs showing changes in ground-water storage in each of six segments of the Lower Platte River Valley, together with the monthly precipitation in inches at the nearest stations of the U. S. Weather Bureau.

In each graph the curve starts at the zero point—this point representing the total amount of water in ground-water storage on January 1, 1946. Points along the curve which are above the zero line represent times at which ground-water storage was in excess of the amount in storage on January 1, 1946. Likewise, points below the zero line represent times at which ground-water storage was less than the amount in storage at the beginning of 1946. The increases or losses in ground-water storage, in thousands of acre-feet, can be determined by reference to the scale on the left margin of the graphs. Actually, the curves are an index of changes in water level in the respective segments.




Figure 6. Hydrographs showing fluctuations of the water levels in six wells in the Lower Platte River Valley between North Platte and Kearney.




Figure 7. Hydrographs showing fluctuations of the water levels in six wells in the Lower Platte River Valley between Kearney and Grand Island.




Figure 6. Hydrographs showing fluctuations of the water levels in six wells in the Lower Platte River Valley between North Platte and Kearney.

Depths to water level below measuring point, in feet.




Figure 7. Hydrographs showing fluctuations of the water levels in six wells in the Lower Platte River Valley between Kearney and Grand Island.

These long-term water-level records are of interest, because most of them started in the fall of 1930 and have continued without interruption until the present time. In 1930 the water table was relatively high throughout the Lower Platte River Valley, largely as a result of unusually favorable precipitation during that year, with resultant lesser withdrawals from the ground-water reservoir. During the period from about 1931 through 1939, when drought conditions were prevalent, the water levels in observation wells showed net declines. After 1939, with a return to a period of more nearly normal precipitation, the water levels in many of the observation wells showed net rises. During the period from 1940 through 1946, the average annual precipitation for the State was below normal only twice, in 1940 and 1943. In each of the remaining years during this same period the average annual precipitation was above normal. The plus departures from normal precipitation slightly more than balanced the minus departures during this period.

It will be noted that although the water level records of several of the observation wells show annual losses and partial recovery, or even gains, the hydrographs of several wells in the critical area on the north side of the river between Kearney and Grand Island show that the long-term trend is downward despite partial recovery in 1947. The hydrographs of wells 265, 249, and 247 substantiate this fact rather conclusively. This situation exists in an area where there is no recharge from irrigation with river water.

In much of the valley area between Gothenburg and Kearney the ground-water levels have recovered from low stages reached in 1939. An exception, however, is the area in the vicinity of Cozad and Lexington, mentioned earlier, where the water levels in wells show net declines of 1 to 3 feet since 1930. The principal reason that ground-water levels west of Kearney have declined less than they have in the area between Kearney and Grand Island is the fact that the ground-water reservoir in the area between Gothenburg and Kearney receives valuable recharge from surface-water irrigation in that area.

Ground water is used for irrigation throughout the Lower Platte River Valley. The locations of irrigation wells are shown in figure 8. It is noteworthy that approximately 3,930 irrigation wells are shown on this map. As this map was prepared in the

spring of 1947, the number of well locations shown on the map is considerably less than the number of irrigation wells existing at present. The map shows that the greatest concentration of irrigation wells is in Hall County, which has a total of 1,045. Dawson County with 983 wells and Buffalo County with 905 wells are close behind. This heavy development of pump irrigation took place during and immediately following the drought of the 1930's and was particularly heavy in the three counties mentioned.

For the sake of comparison, it should be pointed out that a field census made in 1931 and 1932 in connection with the cooperative ground-water survey showed that about 920 irrigation wells were being operated at that time between Gothenburg and Chapman. The total number of irrigation wells in this section had increased to about 3,275 by 1946, as shown by field surveys made by the Bureau of Reclamation in that year. Thus an increase of nearly 260 per cent is indicated for the period 1932-1946, or an average increase of about 19 per cent per year. The rate of increase in the future is, of course, a matter of conjecture.

It is hoped that the maps and charts presented in this paper will result in a better understanding of the methods used for interpreting the vast amount of basic water-level data that has been collected in the Lower Platte River Valley in Nebraska since the beginning of the cooperative ground-water survey in 1930.

Figure 2

# PLATTE RIVER VALLEY

Changes in Ground-water Levels  
from 1900 to December, 1939






Figure 2

# MAP OF THE LOWER PLATEAU

Showing Net Changes in Ground Water  
December, 1930 to December, 1931




Figure 3

# R PLATTE RIVER VALLEY

Changes in Ground-water Levels  
from September to December, 1946




Figure 3

# MAP OF THE LOWER PLATE

Showing Net Changes in Groundwater  
December, 1939 to December, 1940





Figure 4  
 Changes in Ground-water Levels  
 1930 to December, 1946





U.S. Department of the Interior  
 Geological Survey, Water Resources Branch, Division of Ground Water

Figure 4

# MAP OF THE LOWER PLATTE R

Showing Net Changes in Ground-water  
December, 1930 to December, 1931






U. S. GEOLOGICAL SURVEY

**MAP OF THE LOWER PLATTE VALLEY  
SHOWING  
LOCATIONS OF IRRIGATION WELLS**


COMPILED FROM THE FOLLOWING SOURCES:

1. U.S. Geological Survey Water-Supply Paper 779.
2. Field inventory of irrigation wells by the Conservation and Survey Division of the University of Nebraska, 1941.
3. Mail canvass of County Agricultural Agents by Dr. G. E. Condra, 1941.
4. District Supervisors and others of the Soil Conservation Service.
5. Water Facilities Supervisors of the Farm Security Administration.
6. Nebraska Well Drillers and Pump Manufacturers.
7. Field inventory by the U. S. Geological Survey.
8. Field inventory by the Bureau of Reclamation.



1947

Figure 8. Map of the Lower Platte River Valley showing the locations of irrigation wells.



HOWARD

R11W

MERRICK

R12W

R13W

R14W

R15W

R16W

R17W

R18W

R19W

R20W

R21W

R22W

DAWSON

R21W

R20W

R19W

R18W

BUFFALO

R17W

R16W

R15W

R14W

R13W

HALL

HAMILTON

R6W

R7W

R8W

R9W

R10W

R11W

R12W

R13W

R14W

R15W

R16W

R17W

R18W

R19W

R20W

R21W

R22W

ADAMS

R12W

R11W

R10W

R9W

R8W

R7W

R6W

PHELPS

KEARNEY

R19W

R18W

R17W

R16W

R15W

R14W

R13W

R12W

R11W

R10W

R9W

R8W

R7W

R6W

R5W

R4W

R3W

R2W

R1W

R0W

R0E

R1E

R2E

R3E

R4E

R5E

R6E

R7E

R8E

R9E

R10E

R11E

R12E

R13E

R14E

R15E

R16E

R17E

R18E

R19E

R20E

R21E

R22E

R23E

R24E

R25E

R26E

R27E

R28E

R29E

R30E

R31E

R32E

R33E

R34E

R35E

R36E

R37E

R38E

R39E

R40E

R41E

R42E

R43E

R44E

R45E

R46E

R47E

R48E

R49E

R50E

R51E

R52E

R53E

R54E

R55E

R56E

R57E

R58E

R59E

R60E

R61E

R62E

R63E

R64E

R65E

R66E

R67E

R68E

R69E

R70E

R71E

R72E

R73E

R74E

R75E

R76E

R77E

R78E

R79E

R80E

R81E

R82E

R83E

R84E

R85E

R86E

R87E

R88E

R89E

R90E

R91E

R92E

R93E

R94E

R95E

R96E

R97E

R98E

R99E

R100E

R101E

R102E

R103E

R104E

R105E

R106E

R107E

R108E

R109E

R110E

R111E

R112E

R113E

R114E

R115E

R116E

R117E

R118E

R119E

R120E

R121E

R122E

R123E

R124E

R125E

R126E

R127E

R128E

R129E

R130E

R131E

R132E

R133E

R134E

R135E

R136E

R137E

R138E

R139E

R140E

R141E

R142E

R143E

R144E

R145E

R146E

R147E

R148E

R149E

R150E

R151E

R152E

R153E

R154E

R155E

R156E

R157E

R158E

R159E

R160E

R161E

R162E

R163E

R164E

R165E

R166E

R167E

R168E

R169E

R170E

R171E

R172E

R173E

R174E

R175E

R176E

R177E

R178E

R179E

R180E

R181E

R182E

R183E

R184E

R185E

R186E

R187E

R188E

R189E

R190E

R191E

R192E

R193E

R194E

R195E

R196E

R197E

R198E

R199E

R200E

R201E

R202E

R203E

R204E

R205E

R206E

R207E

R208E

R209E

R210E

R211E

R212E

R213E

R214E

R215E

R216E

R217E

R218E

R219E

R220E

R221E

R222E

R223E

R224E

R225E

R226E

R227E

R228E

R229E

R230E

R231E

R232E

R233E

R234E

R235E

R236E

R237E

R238E

R239E

R240E

R241E

R242E

R243E

R244E

R245E

R246E

R247E

R248E

R249E

R250E

R251E

R252E

R253E

R254E

R255E

R256E

R257E

R258E

R259E

R260E

R261E

R262E

R263E

R264E

R265E

R266E

R2





LINCOLN

CUSTER

DAWSON

BUFFALO

FRONTIER

GOSPER

PHELPS

PLATTE RIVER