



Reduction–oxidation (redox) system in radiation-induced normal tissue injury: molecular mechanisms and implications in radiation therapeutics

R. Yahyapour¹ · E. Motevaseli² · A. Rezaeyan³ · H. Abdollahi³ · B. Farhood⁴ · M. Cheki⁵ · S. Rezapoor⁶ · D. Shabeeb^{7,8} · A. E. Musa⁹ · M. Najafi¹⁰  · V. Villa¹¹

Received: 22 October 2017 / Accepted: 27 December 2017 / Published online: 9 January 2018

© Federación de Sociedades Españolas de Oncología (FESEO) 2018

Abstract

Every year, millions of cancer patients undergo radiation therapy for treating and destroying abnormal cell growths within normal cell environmental conditions. Thus, ionizing radiation can have positive therapeutic effects on cancer cells as well as post-detrimental effects on surrounding normal tissues. Previous studies in the past years have proposed that the reduction and oxidation metabolism in cells changes in response to ionizing radiation and has a key role in radiation toxicity to normal tissue. Free radicals generated from ionizing radiation result in upregulation of cyclooxygenases (COXs), nitric oxide synthase (NOSs), lipoxygenases (LOXs) as well as nicotinamide adenine dinucleotide phosphate oxidase (NADPH oxidase), and their effected changes in mitochondrial functions are markedly noticeable. Each of these enzymes is diversely expressed in multiple cells, tissues and organs in a specific manner. Overproduction of reactive oxygen radicals (ROS), reactive hydroxyl radical (ROH) and reactive nitrogen radicals (RNS) in multiple cellular environments in the affected nucleus, cell membranes, cytosol and mitochondria, and other organelles, can specifically affect the sensitive and modifying enzymes of the redox system and repair proteins that play a pivotal role in both early and late effects of radiation. In recent years, ionizing radiation has been known to affect the redox functions and metabolism of NADPH oxidases (NOXs) as well as having destabilizing and detrimental effects on directly and indirectly affected cells, tissues and organs. More noteworthy, chronic free radical production may continue for years, increasing the risk of carcinogenesis and other oxidative stress-driven degenerative diseases as well as pathologies, in addition to late effect complications of organ fibrosis. Hence, knowledge about the mechanisms of chronic oxidative damage and injury in affected cells, tissues and organs following exposure to

M. Najafi and V. Villa both share the senior authorship.

✉ M. Najafi
masoudnajafi67@yahoo.com

¹ School of Medicine, Jiroft University of Medical Sciences, Jiroft, Iran

² Department of Molecular Medicine, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

³ Department of Medical Physics, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

⁴ Departments of Medical Physics and Radiology, Faculty of Paramedical Sciences, Kashan University of Medical Sciences, Kashan, Iran

⁵ Department of Radiologic Technology, Faculty of Paramedicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁶ Department of Radiology, Faculty of Paramedical, Tehran University of Medical Sciences, Tehran, Iran

⁷ Department of Medical Physics and Biomedical Engineering, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

⁸ Department of Physiology, College of Medicine, University of Misan, Amarah, Iraq

⁹ Research Center for Molecular and Cellular Imaging, Tehran University of Medical Sciences, Tehran, Iran

¹⁰ Radiology and Nuclear Medicine Department, School of Paramedical Sciences, Kermanshah University of Medical Science, Kermanshah, Iran

¹¹ Scientific Research Department, Armed Forces Radiobiology Research Institute (AFRRI), Uniformed Services University of Health Sciences (USUHS), Bethesda, MD 20889-5603, USA

ionizing radiation may help in the development of treatment and management strategies of complications associated with radiotherapy (RT) or radiation accident victims. Thus, this medically relevant phenomenon may lead to the discovery of potential antioxidants and inhibitors with promising results in targeting and modulating the ROS/NO-sensitive enzymes in irradiated tissues and organ injury systems.

Keywords Radiation · Redox · Normal tissue injury · Inflammation · NADPH oxidase

Introduction

Annually, more than 12 million new cases of cancer are reported worldwide [1]. Approximately, more than half of these patients require radiation therapy (RT) either alone or in combination with other modalities such as surgery, immunotherapy, hyperthermia, chemotherapy and hormone therapy. In addition to clinical applications, several people are exposed to lethal or sub-lethal doses of ionizing radiation resulting from radiation accident or terrorist activities [2, 3]. So far, several studies have been conducted to understand the mechanisms of minimizing the detrimental effects of exposure to ionizing radiation (IR) on normal tissues.

Exposing cells to IR causes immediate free radical formation with a nanoseconds half-life. For many years, it has been believed that these free radicals, as well as direct radiation interaction with DNA, are responsible for the side effects of exposure to IR. However, the discovery of new phenomena in radiobiology such as radiation-induced bystander effect, non-targeted effect and genomic instability, have challenged this central dogma [4]. In recent years, studies have demonstrated that changes in the normal functions of the reduction/oxidation (redox) systems are involved in several damages following exposure to IR [5]. Free radical production by the redox system begins some few hours after exposure and may continue for several years [6, 7]. The redox system plays a key role in acute radiation syndrome. It is also responsible for several early and late effects of exposure to IR such as bystander effect, out-of-field effect, inflammation, fibrosis and others [8].

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are the main sources of damage to normal tissues after exposure to IR. During normal cell function, ROS and RNS are essential mediators for several cellular processes such as immune responses, cell signaling, microbial defense, differentiation, cell adhesion, apoptosis and others [9]. Antioxidant systems include enzymes such as superoxide dismutase (SOD) and glutathione peroxidase (GPX) as well as peptides such as glutathione (GSH). They neutralize additional free radicals and protect cells against the detrimental effects. Exposure to IR causes excessive production of free radicals over the antioxidant system potency, resulting in oxidative damage to DNA, proteins and lipids. These effects cause damages to normal cell function and

may lead to genomic instability that increases the risk of malignancies [10].

Redox system biology and enzymology, redox chemistry and ionizing radiation interactions

Free radicals including ROS, ROH and RNS are recognized as dual role players which can have both deleterious and beneficial consequences. Free radicals formed via redox system biology can act as messengers in cell signaling and changes in gene expression patterns. For example, several types of protein kinases and transcription factors are stimulated by oxidation reactions, while protein phosphatases are inactivated. These changes result in the activation of several ROS/RNS-producing enzymes which may continue for hours, days, months or years [5]. So far, several types of oxidoreductases with the ability to produce superoxide in cells have been identified. They include cyclooxygenase (COX), lipoxygenase, nitric oxide synthase (NOS), cytochrome P450 enzymes, xanthine oxidase, NADPH oxidase and mitochondrial electron transfer chain. It is generally accepted that mitochondrial ROS production and expression of other genes involved in redox system such as NF- κ B, COX-2, iNOS and NADPH oxidase amplify each other [11].

A large number of studies have been conducted to reveal the roles of interacting redox biology and chemistry systems in oxidative-mediated damages induced by IR. Recent studies have demonstrated that redox activity is involved in both early and late effects of exposure to IR. Moreover, the results of different studies have indicated that the expression of genes involved in the redox system is tissue dependent. Hence, an understanding of the specific functions of such redox biological and chemical systems' interactions with IR is specifically described further.

Radiation-induced inflammation triggers redox activation

Inflammation plays a key role in redox activation. Exposure of normal cells directly to IR or ROS will result in both nucleus and mitochondria DNA damages, which may cause cell death through apoptosis, mitotic catastrophe or necrosis.

Mitotic catastrophe is not immunogenic, while apoptosis and necrosis trigger immune activation through the release of danger signals [12]. Although apoptotic bodies are omitted by macrophages and do not activate inflammatory response, necrotic cells release various signals to immune cells including mast cells and lymphocytes which will result in the secretion of inflammatory cytokines [13]. Apoptosis can trigger the release of anti-inflammatory cytokines such as IL-10 and TGF- β , while necrosis may lead to the release of inflammatory cytokines such as IL-1, IL-6, IL-8, IL-13, IL-33 and TNF- α , as well as other inflammatory mediators [14]. In the absence of suppression of these responses by the immune system, chronic inflammation may continue for a long time after exposure. This is associated with chronic oxidative damage, which lead to genomic instability and damage to the normal function of organs [15] (Fig. 1).


NADPH oxidases (NOXs) functions and interactions

NADPH oxidase enzymes are a group of oxidoreductases that transfer electron from NADPH to oxygen molecules. So far, several subtypes of these enzymes have been detected within cells. NADPH oxidase genes have

been discovered in both phagocytic and non-phagocytic cells and are involved in immune system responses and inflammation. It was suggested that the NADPH oxidase system is involved in signaling pathways that mediate cell growth, cell survival and death. In response to inflammatory stimuli, macrophages and neutrophils produce superoxide via the phagocytic NADPH oxidase (NOX). On the other hand, ROS produced by non-phagocytic isoforms of NADPH oxidase (such as NOX2 in the membrane) are involved in regulating intracellular signaling cascades in various types of non-phagocytic cells [16]. NOX1-5, DUOX1 and DUOX2 are the most important subtypes of NADPH oxidase enzymes involved in respiratory burst following exposure to radiation. In contrast to other sources of ROS, NADPH oxidase (NOX) is a professional and distinct ROS generator through conversion of O₂ to superoxide anion O₂⁻ [17, 18].

These enzymes have high stability, resulting in continuous ROS production following exposure. Each of these enzymes is activated in specific tissue type of cells. In addition, enzymes can be stimulated by a large group of stressors and stimuli, as well as different types of cytokines and growth factors. A large group of inflammatory cytokines, chemokines and hormones such as IL-1, TNF- α , TGF- β , IFN- γ and angiotensin II are implicated in NOX system activation [19].

Fig. 1 Mechanisms of cell death trigger inflammation and free radical production after exposure to ionizing radiation



NOX1

NOX1 is expressed in several types of cells such as endothelial cells, in the placenta, prostate and uterus, and osteoclasts, as well as in some malignancies such as melanoma and colon cancer [20–22]. Hence, overexpression of NOX1 may be involved in some malignancies. In addition to risk of carcinogenesis, the role of NOX1 in radiation fibrosis has been proposed. Choi et al. showed that inhibiting NOX1 but not NOX2 or NOX4 ameliorates collagen deposition and pulmonary fibrosis markers such as α -SMA and fibroblast-specific protein 1 (FSP1) in C57BL/6J mice following lung irradiation. Furthermore, ROS production following NOX1 inhibition decreased significantly [23].

NOX2

There are some evidences that NOX2 expressions in both phagocytic and non-phagocytic cells are regulated by some cytokines and growth factors such as IFN- γ , TGF- β and IL-12 [16, 24]. Studies have demonstrated that some mediators such as TLR-2 have an intermediate role in NOX2 activation. In response to IR, NOX2 has an important role in continuous ROS production. Narayanan et al. showed that irradiating human lung fibroblasts with alpha particle produces O_2^- and H_2O_2 . Analysis showed that the plasma membrane-bound NOX2 is primarily responsible for O_2^- and H_2O_2 production [25]. The upregulation of upstream genes such as NF-KB, Raf-1, ERK 1/2, c-Jun, p38, activator protein 1 (AP-1) and calcium signaling is involved in this process. Scavenging of free radicals by SOD or catalase inhibited these signaling pathways and subsequent NOX2 upregulation [26–28]. The involvement of NOX2 in radiation-induced salivary gland injury was demonstrated by Kim et al. They showed that exposing rats to 18 Gy of X-ray (2 Gy/min) increases NOX2 gene expression in salivary glands at least 7 days after irradiation. Their results indicated that apoptotic genes such as caspase-9 as well as MAPKs including p-38 and JNK are involved in NOX2 signaling cascades [29]. NOX2 is involved in persistent ROS production in the intestine as well. Datta et al. have shown that irradiating mice with gamma rays and high LET ^{56}Fe radiation causes stimulation of ROS production in the intestinal epithelial cells for 1 year after exposure. The result of this study indicated that NOX1, NOX2 and mitochondria malfunction are responsible for persistent oxidative damage [30].

NOX3

Studies conducted to depict the role of NOX3 in radiation damage are very limited. A study by Shin et al. showed that exposure to radiation upregulates the expression of NOX3

in the oral mucosa of rat. Increased NOX3 expression was associated with necrotic inflammatory exudates and ulceration in the oral mucosa [31].

NOX4

NOX4 is one of the most important subtypes of the NOX system in response to IR. Some studies have proposed a role for NOX4 in radiation-induced bone marrow toxicity. Evaluating different ROS/NO producing enzymes in mice bone marrow showed that increase in NOX4 activity following exposure has a central role for ROS production and bone marrow stem cells damage after exposure. Analysis showed persistent ROS production 8 weeks after exposure. Inhibiting the NOX system resulted in better survival and decreased bone marrow damage [32–34]. As TGF- β has a pivotal role in bone marrow toxicity following exposure to radiation, it seems that TGF- β –NOX4 pathway is responsible for the continuous ROS/NO production and subsequent genomic instability in bone marrow following exposure [35]. Moreover, studies have shown that other mediators such as TLR-4 and MyD88 have an intermediate role in NOX4 activation. The C-terminal region of NOX4 has an interaction with the tail of TLR-4. However, it was proposed that other mediators such as MyD88 and IRAK were involved in this pathway. These interactions are essential for ROS production from NOX4 [36].

NOX5

NOX5-induced ROS production is due to Ca^{2+} flux through Ca^{2+} -binding sites [37]. This gene is absent in rodents. Evidences for its role in radiation oxidative damages are limited. A study by Weyemi et al. on human primary fibroblasts have revealed that inhibition of both NOX4 and NOX5 leads to reduced levels of DNA damage associated with increased cell survival. Results showed that the levels of protection by inhibition of these genes are similar to administering two potent radioprotectors: *N*-acetylcysteine (NAC) and fulvene-5 [38].

DUOX1/DUOX2

Some evidences support the role of DUOX1/DUOX2 in chronic oxidative stress and subsequent consequences of IR such as fibrosis. However, data for responses of these genes to IR are very limited. IFN- γ , IL-4 and IL-13 have pivotal roles in the upregulation of DUOX1 and DUOX2. Hassani et al. showed that DUOX1 gene expression is upregulated for several days after exposing human thyrocytes to radiation. Analysis showed that IL-13-p38 MAPK is responsible for persistent DUOX1-induced H_2O_2 production following irradiation [39]. IL-4 and IL-13 induce DUOX2 production as

well as increased ROS production. IL-4–STAT6 pathway is responsible for upregulating DUOX2. The increased expression of these genes may be involved in the development of pancreatic and gastrointestinal malignancies [40]. Further studies are needed to depict possible roles of these genes in radiation-induced carcinogenesis and other side effects.

COX-2

COX-2 has a central role in inflammatory responses which convert arachidonic acid liberated from membrane phospholipids to prostaglandins (PGs). During the synthesis of PGE₂, production of ROS is a common secondary effect of arachidonic acid metabolism [41]. Several studies have indicated that upregulation of COX-2 is involved in different toxicities following exposure to IR [42]. Increased COX-2 expression was reported for its association with radiation toxicity in the gastrointestinal system such as the intestine and colon [43]. Furthermore, upregulated COX-2 gene expression is involved in radiation toxicities in the lung, heart, brain, kidney and others [44]. COX-2 upregulation causes accumulation of immune system cells and appearance of inflammation signs. Inflammatory cells including macrophages and lymphocytes further enhance oxidative damage through secretion of NO and ROS. These changes cause activation of matrix metalloproteinases (MMPs) which change the normal function of tissues through deposition of collagen and fibronectin. Thus, COX-2 mediates pathological damages induced by ionizing radiation such as fibrosis, atherosclerosis and vascular damage [45].

It was suggested that COX-2 can stimulate carcinogenesis through signal modulation involved in cell proliferation and apoptosis [46]. In addition, ROS produced by COX-2 play a key role in mutagenesis and genomic instability. Increased COX-2 expression has been reported for its association with several malignancies such as breast, esophageal, gastric, colorectal and lung cancer [47]. Some studies have also shown the role of COX-2 in oxidative damage in non-irradiated cells. Overexpression of COX-2 in the distant lung and bronchial after pelvis or abdominal irradiation in rats and mice have been reported. These effects lead to increased ROS production and DNA oxidative damage through bystander and non-targeted responses [48, 49].

Lipoxygenases

Lipoxygenases (LOXs) are iron-containing enzymes which catalyze the deoxygenation of unsaturated fatty acids. This process is associated with ROS production and initiating lipoperoxidation of membranes as well as some changes in the cell metabolism [50]. Thus, LOXs can stimulate DNA damage, genomic instability and also cell death, especially apoptosis [51]. In mammals, increased LOXs have been

found to be associated with some inflammatory diseases and cancer. A study by Matyshevskaia et al. showed an increase in LOX activity of lymphocytes within 6 h post-irradiation of Wistar rats with 1 Gy X-ray. They also observed that inhibiting LOX activity leads to a remarkable reduction in DNA damage. The results of this study proposed that LOX activity was involved in ROS production during early hours after exposure [52]. In another study, Grichenko et al. have shown that LOX activity was obvious 1 h after exposure but not at later times [53].

Nitric oxide in DNA damage and inflammation

Nitric oxide (NO) is an important mediator which affects a number of targets within cells. iNOS is the main source of NO during stress conditions such as inflammation and plays a key role in oxidative stress and carcinogenesis. NO is generated by macrophages via iNOS enzyme in response to inflammatory stimulus. NO is highly reactive, interacting with mitochondria-derived superoxide to form higher reactive peroxynitrite. On the other hand, high level of NO competes with O₂ in the ETC and may suppress respiration in the mitochondria. This effect may lead to increased superoxide generation which amplifies oxidative stress [54].

Several studies have indicated that reactive nitrogen species (RNS) are involved in radiation-induced normal tissue injury [55–57]. In addition, NO has a pivotal role in radiation-induced bystander and non-targeted effects [48, 58]. Increased NO level following radiation exposure has shown association with DNA damage and genomic instability [59]. A study by Ohta et al. proposed that increase in NO level has a direct relation to radiation dose. Moreover, they observed that increased serum level of NO occurs in the early hours after exposure [55]. However, it seems that increased NO production by iNOS in exposed tissues continues for several days or months after exposure. This is associated with long-term pathological changes in irradiated tissues [60].

One important mechanism for NO-induced normal tissue damage is nitro-acetylation and subsequent epigenetic changes in some enzymes such as DNA repair enzymes. NO produced by macrophages and neutrophils has a role in inhibiting DNA repair enzymes involved in the base excision repair (BER) and mismatch repair (MMR) pathways [61]. Elevated NO production which can be seen following exposure to radiation may suppress the activity of some DNA repair enzymes in these pathways [62]. Studies have proposed that chronic upregulation of nitric oxide synthases (NOS) especially iNOS, causing nitro-acetylation and decreased half-life of ogg1 and AGT [63, 64]. These effects on DNA repair responses result in incomplete repair of DNA damage which may provide grounds for mutation, chromosomal instability, genomic instability and finally carcinogenesis [65, 66].

Mitochondrial functions as an energy and free radical reservoir

Mitochondria sources energy for the cells through reducing oxygen to water and synthesizing adenosine triphosphate (ATP). This process is known as oxidative phosphorylation. It accounts for the consumption of 90% of all oxygen taken up by mammals [67]. Oxidative phosphorylation occurs within the electron transport chains, a series of enzymes embedded within the inner mitochondrial membrane. During electron transportation, a percentage of the oxygen molecules undergo a one-electron reduction to superoxide [68]. In normal conditions, antioxidant defense systems such as catalase, superoxide dismutase and glutathione peroxidases neutralize superoxide and form free radicals [69]. Thus, this system protects cells from oxidative damage resulting from mitochondria activity. O_2 radicals may be released to the cytosol to generate reactive oxygen species. Furthermore, these radicals may react with NO, producing peroxynitrite [70].

Although several studies have shown that the main source for radiation-induced ROS production is the mitochondria, its activation mechanisms have not been recognized completely. A study by Yamamori et al. indicated that irradiating human lung carcinoma A549 cells increases the mitochondrial contents of the cells. Furthermore, they showed that exposure of cells to radiation increases the mitochondrial membrane potential and also stimulates the mitochondrial electron transport chain (ETC) function [71]. In addition, some studies reported that irradiation leads to an increase in the mitochondrial mass in different cell lines [72, 73].

An investigation by Tulard et al. revealed a persistent and dose-dependent increase in mitochondrial ROS in human colon cells after exposure to gamma radiation [74]. Evaluation of intracellular and mitochondrial ROS after irradiation has shown that a dose-dependent increase in both sources of ROS is due to radiation. However, analysis showed a difference in the intracellular and mitochondrial ROS profile. Intracellular ROS increased soon after exposure, subsided after 24 h, and thereafter increased for another 3 days. In contrast to intracellular ROS, mitochondrial ROS level rises gently and attains its peak 3 days after exposure. This may remain high for a week after irradiation [75]. These results have been confirmed by other studies [76–78]. It was suggested that mitochondrial ROS are involved in DNA damage and tumor development [79]. In different tumor cells, the increased level of mitochondrial ROS was confirmed. For example, it was observed that mtDNA mutations are associated with malignancies such as lung, stomach and breast cancer as well as leukemia and lymphoma [80, 81].

However, the exact mechanisms for increased oxidative phosphorylation in mitochondria remain to be elucidated.

Mitochondrial dysfunction, inflammation and intracellular ROS play a key role. Increased the level of ROS production stimulates mitochondria response through a phenomenon known as ROS-induced ROS. Secretion of calcium ions into cytosol plays a key role in this signaling. Following secretion of Ca^{2+} in the endoplasmic reticulum to cytosol during stress conditions, these ions are accumulated in the mitochondria, leading to the disruption of the normal ETC activity which causes increase in ROS production [82, 83].

Mitochondria malfunction is another reason for ROS production. It was proposed that mutation in mtDNA and ETC are involved in this process. Yoshida et al. have shown that exposure of rats' A7r5 cells to 5 Gy gamma rays causes decreased activity of complex I and increased ROS production. Complex I (NADH dehydrogenase) is the most important complex involved in the release of ROS from the ETC [78]. In another study, Dayal et al. showed that dysfunction of complex II after exposure to 10 Gy X-rays played a key role in ROS production and oxidative stress in GM10115 cells. Irradiation of the heart of mice with 2 Gy has shown that succinate-stimulated respiration decreased significantly compared to the control mice. This study has shown that cytochrome C as well as ETC1 and three activities were reduced [84]. This could be because unstable mitochondria produce more hydrogen peroxide compared to normal cells, resulting from decreased respiratory rate [85]. Evidences show that the NADPH oxidase system and mitochondria have a synergic effect on each other in response to stress signaling [86]. Also, studies proposed that the ROS-derived NOX system is involved in mitochondrial dysfunction and subsequent ROS production in this organelle [87, 88].

Epigenetics of redox activation

Evidences indicate that the redox system is related to epigenetic regulation. On one hand, free radicals including both ROS and NO regulate epigenetic processes such as DNA methylation, histone methylation and acetylation. On the other hand, changes in some miRNAs can increase or decrease oxidative damage [89]. Exposure to IR increases the expression of miRNAs involved in ROS production such as let-7 family, mir-15b, mir-21, mir-128 and mir-636 [90–92]. The best example for redox activation of epigenetic modulators is upregulation of mir-21 following exposure to radiation and bystander cells. mir-21 is activated in oxidative stress conditions, as well as following upregulation of some cytokines such as TGF- β [93]. mir-21 induces oxidative stress via targeting of SOD and TNF [94]. Following exposure to radiation, it can suppress detoxification of superoxide by targeting SOD3. In addition, it subdues the regulation of TNF-a, resulting in reducing SOD2 levels [94].

The role of mir-21 in ROS production in bystander cells has also been investigated. Xu et al. showed that the upregulation of mir-21 in bystander MRC-5 cells is associated with increased oxidative and DNA damage. Their results showed that expression of SOD3 was reduced significantly [95]. Tian et al. [96] achieved similar results for MnSOD or SOD2 in bystander WS1 cells following alpha particle irradiation of HaCaT keratinocytes. TGF-β is responsible for the oxidative damage in bystander cells through mir-21. An investigation by Jiang et al. [97] showed that inhibiting TGF-βR1 via a selective inhibitor resulted in abolishing mir-21 and oxidative stress in bystander cells. Since free radicals can upregulate mir-21 and TGF-β, it is possible that mir-21 via a positive loop feedback plays a role in continuous oxidative damage following exposure to ionizing radiation (Fig. 2).

Hypoxia

Hypoxia results from vascular damage and tissue injury. Some changes associated with hypoxia have a regulatory role in tissue remodeling and wound healing. Following exposure to IR, increased inflammatory cytokines and chemokines stimulate accumulation of macrophages in the injured area. This phenomenon is associated with increased oxygen consumption by activated macrophages leading to a low oxygen state and hypoxia in the injured tissues. This hypoxic state stimulates free radical production of some redox system agents such as Ca²⁺ flux, mitochondria and NADPH oxidase enzymes [98, 99]. Evidences indicate that

hypoxia is involved in chronic oxidative stress and radiation-induced late normal tissue injury such as fibrosis [100, 101]. These and many other factors are responsible for this development.

Bystander/non-targeted effect-induced redox activation

Radiation-induced bystander effect refers to a phenomenon in which irradiated cells secrete signals to adjacent non-irradiated cells that cause damage to them. Studies have shown that ROS and NO play a pivotal role in DNA damage and genomic instability in bystander cells. Results from several studies proposed that upregulating some genes involved in redox system such as COX-2, iNOS, NADPH oxidase and also mitochondria plays a central role in this phenomenon [11]. COX-2 and iNOS have a synergic effect on each other [102]. Furthermore, ROS and NO produced by these enzymes increase the activity of the electron transfer chain (ETC) in the mitochondria. Several experiments have been conducted to reveal the signaling pathways that cause oxidative damage in bystander cells. Exposing cells to radiation cause damages to DNA and other structures such as the membrane and mitochondria. These damages result in the release of various products such as exosomes, miRNAs, oxidized DNA and other danger alarms from irradiated cells. In response to these products, macrophages and lymphocytes release several cytokines such as IL-1, IL-6, IL-8, IL-33,




Fig. 2 Mechanisms of redox system activation following exposure to radiation

TNF- α , TGF β and others [103]. Migration of these signals to other cells leads to long-term detrimental changes that may give rise to secondary effects of ionizing radiation on normal tissues. In vitro studies have revealed that gap junction intercellular communication (GJIC) plays a key role in the transfer of bystander signals to non-irradiated cells [104–106]. Inhibiting GJIC by specific inhibitors resulted in the suppression of DNA damage in bystander cells [107, 108].

Several studies have shown that inflammatory responses to DNA damage, DNA repair and cell death play a key role in the secretion of factors involved in activating redox systems [8, 109]. As regards the ability of several of these factors to migrate to distant organs, it is acceptable that redox system activity increases in non-targeted organs. The released clastogenic signals can migrate to distant tissue and stimulate ROS/NO production [110]. Several in vivo studies have shown that local irradiation of a limited area causes ROS production and oxidative damage in out-of-field tissues [48, 111]. A study by Chai et al. showed that the TGF- β –TGF β R1–COX-2 pathway plays an important role in ROS production and oxidative DNA damage in distant lung tissues. However, non-targeted induced oxidative damage affects non-irradiated tissues in a tissue specific manner. The expression of TGF β R1 and its cascades such as COX-2 and ROS in the lung was obvious, but not for the liver [112]. As oxidative DNA damage causes hypomethylation and other epigenetic changes, this effect may result in a tissue-specific epigenetic change in non-irradiated tissues. For example, cranial irradiation resulted in long-term hypomethylation and changes in miRNAs profile in rat's spleen. These changes were not observed in the skin. A better understanding of the basic mechanisms of this phenomenon may help improvements in the therapeutic ratio of RT [113].

Genomic instability is a phenomenon seen in several types of malignancies. It is associated with chromosomal aberrations such as increased mutation frequency within the genome, damage to DNA repair genes, alteration in mitochondrial function, mutation in mitochondrial DNA, change in energy balance within cells, attenuation of antioxidant enzymes and so on [114–120]. These abnormal changes in cells lead to persistent ROS production, oxidative stress and mutation in chromosome, making cells susceptible to carcinogenesis [121].

Evidences indicated that most cancers have a type of genetic instability. This phenomenon in many tumors causes a large number of genetic alterations. Although, so far, the mechanisms involved in genomic instability remain unknown, the results of several studies have proposed a role for ROS. A study by Limoli et al. has shown that clones derived from cells exposed to ionizing radiation have abnormal increase in ROS levels. In addition, the number of dysfunctional mitochondria was higher compared to normal

clones [122]. A further study observed that using some scavengers such as DMSO and cationic thiol cysteamine reduces genomic instability after irradiation. This implies the role of continuous ROS production following exposure to radiation in the induction of genomic instability. Furthermore, results indicated that the abnormal functions of mitochondria have a key role in chronic oxidative stress [85]. Genomic instability induced by different types of radiations such as low and high LET radiations has been revealed in both in vitro and in vivo studies [123] (Fig. 3).

Targeting of the redox system for mitigation of radiation-induced normal tissue injury

Normal tissue protection during RT is an active area of research in radiation oncology. It is widely known that radiation induces a wide range of clinical disorders which reduce the outcomes of RT. For many years, radiation protectors have attracted a great deal of attention and various agents have been tested for different tissues [124]. It has been observed that the responses of different organs to IR are distinct [49]. On the other hand, results of a large number of studies have indicated that no agent can protect all organs against IR. Moreover, in clinical applications, protection of normal tissues against RT is a complex biological process and finding selective drugs requires advanced biological experiments. In recent years, researchers have investigated a variety of approaches to obtain the best biological mechanism of radiation protection. It was revealed that these mechanisms are tissue dependent and differ based on the structure of organs as well as functions, in addition to the immune system response [125]. Hence, selection of an appropriate radioprotector is based on tissue responses.

In this present radiation biology era, to find the best clinical radioprotector, new approaches such as redox mechanisms of irradiated organs are of particular interest [109]. As a good example, targeting NOX2 and NOX4 genes has a critical role in bone marrow sparing following exposure to radiation. Several studies have indicated that some radiation modifiers cause reduction of NOX4 gene expression and amelioration of bone marrow toxicity following exposure. Guoshun et al. showed that total body irradiation (TBI) of mice results in long-term upregulation of NOX4 and increase in ROS production in bone marrow hematopoietic stem cells (HSCs). These changes were involved in radiation-induced chronic oxidative damage and long-term injury in the bone marrow. Moreover, they showed that treatment with metformin significantly attenuated ROS production and ameliorated micronuclei formation via NOX4 downregulation in HSCs [126]. Similar results were obtained after administering 5-methoxytryptamine- α -lipoic acid and resveratrol [127, 128]. Moreover, inhibition of these genes may




Fig. 3 Mechanisms of radiation-induced bystander/non-targeted effect

protect the lungs and cardiovascular system from fibrosis [129].

NO plays a key role in radiation-induced chronic injury in irradiated cells, as well as bystander cells [130, 131]. iNOS as the main source of NO is expressed in several tissues such as lungs and gastrointestinal organs. Hence, inhibiting this enzyme is an approach for normal tissue protection against inflammatory responses to ionizing radiation. Some inhibitors like *N*-nitro-*L*-arginine methyl ester and amino guanidine have shown ability to ameliorate radiation injury in the lung [132, 133]. In another study by Erbil et al., it was observed that treatment with *N*(omega)-nitroarginine methyl ester (L-NAME), which is another type of iNOS inhibitor, can ameliorate radiation-induced enteritis [134]. However, in the gastrointestinal system, NOX1, NOX2, cyclooxygenase-2 (COX-2) and mitochondria are other sources of ROS [109, 135].

Targeting mitochondria-induced ROS has been proposed by some studies for mitigating radiation injury. Rwigema et al. evaluated three types of mitochondrial targeting drugs including JP4-039, MCF201-89 and BEB55. The study was conducted as both in vitro for mice hematopoietic progenitor cell line and in vivo for mice. Their results showed that administering these drugs after exposure to a lethal dose of radiation (9.5 Gy) can reduce death of progenitor cells as well as increase survival of irradiated mice [136]. It has been shown that targeting mitochondria is associated with arresting cell cycle in G1, leading to increased DNA damage repair [137]. Reducing apoptosis, especially

in apoptosis-prone cells like hematopoietic system cells is another property of mitochondria-targeting agents, which can prolong survival [138]. Similar results have been confirmed for other mitochondria-targeting agents [139, 140] (Table 1).

Summary and conclusion

This review has presented the mechanisms of ROS production and oxidative damage derived from some enzymes and mitochondria. It is clear that these systems contribute to acute and late deleterious effects of ionizing radiation. In addition, results from several studies propose that ROS as well as NO production by mitochondria and inflammatory cells is involved in oxidative damage to bystander cells and non-targeted tissues. Although the roles of some inflammatory mediators such as lipoxygenases, COX-2 and iNOS have been confirmed in radiation toxicity, recent studies indicate that mitochondria malfunction and upregulation of NADPH oxidase enzymes have pivotal roles. NADPH oxidase enzymes including NOXs and DUOX1-2 are H₂O₂-producing enzymes with abilities to stimulate continuous oxidative damage and genomic instability for a long time after exposure. However, recent studies have highlighted the roles of these enzymes in other consequences of exposure to radiation such as pneumonitis, fibrosis and vascular injury. With regard to the role of NADPH oxidase in radiation-induced oxidative stress, it is

Table 1 Redox enzyme responses in radiation-induced normal tissue injury

Target	Organs	Effects	Inhibitors	Refs.
NOX1	Lung, intestine	Fibrosis, enteritis, oxidative damage	Apocynin, diphenyleneiodonium (DPI)	[23]
NOX2	Salivary gland, intestine	Mucositis, enteritis, oxidative damage	Apocynin, DPI	[29]
NOX3	Oral mucosa	Mucositis, ulceration	Apocynin, DPI	[31]
NOX4	Bone marrow, fibroblast cells	Genomic instability, senescence of stem cells, hematopoietic system syndrome	Metformin, resveratrol, melatonin, <i>N</i> -acetylcysteine, fulvene-5, DPI	[126]
NOX5	Fibroblast cells	Genomic instability	<i>N</i> -acetylcysteine, fulvene-5	[38]
DUOX1&2	Thyroid	Chronic oxidative stress, fibrosis	IC ₅₀	[141]
COX-2	Joints, heart, lung	Arthritis, pneumonitis, DNA damage	Celecoxib, resveratrol	[142]
iNOS	Lung, intestine	Enteritis, pneumonitis	<i>N</i> -nitro- <i>L</i> -arginine methyl ester, amino guanidine, <i>L</i> -NAME	[109, 135]
Mitochondria	Bone marrow, gastrointestinal system, pulmonary epithelial cells	Enteritis, oxidative DNA damage, fibrosis	Metformin, JP4-039, MCF201-89, BEB55, cationic rhodamine 19	[143, 144]

possible to detect involvement of some other sub-families of NADPH oxidase in radiation injury after some years. This is very vital for normal tissue regeneration in the abdominal or cardiovascular systems.

In recent years, numerous studies have indicated that epigenetic modulators play a key role in redox induced normal tissue injury following exposure to ionizing radiation. Upregulation of some immune mediators such as TGF- β and free radicals stimulates the expression of various miRNAs. For example, TGF- β , which has a direct relationship with radiation dose and oxidative damage, upregulates the expression of miRNA21. On the other hand, miRNA21 can suppress SOD activity, resulting in the amplification of oxidative damage induced by inflammatory mediators. The interaction of ionizing radiation, inflammation and epigenetic modulators is very complicated and requires further studies to illustrate this complicated interrelationship between them.

Despite the complicated interrelationships between these factors, it has been confirmed that all mentioned inflammatory mediators, free radicals and some epigenetic modulators amplify each other, resulting in a positive feedback loop. During these interactions, free radicals continually attack genome and cell structure. In addition, oxidative damage changes the expression of various genes involved in the long-term detrimental effects of radiation. Modulation of ROS production from these interactions has been shown to ameliorate lethality, pathological damages as well as genomic instability. Hence, the management of each of these enzymes depends on the irradiated organs. Moreover, knowledge of the mechanisms of redox activation in each organ can help in the production of novel radioprotectors and mitigators with higher efficacy.

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Disclaimer All opinions are the personal and professional opinions of the authors and are not the opinions of their respective academic affiliations and agencies. Masoud Najafi and Vilmar Villa share both senior authorship.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA Cancer J Clin*. 2017;67(1):7–30.
2. Mettler FA Jr, Voelz GL. Major radiation exposure—what to expect and how to respond. *N Engl J Med*. 2002;346(20):1554–61.
3. Ring JP. Radiation risks and dirty bombs. *Health Phys*. 2004;86(2 Suppl):S42–7.
4. Mothersill C, Seymour CB. Radiation-induced bystander effects and the DNA paradigm: an “out of field” perspective. *Mutat Res*. 2006;597(1–2):5–10.
5. Spitz DR, Azzam EI, Li JJ, Gius D. Metabolic oxidation/reduction reactions and cellular responses to ionizing radiation: a unifying concept in stress response biology. *Cancer Metastasis Rev*. 2004;23(3–4):311–22.
6. Sieber F, Muir SA, Cohen EP, North PE, Fish BL, Irving AA, et al. High-dose selenium for the mitigation of radiation injury: a pilot study in a rat model. *Radiat Res*. 2009;171(3):368–73.
7. Zhao W, Robbins ME. Inflammation and chronic oxidative stress in radiation-induced late normal tissue injury: therapeutic implications. *Curr Med Chem*. 2009;16(2):130–43.
8. Najafi M, Shirazi A, Motevaseli E, Geraily Gh, Norouzi F, Heidari M, Rezapoor S. The melatonin immunomodulatory actions in radiotherapy. *Biophys Rev*. 2017;9(2):139–48.

9. Bae YS, Oh H, Rhee SG, Yoo YD. Regulation of reactive oxygen species generation in cell signaling. *Mol Cells*. 2011;32(6):491–509.
10. Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative stress and antioxidant defense. *World Allergy Organ J*. 2012;5(1):9–19.
11. Najafi M, Fardid R, Hadadi G, Fardid M. The mechanisms of radiation-induced bystander effect. *J Biomed Phys Eng*. 2014;4(4):163–72.
12. Pugin J. How tissue injury alarms the immune system and causes a systemic inflammatory response syndrome. *Ann Intensive Care*. 2012;2:27.
13. Frey B, Hehlgans S, Rodel F, Gaipf US. Modulation of inflammation by low and high doses of ionizing radiation: implications for benign and malign diseases. *Cancer Lett*. 2015;368(2):230–7.
14. Frey B, Ruckert M, Deloch L, Ruhle PF, Derer A, Fietkau R, et al. Immunomodulation by ionizing radiation-impact for design of radio-immunotherapies and for treatment of inflammatory diseases. *Immunol Rev*. 2017;280(1):231–48.
15. Yahyapour R, Amini P, Rezapoor S, Rezaeyan A, Farhood B, Cheki M, et al. Targeting of inflammation for radiation protection and mitigation. *Curr Mol Pharmacol*. 2018. <https://doi.org/10.2174/1874467210666171108165641>.
16. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev*. 2007;87(1):245–313.
17. Drummond GR, Selemidis S, Griendling KK, Sobey CG. Combating oxidative stress in vascular disease: NADPH oxidases as therapeutic targets. *Nat Rev Drug Discov*. 2011;10(6):453–71.
18. Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol*. 2004;4(3):181–9.
19. Panday A, Sahoo MK, Osorio D, Batra S. NADPH oxidases: an overview from structure to innate immunity-associated pathologies. *Cell Mol Immunol*. 2015;12(1):5–23.
20. Fu X-J, Peng Y-B, Hu Y-P, Shi Y-Z, Yao M, Zhang X. NADPH oxidase 1 and its derived reactive oxygen species mediated tissue injury and repair. *Oxid Med Cell Longev*. 2014;2014:282854. <https://doi.org/10.1155/2014/282854>.
21. Sun Z, Liu F. Association of Nox1 and vinculin with colon cancer progression. *Cancer Invest*. 2013;31(4):273–8.
22. Liu F, Garcia AMG, Meyskens FL. NADPH oxidase 1 overexpression enhances invasion via matrix metalloproteinase-2 and epithelial–mesenchymal transition in melanoma cells. *J Invest Dermatol*. 2012;132(8):2033–41.
23. Choi S-H, Kim M, Lee H-J, Kim E-H, Kim C-H, Lee Y-J. Effects of NOX1 on fibroblastic changes of endothelial cells in radiation-induced pulmonary fibrosis. *Mol Med Rep*. 2016;13(5):4135–42.
24. Jendrysik MA, Vasilevsky S, Yi L, Wood A, Zhu N, Zhao Y, et al. NADPH oxidase-2 derived ROS dictates murine DC cytokine-mediated cell fate decisions during CD4 T helper-cell commitment. *PLoS ONE*. 2011;6(12):e28198.
25. Narayanan P, Goodwin EH, Lehnert B. α particles initiate biological production of superoxide anions and hydrogen peroxide in human cells. *Cancer Res*. 1997;57(18):3963–71.
26. Azzam EI, de Toledo SM, Spitz DR, Little JB. Oxidative metabolism modulates signal transduction and micronucleus formation in bystander cells from α -particle-irradiated normal human fibroblast cultures. *Cancer Res*. 2002;62(19):5436–42.
27. Little J, Azzam E, De Toledo S, Nagasawa H. Bystander effects: intercellular transmission of radiation damage signals. *Radiat Prot Dosim*. 2002;99(1–4):159–62.
28. Paillas S, Ladjohounlou R, Lozza C, Pichard A, Boudousq V, Jarlier M, et al. Localized irradiation of cell membrane by auger electrons is cytotoxic through oxidative stress-mediated nontargeted effects. *Antioxid Redox Signal*. 2016;25(8):467–84.
29. Kim JH, Kim KM, Jung MH, Jung JH, Kang KM, Jeong BK, et al. Protective effects of alpha lipoic acid on radiation-induced salivary gland injury in rats. *Oncotarget*. 2016;7(20):29143–53.
30. Datta K, Suman S, Kallakury BV, Fornace AJ Jr. Exposure to heavy ion radiation induces persistent oxidative stress in mouse intestine. *PLoS ONE*. 2012;7(8):e42224.
31. Shin YS, Shin HA, Kang SU, Kim JH, Oh YT, Park KH, et al. Effect of epicatechin against radiation-induced oral mucositis: in vitro and in vivo study. *PLoS ONE*. 2013;8(7):e69151.
32. Pazhanisamy SK, Li H, Wang Y, Batinic-Haberle I, Zhou D. NADPH oxidase inhibition attenuates total body irradiation-induced haematopoietic genomic instability. *Mutagenesis*. 2011;26(3):431–5.
33. Wang Y, Liu L, Pazhanisamy SK, Li H, Meng A, Zhou D. Total body irradiation causes residual bone marrow injury by induction of persistent oxidative stress in murine hematopoietic stem cells. *Free Radic Biol Med*. 2010;48(2):348–56. <https://doi.org/10.1016/j.freeradbiomed.2009.11.005>.
34. Chang J, Feng W, Wang Y, Luo Y, Allen AR, Koturbash I, et al. Whole-body proton irradiation causes long-term damage to hematopoietic stem cells in mice. *Radiat Res*. 2015;183(2):240–8.
35. Zhang H, Y-a Wang, Meng A, Yan H, Wang X, Niu J, et al. Inhibiting TGF β 1 has a protective effect on mouse bone marrow suppression following ionizing radiation exposure in vitro. *J Radiat Res*. 2013;54(4):630–6.
36. Deng S, Yu K, Zhang B, Yao Y, Wang Z, Zhang J, et al. Toll-like receptor 4 promotes NO synthesis by upregulating GCHI expression under oxidative stress conditions in sheep monocytes/macrophages. *Oxid Med Cell Longev*. 2015;2015:359315. <http://doi.org/10.1155/2015/359315>.
37. Banfi B, Molnar G, Maturana A, Steger K, Hegedus B, Demareux N, et al. A Ca(2+)-activated NADPH oxidase in testis, spleen, and lymph nodes. *J Biol Chem*. 2001;276(40):37594–601.
38. Weyemi U, Redon CE, Aziz T, Choudhuri R, Maeda D, Parekh PR, et al. Inactivation of NADPH oxidases NOX4 and NOX5 protects human primary fibroblasts from ionizing radiation-induced DNA damage. *Radiat Res*. 2015;183(3):262–70.
39. Ameziane-El-Hassani R, Talbot M, de Souza Dos Santos MC, Al Ghuzlan A, Hartl D, Bidart J-M, et al. NADPH oxidase DUOX1 promotes long-term persistence of oxidative stress after an exposure to irradiation. *Proc Natl Acad Sci USA*. 2015;112(16):5051–6.
40. Wu Y, Doroshov JH. Abstract 5358: IL-4/IL-13 induce Duox2/DuoxA2 expression and reactive oxygen production in human pancreatic and colon cancer cells. *Cancer research*. 2014;74(19 Supp):5358.
41. Song J, Wei Y, Chen Q, Xing D. Cyclooxygenase 2-mediated apoptotic and inflammatory responses in photodynamic therapy treated breast adenocarcinoma cells and xenografts. *J Photochem Photobiol B*. 2014;134:27–36.
42. Steinauer KK, Gibbs I, Ning S, French JN, Armstrong J, Knox SJ. Radiation induces upregulation of cyclooxygenase-2 (COX-2) protein in PC-3 cells. *Int J Radiat Oncol Biol Phys*. 2000;48(2):325–8.
43. Wang D, DuBois RN. The role of COX-2 in intestinal inflammation and colorectal cancer. *Oncogene*. 2010;29(6):781–8.
44. Laube M, Kniess T, Pietzsch J. Development of antioxidant COX-2 inhibitors as radioprotective agents for radiation therapy—a hypothesis-driven review. *Antioxidants (Basel)*. 2016;5(2):14.
45. Rezaeyan A, Haddadi GH, Hosseinzadeh M, Moradi M, Najafi M. Radioprotective effects of hesperidin on oxidative damages and histopathological changes induced by X-irradiation in rats heart tissue. *J Med Phys*. 2016;41(3):182–91.

46. Lee Y-K, Park SY, Kim Y-M, Lee WS, Park OJ. AMP kinase/cyclooxygenase-2 pathway regulates proliferation and apoptosis of cancer cells treated with quercetin. *Exp Mol Med*. 2009;41(3):201–7.
47. Sobolewski C, Cerella C, Dicato M, Ghibelli L, Diedrich M. The role of cyclooxygenase-2 in cell proliferation and cell death in human malignancies. *Int J Cell Biol*. 2010;2010:215158.
48. Yahyapour R, Motevaseli E, Rezaeyan A, Abdollahi H, Farhood B, Cheki M, et al. Mechanisms of radiation bystander and non-targeted effects: implications to radiation carcinogenesis and radiotherapy. *Curr Radiopharm*. 2017. <https://doi.org/10.2174/1874471011666171229123130>.
49. Chai Y, Calaf GM, Zhou H, Ghandhi SA, Elliston CD, Wen G, et al. Radiation induced COX-2 expression and mutagenesis at non-targeted lung tissues of gpt delta transgenic mice. *Br J Cancer*. 2013;108(1):91–8. <https://doi.org/10.1038/bjc.2012.498>.
50. Maccarrone M. Lipoxygenases, apoptosis, and the role of antioxidants. In: Demmig-Adams B, Adams WW, Mattoo AK, editors. *Photoprotection, photoinhibition, gene regulation, and environment*. Dordrecht: Springer; 2008. p. 321–32.
51. Grichenko O, Shaposhnikova V, Levitman M, Kudriavtsev A, Korystov I. A study of the role of lipoxygenases in radiation-induced apoptosis of thymocytes. Inhibitory analysis. *Radiatsionnaia biologiiia, radioecologiiia/Rossiiskaia akademiia nauk*. 2003;44(1):27–31.
52. Matyshevskaya OP, Pastukh VN, Solodushko VA. Inhibition of lipoxygenase activity reduces radiation-induced DNA fragmentation in lymphocytes. *Radiats Biol Radioecol*. 1999;39(2–3):282–6.
53. Grichenko OE, Pushin AC, Shaposhnikova VV, Levitman M, Korystov IuN. Analysis of 15-lipoxygenase activity in irradiated thymocytes. *Izv Akad Nauk Ser Biol*. 2004;5:517–21.
54. Aktan F. iNOS-mediated nitric oxide production and its regulation. *Life Sci*. 2004;75(6):639–53.
55. Ohta S, Matsuda S, Gunji M, Kamogawa A. The role of nitric oxide in radiation damage. *Biol Pharm Bull*. 2007;30(6):1102–7.
56. Kiang JG, Smith JT, Agravante NG. Geldanamycin analog 17-DMAG inhibits iNOS and caspases in gamma-irradiated human T cells. *Radiat Res*. 2009;172(3):321–30.
57. Malaviya R, Gow AJ, Francis M, Abramova EV, Laskin JD, Laskin DL. Radiation-induced lung injury and inflammation in mice: role of inducible nitric oxide synthase and surfactant protein D. *Toxicol Sci*. 2015;144(1):27–38.
58. Yakovlev VA. Role of nitric oxide in the radiation-induced bystander effect. *Redox Biol*. 2015;6:396–400.
59. Yakovlev VA. Nitric oxide dependent downregulation of BRCA1 expression promotes genetic instability. *Cancer Res*. 2013;73(2):706–15.
60. Gaiad A, Lehnert SM, Chehayeb B, Chehayeb D, Kaplan I, Sheinouda G. Inducible nitric oxide synthase and nitrotyrosine in mice with radiation-induced lung damage. *Am J Clin Oncol*. 2003;26(4):e67–72.
61. Lahtz C, Pfeifer GP. Epigenetic changes of DNA repair genes in cancer. *J Mol Cell Biol*. 2011;3(1):51–8.
62. Kidane D, Chae WJ, Czochor J, Eckert KA, Glazer PM, Bothwell ALM, et al. Interplay between DNA repair and inflammation, and the link to cancer. *Crit Rev Biochem Mol Biol*. 2014;49(2):116–39.
63. Wink DA, Laval J. The Fpg protein, a DNA repair enzyme, is inhibited by the biomediator nitric oxide in vitro and in vivo. *Carcinogenesis*. 1994;15(10):2125–9.
64. Rezapoor S, Shirazi A, Abbasi S, Bazzaz JT, Izadi P, Rezaeejam H, et al. Modulation of radiation-induced base excision repair pathway gene expression by melatonin. *J Med Phys*. 2017;42(4):245–50.
65. Shinmura K, Kohno T, Kasai H, Koda K, Sugimura H, Yokota J. Infrequent mutations of the hOGG1 gene, that is involved in the excision of 8-hydroxyguanine in damaged DNA, in human gastric cancer. *Cancer Sci*. 1998;89(8):825–8.
66. Chevillard S, Radicella JP, Levalois C, Lebeau J, Poupon M-F, Oudard S, et al. Mutations in OGG1, a gene involved in the repair of oxidative DNA damage, are found in human lung and kidney tumours. *Oncogene*. 1998;16(23):3083–6.
67. Rolfe DF, Brown GC. Cellular energy utilization and molecular origin of standard metabolic rate in mammals. *Physiol Rev*. 1997;77(3):731–58.
68. Brand MD, Affourtit C, Esteves TC, Green K, Lambert AJ, Miwa S, et al. Mitochondrial superoxide: production, biological effects, and activation of uncoupling proteins. *Free Radic Biol Med*. 2004;37(6):755–67.
69. Tian L, Cai Q, Wei H. Alterations of antioxidant enzymes and oxidative damage to macromolecules in different organs of rats during aging. *Free Radic Biol Med*. 1998;24(9):1477–84.
70. Packer MA, Murphy MP. Peroxynitrite formed by simultaneous nitric oxide and superoxide generation causes cyclosporin-A-sensitive mitochondrial calcium efflux and depolarisation. *Eur J Biochem*. 1995;234(1):231–9.
71. Yamamori T, Yasui H, Yamazumi M, Wada Y, Nakamura Y, Nakamura H, et al. Ionizing radiation induces mitochondrial reactive oxygen species production accompanied by upregulation of mitochondrial electron transport chain function and mitochondrial content under control of the cell cycle checkpoint. *Free Radic Biol Med*. 2012;53(2):260–70.
72. Nugent SME, Mothersill CE, Seymour C, McClean B, Lyng FM, Murphy JEJ. Increased mitochondrial mass in cells with functionally compromised mitochondria after exposure to both direct γ radiation and bystander factors. *Radiat Res*. 2007;168(1):134–42.
73. Wang L, Kuwahara Y, Li L, Baba T, Shin R-W, Ohkubo Y, et al. Analysis of common deletion (CD) and a novel deletion of mitochondrial DNA induced by ionizing radiation. *Int J Radiat Biol*. 2007;83(7):433–42.
74. Tulard A, Hoffschir F, de Boisferon FH, Luccioni C, Bravard A. Persistent oxidative stress after ionizing radiation is involved in inherited radiosensitivity. *Free Radic Biol Med*. 2003;35(1):68–77.
75. Kobashigawa S, Suzuki K, Yamashita S. Ionizing radiation accelerates Drp1-dependent mitochondrial fission, which involves delayed mitochondrial reactive oxygen species production in normal human fibroblast-like cells. *Biochem Biophys Res Commun*. 2011;414(4):795–800.
76. Hosoki A, Yonekura S-I, Qing-Li Z, Zheng-Li W, Takasaki I, Tabuchi Y, et al. Mitochondria-targeted superoxide dismutase (SOD2) regulates radiation resistance and radiation stress response in HeLa cells. *J Radiat Res*. 2012;53(1):58–71.
77. Ogura A, Oowada S, Kon Y, Hirayama A, Yasui H, Meike S, et al. Redox regulation in radiation-induced cytochrome *c* release from mitochondria of human lung carcinoma A549 cells. *Cancer Lett*. 2009;277(1):64–71.
78. Yoshida T, Goto S, Kawakatsu M, Urata Y, Li T-S. Mitochondrial dysfunction, a probable cause of persistent oxidative stress after exposure to ionizing radiation. *Free Radic Res*. 2012;46(2):147–53.
79. Choi KM, Kang CM, Cho ES, Kang SM, Lee SB, Um HD. Ionizing radiation-induced micronucleus formation is mediated by reactive oxygen species that are produced in a manner dependent on mitochondria, Nox1, and JNK. *Oncol Rep*. 2007;17(5):1183–8.
80. Copeland WC, Wachsmann JT, Johnson FM, Penta JS. Mitochondrial DNA alterations in cancer. *Cancer Invest*. 2002;20:557–69.

81. Laurent A, Nicco C, Chereau C, et al. Controlling tumor growth by modulating endogenous production of reactive oxygen species. *Cancer Res.* 2005;65:948–56.
82. Hajnoczky G, Csordas G, Das S, Garcia-Perez C, Saotome M, Sinha Roy S, et al. Mitochondrial calcium signalling and cell death: approaches for assessing the role of mitochondrial Ca^{2+} uptake in apoptosis. *Cell Calcium.* 2006;40(5–6):553–60.
83. Murphy MP. Mitochondrial dysfunction indirectly elevates ROS production by the endoplasmic reticulum. *Cell Metab.* 2013;18(2):145–6.
84. Barjaktarovic Z, Schmaltz D, Shyla A, Azimzadeh O, Schulz S, Haagen J, et al. Radiation-induced signaling results in mitochondrial impairment in mouse heart at 4 weeks after exposure to X-rays. *PLoS ONE.* 2011;6(12):e27811.
85. Kim GJ, Fiskum GM, Morgan WF. A role for mitochondrial dysfunction in perpetuating radiation-induced genomic instability. *Cancer Res.* 2006;66(21):10377–83.
86. Dikalov S. Cross talk between mitochondria and NADPH oxidases. *Free Radic Biol Med.* 2011;51(7):1289–301.
87. Ago T, Kuroda J, Pain J, Fu C, Li H, Sadoshima J. Upregulation of Nox4 by hypertrophic stimuli promotes apoptosis and mitochondrial dysfunction in cardiac myocytes. *Circ Res.* 2010;106(7):1253–64.
88. Kowluru RA, Mishra M. Oxidative stress, mitochondrial damage and diabetic retinopathy. *Biochim Biophys Acta.* 2015;1852(11):2474–83.
89. Afanas'ev I. Mechanisms of superoxide signaling in epigenetic processes: relation to aging and cancer. *Aging Dis.* 2015;6(3):216–27.
90. Simone NL, Soule BP, Ly D, Saleh AD, Savage JE, Degraff W, et al. Ionizing radiation-induced oxidative stress alters miRNA expression. *PLoS ONE.* 2009;4(7):e6377.
91. Ye L, Yu G, Wang C, Du B, Sun D, Liu J, et al. MicroRNA128a, BMI1 polycomb ring finger oncogene, and reactive oxygen species inhibit the growth of U87 MG glioblastoma cells following exposure to X-ray radiation. *Mol Med Rep.* 2015;12(4):6247–54.
92. Chaudhry MA, Omaruddin RA, Brumbaugh CD, Tariq MA, Pourmand N. Identification of radiation-induced microRNA transcriptome by next-generation massively parallel sequencing. *J Radiat Res.* 2013;54(5):808–22.
93. Ling M, Li Y, Xu Y, Pang Y, Shen L, Jiang R, et al. Regulation of miRNA-21 by reactive oxygen species-activated ERK/NF-kappaB in arsenite-induced cell transformation. *Free Radic Biol Med.* 2012;52(9):1508–18.
94. Zhang X, Ng W-L, Wang P, Tian L, Werner E, Wang H, et al. MicroRNA-21 modulates the levels of reactive oxygen species levels by targeting SOD3 and TNF α . *Can Res.* 2012;72(18):4707–13.
95. Xu S, Ding N, Pei H, Hu W, Wei W, Zhang X, et al. MiR-21 is involved in radiation-induced bystander effects. *RNA Biol.* 2014;11(9):1161–70.
96. Tian W, Yin X, Wang L, Wang J, Zhu W, Cao J, et al. The key role of miR-21-regulated SOD2 in the medium-mediated bystander responses in human fibroblasts induced by alpha-irradiated keratinocytes. *Mutat Res.* 2015;780:77–85.
97. Jiang Y, Chen X, Tian W, Yin X, Wang J, Yang H. The role of TGF- β 1–miR-21–ROS pathway in bystander responses induced by irradiated non-small-cell lung cancer cells. *Br J Cancer.* 2014;111(4):772–80.
98. Rathore R, Zheng YM, Niu CF, Liu QH, Korde A, Ho YS, et al. Hypoxia activates NADPH oxidase to increase [ROS] i and [Ca $^{2+}$] through the mitochondrial ROS-PKCepsilon signaling axis in pulmonary artery smooth muscle cells. *Free Radic Biol Med.* 2008;45(9):1223–31.
99. Marshall C, Mamary AJ, Verhoeven AJ, Marshall BE. Pulmonary artery NADPH-oxidase is activated in hypoxic pulmonary vasoconstriction. *Am J Respir Cell Mol Biol.* 1996;15(5):633–44.
100. Halberg N, Khan T, Trujillo ME, Wernstedt-Asterholm I, Attie AD, Sherwani S, et al. Hypoxia-inducible factor 1alpha induces fibrosis and insulin resistance in white adipose tissue. *Mol Cell Biol.* 2009;29(16):4467–83.
101. Vujaskovic Z, Anscher MS, Feng QF, Rabbani ZN, Amin K, Samulski TS, et al. Radiation-induced hypoxia may perpetuate late normal tissue injury. *Int J Radiat Oncol Biol Phys.* 2001;50(4):851–5.
102. Choi YJ, Kim HS, Lee J, Chung J, Lee JS, Choi JS, et al. Down-regulation of oxidative stress and COX-2 and iNOS expressions by dimethyl lithospermate in aged rat kidney. *Arch Pharm Res.* 2014;37(8):1032–8.
103. Hei TK, Zhou H, Ivanov VN, Hong M, Lieberman HB, Brenner DJ, et al. Mechanism of radiation-induced bystander effects: a unifying model. *J Pharm Pharmacol.* 2008;60(8):943–50.
104. Edwards GO, Botchway SW, Hirst G, Wharton CW, Chipman JK, Meldrum RA. Gap junction communication dynamics and bystander effects from ultrasoft X-rays. *Br J Cancer.* 2004;90(7):1450–6.
105. Suzuki M, Tsuruoka C. Heavy charged particles produce a bystander effect via cell–cell junctions. *Biol Sci Space.* 2004;18(4):241–6.
106. Shao C, Furusawa Y, Aoki M, Ando K. Role of gap junctional intercellular communication in radiation-induced bystander effects in human fibroblasts. *Radiat Res.* 2003;160(3):318–23.
107. Bishayee A, Rao DV, Howell RW. Evidence for pronounced bystander effects caused by nonuniform distributions of radioactivity using a novel three-dimensional tissue culture model. *Radiat Res.* 1999;152(1):88–97.
108. Azzam EI, de Toledo SM, Little JB. Direct evidence for the participation of gap junction-mediated intercellular communication in the transmission of damage signals from α -particle irradiated to nonirradiated cells. *Proc Natl Acad Sci USA.* 2001;98(2):473–8.
109. Najafi M, Shirazi A, Motevaseli E, Rezaeyan A, Salajegheh A, Rezapoor S. Melatonin as an anti-inflammatory agent in radiotherapy. *Inflammopharmacology.* 2017;25(4):403–13.
110. Morgan WF. Non-targeted and delayed effects of exposure to ionizing radiation: I. Radiation-induced genomic instability and bystander effects in vitro. 2003. *Radiat Res.* 2012;178(2):Av223–36.
111. Ghobadi A, Shirazi A, Najafi M, Kahkesh MH, Rezapoor S. Melatonin ameliorates radiation-induced oxidative stress at targeted and nontargeted lung tissue. *J Med Phys.* 2017;42(4):241.
112. Chai Y, Lam RK, Calaf GM, Zhou H, Amundson S, Hei TK. Radiation-induced non-targeted response in vivo: role of the TGFbeta-TGFBR1-COX-2 signalling pathway. *Br J Cancer.* 2013;108(5):1106–12.
113. Najafi MSA, Rezaeyan A. Bystander effect and second primary cancers following radiotherapy: what are its significances? *J Med Phys.* 2017;42:55–6.
114. Negrini S, Gorgoulis VG, Halazonetis TD. Genomic instability—an evolving hallmark of cancer. *Nat Rev Mol Cell Biol.* 2010;11(3):220–8.
115. Ferguson LR, Chen H, Collins AR, Connell M, Damia G, Dasgupta S, et al. Genomic instability in human cancer: Molecular insights and opportunities for therapeutic attack and prevention through diet and nutrition. *Semin cancer biol.* 2015;35(Suppl):S5–24.
116. Goel A, Nagasaka T, Arnold CN, Inoue T, Hamilton C, Niedzwiecki D, et al. The CpG island methylator phenotype and chromosomal instability are inversely correlated in sporadic colorectal cancer. *Gastroenterology.* 2007;132(1):127–38.

117. Chatterjee A, Dasgupta S, Sidransky D. Mitochondrial subversion in cancer. *Cancer Prev Res*. 2011;4(5):638–54.
118. Cook CC, Kim A, Terao S, Gotoh A, Higuchi M. Consumption of oxygen: a mitochondrial-generated progression signal of advanced cancer. *Cell Death Dis*. 2012;3(1):e258.
119. Konki M, Pasumarthy K, Malonzo M, Sainio A, Valensisi C, Söderström M, et al. Epigenetic silencing of the key antioxidant enzyme catalase in karyotypically abnormal human pluripotent stem cells. *Sci rep*. 2016;6:22190.
120. Rooney S, Alt FW, Lombard D, Whitlow S, Eckersdorff M, Fleming J, et al. Defective DNA repair and increased genomic instability in Artemis-deficient murine cells. *J Exp Med*. 2003;197(5):553–65.
121. Ziech D, Franco R, Pappa A, Panayiotidis MI. Reactive oxygen species (ROS)—induced genetic and epigenetic alterations in human carcinogenesis. *Mutat Res*. 2011;711(1–2):167–73.
122. Limoli CL, Giedzinski E, Morgan WF, Swartz SG, Jones GD, Hyun W. Persistent oxidative stress in chromosomally unstable cells. *Cancer Res*. 2003;63(12):3107–11.
123. Morgan WF, Day JP, Kaplan MI, McGhee EM, Limoli CL. Genomic instability induced by ionizing radiation. *Radiat Res*. 1996;146(3):247–58.
124. Abdollahi H, Shiri I, Atashzar M, Sarebani M, Moloudi K, Samadian H. Radiation protection and secondary cancer prevention using biological radioprotectors in radiotherapy. *Int J Cancer Ther Oncol*. 2015;3(3):335.
125. Kim K, Damoiseaux R, Norris AJ, Rivina L, Bradley K, Jung ME, et al. High throughput screening of small molecule libraries for modifiers of radiation responses. *Int J Radiat Biol*. 2011;87(8):839–45.
126. Xu G, Wu H, Zhang J, Li D, Wang Y, Wang Y, et al. Metformin ameliorates ionizing irradiation-induced long-term hematopoietic stem cell injury in mice. *Free Radic Biol Med*. 2015;87:15–25.
127. Zhang H, Zhai Z, Wang Y, Zhang J, Wu H, Wang Y, et al. Resveratrol ameliorates ionizing irradiation-induced long-term hematopoietic stem cell injury in mice. *Free Radic Biol Med*. 2013;54:40–50.
128. Li D, Tian Z, Tang W, Zhang J, Lu L, Sun Z, et al. The protective effects of 5-methoxytryptamine- α -lipoic acid on ionizing radiation-induced hematopoietic injury. *Int J Mol Sci*. 2016;17(6):935.
129. Van Buul JD, Fernandez-Borja M, Anthony EC, Hordijk PL. Expression and localization of NOX2 and NOX4 in primary human endothelial cells. *Antioxid Redox Signal*. 2005;7(3–4):308–17.
130. Han W, Wu L, Chen S, Bao L, Zhang L, Jiang E, et al. Constitutive nitric oxide acting as a possible intercellular signaling molecule in the initiation of radiation-induced DNA double strand breaks in non-irradiated bystander cells. *Oncogene*. 2007;26(16):2330–9.
131. Malaviya R, Gow AJ, Francis M, Abramova EV, Laskin JD, Laskin DL. Radiation-induced lung injury and inflammation in mice: role of inducible nitric oxide synthase and surfactant protein D. *Toxicol Sci*. 2014;144(1):27–38.
132. Nozaki Y, Hasegawa Y, Takeuchi A, Fan Z, Isoke K, Nakashima I, et al. Nitric oxide as an inflammatory mediator of radiation pneumonitis in rats. *Am J Physiol Lung Cell Mol Physiol*. 1997;272(4):L651–8.
133. Tsuji C, Shioya S, Hirota Y, Fukuyama N, Kurita D, Tanigaki T, et al. Increased production of nitrotyrosine in lung tissue of rats with radiation-induced acute lung injury. *Am J Physiol Lung Cell Mol Physiol*. 2000;278(4):L719–25.
134. Erbil Y, Dibekoglu C, Turkoglu U, Ademoglu E, Berber E, Kizir A, et al. Nitric oxide and radiation enteritis. *Eur J Surg*. 1998;164(11):863–8.
135. Abdollahi H, Atashzar M, Amini M. The potential use of biogas producing microorganisms in radiation protection. *J Med Hypotheses Idea*. 2015;9(2):67–71.
136. Rwigema J-CM, Beck B, Wang W, Doemling A, Epperly MW, Shields D, et al. Two strategies for the development of mitochondrion-targeted small molecule radiation damage mitigators. *Int J Radiat Oncol Biol Phys*. 2011;80(3):860–8.
137. Rajagopalan, Gupta K, Epperly MW, Franicola D, Zhang X, Wang H, et al. The mitochondria-targeted nitroxide JP4-039 augments potentially lethal irradiation damage repair. In *Vivo* (Athens, Greece). 2009;23(5):717–26.
138. Stoyanovsky DA, Huang Z, Jiang J, Belikova NA, Tyurin V, Epperly MW, et al. A manganese-porphyrin complex decomposes H₂O₂, inhibits apoptosis, and acts as a radiation mitigator in vivo. *ACS Med Chem Lett*. 2011;2(11):814–7.
139. Stoyanovsky D, Jiang J, Murphy M, Epperly M, Li S, Greenberger JS, et al. Mitigation of irradiation damage in vitro and in vivo by mitochondrial targeted glutathione peroxidase 4 mimic mito-ehselen. *Int J Radiat Oncol Biol Phys*. 2015;93(3):E541.
140. Stoyanovsky DA, Jiang J, Murphy MP, Epperly M, Zhang X, Li S, et al. Design and synthesis of a mitochondria-targeted mimic of glutathione peroxidase, MitoEhselen-2, as a radiation mitigator. *ACS Med Chem Lett*. 2014;5(12):1304–7.
141. Aoyama T, Paik YH, Watanabe S, Laleu B, Gaggini F, Fioraso-Cartier L, et al. Nicotinamide adenine dinucleotide phosphate oxidase in experimental liver fibrosis: GKT137831 as a novel potential therapeutic agent. *Hepatology*. 2012;56(6):2316–27.
142. Zykova TA, Zhu F, Zhai X, Ma WY, Ermakova SP, Lee KW, et al. Resveratrol directly targets COX-2 to inhibit carcinogenesis. *Mol Carcinog*. 2008;47(10):797–805.
143. Rwigema J-CM, Beck B, Wang W, Doemling A, Epperly MW, Shields D, et al. Two strategies for the development of mitochondrion-targeted small molecule radiation damage mitigators. *Int J Radiat Oncol Biol Phys*. 2011;80(3):860–8.
144. Fetisova EK, Antoschina MM, Cherepanynets VD, Izumov DS, Kireev II, Kireev RI, et al. Radioprotective effects of mitochondria-targeted antioxidant SkQR1. *Radiat Res*. 2015;183(1):64–71.