

Medication management – the missing link in dementia interventions

Running head – Medication management in dementia

Ian D Maidment (corresponding author), Senior Pharmacist, Kent and Medway NHS and Social Care Partnership Trust, Honorary Research Fellow, Centre for Health Services Studies, University of Kent.
Corresponding address – Pharmacy Department, Eastern and Area Coastal Office, St Martin's Hospital, Littlebourne Rd., Canterbury, Kent, CT1 1AZ, UK
Telephone / fax – 01227 812203. Email – ian.maidment@nhs.net

Dr. Chris Fox, Senior Clinical Lecturer / Hon Consultant Psychiatrist
School of Medicine, Health Policy and Practice, University of East Anglia

Dr. Malaz Boustani, Associate Professor of Medicine,
Associate Director of Indiana University Center for Aging Research,
Regenstrief Institute

Professor Cornelius Katona, Honorary Professor of Old Age Psychiatry,
University College London

Key words – Medication management, dementia, adverse events, adherence

Total word count - 1422

Dementia has an estimated global prevalence of 35 million individuals (Alzheimer's Disease International, 2009). In the UK 570,000 people live with dementia - a figure which will double over the next 30 years, reflecting global predictions (Alzheimer's Disease International, 2009; DoH, 2009a). People with dementia are commonly prescribed complex medication regimens, containing both physical and psychotropic medication (Schubert et al, 2006).

Medication related adverse events due to poor medication management are associated with significant morbidity and admission to secondary care (Pirmohamed et al, 2004). Medication management is the entire way in which medicines are selected, procured, delivered, prescribed, administered and reviewed to optimise the contribution that they make to producing informed and desired outcomes of patient care (Audit Commission, 2001). The importance of safe and effective medication management for people with dementia is recognised nationally in the UK, by both the National Dementia Strategy (DoH, 2010) and NICE guidance (National Collaborating Centre for Mental Health, 2007), and internationally (Ministry of Health Services, 2007; Agency for Healthcare Research and Quality, 2009; see table 1 for a summary of the key factors). However, medication management care pathways in dementia are often complex. In addition, there is an inherent tension between ensuring that the symptoms of dementia are treated effectively and limiting iatrogenic disease associated with medication.

Please insert table 1 here.

There has been relatively little research on the impact of dementia on the prevalence of adverse drug events, in terms of adverse drug reactions (ADRs) and/or medication errors (Maidment et al, 2008; Gomez-Pavon et al, 2010). Dementia has, however, been identified as the main disease state which increases the likelihood that other risk factors for ADRs, including inappropriate prescribing, old age, adherence issues, drug interactions, comorbidity and polypharmacy, are present (Gomez-Pavon et al, 2010).

Medication errors (any incident where there has been an error in the process of prescribing, dispensing, preparing, administering, monitoring or providing medicines advice, regardless of whether any harm occurred; NPSA, 2009) may also be more common in people with dementia due to the involvement of multiple health and social care professionals (Maidment et al, 2008). The direct impact of cognitive impairment is of course also important (Nirodi et al, 2002). Within the general population, patients identify and thereby avoid nearly a quarter of errors, but both dementia and medication impair cognition and decision-making facilities (Boustani et al, 2003; Warner et al, 2005; Schubert et al, 2006; Maidment et al, 2008). Patients with dementia may be less likely to question a prescription or a change in medicine, be less aware of potential adverse events, and of whether monitoring is required, and therefore more likely to fail to identify a potential error (Maidment et al, 2006; Barber et al, 2009).

Both adverse drug reactions and medication errors may lead to admission to resource-intensive care facilities. One of the largest studies of its type estimated that adverse drug events caused 6.5% of all admissions to secondary care facilities (Pirmohamed et al, 2004). The same study found that such events might account for as many as 6,000 deaths each year in England. In a study of 13,000 unplanned admissions, of which 714 (5.6%) were related to medication, impaired cognition was identified as the main predictor of preventable medication-related admission to hospital (Odds ratio, 11.9; 95% confidence interval, 3.9 to 36.3; Leendertse et al, 2008).

The impact of adverse drug events on admission to residential care facilities is less well studied. However, it is clear that people with dementia are frequently prescribed medicines with both anti-cholinergic and sedative activity, both of which impair cognition (Boustani et al, 2003; Mulsant et al, 2003). It has been estimated that 20 to 50% of people with dementia take at least one medication with anticholinergic activity (Boustani et al, 2003; Schubert et al, 2006). Admission to residential care facilities is linked with impaired cognition and central ADRs could be associated with admission to residential care facilities via their impact on cognition.

Informal carers of older people, such as family members, may conduct as many as 10 activities in relation to medication management, such as noticing and managing side-effects, deciding when to administer the medication and supplying appropriate information (Smith et al, 2003; Francis et al, 2002). They have a key role in ensuring safe medication management (Smith et al, 2003; Francis et al, 2002). People with dementia often rely upon informal carers to manage their medication particularly as the dementia progresses

(Cotrell et al, 2006; Arlt et al, 2008). This role places significant strain on such carers (Francis et al, 2002). The greater the number of medication management related activities, the worse the social functioning and the mental health of the carer (Francis et al, 2002). People with dementia may over-estimate their ability to manage their own medication safely and there is a lack of research on both the causes, and the impact, of switching from self-administration to carer-led administration (Cotrell et al, 2006; Arlt et al, 2008). One study, however, found that cognitive impairment was only associated with a difficulty in self-medicating safely in people prescribed complex medication regimens (Maddigan et al, 2003).

Adherence to, and concordance with, prescribed medication may be particularly problematic for people with dementia (Mackin et al, 2007). An inability to manage complex medication regimens is a predictor for admission to residential care facilities (Lieto et al, 2005). A successful self-medication programme that includes adequate support, teaching the older person about the medication and other measures to improve compliance, can reduce the risk of medication errors and help maintain independence (Maddigan et al, 2003). However, simply improving adherence may be counter-productive and an over-emphasis on adherence may increase the incidence of medication related iatrogenic disease (Holland et al, 2005; Holmes et al, 2006). In the HOMER trial - an intervention designed to improve medication management - a pharmacist-led, home-based medication review actually *increased* readmission rates in the intervention group compared to treatment as usual (234 vs. 178; OR 1.30, 95% confidence interval 1.07 to 1.58; P = 0.009) - possibly by over-encouraging improved adherence and thereby precipitating iatrogenic disease (Holland et al, 2005). Other trials of pharmacist-led medication review have produced variable results; though some studies have reduced the number of medicines prescribed, improving real life outcomes has been problematic (Zermansky et al, 2001; MEDMAN Team, 2007; Patterson et al, 2010; RESPECT Trial Team, 2010).

Research to date in this area has tended to focus on the use of anti-psychotics to treat behaviour that challenges. Such behaviour is amongst the main drivers for admission to residential care facilities (Holmes et al, 1998; Souetre et al, 1999; Gaugler et al, 2009; Ballard et al, 2010). Concern about the use of anti-psychotics in dementia has been highlighted for a long time (DoH, 2001). Recently the National Dementia Strategy targeted a two-thirds reduction in the use of anti-psychotics to treat behaviour that challenges (DoH, 2009b & 2010). While a significant proportion of anti-psychotic prescribing (in particular for long-term use) is inappropriate, this is an ambitious target and may prove unrealistic. The US 1987 Omnibus Budget Reconciliation Act (OBRA) restricted the use of anti-psychotics to certain approved indications, but only achieved a reduction in antipsychotic use of between 28 and 36% (Furniss, 2002; National Long Term Care Ombudsman Resource Center, 2011). Pharmacological methods will still be needed to treat acute behaviour that challenges; such a rigid and ambitious target may have adverse consequences such as the use of alternative unlicensed therapies for example, benzodiazepines, which are themselves associated with serious side-effects (Ancill et al, 1991; Guthrie et al, 2010). Indeed, one large study

involving 4850 nursing home residents, which examined the impact of the OBRA guidelines, noted a 48.6% increase in regular and a 27.5% increase in 'as required' prescriptions for anxiolytics such as benzodiazepines and buspirone (Borson et al, 1997).

We conclude that medication management in dementia is a broad concept that should encompass a complete review of medication including assessment of indication, dosage, interactions and continued need. Optimising the management of all medication in dementia offers significant potential to improve dementia care in several ways:

- (a) improving the quality of care for people with dementia and their carers;
- (b) improving key outcomes, in dementia by reducing iatrogenic disease;
- (c) enhancing the healthcare professional/patient relationship, by for example empowering people with dementia and their carers;
- (d) reducing inappropriate medication and the need for resource intensive interventions consequent to mismanagement;
- (e) enhancing non-medication management pathways for people with dementia.

Research is urgently needed to identify the most effective ways of delivering effective medication management in dementia.

Conflict of Interest

No conflicts of interest were identified.

References

Agency for Healthcare Research and Quality. 2009. AHRQ – Health Information Technology Portfolio's 2009 Annual Report. Available on www.ahrq.gov (accessed 6 January 2011).

Alzheimer's Disease International. 2009. World Alzheimer Report 2009. Available on: www.alz.co.uk/research/files/WorldAlzheimerReport.pdf (Accessed 1 December 2010)

Ancill RJ, Carlyle WW, Liang RA et al. 1991. Agitation in the Demented Elderly: A role for benzodiazepines? *Int Clin Psychopharmacol* **6**: 141-146.

Arlt S, Lindner R, Rosler A et al. 2008. Adherence to medication in patients with dementia. *Drugs Aging* **25**: 1033-1047.

Audit Commission 2001. A spoonful of sugar – medicines management in NHS hospitals. Holbrooks Printers, Portsmouth. Available on <http://www.audit-commission.gov.uk/SiteCollectionDocuments/AuditCommissionReports/NationalStudies/nrspoonfulsugar.pdf> (accessed 6 April 2011)

Ballard C, Creese B, Corbett A, Aarsland D. 2010. Atypical antipsychotics for the treatment of behavioral and psychological symptoms in dementia, with a

particular focus on longer term outcomes and mortality. *Expert Opin Drug Saf* **10**: 35-43

Barber ND, Alldred DP, Raynor DK et al. 2009. Care homes' use of medicines study: prevalence and potential harm of medication errors in care homes for older people. *Qual Saf Health Care* **18**: 341-346.

Borson S, Doane K. (1997). The Impact of OBRA-87 on Psychotropic Drug Prescribing in Skilled Nursing Facilities. *Psych Ser* **48**: 1289-296.

Boustani MA, Peterson B, Hanson L, Harris R, Krasnov C. 2003. *Screening for Dementia*. Systematic Evidence Review. Agency for Healthcare Research and Quality, Rockville, MD. Available on www.ahrq.gov/clinic/uspstfix.htm (accessed 6 January 2011).

Cotrell V, Wild K, Bader T. 2006. Medication management and adherence among cognitively impaired older adults. *J Gerontol Soc Work* **47**: 31-46.

Department of Health. 2001. National Service Framework for Older People. London, Stationary Office. Available on http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4071283.pdf (accessed 6 April 2011).

Department of Health. 2009a. Living well with dementia: a national dementia strategy. London, Stationary Office. Available on http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_094051.pdf (accessed 6 January 2011).

Department of Health. 2009b. The use of antipsychotic medication for people with dementia: Time for action Living well with dementia: A National Dementia Strategy. London, Stationary Office. Available on www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_108303 (accessed 7 April 2011)

Department of Health. 2010. Quality outcomes for people with dementia: building on the work of the National Dementia Strategy. London, Stationary Office. Available on http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_119827 (accessed 6 January 2011).

Francis SA, Smith F, Gray N et al. 2002. The roles of informal carers in the management of medication for older-care recipients. *Int J Pharm Pract* **3**: 1-10.

Furniss L. 2002. Use of medicines in nursing homes for older people. *Advances Psychiatr Treatment* **8**: 198-204

Gaugler JE, Yu F, Krichbaum K, Wyman JF. 2009. Predictors of nursing home admission for person with dementia. *Med Care* **47**: 191-198.

Gomez-Pavon J, Gonzalez Garcia P, Frances Roman I et al. 2010. Recommendations for the prevention of adverse drug reactions in older adults with dementia. *Rev Esp Geriatr Gerontol* **45**: 89-96.

Guthrie B, Clark A, McGowan G. 2010. The burden of psychotropic drug prescribing in people with dementia: a population database study. *Age Ageing* **39**: 637-642.

Holland R, Lenaghan E, Harvey I et al. 2005. Does home based medication review keep older people out of hospital? The HOMER RCT. *BMJ* **330**: 293
doi: 10.1136/bmj.38338.674583.AE

Holmes J, Pugner K, Phillips R et al. 1998. Managing Alzheimer's Disease: the cost of care per patient. *British Journal Health Care Management* **4**: 332-337.

Holmes HM, Hayley DC, Caleb Alexander GC et al. 2006. Reconsidering Medication Appropriateness for Patients Late in Life. *Arch Intern Med* **166**: 605-609.

Leendertse AJ, Egberts ACG, Stoker LJ et al. 2008. Frequency of and risk factors for preventable medication-related hospital admissions in the Netherlands. *Arch Intern Med* **168**: 1890-1896.

Lieto JM, Schmidt KS. 2005. Reduced ability to self-administer medication is associated with assisted living placement in a continuing care retirement community. *J Am Med Dir Assoc* **6**: 246-249.

Mackin RS, Arean PA. 2007. Cognitive and psychiatric predictors of medical treatment adherence among older adults in primary care clinics. *Int J Geriatr Psychiatry* **22**: 55-60.

Maddigan SL, Farris KB, Keating N et al. 2003. Predictors of older adults' capacity for medication management in a self-medication program: a retrospective chart review. *J Aging Health* **15**: 332-352.

Maidment ID, Paton C, Lelliott P. 2006. A Review of Medication Errors in Mental Health Care. *Qual Saf Health Care* **15**: 409-413.

Maidment ID, Haw C, Stubbs J et al. 2008. Medication Errors in Older People with mental health problems: a review. *Int J Geriatr Psychiatry* **23**: 564-73.

MEDMAN study team. 2007. The MEDMAN study: a randomised controlled trial of community pharmacy-led medicines management for patients with coronary heart disease. *Family Practice* **24**: 189-200.

Ministry of Health Services. 2007. British Columbia – Guidelines and Protocols Advisory Committee. Cognitive Impairment in the Elderly: recognition, diagnosis and management. Available on http://www.bcguidelines.ca/gpac/guideline_cognitive.html#recommendation_9 (accessed 6 January 2011)

Mulsant BH, Pollock BG, Kirshner M, Shen C, Hiroko D, Ganguli M. 2003. Serum Anticholinergic Activity in a community-based sample of older adults. *Arch Gen Psychiatry* **60**: 198-203.

National Long Term Care Ombudsman Resource Center. 2011. OBRA '87 SUMMARY. Available on <http://www.allhealth.org/briefingmaterials/OBRA87Summary-984.pdf> (accessed 7 January 2011)

National Collaborating Centre for Mental Health. 2007. The NICE-SCIE guideline on supporting people with dementia and their carers in health and social care. Available on <http://www.nice.org.uk/nicemedia/live/10998/30320/30320.pdf> (accessed 10 January 2011)

Nirodi P, Mitchell AJ. 2002. The quality of psychotropic drug prescribing in patients in psychiatric units for the elderly. *Aging Ment Health* **6**: 191-196.

NPSA. 2009. Safety in doses – improving the use of medicines in the NHS. Learning from National Reporting 2007. National Reporting and Learning System. Available on <http://www.nrls.npsa.nhs.uk/resources/?entryid45=61625> (accessed 8 April 2011)

Patterson SM, Hughes CM, Crealey G et al. 2010. An evaluation of an adopted US model of pharmaceutical care to improve psychoactive prescribing for nursing home residents in Northern Ireland (Fleetwood Northern Ireland Study). *Journal of the American Geriatric Society* **58**:44-53.

Pirmohamed M, James S, Meakin S, et al. 2004. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* **329**: 15-19.

RESPECT trial team. 2010 Effectiveness of shared pharmaceutical care for older patients: RESPECT trial findings. *British Journal of General Practice* **60**: e10-9.

Schubert CC, Boustani M, Callahan CM et al., 2006. Comorbidity profile of dementia patients in primary care: Are they sicker? *J Am Geriatr Soc* **54**: 104-109.

Smith F, Francis SA, Gray N, Denham M, Graffy J. (2003). A multi-centre survey among informal carers who manage medication for older care

recipients: problems experienced and development of services. *Health Soc Care Community* **11**: 138-45.

Souetre E, Thwaites RMA, Yearley HL. 1999. Economic impact of Alzheimer's disease in the United Kingdom: Cost of care and disease severity for non-institutionalized patients with Alzheimer's disease. *Br J Psychiatry* **174**: 51-55.

Warner B, Gerrett D. 2005. Identification of medication errors through community pharmacies. *Int J Pharm Pract* **13**: 1-6.

Zermansky A, Petty DR, Raynor DK et al. 2001. Randomised controlled trial of clinical medication review by a pharmacist of elderly patients receiving repeat prescriptions in general practice. *BMJ* **323**: 1-5.

Table 1 – Key Factors in Medication Management in Dementia

Ensuring effective evidence-based treatments are utilised.

Limiting adverse events associated with medication, such as adverse reactions and medication errors.

Ensuring appropriate adherence to medication.

Supporting the role of carers in safe and effective medication management.

Ensuring that medication can facilitate independent living.