

Disordered eating: the young male side

Sabine Ammann¹, BA

André Berchtold², PhD

Yara Barrense-Dias¹, MA

Christina Akre¹, PhD

Joan-Carles Surís¹, MD, MPH, PhD

¹ Research Group on Adolescent Health (GRSA), Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, 1010 Lausanne, Switzerland.

² Institute of Social Sciences & NCCR LIVES, University of Lausanne, 1015 Lausanne, Switzerland

Corresponding author: Prof. Joan-Carles Surís
Institute of Social and Preventive Medicine (IUMSP)
Route de la Corniche 10
1010 Lausanne / Switzerland
Phone : +4121 3147375 / Fax : +4121 3147373
Email : joan-carles.suris@chuv.ch

DISCLOSURE

Conflicts of Interest. The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

Funding

TheGenerationFree study was financed by the *Programme Intercantonal de Lutte contre la Dépendance au Jeu* (PILDJ) and the canton of Fribourg. The funding bodies had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

ABSTRACT

Disordered eating (DE) is known to predominate among girls, and historically studies have focused primarily on them. This cross-sectional study aimed to investigate the characteristics of adolescent and young adult males at risk of DE. Participants (N=2269; 15-24 years) in a study assessing their lifestyle were divided into two groups based on the SCOFF five-question screening tool: a SCOFF[+] group for participants at risk of disordered eating, and a SCOFF[-] group for everyone else. SCOFF[+] participants were more likely to be overweight and obese; have a lower socio-economic status and non-Swiss parents; experience weight issues; and report a chronic condition, poorer emotional well-being and a less positive attitude towards life. With 10% of males at risk of DE, clinicians should be aware of their specific characteristics, which should be considered red flags. DE is no longer a female-specific issue.

Keywords. male; disordered eating; adolescent; young adult; Switzerland; cross-sectional study

INTRODUCTION

There is a rising incidence of eating disorders (EDs) during adolescence. A study in six European countries analyzing lifetime ED prevalence showed that the majority of cases started between ten and twenty years of age ¹. ED cover a range of serious disturbances in eating habits or weight-control behavior ², which are complex issues linked to predisposing biological, psychological and social factors. Various hypotheses influencing the onset of EDs during adolescence have been suggested, such as: an increase of adipose tissue in girls, the importance of peer influence on body image, exposure to media-idealized images³, and difficulty in achieving independence from parents ⁴.

EDs are known to predominate among girls, and historically studies have focused primarily on them. However, research is increasing among males in the light of growing awareness that they, too, suffer from this condition. The usually accepted male vs. female ratio of 1:10 among adults shows fewer disparities among adolescents ⁵. In fact, the gender ratio seems to be an increasing function of age as it is 1:4 in childhood (5-13 years) ⁶. Moreover, EDs are even less sex-specific when partial eating disorders are included ⁷. Depending on age and the criteria used, ED prevalence among males varies greatly from 0.5% to 16.6% ^{8,9}.

Males find it difficult to identify themselves as having an ED and are reluctant to see their doctor because of the associated gender stereotype of an illness affecting only females ¹⁰⁻¹³. Since they are under-diagnosed, the prevalence among males is likely underestimated ¹⁴. There is a need to improve detection of ED in males and increase awareness among healthcare providers ¹⁰.

Recently, studies on ED exclusively focusing on males have increased and included larger sample sizes than previously ¹⁵. Indeed, different aspects of ED among males are being

analyzed, for example forecasting biological, social and individual factors associated with the onset¹⁶ and persistence of ED¹⁷, prevalence of each type of ED among males with the introduction of DSM-5¹⁸, and behavioral symptoms associated with ED^{19,20}. Nevertheless, there are few studies specifically concerned with adolescent and young adult (AYA) males.

One of these studies analyzed disordered eating among adolescent males in Switzerland²¹: they were more likely to be overweight or obese, to report depression, and to have a history of sexual or physical abuse and delinquent behavior. In this paper we will take the research of Dominé et al.²¹ a step further by assessing associations with socioeconomic status, self-perceived school performance, and the relationship with both parents.

The link between EDs and socioeconomic status is unclear in the literature with some research indicating that EDs are more common in higher income levels for males and in both the highest and lowest for females²². Others²³ report overall that EDs are more frequent in the lowest income quarter.

The association between ED and school performance is also unclear. Some authors²² report lower academic achievement in both genders, while others describe it mainly among girls^{24,25}. The literature on the influence of the parent-adolescent relationship on ED is scarce. Krug et al. found that low parental warmth was associated with EDs in girls but not in boys, while Berge et al.²⁶ concluded that higher family functioning and parental connection were protective factors against ED.

Disordered eating and EDs result from a complex interaction of biological processes, sociocultural factors, and psychological and individual traits like low self-esteem, perfectionism and negative affect^{27,28}. Other evidence suggests that the life context of AYAs, be it the family or scholastic environment, might act as a potential correlate of disordered eating^{29,30}.

The timing of pubertal transition has been included as a variable in this study because it is known to have an impact on social development and psychological adjustment among adolescents³¹. There is evidence that pubertal timing affects females and males in different ways³². Early-maturing females are more likely to be depressed and tend to be less popular with peers³³. In contrast, early-maturing males are viewed as more attractive and demonstrate higher level of body satisfaction³⁴.

Most studies analyze boys suffering from ED through the prism of girls with EDs, which implies that both genders present the same symptoms. However, males seem to present different criteria regarding EDs. Body concern among adolescent males tends to focus more on muscularity than on a drive for thinness, which is characteristic among girls^{35,36}. Males before obvious signs of ED tend to be overweight, which clinicians can overlook as a sign of being at risk^{36,37}. In fact, males can be motivated to lose weight and increase muscle mass to achieve their ideal body: large chest and shoulders and a slim waist^{14,38}. Although some studies show that athletes of both genders are at risk of ED, male athletes are more likely to exhibit disturbed eating at sub-threshold diagnostic levels³⁹.

Early detection of disturbed eating is important. Treatment tends to be more successful when initiated early and while preventing harmful consequences in the long run⁴⁰. For that reason, being aware of the distinguishing characteristics of males who are at risk of developing disordered eating could help clinicians identify them and offer a targeted prevention. The aim of this study is to describe the characteristics of males at risk of disordered eating while controlling for potential confounding factors.

METHODS

Data were drawn from the baseline wave of the GenerationFRee study which aims to assess the lifestyles of AYAs in the canton of Fribourg, Switzerland. Data used for this study were collected during the 2014-2015 school year. AYAs in the canton's eleven post-mandatory schools (five high schools and six professional schools) completed an anonymous Web-based self-administrated questionnaire. The mandatory school age in Switzerland is 15; afterwards, about one third of adolescents follow high school and two thirds vocational training. The latter are enrolled by companies to train for their future profession and attend classes at vocational school 1-2 days a week. Questionnaires were filled in online in the schools' computer facility.

Out of 5 834 AYAs, 200 were unwilling to participate, 211 were not in the target age group, and 244 failed to complete the questionnaire reliably and were thus excluded. The final sample consisted of 5 179 AYAs of which only male respondents were included in the analysis (n= 2 269).

Dependent variable

We divided the surveyed males into two groups using the SCOFF questionnaire, a validated screening tool for identifying subjects at-risk of disordered eating ⁴¹. The five items of the questionnaire (*Do you make yourself sick because you feel uncomfortably full? / Do you worry you have lost control over how much you eat? / Have you recently lost more than 14 pounds in a three-month period? / Do you believe yourself to be fat when others say you are too thin? / Would you say that food dominates your life?*) are each answered as "yes" or "no". The cut-off point for the SCOFF is set at two or more positive answers, with a sensitivity of 85% and a specificity of 90%. Regarding eating disorders, the SCOFF is excellent at detecting all true cases of anorexia nervosa and bulimia nervosa and most cases of eating disorders not otherwise

specified (EDNOS)⁴¹. The SCOFF questionnaire has been translated into French and validated among adolescents⁴².

We included subjects who answered positively to at least two of the five questions in the group with a positive SCOFF (SCOFF[+]; n=230, 10.1%). Participants with one or no positive answers formed the group with a negative SCOFF (SCOFF[-]; n=2 039, 89.9%).

Independent variables

Personal variables included age, height, weight, pubertal timing, AYA's birthplace (Switzerland/abroad), and residence (urban/rural).

BMI was calculated from self-reported height and weight and we used the cut-offs for underweight, overweight and obesity defined by Cole et al.^{43,44}. Self-assessment of pubertal timing was determined by the question "If you think about the age you began puberty, compared with your same age peers, would you say that you were..." trichotomized into 'early pubertal timing', 'average pubertal timing' and 'late pubertal timing'. This method has been shown to be a valid approximation to pubertal timing with a 94% correlation with physician evaluations⁴⁵.

Family variables included family structure (parents living together or not), parents' birthplace (at least one born in Switzerland/both abroad), socioeconomic status (SES), and mother-AYA and father-AYA relationship, which is a good indicator of adolescent functioning⁴⁶.

Self-assessment of SES was determined by the question used in the ESPAD project⁴⁷: "Compared to the financial situation of other families in Switzerland, would you say that your family is..." with seven possible answers ranging from 'very well below average' to 'very well above average' and trichotomized into 'above average', 'average' and 'below average'. Father and mother relationship with the AYA were rated on a scale from 1 [very poor] to 10 [excellent].

School variables included academic track (student/apprentice) and self-reported school performance (above average, average or below average).

We controlled for variables likely to have a relationship with disordered eating such as the average weekly number of days of physical activity lasting a minimum of 60 minutes, having faced weight issues the past year, having a chronic condition (disability or chronic disease)⁴⁸, emotional well-being⁴⁹, self-reported health (good/poor)⁵⁰ and positive attitude towards life.

To measure emotional well-being we used the WHO-Five Well-Being Index (WHO-5), whose validity in adolescents has been proved⁵¹. The WHO-5 index includes five items and each one is rated on a six-point Likert scale ranging from 0 (at no time) to 5 (all of the time)⁵². Scores are added and a result below 13/25 indicates poor well-being. Cronbach's alpha in the present study was .81. We measured their vision of the future using the Positive attitude to life scale, which includes six items ranging from 1 (absolutely untrue) to 6 (absolutely true), a higher score meaning that they had a more positive attitude to life⁵³. Cronbach's alpha in the present study was .87.

The study protocol was approved by the Ethics Committee of the canton of Vaud.

Statistical analyses were performed with STATA 13.0 (StataCorp, College Station, Texas). In a first step, we used the Chi-square test and Student's t-test to compare categorical and continuous variables between the two groups. Results are given as point prevalence and means. In a second step, all statistically significant variables ($p < 0.05$) at the bivariate level were included in a backward logistic regression analysis. Because the SCOFF and the variable "weight issues during the past year" might interact with each other, we reran the analysis without the variable "weight issue during the past year". However, the results did not change significantly (data not shown). To avoid setting aside potentially important variables that could interact with

others, we also tried to include in the regression model variables significant at the 10% level at the bivariate level, but again without there being any change. Finally, the specification of the final multivariate model was checked using the linktest, but no problem was found. Results are given as adjusted odds ratio (aOR) with 95% confidence intervals (95%CI).

RESULTS

At the bivariate level AYAs at risk of disordered eating were significantly more likely to be overweight or obese, to report early pubertal timing, to be born abroad and to have a low socio-economic status. Regarding family-related variables, SCOFF[+] males were significantly more likely to have non-Swiss-born parents not living together, and to have a poorer relationship both with their father and their mother. Considering health-related variables, SCOFF[+] males were significantly more likely to report poor health, to have faced weight issues during the past year, and to have a chronic condition, poor emotional wellbeing and a less positive attitude towards life (Table 1).

The results of the backward logistic regression are reported in Table 2. After controlling for confounding factors and comparing to the control group, SCOFF[+] males were more likely to be overweight (aOR: 2.58 [1.78:3.75]) or obese (aOR: 3.12 [1.70:5.71]). For their social background, SCOFF[+] males were more likely to have a lower socio-economic status (aOR: 2.20 [1.39:3.48]) and non-Swiss parents (aOR: 1.66 [1.19:2.31]). They were also more likely to have faced weight issues during the past year (aOR: 5.79 [2.98:11.27]), and to report a chronic condition (aOR: 1.63 [1.11:2.40]), poorer emotional well-being (aOR: 1.79 [1.19:2.69]) and a less positive attitude towards life (aOR: 0.95 [0.93:0.98]).

DISCUSSION

Overall 10% of our sample was at risk of disordered eating. They were more likely to be overweight or obese, to be of lower socio-economic status, to have both parents of foreign origin, to suffer from a chronic condition and to have a less positive attitude towards life.

Dominé et al.²¹ had also found an association between unhealthy eating behaviors and overweight or obesity among males. The reverse is also true as overweight youth have an increased risk for disordered eating behaviors. Indeed, dieting, for example, is a common risk factor for overweight and ED, as it might trigger a cycle of restriction and overeating, thus possibly leading to binge eating and weight gain⁵⁴. Increased body mass contributes to body dissatisfaction for not fitting with the leanness ideal, which in turn is a risk factor for eating pathology⁵⁵⁻⁵⁷.

Regarding associations between SES and disordered eating, the literature is inconclusive, especially for males⁵⁸. Nevertheless, the link between low socio-economic status and overweight/obesity has been well established in numerous studies even though the underlying mechanisms are still not well understood⁵⁹. We can hypothesize that eating healthily is more financially demanding but also needs nutrition knowledge and time for cooking, which can be lacking when both parents work⁶⁰. Overweight/obesity and their psychosocial consequences such as peer teasing and repetitive dieting are known risk factors for disordered eating^{3,19,54,61}. Moreover, the dysfunctional family environment often observed in low-income households might lead to chaotic behaviors, among them disordered eating⁶². Further research is required to assess the link between socio-economic status and disordered eating, especially among males.

Our results also demonstrate that SCOFF[+] males were more likely to self-report early puberty. There is a controversy in the literature regarding whether early or late puberty among males is associated with disordered eating, highlighting the need for further research^{4,63}.

Interestingly, Dominé et al.²¹ also found an association between early puberty and disordered eating behaviors or concerns. Psychological mechanisms could account in this study for the relationship between early puberty and disordered eating. Differing development among males might foster feelings of alienation, especially during a development phase where comparison to peers is a major concern. More important still, experiencing puberty before peers might lead to greater psychological disturbance: early maturing males are less likely to be prepared to face bodily changes and might focus more on their body. While females move further away from the thinness ideal with pubertal development, males tend to move closer to the male ideal of acquiring muscular mass and developing wider shoulders. McCabe showed in her study that females experienced more body dissatisfaction with an increase in body fat linked to puberty and tended to adopt strategies to reduce weight⁶⁴. For adolescent males in the study however, pubertal changes led to an increased use of strategies to enhance muscle tone, which in turn was associated with greater anxiety⁶⁴. Therefore it seems that a different mechanism underlies the association of pubertal timing and puberty onset with disordered eating pathology between females and males.

The risk of disordered eating was significantly more predominant among males with foreign-born parents. Regarding ethnicity, the literature highlights an emergence of disordered eating in non-Western cultures⁶⁵. Furthermore, some authors demonstrated an association between culture change and disordered eating that depends on the studied group⁶⁶. A possible explanation is that children of foreign-born parents grow up between two different cultures and thus can feel torn between original and local values. Disordered eating can be a way to cope with the identity issues related to the culture shock. However, this cultural turmoil may express itself in a different manner. Some cultures look at overweight or obesity positively as a sign of wealth,

preferring a larger body size. A possible explanation is that, when confronted with local body standards, AYAs of foreign-born parents might experience conflicting expectations, possibly leading to eating disorders.

We also found that SCOFF[+] males were more likely to have a chronic condition. Chronic conditions that require dietary management like diabetes or cystic fibrosis might increase preoccupation with food. Moreover, physical stigma due to illness or treatment can foster body dissatisfaction, a risk factor of disordered eating⁴⁸. AYAs suffering from a chronic condition are striving to be like their healthy peers and report lower self-esteem⁶⁷. Disordered eating may be a functional response to confronting an identity crisis.

Research focusing on the relationship between disordered eating and attitude towards life is rare. In this study, SCOFF[+] males reported a less positive attitude towards life. The thoughts of disordered eating subjects revolve mostly around food and weight, thus not leaving much room for other concerns. Disordered eating might thus serve to mask existential concerns related to purpose in life and self-identity. Furthermore, depression is a common psychiatric comorbidity in patients with an eating pathology that is associated with feelings of hopelessness and worthlessness which darken one's future^{68,69}.

As expected, our results showed that SCOFF[+] males faced weight issues during the past year. This concurs with the SCOFF evaluation, thus underscoring its relevance. Additionally, in line with earlier studies, we found that SCOFF[+] males had a reduced emotional well-being⁷⁰.

The main strengths of this study are that it is based on a large representative sample of AYAs in post-mandatory education and that it focuses exclusively on males. However, there are also some limitations. First, because of the study's cross-sectional design no causal relationship

can be drawn. Second, as the survey results are based on self-reported questionnaire responses, bias cannot be excluded. Regarding weight status in particular, overweight or obese youth tend to underestimate their actual weight ⁷¹, which could lead to our underestimating results (although use of anonymous data collection should have minimized this problem). Third, an important limitation of the body mass index (BMI) should be considered. Indeed, it does not distinguish between body fat, muscle mass or bone density. SCOFF[+] males considered as overweight or obese in terms of their BMI could in fact be muscular. That possibility would correspond to the muscularity concern common to males. Fourth, the GenerationFree study did not include more specific male assessment such as a drive for muscularity that could have improved the present analysis ⁷². However, the initial aim of the GenerationFree study was to focus on AYAs' lifestyle independently of gender. Fifth, the study did not assess disordered eating oriented toward gaining muscularity rather than losing body fat (termed muscularity-oriented disordered eating), although, at present, there is no validated and published scale that assesses this construct⁷³. Sixth, the SCOFF questionnaire is only a screening tool for disordered eating and cannot replace a clinician's diagnosis. However, the SCOFF questionnaire has a high sensitivity with few false positives ⁴¹.

Given the public health significance of disordered eating, further research specifically focused on males is warranted to better define their characteristics. Furthermore, the success of enhanced eating disorder prevention depends on longitudinal data to understand the risk factors of disordered eating among males.

CONCLUSIONS

Based on this study's results, 10% of young males are at risk of disordered eating. This finding provides further evidence suggesting that disordered eating is no longer a specific female

issue (even though females in our study report a much higher rate, 31% [data not shown]) and tends to affect males too. Besides, although males show some similarities with females at risk of disordered eating, they also present important differences. Therefore, looking at males through a female prism (academic overachievers, higher socioeconomic status) would be much too restrictive. From a clinical standpoint, it is important for practitioners to consider the specificities of males at risk of disordered eating, especially overweight and obesity. Healthcare providers should bear in mind these characteristics when consulting with AYA males and should be encouraged to discuss disordered eating with them.

DISCLOSURE

Conflicts of Interest. The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

Funding

The GenerationFree study was financed by the *Programme Intercantonal de Lutte contre la Dépendance au Jeu* (PILDJ) and the canton of Fribourg. The funding body had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

AUTHOR CONTRIBUTIONS

Sabine Ammann analyzed and interpreted the data, drafted and revised the manuscript, and approved the final version of the manuscript as submitted.

André Berchtold conceptualized and designed the study, obtained funding, analyzed and interpreted the data, critically revised the manuscript, and approved the final version of the manuscript as submitted.

Yara Barrense-Dias conceptualized and designed the study, coordinated data collection, critically revised the manuscript, and approved the final version of the manuscript as submitted.

Christina Akre conceptualized and designed the study, coordinated data collection, critically revised the manuscript, and approved the final version of the manuscript as submitted.

Joan-Carles Surís conceptualized and designed the study, obtained funding, acquired, analyzed and interpreted the data, critically revised the manuscript, and approved the final version of the manuscript as submitted.

REFERENCES

1. Preti A, Girolamo Gd, Vilagut G, et al. The epidemiology of eating disorders in six European countries: Results of the ESEMeD-WMH project. *Journal of Psychiatric Research*. 2009;43(14):1125-1132.
2. Fairburn CG, Harrison PJ. Eating disorders. *The Lancet*. 2003;361:407-416.
3. Dakanalis A, Carra G, Calogero R, et al. The developmental effects of media-ideal internalization and self-objectification processes on adolescents' negative body-feelings, dietary restraint, and binge eating. *Eur Child Adolesc Psychiatry*. 2015;24(8):997-1010.
4. Klump KL. Puberty as a critical risk period for eating disorders: a review of human and animal studies. *Horm Behav*. 2013;64(2):399-410.
5. Kjelsas E, Bjornstrom C, Gotestam KG. Prevalence of eating disorders in female and male adolescents (14-15 years). *Eat Behav*. 2004;5(1):13-25.
6. Madden S, Morris A, Zurynski YA, Kohn M, Elliot EJ. Burden of eating disorders in 5-13-year-old children in Australia. *The Medical Journal of Australia*. 2009;190(8):410-414.
7. Woodside DB, Garfinkel PE, Lin E, et al. Comparisons of men with full or partial eating disorders, men without eating disorders, and women with eating disorders in the community. *The American Journal of Psychiatry*. 2001;158(4):570-574.
8. Smink FR, van Hoeken D, Oldehinkel AJ, Hoek HW. Prevalence and severity of DSM-5 eating disorders in a community cohort of adolescents. *Int J Eat Disord*. 2014;47(6):610-619.
9. Valls M, Callahan S, Rousseau A, Chabrol H. Troubles du comportement alimentaire et symptomatologie dépressive : étude épidémiologique chez les jeunes hommes. *L'Encéphale*. 2014;40(3):223-230.
10. Raisanen U, Hunt K. The role of gendered constructions of eating disorders in delayed help-seeking in men: a qualitative interview study. *BMJ Open*. 2014;4(4):e004342-e004342.
11. Griffiths S, Mond JM, Li Z, et al. Self-stigma of seeking treatment and being male predict an increased likelihood of having an undiagnosed eating disorder. *Int J Eat Disord*. 2015;48(6):775-778.
12. Griffiths S, Mond JM, Murray SB, Touyz S. The prevalence and adverse associations of stigmatization in people with eating disorders. *Int J Eat Disord*. 2015;48(6):767-774.
13. Mitchison D, Mond J. Epidemiology of eating disorders, eating disordered behaviour, and body image disturbance in males: a narrative review. *Journal of eating disorders*. 2015;3(1):20.
14. Strother E, Lemberg R, Stanford SC, Turberville D. Eating Disorders in Men: Underdiagnosed, Undertreated, and Misunderstood. *Eating Disorders*. 2012;20(5):346-355.
15. Coelho JS, Kumar A, Kilvert M, Kunkel L, Lam P-Y. Male Youth With Eating Disorders: Clinical and Medical Characteristics of a Sample of Inpatients. *Eating Disorders*. 2015:1-7.
16. Dakanalis A, Favagrossa L, Clerici M, et al. Body Dissatisfaction and Eating Disorder Symptomatology: A Latent Structural Equation Modeling Analysis of Moderating Variables in 18-to-28-Year-Old Males. *The Journal of Psychology*. 2015;149(1):85-112.

17. Dakanalis A, Clerici M, Caslini M, et al. Predictors of initiation and persistence of recurrent binge eating and inappropriate weight compensatory behaviors in college men. *Int J Eat Disord*. 2016;49(6):581-590.
18. Allen KL, Byrne SM, Oddy WH, Crosby RD. DSM-IV-TR and DSM-5 eating disorders in adolescents: prevalence, stability, and psychosocial correlates in a population-based sample of male and female adolescents. *Journal of abnormal psychology*. 2013;122(3):720-732.
19. Culbert KM, Racine SE, Klump KL. Research Review: What we have learned about the causes of eating disorders - a synthesis of sociocultural, psychological, and biological research. *Journal of child psychology and psychiatry, and allied disciplines*. 2015;56(11):1141-1164.
20. Wyssen A, Bryjova J, Meyer AH, Munsch S. A model of disturbed eating behavior in men: The role of body dissatisfaction, emotion dysregulation and cognitive distortions. *Psychiatry research*. 2016;246:9-15.
21. Dominé F, Berchtold A, Akre C, Michaud P-A, Suris J-C. Disordered eating behaviors: what about boys? *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*. 2009;44(2):111-117.
22. Lee HJ, Park S, Kim CI, et al. The association between disturbed eating behavior and socioeconomic status: the Online Korean Adolescent Panel Survey (OnKAPS). *PLoS One*. 2013;8(3):e57880.
23. Ahren JC, Chiesa F, Koupil I, Magnusson C, Dalman C, Goodman A. We are family--parents, siblings, and eating disorders in a prospective total-population study of 250,000 Swedish males and females. *Int J Eat Disord*. 2013;46(7):693-700.
24. Veses AM, Gomez-Martinez S, de Heredia FP, et al. Cognition and the risk of eating disorders in Spanish adolescents: the AVENA and AFINOS studies. *Eur J Pediatr*. 2015;174(2):229-236.
25. Tabler J, Utz RL. The influence of adolescent eating disorders or disordered eating behaviors on socioeconomic achievement in early adulthood. *Int J Eat Disord*. 2015;48(6):622-632.
26. Berge JM, Wall M, Larson N, Eisenberg ME, Loth KA, Neumark-Sztainer D. The unique and additive associations of family functioning and parenting practices with disordered eating behaviors in diverse adolescents. *J Behav Med*. 2014;37(2):205-217.
27. Jacobi C, Hayward C, de Zwaan M, Kraemer HC, Agras WS. Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychological bulletin*. 2004;130(1):19-65.
28. Ricciardelli LA, McCabe MP. A biopsychosocial model of disordered eating and the pursuit of muscularity in adolescent boys. *Psychological bulletin*. 2004;130(2):179-205.
29. Croll J, Neumark-Sztainer D, Story M, Ireland M. Prevalence and risk and protective factors related to disordered eating behaviors among adolescents: relationship to gender and ethnicity. *J Adolesc Health*. 2002;31(2):166-175.
30. Haines J, Rifas-Shiman SL, Horton NJ, et al. Family functioning and quality of parent-adolescent relationship: cross-sectional associations with adolescent weight-related behaviors and weight status. *The international journal of behavioral nutrition and physical activity*. 2016;13:68.
31. Alsaker FD. Is puberty a critical period for socialization? *Journal of adolescence*. 1995;18(4):427-444.

32. Graber JA, Seeley Jr Fau - Brooks-Gunn J, Brooks-Gunn J Fau - Lewinsohn PM, Lewinsohn PM. Is pubertal timing associated with psychopathology in young adulthood. (0890-8567 (Print)).
33. Stice E, Presnell K, Bearman SK. Relation of early menarche to depression, eating disorders, substance abuse, and comorbid psychopathology among adolescent girls. *Developmental psychology*. 2001;37(5):608-619.
34. McCabe MP, Ricciardelli LA. A longitudinal study of pubertal timing and extreme body change behaviors among adolescent boys and girls. *Adolescence*. 2004;39(153):145-166.
35. Neumark-Sztainer D, Eisenberg ME. Body image concerns, muscle-enhancing behaviors, and eating disorders in males. *JAMA*. 2014;312(20):2156-2157.
36. Campbell K, Peebles R. Eating disorders in children and adolescents: state of the art review. *Pediatrics*. 2014;134(3):582-592.
37. Raevuori A, Keski-Rahkonen A, Hoek HW. A review of eating disorders in males. *Current Opinion in Psychiatry*. 2014;27(6):426-430.
38. McCabe MP, Ricciardelli LA. Body image and body change techniques among young adolescent boys. *European Eating Disorders Review*. 2001;9(5):335-347.
39. Chatterton JM, Petrie TA. Prevalence of disordered eating and pathogenic weight control behaviors among male collegiate athletes. *Eat Disord*. 2013;21(4):328-341.
40. Herpertz-Dahlmann B, Dempfle A, Konrad K, Klasen F, Ravens-Sieberer U, The Bsg. Eating disorder symptoms do not just disappear: the implications of adolescent eating-disordered behaviour for body weight and mental health in young adulthood. *European Child & Adolescent Psychiatry*. 2014.
41. Hill LS, Reid F, Morgan JF, Lacey JH. SCOFF, the development of an eating disorder screening questionnaire. *Int J Eat Disord*. 2010;43(4):344-351.
42. Garcia FD, Grigioni S, Chelali S, Meyrignac G, Thibaut F, Dechelotte P. Validation of the French version of SCOFF questionnaire for screening of eating disorders among adults. *World J Biol Psychiatry*. 2010;11(7):888-893.
43. Cole TJ, Flegal KM, Nicholls D, Jackson AA. Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ*. 2007;335(7612):194-194.
44. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ (Clinical research ed)*. 2000;320(7244):1240-1243.
45. Mohnke S, Warschburger P. [Body dissatisfaction among female and male adolescents: comparing prevalence, predictors, and consequences between the sexes]. *Praxis Der Kinderpsychologie Und Kinderpsychiatrie*. 2011;60(4):285-303.
46. Moretti MM, Peled M. Adolescent-parent attachment: Bonds that support healthy development. *Paediatr Child Health*. 2004;9(8):551-555.
47. Hibell B, Guttormsson U, Ahlström S, et al. *The 2007 ESPAD Report. Substance Use Among Students in 35 European Countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN);2009.
48. Piquart M. Body image of children and adolescents with chronic illness: a meta-analytic comparison with healthy peers. *Body image*. 2013;10(2):141-148.
49. Fursland A, Watson HJ. Eating disorders: A hidden phenomenon in outpatient mental health?: Eating Disorder Screening. *International Journal of Eating Disorders*. 2014;47(4):422-425.

50. Norris ML, Apsimon M, Harrison M, et al. An examination of medical and psychological morbidity in adolescent males with eating disorders. *Eating Disorders*. 2012;20(5):405-415.
51. Allgaier A-K, Pietsch K, Frühe B, Prast E, Sigl-Glöckner J, Schulte-Körne G. Depression in pediatric care: is the WHO-Five Well-Being Index a valid screening instrument for children and adolescents? *General Hospital Psychiatry*. 2012;34(3):234-241.
52. Henkel V, Mergl R, Kohnen R, Maier W, Möller H-J, Hegerl U. Identifying depression in primary care: a comparison of different methods in a prospective cohort study. *BMJ*. 2003;326(7382):200-201.
53. Grob A, Lüthi R, Kaiser FG, Flammer A, Mackinnon A, Wearing AJ. Berner Fragebogen zum Wohlbefinden Jugendlicher (BFW). *Diagnostica*. 1991;37(1):66-75.
54. Neumark-Sztainer DR, Wall MM, Haines JI, Story MT, Sherwood NE, van den Berg PA. Shared risk and protective factors for overweight and disordered eating in adolescents. *Am J Prev Med*. 2007;33(5):359-369.
55. Stice E. Risk and maintenance factors for eating pathology: a meta-analytic review. *Psychol Bull*. 2002;128(5):825-848.
56. Riva G, Gaggioli A, Dakanalis A. From body dissatisfaction to obesity: how virtual reality may improve obesity prevention and treatment in adolescents. *Studies in health technology and informatics*. 2013;184:356-362.
57. Dakanalis A, Carra G, Clerici M, Riva G. Efforts to make clearer the relationship between body dissatisfaction and binge eating. *Eating and weight disorders : EWD*. 2015;20(1):145-146.
58. Mitchison D, Hay P, Slewa-Younan S, Mond J. The changing demographic profile of eating disorder behaviors in the community. *BMC Public Health*. 2014;14(1):943.
59. Ball K, Crawford D. Socio-economic factors in obesity: a case of slim chance in a fat world? *Asia Pac J Clin Nutr*. 2006;15 Suppl:15-20.
60. Drewnowski A, Darmon N. The economics of obesity: dietary energy density and energy cost. *Am J Clin Nutr*. 2005;82(1 Suppl):265s-273s.
61. Dakanalis A, Pla-Sanjuanelo J, Caslini M, et al. Predicting onset and maintenance of men's eating disorders. *International Journal of Clinical and Health Psychology*. 2016;16(3):247-255.
62. Schreier HM, Chen E. Socioeconomic status and the health of youth: a multilevel, multidomain approach to conceptualizing pathways. *Psychol Bull*. 2013;139(3):606-654.
63. Zehr JL, Culbert KM, Sisk CL, Klump KL. An association of early puberty with disordered eating and anxiety in a population of undergraduate women and men. *Horm Behav*. 2007;52(4):427-435.
64. McCabe MP, Ricciardelli LA, Banfield S. Body image, strategies to change muscles and weight, and puberty: do they impact on positive and negative affect among adolescent boys and girls? *Eat Behav*. 2001;2(2):129-149.
65. Pike KM, Hoek HW, Dunne PE. Cultural trends and eating disorders. *Current Opinion in Psychiatry*. 2014;27(6):436-442.
66. Doris E, Shekriladze I, Javakhishvili N, Jones R, Treasure J, Tchanturia K. Is cultural change associated with eating disorders? A systematic review of the literature. *Eat Weight Disord*. 2015;20(2):149-160.
67. Pinquart M. Self-esteem of children and adolescents with chronic illness: a meta-analysis. *Child Care Health Dev*. 2013;39(2):153-161.

68. Jenkins PE, Hoste RR, Doyle AC, et al. Health-related quality of life among adolescents with eating disorders. *Journal of Psychosomatic Research*. 2014;76(1):1-5.
69. Ferreiro F, Wichstrom L, Seoane G, Senra C. Reciprocal associations between depressive symptoms and disordered eating among adolescent girls and boys: a multiwave, prospective study. *J Abnorm Child Psychol*. 2014;42(5):803-812.
70. Engel SG, Adair CE, Las Hayas C, Abraham S. Health-related quality of life and eating disorders: a review and update. *Int J Eat Disord*. 2009;42(2):179-187.
71. Sherry B, Jefferds ME, Grummer-Strawn LM. Accuracy of adolescent self-report of height and weight in assessing overweight status: a literature review. *Arch Pediatr Adolesc Med*. 2007;161(12):1154-1161.
72. Dakanalis A, Zanetti AM, Riva G, et al. Male body dissatisfaction and eating disorder symptomatology: moderating variables among men. *J Health Psychol*. 2015;20(1):80-90.
73. Griffiths S, Murray SB, Touyz S. Disordered eating and the muscular ideal. *Journal of eating disorders*. 2013;1:15.