

Langmuir, 2017, 33 (38), pp 10106–10106

DOI: 10.1021/acs.langmuir.7b02999

Accepted for publication

26/ 8/ 17

**Correction to “Lipid spontaneous curvatures estimated from
temperature-dependent changes in inverse hexagonal phase lattice
parameters: the effects of metal cations”**

Marcus K. Dymond^{1*}, Richard J. Gillams², Duncan J. Parker², Jamie Burrell², Ana Labrador³, Tommy Nylander⁴, George S. Attard^{2*}

¹ Division of Chemistry, School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, BN2 4GJ, UK

² Chemistry, Faculty of Natural & Environmental Sciences, University of Southampton, Southampton SO17 1BJ, UK

³ MAX IV Laboratory, Lund University, PO Box 118, SE-221 00, Lund, Sweden

⁴ Physical Chemistry, Lund University, PO Box 124, SE-221 00, Lund, Sweden

(*) Authors for correspondence: M.Dymond@brighton.ac.uk; gza@soton.ac.uk

Correction:

It has come to our attention that there are two typos in the original manuscript concerning equations 7 and 8. These should read as;

We define the ‘curvature power’ of lipid j (χ_j) as:

$$\chi_j = -(L_{p,PE(37)} - L_{p,j(37)}) / x_j \quad (7).$$

where $L_{p,j(37)}$ is the lattice parameter, at 37 °C, of lipid j , $L_{p,PE(37)}$ is the lattice parameter of DOPE and x_j is the mole fraction of the lipid j .

The universal curve can be fitted with the linear equation:

$$c_0 = 0.0731 \chi - 0.3694 . \quad (8)$$