

New Triterpene from *Conyza aegyptiaca* L.*

Mohamed Abbas Metwally

Department of Chemistry, Faculty of Science, University of Mansoura, Mansoura, Egypt

Z. Naturforsch. **44b**, 1584 (1989); received July 26, 1989

Triterpene, *Conyza aegyptiaca* L.

A reinvestigation of *C. aegyptiaca* L. gave in addition to phytol a new triterpene. Its structure was determined by IR, ¹H NMR and mass spectral data.

Introduction


From the large genus *Conyza* (tribe Astereae, family Compositae) about 10% of the approximately 2000 species in this tribe have been subjected to chemical scrutiny. However, work has been confined largely to root constituents. Among these, polyacetlenes, polyenes and related substances and, in certain groups, coumarins are very characteristic. Less widely distributed, perhaps because they are less intensively searched for, are diterpenoids, so far largely of the labdane and clerodane type.

Our previous investigation [1] of *C. aegyptiaca* L. resulted in isolation of α -curcumene, germacrene D, β -farnesene, caryophyllenepoxide, lachnophyllum ester, matricaria ester, stigmasterol, squalene and some acetylenic compounds from the plant material collected in August. In continuation of our study on the Egyptian Compositae [2] the present work deals with investigation of the same plant collected in April.

A reinvestigation of *C. aegyptiaca* L. afforded a phytol and a new triterpene. The molecular formula of the triterpene is C₃₀H₅₀O, was deduced from the chemical ionization mass spectrum, while the IR spectrum showed a well-defined absorption due to OH (3620 cm⁻¹). The ¹H NMR spectrum showed the presence of seven methyl groups, H-3 (3.73, dd, 1H, *J* = 5 Hz), with a (1H, dddd) centered at 0.35 (*J* = 7 Hz) suggesting the attachment of a cyclopropane ring to C-21 in epifriedelinol and the absence of the methyl group at C-30 (see Experimental). The triterpene gave molecular ion [M]⁺ at *m/z* 426 corresponding to C₃₀H₅₀O. The fragment *m/z* 411, produced from [M]⁺ after removal of a methyl group. The appearance of fragments *m/z* 394, 341, 275, 205 and 109 can be explained on the basis of elimination of OH,

C₄H₅, C₅H₆, C₅H₁₀ and C₇H₁₂ from the fragment *m/z* 411. The *m/z* 109 ion is converted into the fragment ion *m/z* 55 (base peak) by loss of C₄H₆. Thus, the structure of the triterpene was settled to be 1.

The seasonal difference in the constituents of *C. aegyptiaca* L. is worthy of comment. The sample collected in August from the same locality did not contain this type of triterpene.



Experimental

The air dried plant material, collected in April, from the garden of Mansoura University, was extracted with C₂H₅OC₂H₅/petroleum ether/CH₃OH (1:3:1). The extract of the aerial parts of *C. aegyptiaca* (150 g) was first treated with CH₃OH to remove long chain hydrocarbons and then partially separated by CC (SiO₂) with petroleum ether and increasing amounts of C₂H₅OC₂H₅ and finally C₂H₅OC₂H₅/CH₃OH (10:1). The fraction obtained with petroleum ether/C₂H₅OC₂H₅ (9:1) afforded 10 mg phytol. The fraction obtained with petroleum ether/C₂H₅OC₂H₅ (3:1) afforded (1) 5 mg colourless oil; IR ν CCl₄/max. cm⁻¹: 3620 (OH); MS *m/z* (rel. int.): 426.386 [M]⁺ (2) (calcd for C₃₀H₅₀O: 426.386), 411 [M-CH₃]⁺ (45), 394 [411-OH]⁺ (8), 341 [394-C₂H₅]⁺ (2), 275 [341-C₅H₆]⁺ (35), 205 [275-C₅H₁₀]⁺ (22), 109 [205-C₇H₁₂]⁺ (61), 55 [109-C₄H₆]⁺ (100); ¹H NMR (400 MHz) CDCl₃: δ 0.7, 0.83, 0.93, 0.94, 0.98, 0.99, 1.01 (7CH₃), 3.73 (dd, 1H, H-3, *J* = 5 Hz) and 0.35 (dddd, 1H, C-30 CH₂, *J* = 7 Hz).

We thank Prof. F. Bohlmann, TU Berlin (West) for the spectral measurements.

* Part 18 in the series of Egyptian Compositae, for part 17, see M. A. Metwally, Pharmazie 1989.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen 0932-0776/89/1200-1584/\$ 01.00/0

[1] M. A. Metwally and A. M. Dawidar, Pharmazie **39**, H8, 575 (1984).

[2] M. A. Metwally, Pharmazie, communicated.