

THE BOUNDED APPROXIMATION PROPERTY OF VARIABLE LEBESGUE SPACES AND NUCLEARITY

JULIO DELGADO AND MICHAEL RUZHANSKY

ABSTRACT. In this paper we prove the bounded approximation property for variable exponent Lebesgue spaces, study the concept of nuclearity on such spaces and apply it to trace formulae such as the Grothendieck-Lidskii formula. We apply the obtained results to derive criteria for nuclearity and trace formulae for periodic operators on \mathbb{R}^n in terms of global symbols.

1. INTRODUCTION

The approximation property on a Banach space arises in the study of the concept of trace and was first introduced in its current shape by Grothendieck in his monumental work [Gro55]. A particular importance for a Banach space enjoying this property is that the trace can be defined and consequently the Fredholm's determinant leading to numerous further developments. Indeed, this problematic finds itself closely related to a wide range of analysis areas: operator theory, spectral analysis, harmonic analysis, functional analysis, PDEs.

In [Enf73] Enflo constructed a counterexample to the approximation property in Banach spaces. A more natural counterexample was then found by Szankowski [Sza81] who proved that $B(H)$ does not have the approximation property. More recently these properties have been intensively investigated by Figiel, Johnson, Pelczyński and Szankowski in [FJP11], [JS12]. Alberti, Csörnyei, Pelczyński and Preiss [ACPP05] established the bounded approximation property (BAP) for functions of bounded variations, and Roginskaya and Wojciechowski [RW14] for Sobolev spaces $W^{1,1}$. The authors have recently established the metric approximation property for mixed-norm L^p , modulation and Wiener amalgam spaces in [DRW16a], see also [DRW16b]. Other works on the bounded approximation property can be found in [LLO10], [LLO14]. A weak approximation property has been introduced and investigated in [LO05]. The fact that the approximation property does not imply the bounded approximation property was proved in [FJ73]. For a historical perspective and an introduction to the subject the reader can be referred to Pietsch's book [Pie07, Section 5.7.4] and the recent revisited presentation on the Grothendieck's classical work by Diestel, Fourie and Swart [DFS08]. The monograph [Rya02] contains a more accessible introduction to the topic as well as several examples of spaces enjoying

Date: October 17, 2018.

2010 Mathematics Subject Classification. Primary 46B28, 47B10; Secondary 47G10, 47B06.

Key words and phrases. Variable exponent Lebesgue spaces, approximation property, nuclearity, trace formulae.

The first author was supported by the Leverhulme Research Grant RPG-2014-02. The second author was supported by the EPSRC Grant EP/K039407/1. No new data was collected or generated during the course of research.

approximation properties. An introductory survey to the concept of trace on Banach spaces appeared in [Rob14] by Robert.

To formulate the notions more precisely, let $\mathcal{B}_1, \mathcal{B}_2$ be Banach spaces. A linear operator T from \mathcal{B}_1 to \mathcal{B}_2 is called *nuclear* if there exist sequences (x'_n) in \mathcal{B}'_1 and (y_n) in \mathcal{B}_2 such that

$$Tx = \sum_{n=1}^{\infty} \langle x, x'_n \rangle y_n \quad \text{and} \quad \sum_{n=1}^{\infty} \|x'_n\|_{\mathcal{B}'_1} \|y_n\|_{\mathcal{B}_2} < \infty.$$

This definition agrees with the concept of a trace class operator in the setting of Hilbert spaces. The set of nuclear operators from \mathcal{B}_1 into \mathcal{B}_2 forms the ideal of nuclear operators $\mathcal{N}(\mathcal{B}_1, \mathcal{B}_2)$ endowed with the norm

$$N(T) = \inf \left\{ \sum_{n=1}^{\infty} \|x'_n\|_{\mathcal{B}'_1} \|y_n\|_{\mathcal{B}_2} : T = \sum_{n=1}^{\infty} x'_n \otimes y_n \right\}.$$

If $\mathcal{B} = \mathcal{B}_1 = \mathcal{B}_2$, it is natural to attempt to define the trace of $T \in \mathcal{N}(\mathcal{B})$ by

$$\text{Tr}(T) := \sum_{n=1}^{\infty} x'_n(y_n), \quad (1.1)$$

where $T = \sum_{n=1}^{\infty} x'_n \otimes y_n$ is a representation of T . Grothendieck [Gro55] proved that the trace $\text{Tr}(T)$ is well defined for all nuclear operators $T \in \mathcal{N}(\mathcal{B})$ if and only if the Banach space \mathcal{B} has the *approximation property* (see also Pietsch [Pie87] or Defant and Floret [DF93]), which means that for every compact set K in \mathcal{B} and for every $\epsilon > 0$ there exists $F \in \mathcal{F}(\mathcal{B})$ such that

$$\|x - Fx\| < \epsilon \quad \text{for all } x \in K,$$

where we have denoted by $\mathcal{F}(\mathcal{B})$ the space of all finite rank bounded linear operators on \mathcal{B} . We denote by $\mathcal{L}(\mathcal{B})$ the Banach algebra of bounded linear operators on \mathcal{B} .

There are more related approximation properties, e.g. if in the definition above the operator F satisfies $\|F\| \leq M$ for a fixed $M > 0$ one says that \mathcal{B} possesses the *bounded approximation property*. In the case $M = 1$ one says that \mathcal{B} has the *metric approximation property*. The fact that the classical spaces $C(X)$, where X is a compact topological space and $L^p(\mu)$ for $1 \leq p < \infty$ satisfy the metric approximation property can be found in [Pie80].

As we know from Lidskii [Lid59], in Hilbert spaces the operator trace is equal to the sum of the eigenvalues of the operator counted with multiplicities. This property is nowadays called the Lidskii formula. An important feature on Banach spaces even endowed with the approximation property is that the Lidskii formula does not hold in general for nuclear operators. Thus, in the setting of Banach spaces, Grothendieck [Gro55] introduced a more restricted class of operators where Lidskii formula holds, this fact motivating the following definition.

Let $\mathcal{B}_1, \mathcal{B}_2$ be Banach spaces and let $0 < r \leq 1$. A linear operator T from \mathcal{B}_1 into \mathcal{B}_2 is called *r-nuclear* if there exist sequences (x'_n) in \mathcal{B}'_1 and (y_n) in \mathcal{B}_2 so that

$$Tx = \sum_{n=1}^{\infty} \langle x, x'_n \rangle y_n \quad \text{and} \quad \sum_{n=1}^{\infty} \|x'_n\|_{\mathcal{B}'_1}^r \|y_n\|_{\mathcal{B}_2}^r < \infty. \quad (1.2)$$

We associate a quasi-norm $n_r(T)$ by

$$n_r(T)^r := \inf \left\{ \sum_{n=1}^{\infty} \|x'_n\|_{\mathcal{B}_1}^r \|y_n\|_{\mathcal{B}_2}^r \right\},$$

where the infimum is taken over the representations of T as in (1.2). When $r = 1$ the 1-nuclear operators agree with the nuclear operators, and as already mentioned, in that case this definition also agrees with the concept of trace class operators in the setting of Hilbert spaces ($\mathcal{B}_1 = \mathcal{B}_2 = H$). More generally, Oloff proved in [Olo72] that the class of r -nuclear operators coincides with the Schatten class $S_r(H)$ when $\mathcal{B}_1 = \mathcal{B}_2 = H$ is a Hilbert space and $0 < r \leq 1$. Moreover, Oloff proved that

$$\|T\|_{S_r} = n_r(T), \quad (1.3)$$

where $\|\cdot\|_{S_r}$ denotes the classical Schatten quasi-norms in terms of singular values.

In [Gro55] Grothendieck proved that if T is $\frac{2}{3}$ -nuclear from \mathcal{B} into \mathcal{B} for a Banach space \mathcal{B} , then

$$\mathrm{Tr}(T) = \sum_{j=1}^{\infty} \lambda_j, \quad (1.4)$$

where λ_j ($j = 1, 2, \dots$) are the eigenvalues of T with multiplicities taken into account, and $\mathrm{Tr}(T)$ is as in (1.1). Grothendieck also established its applications to the distribution of eigenvalues of operators in Banach spaces. We refer to [DR14c] for several conclusions in the setting of compact Lie groups concerning summability and distribution of eigenvalues of operators on L^p -spaces once we have information on their r -nuclearity. See also [DRT] for applications of the notion of nuclearity to boundary value problems. Kernel conditions on compact manifolds have been investigated in [DR14d], [DR14b].

On the other hand, the variable exponent Lebesgue spaces are a generalisation of the classical Lebesgue spaces, replacing the constant exponent p by a variable exponent function $p(x)$. Variable exponent Lebesgue spaces were introduced by Orlicz [Orl31] in 1931 and some properties were further developed by Nakano in the 1950s [Nak50], [Nak51] within the more general framework of modular spaces. Subsequently developments of modular spaces were carried out in the 1970s and 1980s by Hudzik, Musielak, Portnov [Hud76a], [Hud76b], [Hud76c], [Mus83], [Por66].

A more specific study of variable Lebesgue spaces only appears in 1961 with the work of Tsenov [Tse61] who independently discovered those spaces and later in the works of Sharapudinov [Sha79], [Sha83], [Sha86], [Sha96] and Zhikov [Zhi82], [Zhi92], [Zhi97]. Further, the development of the analysis of many problems on those spaces has been of great interest in the last decades as has been exhibited in the recent book [DHHR11], [CUF13], [CUFRW14] and the literature therein.

We now briefly recall the definition of variable exponent Lebesgue spaces and we refer the reader to [DHHR11] and [CUF13] for the basic properties of such spaces. Let $(\Omega, \mathcal{M}, \mu)$ be a σ -finite, complete measure space. We define $\mathcal{P}(\Omega, \mu)$ to be the set of all μ -measurable functions $p : \Omega \rightarrow [1, \infty]$. The functions in $\mathcal{P}(\Omega, \mu)$ are called variable exponents on Ω . We define

$$p^+ = p_{\Omega}^+ := \operatorname{ess\,sup}_{x \in \Omega} p(x), \quad p^- = p_{\Omega}^- := \operatorname{ess\,inf}_{x \in \Omega} p(x).$$

If $p^+ < \infty$, then p is called a *bounded variable exponent*. If $f : \Omega \rightarrow \mathbb{R}$ is a measurable function we define the *modular* associated with $p = p(\cdot)$ by

$$\rho_{p(\cdot)}(f) := \int_{\Omega} |f(x)|^{p(x)} d\mu(x),$$

and

$$\|f\|_{L^{p(\cdot)}(\mu)} := \inf\{\lambda > 0 : \rho_{p(\cdot)}(f/\lambda) \leq 1\}.$$

The resulting spaces $L^{p(\cdot)}(\mu)$ of measurable functions such that $\|f\|_{L^{p(\cdot)}(\mu)} < \infty$ are Banach spaces and enjoy many properties similar to the classical Lebesgue L^p spaces. For example, we will often make use of the following modification of Hölder inequality which becomes affected by factor 2: let $p, q, s \in \mathcal{P}(\Omega, \mu)$ be such that

$$\frac{1}{s(x)} = \frac{1}{p(x)} + \frac{1}{q(x)}$$

holds for μ -almost every $x \in \Omega$. Then we have

$$\|fg\|_{L^s(\cdot)(\mu)} \leq 2 \|f\|_{L^p(\cdot)(\mu)} \|g\|_{L^q(\cdot)(\mu)}. \quad (1.5)$$

We refer to [DHHR11, Lemma 3.2.20] for a more detailed statement.

At the same time, there are some exceptions and differences to the classical theory, for instance the Young inequality fails in the variable exponent case, a fact proved in 1991 by Kováčik and Rákosník (cf. [KR91]) and essentially due to the loss of boundedness of translation operators on $L^{p(\cdot)}$ spaces (see also [CUF13], Theorem 5.19). If the variable exponent $p(\cdot)$ is bounded the space $L^{p(\cdot)}(\mu)$ is separable and if we denote by $p'(\cdot)$ the variable exponent defined pointwise by

$$\frac{1}{p(x)} + \frac{1}{p'(x)} = 1,$$

then $(L^{p(\cdot)}(\mu))' = L^{p'(\cdot)}(\mu)$, where the identity refers to the associate space and not necessarily to the isometric dual space. Moreover, if $1 < p^- \leq p^+ < \infty$ the space $L^{p(\cdot)}(\mu)$ is reflexive. For the study of the approximation property we will restrict to consider bounded variable exponents due to the density of the simple functions in $L^{p(\cdot)}$ in that case.

In this work we are going to establish the bounded approximation property for variable exponent Lebesgue spaces, study the concept of nuclearity on such spaces and apply it to trace formulae such as the Grothendieck-Lidskii formula and the analysis of pseudo-differential operators on the torus.

2. BOUNDED APPROXIMATION PROPERTY FOR VARIABLE EXPONENT LEBESGUE SPACES

In this section we will prove that the variable exponent spaces $L^{p(\cdot)}(\mu)$ satisfy the bounded approximation property.

In the rest of this section we will assume that our measure space $(\Omega, \mathcal{M}, \mu)$ is σ -finite and complete. We will also assume that the exponent $p(\cdot)$ is bounded since only in such case the simple functions are dense in $L^{p(\cdot)}(\mu)$ (cf. [DHHR11], Corollary 3.4.10).

We shall now formulate some preparatory lemmata useful for the proof of the bounded approximation property. Let I be a countable set of indices endowed with the counting measure ν . For $p \in \mathcal{P}(I, \nu)$, we will denote by $\ell^{p(\cdot)}(I)$ or simply by $\ell^{p(\cdot)}$ the corresponding variable exponent Lebesgue space whose norm is given by

$$\|h\|_{\ell^{p(\cdot)}} = \inf\{\lambda > 0 : \sum_{k \in I} \left| \frac{h_k}{\lambda} \right|^{p_k} \leq 1\}.$$

Given a Banach space \mathcal{B} and $u \in \mathcal{B}, z \in \mathcal{B}'$, we will also denote by $\langle u, z \rangle_{\mathcal{B}, \mathcal{B}'}$, or simply by $\langle u, z \rangle$, the valuation $z(u)$.

Lemma 2.1. *Let \mathcal{B} be a Banach space and $q \in \mathcal{P}(I, \nu)$. Let $(u_i)_{i \in I}, (v_i)_{i \in I}$ be sequences in $\mathcal{B}', \mathcal{B}$ respectively such that*

$$\|\langle x, u_i \rangle\|_{\ell^{q(\cdot)}}, \|\langle v_i, z \rangle\|_{\ell^{q'(\cdot)}} \leq 1, \quad \text{for } \|x\|_{\mathcal{B}}, \|z\|_{\mathcal{B}'} \leq 1.$$

Then the operator $T = \sum_{i \in I} u_i \otimes v_i$ from \mathcal{B} into \mathcal{B} is well defined, bounded and satisfies $\|T\|_{\mathcal{L}(\mathcal{B})} \leq 2$.

Proof. Let $N \subset I$ be a finite subset of I . Let us write $T_N := \sum_{i \in N} u_i \otimes v_i$. It is clear that T_N is well defined. Moreover T_N is a bounded finite rank operator. Now, since $T_N x = \sum_{i \in N} \langle x, u_i \rangle v_i$, we observe that for $x \in \mathcal{B}, z \in \mathcal{B}'$ such that $\|x\|_{\mathcal{B}}, \|z\|_{\mathcal{B}'} \leq 1$, applying the Hölder inequality (1.5) for variable exponent spaces we obtain

$$|\langle T_N x, z \rangle| \leq \sum_{i \in N} |\langle x, u_i \rangle| |\langle v_i, z \rangle| \leq 2 \|\langle x, u_i \rangle\|_{\ell^{q(\cdot)}} \|\langle v_i, z \rangle\|_{\ell^{q'(\cdot)}} \leq 2.$$

Therefore $T = \lim_N T_N$ exists in $\mathcal{L}(\mathcal{B})$ and $\|T\|_{\mathcal{L}(\mathcal{B})} \leq 2$. \square

Lemma 2.2. *Let $\mathcal{B}_1, \mathcal{B}_2$ be Banach spaces and $(L_i)_i$ a net contained in $\mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)$ such that for every $x \in \mathcal{B}_1$, $\lim_i L_i x = Lx$ for some $Lx \in \mathcal{B}_2$. Then $L \in \mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)$, $\|L_i\|_{\mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)} \leq M$ for some $M > 0$ and L_i converge to L in the topology of uniform convergence on compact sets.*

Proof. The fact that $\|L_i\|_{\mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)} \leq M$ and $L \in \mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)$ follows from the uniform boundedness principle. For the rest, let $K \subset \mathcal{B}_1$ be compact, $\epsilon > 0$ and $M \geq 1$ such that $\|L_i\|_{\mathcal{L}(\mathcal{B}_1, \mathcal{B}_2)} \leq M$. Let $\{x_1, \dots, x_n\} \subset K$ be such that $K \subset \bigcup_{j=1}^n B(x_j, \frac{\epsilon}{3M})$. If i is large enough we have $\|Lx_j - L_i x_j\|_{\mathcal{B}_2} < \frac{\epsilon}{3M}$ for all $1 \leq j \leq n$. Let $x \in K$ and we pick j_0 such that $\|x - x_{j_0}\|_{\mathcal{B}_1} < \frac{\epsilon}{3M}$. Then

$$\|Lx - L_i x\|_{\mathcal{B}_2} \leq \|Lx - Lx_{j_0}\|_{\mathcal{B}_2} + \|Lx_{j_0} - L_i x_{j_0}\|_{\mathcal{B}_2} + \|L_i x_{j_0} - L_i x\|_{\mathcal{B}_2} < \epsilon.$$

Therefore L_i converge to L uniformly on compact sets. \square

As a consequence we obtain:

Corollary 2.3. *Let \mathcal{B} be a Banach space. If there is a net $(L_i)_i$ contained in $\mathcal{F}(\mathcal{B})$ such that $\sup_i \|L_i\| \leq M < \infty$ and $\lim_i L_i x = x$ for every $x \in \mathcal{B}$, then \mathcal{B} has the bounded approximation property with constant M .*

We can now prove the main result of this section:

Theorem 2.4. *Let $p \in \mathcal{P}(\Omega, \mu)$ be a bounded variable exponent. Then, the variable exponent Lebesgue space $L^{p(\cdot)}(\mu)$ has the bounded approximation property.*

Since the Hölder inequality (1.5) in variable Lebesgue spaces $L^{p(\cdot)}$ holds with constant 2, we obtain the bounded approximation property in this setting, rather than the metric approximation property valid for the usual L^p -spaces.

However, the bounded approximation property implies the metric approximation property if the space is reflexive. In our case, this happens if $\Omega \subset \mathbb{R}^n$ is open and $1 < p^-, p^+ < \infty$ (cf. [DHHR11]), in which case $L^{p(\cdot)}(\Omega)$ has the metric approximation property. This gives a small change to Theorem 3.1 in [DRW16b], with the rest of [DRW16b] unchanged.

Proof of Theorem 2.4. We first consider the case when $p \in \mathcal{P}(\Omega, \mu)$ is a simple function and we write

$$p(x) = \sum_{j=1}^l p_j 1_{\Omega_j}(x),$$

where $p_j > 0$, the sets Ω_j are disjoint of finite measure, 1_{Ω_j} denotes the characteristic function of the set Ω_j .

Let $\mathfrak{F} = \{\Omega_1, \dots, \Omega_l\}$ be a finite family of disjoint measurable sets of finite positive measure. We denote by \mathbf{P} the collection of such families. To a $\mathfrak{F} \in \mathbf{P}$ we associate a finite rank operator $L_{\mathfrak{F}}$ from $L^{p(\cdot)}$ into $L^{p(\cdot)}$ defined by

$$L_{\mathfrak{F}} f := \sum_{k=1}^l \mu(\Omega_k)^{-1} \langle f, 1_{\Omega_k} \rangle_{L^{p(\cdot)}, L^{p'(\cdot)}} 1_{\Omega_k}, \quad (2.1)$$

where 1_{Ω_k} denotes the characteristic function of the set Ω_k .

We observe that $L_{\mathfrak{F}} f$ is well defined since $0 < \mu(\Omega_k) < \infty$ for $1 \leq k \leq l$ and the duality $\langle \cdot, \cdot \rangle_{L^{p(\cdot)}, L^{p'(\cdot)}}$ is well defined by using the Hölder inequality for variable exponent spaces, see (1.5).

In the collection \mathbf{P} we define the partial order $\mathfrak{F}_1 \leq \mathfrak{F}_2$ if any set in \mathfrak{F}_1 is the union of sets in \mathfrak{F}_2 . We also say that \mathfrak{F}_2 is *finer* than \mathfrak{F}_1 if $\mathfrak{F}_1 \leq \mathfrak{F}_2$. This order begets a directed set.

Let \mathfrak{F}_1 be the family of sets associated to the exponent $p(\cdot)$. By choosing a finer \mathfrak{F} we can rewrite the operator $L_{\mathfrak{F}}$ given by (2.1) in different ways which will be useful later on: we define

$$u_k := \frac{1_{\Omega_k}}{\mu(\Omega_k)^{\frac{1}{p_k}}}, \quad v_k := \frac{1_{\Omega_k}}{\mu(\Omega_k)^{\frac{1}{p_k}}},$$

so that we can write

$$\begin{aligned} L_{\mathfrak{F}} &= \sum_{k=1}^l \frac{1_{\Omega_k}}{\mu(\Omega_k)^{\frac{1}{p_k}}} \otimes \frac{1_{\Omega_k}}{\mu(\Omega_k)^{\frac{1}{p_k}}} \\ &= \sum_{k=1}^l u_k \otimes v_k \\ &= \sum_{k=1}^l \frac{1}{\mu(\Omega_k)} (1_{\Omega_k} \otimes 1_{\Omega_k}). \end{aligned}$$

We will prove that $\|L_{\mathfrak{P}}\|_{\mathcal{L}(L^{p(\cdot)})} \leq 2$ by applying Lemma 2.1 in the case $\mathcal{B} = L^{p(\cdot)}$, the finite families u_k, v_k and $q = p(k) = p_k$. Let $f \in L^{p(\cdot)}$, $g \in L^{p'(\cdot)}$ be such that $\|f\|_{L^{p(\cdot)}}, \|g\|_{L^{p'(\cdot)}} \leq 1$. Then we have to show that

$$\|\langle f, u_k \rangle\|_{\ell^{p(\cdot)}} \leq 1 \quad \text{and} \quad \|\langle v_k, g \rangle\|_{\ell^{p'(\cdot)}} \leq 1.$$

In order to prove the corresponding property for $f \in L^{p(\cdot)}$, it is enough to consider a simple function $f \in L^{p(\cdot)}$ such that $\|f\|_{L^{p(\cdot)}} \leq 1$. The general case follows then by a standard density argument. By redefining partitions, we can assume that f can be written in the form

$$f(x) = \sum_{k=1}^l \beta_k 1_{\Omega_k}(x).$$

Now, for $\lambda > 0$ we have

$$\rho_{p(\cdot)}(f/\lambda) = \int_{\Omega} \left| \frac{f(x)}{\lambda} \right|^{p(x)} dx = \sum_{k=1}^l \int_{\Omega_k} \left| \frac{\beta_k}{\lambda} \right|^{p_k} dx = \sum_{k=1}^l \left| \frac{\beta_k}{\lambda} \right|^{p_k} \mu(\Omega_k).$$

We also observe that

$$\langle f, u_k \rangle = \beta_k \frac{\mu(\Omega_k)}{\mu(\Omega_k)^{\frac{1}{p_k}}} = \beta_k \mu(\Omega_k)^{\frac{1}{p_k}}.$$

Hence

$$\begin{aligned} \|\langle f, u_k \rangle\|_{\ell^{p(\cdot)}} &= \inf \left\{ \lambda > 0 : \sum_{k=1}^l \left| \frac{\beta_k}{\lambda} \right|^{p_k} \mu(\Omega_k) \leq 1 \right\} \\ &= \inf \left\{ \lambda > 0 : \rho_{p(\cdot)}(f/\lambda) \leq 1 \right\} \\ &= \|f\|_{L^{p(\cdot)}} \leq 1. \end{aligned}$$

We have shown that $\|\langle f, u_k \rangle\|_{\ell^{p(\cdot)}} \leq 1$, the proof of $\|\langle v_k, g \rangle\|_{\ell^{p'(\cdot)}} \leq 1$ is similar and we omit it. Hence $\|L_{\mathfrak{P}}\|_{\mathcal{L}(L^{p(\cdot)})} \leq 2$.

We now consider the net of finite rank operators $(L_{\mathfrak{P}})_{\mathfrak{P} \geq \mathfrak{P}_1}$ and prove that

$$\lim_{\mathfrak{P}} L_{\mathfrak{P}} f = f$$

for every $f \in L^{p(\cdot)}$. It is enough to see this for f simple by the density of simple functions in $L^{p(\cdot)}$ (cf. [DHHR11], Corollary 3.4.10). Indeed, let us write f in the form

$$f(x) := \sum_{m=1}^s \alpha_m 1_{\tilde{\Omega}_m}(x).$$

If we chose \mathfrak{P} finer than $\mathcal{Q}_1 = \{\tilde{\Omega}_m : 1 \leq m \leq s\}$, then $L_{\mathfrak{P}}f = f$. Indeed, since the sets Ω_k are disjoint we have

$$\begin{aligned} L(1_{\Omega_j}) &= \sum_{k=1}^l \frac{1}{\mu(\Omega_k)} (1_{\Omega_k} \otimes 1_{\Omega_k})(1_{\Omega_j}) \\ &= \sum_{k=1}^l \frac{1}{\mu(\Omega_k)} \langle 1_{\Omega_j}, 1_{\Omega_k} \rangle_{L^{p(\cdot)}, L^{p(\cdot)}} (1_{\Omega_k}) \\ &= \frac{1}{\mu(\Omega_j)} \langle 1_{\Omega_j}, 1_{\Omega_j} \rangle_{L^{p(\cdot)}, L^{p(\cdot)}} (1_{\Omega_j}) \\ &= 1_{\Omega_j}. \end{aligned}$$

Therefore, $L_{\mathfrak{P}}f = f$ for $\mathfrak{P} \geq \mathcal{Q}_1$ and thus $\lim_{\mathfrak{P}} L_{\mathfrak{P}}f = f$ in $L^{p(\cdot)}$. We have actually proved that $L^{p(\cdot)}$ satisfied the bounded approximation property by an application of Corollary 2.3.

By an additional argument we will obtain the desired property in the general case. We now consider a variable exponent $p(\cdot)$ such that $p^+ < \infty$. Then, there exists an increasing sequence of simple functions $p^j(\cdot)$ such that $\lim_j p^j(\cdot) = p(\cdot)$ a.e. For each j we associate to a $\mathfrak{P}^j \in \mathbf{P}$ an operator $L_{\mathfrak{P}^j}$ as in (2.1) which due to its form is also defined from $L^{p(\cdot)}(\mu)$ into $L^{p(\cdot)}(\mu)$. The fact that $\lim_j L_{\mathfrak{P}^j}f = f$ in $L^{p(\cdot)}$ follows as in the previous case. We claim that $\|L_{\mathfrak{P}^j}\|_{\mathcal{L}(L^{p(\cdot)})} \leq 2$ which by an application of Corollary 2.3 with $M = 2$ will conclude the proof. Indeed, if f is a simple function such that $\|f\|_{L^{p(\cdot)}} \leq 1$ and \mathfrak{P}_1 is its corresponding family in \mathbf{P} , we observe that by choosing $\mathfrak{P}^j \geq \mathfrak{P}_1$ we obtain $L_{\mathfrak{P}^j}f = f$. \square

3. NUCLEARITY ON VARIABLE EXPONENT LEBESGUE SPACES

In this section we establish some basic properties for the kernels of nuclear operators on $L^{p(\cdot)}$ spaces. We also prove a characterisation of nuclear operators on $L^{p(\cdot)}$.

We start by proving a lemma giving basic properties of a kernel corresponding to a nuclear operator on $L^{p(\cdot)}$ spaces when μ is a finite measure. In the rest of this section we shall consider two variable exponents $p(\cdot) \in \mathcal{P}(\Omega, \mu)$, $q(\cdot) \in \mathcal{P}(\Xi, \nu)$ and the variable exponent conjugate $p'(\cdot)$ of $p(\cdot)$ such that

$$\frac{1}{p(\cdot)} + \frac{1}{p'(\cdot)} = 1.$$

Lemma 3.1. *Let $(\Omega, \mathcal{M}, \mu)$ and (Ξ, \mathcal{M}', ν) be two finite and complete measure spaces. Let $f \in L^{p(\cdot)}(\mu)$, and $(g_n)_n, (h_n)_n$ be sequences in $L^{q(\cdot)}(\nu)$ and $L^{p'(\cdot)}(\mu)$, respectively, such that $\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}} \|h_n\|_{L^{p'(\cdot)}} < \infty$. Then*

(a) *The series $\sum_{j=1}^{\infty} g_j(x)h_j(y)$ converges absolutely for a.e. (x, y) and, consequently,*

$$\lim_n \sum_{j=1}^n g_j(x)h_j(y) \text{ is finite for a.e. } (x, y).$$

(b) For $k(x, y) := \sum_{j=1}^{\infty} g_j(x)h_j(y)$, we have $k \in L^1(\nu \otimes \mu)$.

(c) If $k_n(x, y) = \sum_{j=1}^n g_j(x)h_j(y)$ then $\|k_n - k\|_{L^1(\nu \otimes \mu)} \rightarrow 0$.

(d) $\lim_n \int_{\Omega} \left(\sum_{j=1}^n g_j(x)h_j(y) \right) f(y) d\mu(y) = \int_{\Omega} \left(\sum_{j=1}^{\infty} g_j(x)h_j(y) \right) f(y) d\mu(y)$,

for a.e. x .

Proof. We first write $\tilde{k}_n(x, y) := \sum_{j=1}^n g_j(x)h_j(y)f(y)$ and note that since ν is finite then $\|1\|_{L^{q'(\cdot)}(\nu)} < \infty$. Indeed, $\int_{\Xi} |1/\lambda|^{q'(x)} d\nu(x) = \nu(\Xi) < \infty$, for $\lambda = 1$. Now, by applying the Hölder inequality (1.5) which is affected by the factor 2 in the setting of variable exponents we obtain

$$\begin{aligned} \int_{\Omega} \int_{\Xi} |\tilde{k}_n(x, y)| d\nu(x) d\mu(y) &\leq \int_{\Omega} \int_{\Xi} \sum_{j=1}^n |g_j(x)h_j(y)f(y)| d\nu(x) d\mu(y) \\ &\leq \sum_{j=1}^n \int_{\Xi} |g_j(x)| d\nu(x) \int_{\Omega} |h_j(y)| |f(y)| d\mu(y) \\ &\leq 2 \|1\|_{L^{q'(\cdot)}(\nu)} \|f\|_{L^{p(\cdot)}(\mu)} \sum_{j=1}^n \|g_j\|_{L^{q(\cdot)}(\nu)} \|h_j\|_{L^{p'(\cdot)}(\mu)} \\ &\leq M < \infty \text{ for all } n. \end{aligned}$$

Hence $\|\tilde{k}_n\|_{L^1(\nu \otimes \mu)} \leq M$ for all n . We now consider a sequence (s_n) defined by

$$s_n(x, y) := \sum_{j=1}^n |g_j(x)h_j(y)f(y)|,$$

which is increasing in $L^1(\nu \otimes \mu)$ and satisfies

$$\sup_n \int \int |s_n(x, y)| d\mu(x) d\mu(y) \leq M < \infty.$$

By an application of the monotone convergence theorem, the limit

$$s(x, y) = \lim_n s_n(x, y)$$

exists for a.e. (x, y) and $s \in L^1(\nu \otimes \mu)$. Moreover, since $f = 1 \in L^{p(\cdot)}(\mu)$ and the fact that $|k(x, y)| \leq s(x, y)$ we deduce (a) and (b).

The part (c) can be deduced by using the Lebesgue dominated convergence theorem applied to the sequence (\tilde{k}_n) dominated by $s(x, y)$, and setting $f \equiv 1$, in which case $\tilde{k}_n = k_n$.

For the part (d) we observe that with $\tilde{k}_n(x, y) = \sum_{j=1}^n g_j(x)h_j(y)f(y)$, we have $|\tilde{k}_n(x, y)| \leq s(x, y)$ for all n and every (x, y) . From the fact that $s \in L^1(\nu \otimes \mu)$

we obtain that $s(x, \cdot) \in L^1(\mu)$ for a.e x . Then (d) is obtained from Lebesgue dominated convergence theorem. \square

Remark 3.2. We observe that the condition of finiteness of the measures in the lemma above is crucial to obtain $k = k(x, y) \in L^1(\nu \otimes \mu)$. For instance, let $\Omega = \Xi = \mathbb{R}^n$, $\mu = \nu$ be the Lebesgue measure and $p = p(\cdot), q = q(\cdot)$ constant exponents such that $1 \leq p, q < \infty$. Then by using the fact that $p' > 1$, we define $k(x, y) := g(x)h(y)$, with $g \in L^q(\mu) \setminus \{0\}, h \in L^{p'}(\mu) \setminus L^1(\mu)$. Then

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |k(x, y)| d\mu(x) d\mu(y) = \int_{\mathbb{R}^n} |g(x)| d\mu(x) \int_{\mathbb{R}^n} |h(y)| d\mu(y) = \infty.$$

We can now formulate a characterisation of r -nuclear operators on variable exponent Lebesgue spaces for finite measure spaces.

Theorem 3.3. *Let $(\Omega, \mathcal{M}, \mu)$ and (Ξ, \mathcal{M}', ν) be two complete and finite measure spaces. Let $0 < r \leq 1$. Then T is r -nuclear operator from $L^{p(\cdot)}(\mu)$ into $L^{q(\cdot)}(\nu)$ if and only if there exist a sequence (g_n) in $L^{q(\cdot)}(\nu)$, and a sequence (h_n) in $L^{p'(\cdot)}(\mu)$ such that $\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)}^r \|h_n\|_{L^{p'(\cdot)}(\mu)}^r < \infty$, and such that for all $f \in L^{p(\cdot)}(\mu)$, we have*

$$Tf(x) = \int_{\Omega} \left(\sum_{n=1}^{\infty} g_n(x) h_n(y) \right) f(y) d\mu(y), \text{ for a.e } x.$$

Proof. We will assume that $r = 1$. The case $0 < r < 1$ follows by inclusion. Let T be a nuclear operator from $L^{p(\cdot)}(\mu)$ into $L^{q(\cdot)}(\nu)$. Then there exist sequences (g_n) in $L^{q(\cdot)}(\nu)$, (h_n) in $L^{p'(\cdot)}(\mu)$ such that $\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)} \|h_n\|_{L^{p'(\cdot)}(\mu)} < \infty$ and

$$Tf = \sum_n \langle f, h_n \rangle g_n.$$

Now

$$Tf = \sum_n \langle f, h_n \rangle g_n = \sum_n \left(\int_{\Omega} h_n(y) f(y) d\mu(y) \right) g_n,$$

where the sums converge with respect to the $L^{q(\cdot)}(\nu)$ -norm. There exist (cf. [DHHR11], Lemma 3.2.10) two sub-sequences (\tilde{g}_n) and (\tilde{h}_n) of (g_n) and (h_n) respectively such that

$$(Tf)(x) = \sum_n \langle f, \tilde{h}_n \rangle \tilde{g}_n(x) = \sum_n \left(\int_{\Omega} \tilde{h}_n(y) f(y) d\mu(y) \right) \tilde{g}_n(x), \text{ for a.e } x.$$

Now taking into account that the pair $((\tilde{g}_n), (\tilde{h}_n))$ satisfies

$$\sum_{n=1}^{\infty} \|\tilde{g}_n\|_{L^{q(\cdot)}(\nu)} \|\tilde{h}_n\|_{L^{p'(\cdot)}(\mu)} < \infty,$$

and by applying Lemma 3.1 (d), it follows that

$$\begin{aligned} \sum_{n=1}^{\infty} \left(\int_{\Omega} \tilde{h}_n(y) f(y) d\mu(y) \right) \tilde{g}_n(x) &= \lim_n \sum_{j=1}^n \left(\int_{\Omega} \tilde{h}_j(y) f(y) d\mu(y) \right) \tilde{g}_j(x) \\ &= \lim_n \int_{\Omega} \left(\sum_{j=1}^n \tilde{g}_j(x) \tilde{h}_j(y) f(y) \right) d\mu(y) \\ &= \int_{\Omega} \left(\sum_{n=1}^{\infty} \tilde{g}_n(x) \tilde{h}_n(y) \right) f(y) d\mu(y), \text{ for a.e. } x. \end{aligned}$$

Conversely, let us assume that there exist sequences $(g_n)_n$ in $L^{q(\cdot)}(\nu)$, and $(h_n)_n$ in $L^{p'(\cdot)}(\mu)$ such that $\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)} \|h_n\|_{L^{p'(\cdot)}(\mu)} < \infty$, and for all $f \in L^{p(\cdot)}(\mu)$

$$Tf(x) = \int_{\Omega} \left(\sum_{n=1}^{\infty} g_n(x) h_n(y) \right) f(y) d\mu(y), \text{ for a.e. } x.$$

The Lemma 3.1 (d) gives

$$\begin{aligned} \int_{\Omega} \left(\sum_{n=1}^{\infty} g_n(x) h_n(y) \right) f(y) d\mu(y) &= \lim_n \int_{\Omega} \left(\sum_{j=1}^n g_j(x) h_j(y) f(y) \right) d\mu(y) \\ &= \lim_n \sum_{j=1}^n \left(\int_{\Omega} h_j(y) f(y) d\mu(y) \right) g_j(x) \\ &= \sum_n \left(\int_{\Omega} h_n(y) f(y) d\mu(y) \right) g_n(x) \\ &= \sum_n \langle f, h_n \rangle g_n(x) = (Tf)(x), \text{ a.e. } x. \end{aligned}$$

To prove that $Tf = \sum_n \langle f, h_n \rangle g_n$ in $L^{q(\cdot)}(\nu)$ we let $s_n := \sum_{j=1}^n \langle f, h_j \rangle g_j$, then $(s_n)_n$ is a sequence in $L^{q(\cdot)}(\nu)$ and

$$\begin{aligned} |s_n(x)| &\leq \|f\|_{L^{p(\cdot)}(\mu)} \sum_{j=1}^n \|h_j\|_{L^{p'(\cdot)}(\mu)} |g_j(x)| \\ &\leq \|f\|_{L^{p(\cdot)}(\mu)} \sum_{j=1}^{\infty} \|h_j\|_{L^{p'(\cdot)}(\mu)} |g_j(x)| =: \gamma(x), \text{ for all } n. \end{aligned}$$

Moreover, γ is well defined and $\gamma \in L^{q(\cdot)}(\nu)$ since it is the increasing limit of the sequence $(\gamma_n)_n = (\|f\|_{L^{p(\cdot)}(\mu)} \sum_{j=1}^n \|h_j\|_{L^{p'(\cdot)}(\mu)} |g_j(x)|)_n$ of $L^{q(\cdot)}(\nu)$ functions and

$$\|\gamma_n\|_{L^{q(\cdot)}(\nu)} \leq \|f\|_{L^{p(\cdot)}(\mu)} \sum_{j=1}^{\infty} \|h_j\|_{L^{p'(\cdot)}(\mu)} \|g_j\|_{L^{q(\cdot)}(\nu)} \leq M < \infty.$$

By the monotone convergence theorem we see that $\gamma \in L^{q(\cdot)}(\nu)$. Finally, applying the Lebesgue dominated convergence theorem we deduce that $s_n \rightarrow Tf$ in $L^{q(\cdot)}(\nu)$. \square

In the sequel we also establish a characterisation of r -nuclear operators for σ -finite measures. In order to get an analogue of the finite measures setting we first generalise Lemma 3.1.

Lemma 3.4. *Let $(\Omega, \mathcal{M}, \mu)$ and (Ξ, \mathcal{M}', ν) be two σ -finite, complete measure spaces. Let $f \in L^{p(\cdot)}(\mu)$, and $(g_n)_n, (h_n)_n$ be sequences in $L^{q(\cdot)}(\nu)$ and $L^{p'(\cdot)}(\mu)$, respectively, such that $\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)} \|h_n\|_{L^{p'(\cdot)}(\mu)} < \infty$. Then the parts (a) and (d) of Lemma 3.1 hold.*

Proof. (a) There exist two sequences $(\Omega_k)_k$ and $(\Xi_j)_j$ of disjoint measurable subsets of Ω and Ξ respectively such that $\bigcup_k \Omega_k = \Omega$, $\bigcup_j \Xi_j = \Xi$ and for all j, k

$$\mu(\Omega_k), \nu(\Xi_j) < \infty.$$

We now consider the respective restricted measure spaces $(\Omega_k, \mathcal{M}_k, \mu_k)$ and also $(\Xi_j, \mathcal{M}'_j, \nu_j)$ that we obtain by restricting Ω to Ω_k , and Ξ to Ξ_j for every k, j , and restricting the functions g_n to Ξ_j , and h_n to Ω_k for each n . Then, for all k, j

$$\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu_j)} \|h_n\|_{L^{p'(\cdot)}(\mu_k)} < \infty.$$

By Lemma 3.1 (a) it follows that $\sum_{j=1}^{\infty} g_j(x)h_j(y)$ converges absolutely for a.e. $(x, y) \in$

$\Xi^j \times \Omega^k$. Hence $\sum_{j=1}^{\infty} g_j(x)h_j(y)$ converges absolutely for almost every $(x, y) \in \Xi \times \Omega$.

This proves part (a).

From the part (a) the series $\sum_{j=1}^{\infty} g_j(x)h_j(y)f(y)$ converges absolutely for a.e. $(x, y) \in$

$\Xi \times \Omega$, the part (d) follows from the Lebesgue dominated convergence theorem applied as in the ‘‘only if’’ part of the proof of Theorem 3.3. \square

We are now ready to give the main result of this section, the extension of Theorem 3.3 to the setting of σ -finite measures.

Theorem 3.5. *Let $(\Omega, \mathcal{M}, \mu)$ and (Ξ, \mathcal{M}', ν) be σ -finite complete measure spaces. Let $0 < r \leq 1$. Then T is r -nuclear operator from $L^{p(\cdot)}(\mu)$ into $L^{q(\cdot)}(\nu)$ if and only if there exist a sequence (g_n) in $L^{q(\cdot)}(\nu)$, and a sequence (h_n) in $L^{p'(\cdot)}(\mu)$ such that*

$\sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)}^r \|h_n\|_{L^{p'(\cdot)}(\mu)}^r < \infty$, and such that for all $f \in L^{p(\cdot)}(\mu)$ we have

$$Tf(x) = \int_{\Omega} \left(\sum_{n=1}^{\infty} g_n(x)h_n(y) \right) f(y) d\mu(y), \text{ for a.e. } x.$$

Moreover, if $\Omega = \Xi$, $\mu = \nu$, $p(\cdot) = q(\cdot)$, $p^+ < \infty$, and T is r -nuclear in $\mathcal{L}(L^{p(\cdot)}(\mu))$, then

$$\text{Tr}(T) = \sum_{n=1}^{\infty} \langle g_n, h_n \rangle = \int_{\Omega} \sum_{n=1}^{\infty} g_n(x)h_n(x) d\mu.$$

Proof. For the characterisation it is enough to consider the case $r = 1$. But that characterisation now follows from the same lines of the proof of Theorem 3.3 by replacing the references to part (d) of Lemma 3.1 by part (d) of Lemma 3.4. On the other hand, since $p^+ < \infty$ the bounded approximation property holds and the trace is well defined. We observe that by definition (1.1) we can use the sequences g_n, h_n to calculate the trace, which gives $\text{Tr}(T) = \sum_{n=1}^{\infty} \langle g_n, h_n \rangle$. Moreover, the kernel $k(x, y) = \sum_{n=1}^{\infty} g_n(x)h_n(y)$ is well defined on the diagonal since for $p(\cdot) = q(\cdot)$, we have

$$\begin{aligned} |k(x, x)| &\leq \int_{\Omega} \sum_{n=1}^{\infty} |g_n(x)h_n(x)| d\mu \\ &= \sum_{n=1}^{\infty} \int_{\Omega} |g_n(x)h_n(x)| d\mu \\ &\leq \sum_{n=1}^{\infty} \|g_n\|_{L^{q(\cdot)}(\nu)} \|h_n\|_{L^{p'(\cdot)}(\mu)} < \infty. \end{aligned}$$

Therefore, $k(x, x) \in L^1(\mu)$, $k(x, x)$ is finite for a.e. x and

$$\sum_{n=1}^{\infty} \langle g_n, h_n \rangle = \int_{\Omega} \sum_{n=1}^{\infty} g_n(x)h_n(x) d\mu,$$

completing the proof. \square

4. NUCLEARITY ON $L^{p(\cdot)}$ OF OPERATORS ON THE TORUS

In practice the application of the concept of nuclearity requires an underlying discrete analysis. A source of problems where this situation arises in a natural way is the analysis of operators on compact Lie groups due to the discreteness of the unitary dual. More generally, a discrete Fourier analysis can be associated to a compact manifold as well as a notion of global symbol as developed in [DR14a], [DR14d].

In this section we apply the concept of r -nuclearity on variable exponent Lebesgue spaces to the study of periodic operators on \mathbb{R}^n which we can realise as operators on the torus \mathbb{T}^n , and we point out that all the results in this section have suitable extensions to the setting of compact Lie groups. Recent results on the nuclearity

on Lebesgue spaces on compact Lie groups and Grothendieck-Lidskii formulae have been obtained in [DR14c]. The trace formulas that we establish here are expressed in terms of global toroidal symbols. We first recall some notations and definitions for the Fourier analysis on the torus and the toroidal quantization. The toroidal quantization has been analysed extensively in [RT10b] and [RT09, RT10a], following the initial analysis in [RT07].

We denote the n -dimensional torus by $\mathbb{T}^n = \mathbb{R}^n/\mathbb{Z}^n$. Its unitary dual can be described as $\widehat{\mathbb{T}^n} \simeq \mathbb{Z}^n$, and the collection $\{\xi_k(x) = e^{2\pi i x \cdot k}\}_{k \in \mathbb{Z}^n}$ is an orthonormal basis of $L^2(\mathbb{T}^n)$. We will use the notation $\langle \xi \rangle := 1 + |\xi|$, where $|\cdot|$ denotes the euclidean norm.

Definition 4.1. Let us denote by $\mathcal{S}(\mathbb{Z}^n)$ the space of *rapidly decaying* functions $\phi : \mathbb{Z}^n \rightarrow \mathbb{C}$. That is, $\varphi \in \mathcal{S}(\mathbb{Z}^n)$ if for any $M > 0$ there exists a constant $C_{\varphi, M}$ such that

$$|\varphi(\xi)| \leq C_{\varphi, M} \langle \xi \rangle^{-M}$$

holds for all $\xi \in \mathbb{Z}^n$. The topology on $\mathcal{S}(\mathbb{Z}^n)$ is defined by the seminorms p_k , where $k \in \mathbb{N}_0$ and $p_k(\varphi) = \sup_{\xi \in \mathbb{Z}^n} \langle \xi \rangle^k |\varphi(\xi)|$.

In order to define the class of symbols that we will use, let us recall the definition of the Fourier transform on the torus for a function f in $C^\infty(\mathbb{T}^n)$ given by

$$(\mathcal{F}_{\mathbb{T}^n} f)(\xi) = \widehat{f}(\xi) = \int_{\mathbb{T}^n} e^{-2\pi i x \cdot \xi} f(x) dx.$$

One can prove that

$$\mathcal{F}_{\mathbb{T}^n} : C^\infty(\mathbb{T}^n) \rightarrow \mathcal{S}(\mathbb{Z}^n)$$

is a continuous bijection. The reconstruction formula of f in the form of a discrete integral or sum over the dual group \mathbb{Z}^n is the Fourier series

$$f(x) = \sum_{\xi \in \mathbb{Z}^n} e^{2\pi i x \cdot \xi} (\mathcal{F}_{\mathbb{T}^n} f)(\xi).$$

A corresponding operator is associated to a symbol $\sigma(x, \xi)$ which will be called a periodic pseudo-differential operator or the operator given by the toroidal quantization:

$$T_\sigma f(x) = \sum_{\xi \in \mathbb{Z}^n} e^{2\pi i x \cdot \xi} \sigma(x, \xi) (\mathcal{F}_{\mathbb{T}^n} f)(\xi), \quad (4.1)$$

which can also be written as

$$T_\sigma f(x) = \sum_{\xi \in \mathbb{Z}^n} \int_{\mathbb{T}^n} e^{2\pi i (x-y) \cdot \xi} \sigma(x, \xi) f(y) dy. \quad (4.2)$$

We refer to [RT10b] for an extensive analysis of such toroidal quantization.

In the rest of this section we will consider \mathbb{T}^n endowed with the Borel σ -algebra and the Lebesgue measure so that we will just write $\mathcal{P}(\mathbb{T}^n)$ to denote the corresponding class of variable exponents.

Theorem 4.2. *Let $p(\cdot) \in \mathcal{P}(\mathbb{T}^n)$ and $0 < r \leq 1$. Let $\sigma(x, \xi)$ be a symbol such that*

$$\sum_{\xi \in \mathbb{Z}^n} \|\sigma(\cdot, \xi)\|_{L^{p'(\cdot)}}^r < \infty.$$

Then T_σ is r -nuclear from $L^{p(\cdot)}$ to $L^{q(\cdot)}$ for all $q(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. If $p^+ < \infty$ and $q(\cdot) = p(\cdot)$, then T_σ is r -nuclear on $L^{p(\cdot)}(\mathbb{T}^n)$ and

$$\mathrm{Tr}(T_\sigma) = \int_{\mathbb{T}^n} \sum_{\xi \in \mathbb{Z}^n} \sigma(x, \xi) dx. \quad (4.3)$$

In particular, if additionally $r \leq \frac{2}{3}$, then

$$\mathrm{Tr}(T_\sigma) = \int_{\mathbb{T}^n} \sum_{\xi \in \mathbb{Z}^n} \sigma(x, \xi) dx = \sum_{j=1}^{\infty} \lambda_j, \quad (4.4)$$

where λ_j ($j = 1, 2, \dots$) are the eigenvalues of T_σ on $L^{p(\cdot)}(\mathbb{T}^n)$ with multiplicities taken into account.

Proof. We observe that for a pseudo-differential operator T_σ of the form (4.2) its kernel can be formally written in the form

$$k(x, y) = \sum_{\xi \in \mathbb{Z}^n} e^{2\pi i(x-y)\cdot\xi} \sigma(x, \xi).$$

We write $g_\xi(x) = e^{2\pi i x \cdot \xi} \sigma(x, \xi)$, $h_\xi(y) = e^{-2\pi i y \cdot \xi}$. Now, $\|h_\xi(\cdot)\|_{L^{q(\cdot)}} = \|1\|_{L^{q(\cdot)}} < \infty$ since the measure is finite. Hence

$$\sum_{\xi \in \mathbb{Z}^n} \|\sigma(\cdot, \xi)\|_{L^{p'(\cdot)}}^r \|h_\xi(\cdot)\|_{L^{q(\cdot)}}^r = \|h_\xi(\cdot)\|_{L^{q(\cdot)}}^r \sum_{\xi \in \mathbb{Z}^n} \|\sigma(\cdot, \xi)\|_{L^{p'(\cdot)}}^r < \infty,$$

for all $q(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. An application of Theorem 3.5 yields the r -nuclearity of T_σ from $L^{p(\cdot)}$ to $L^{q(\cdot)}$ for all $q(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. The formula (4.3) for the trace also follows from Theorem 3.5 since $g_\xi(x)h_\xi(x) = \sigma(x, \xi)$. The formula (4.4) follows from (4.3) and Grothendieck's Theorem. \square

As an application, we will consider the composition of a multiplication operator with a multiplier (an operator with symbol depending only on ξ) on the torus \mathbb{T}^n . Given a measurable function α on \mathbb{T}^n , we take the symbols $\alpha(x)$ and $\sigma(\xi)$, the corresponding multiplication is the operator denoted by αT_σ given by $\alpha T_\sigma f = \alpha \sigma(D)f$ on \mathbb{T}^n .

Corollary 4.3. *Let $p(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. Let $0 < r \leq 1$, $\alpha \in L^{p'(\cdot)}$, and let $\sigma(\xi)$ be a symbol such that*

$$\sum_{\xi \in \mathbb{Z}^n} |\sigma(\xi)|^r < \infty.$$

Then αT_σ is r -nuclear from $L^{p(\cdot)}$ to $L^{q(\cdot)}$ for all $q(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. If $p^+ < \infty$ and $q(\cdot) = p(\cdot)$, then αT_σ is r -nuclear on $L^{p(\cdot)}(\mathbb{T}^n)$ and

$$\mathrm{Tr}(\alpha T_\sigma) = \int_{\mathbb{T}^n} \alpha(x) dx \cdot \sum_{\xi \in \mathbb{Z}^n} \sigma(\xi).$$

If additionally $r \leq \frac{2}{3}$, then

$$\mathrm{Tr}(\alpha T_\sigma) = \int_{\mathbb{T}^n} \alpha(x) dx \cdot \sum_{\xi \in \mathbb{Z}^n} \sigma(\xi) = \sum_{j=1}^{\infty} \lambda_j,$$

where λ_j ($j = 1, 2, \dots$) are the eigenvalues of αT_σ with multiplicities taken into account.

Proof. Note that $\|\alpha(\cdot)\sigma(\xi)\|_{L^{p'(\cdot)}}^r = \|\alpha\|_{L^{p'(\cdot)}}^r |\sigma(\xi)|^r$, hence

$$\sum_{\xi \in \mathbb{Z}^n} \|\alpha(\cdot)\sigma(\xi)\|_{L^{p'(\cdot)}}^r = \|\alpha\|_{L^{p'(\cdot)}}^r \sum_{\xi \in \mathbb{Z}^n} |\sigma(\xi)|^r < \infty.$$

An application of Theorem 3.5 concludes the proof. \square

In particular, let us consider the symbol $\sigma(\xi) = (1 + 4\pi^2|\xi|^2)^{-\frac{\tau}{2}}$ for $\tau > 0$. The corresponding multiplication yields the operator $\alpha T_\sigma f = \alpha(I - \Delta)^{-\frac{\tau}{2}} f$ on \mathbb{T}^n . We observe that $\sum_{\xi \in \mathbb{Z}^n} (1 + 4\pi^2|\xi|^2)^{-\frac{r\tau}{2}} < \infty$ if and only if $r\tau > n$. Consequently we obtain:

Corollary 4.4. *Let $p(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. If $0 < r \leq 1$, $\alpha \in L^{p'(\cdot)}$, and $r\tau > n$, then $\alpha T_\sigma = \alpha(I - \Delta)^{-\frac{\tau}{2}}$ is r -nuclear from $L^{p(\cdot)}$ to $L^{q(\cdot)}$ for all $q(\cdot) \in \mathcal{P}(\mathbb{T}^n)$. If additionally $p^+ < \infty$ and $q(\cdot) = p(\cdot)$, then $\alpha(I - \Delta)^{-\frac{\tau}{2}}$ is r -nuclear on $L^{p(\cdot)}(\mathbb{T}^n)$ and*

$$\mathrm{Tr}(\alpha(I - \Delta)^{-\frac{\tau}{2}}) = \int_{\mathbb{T}^n} \alpha(x) dx \cdot \sum_{\xi \in \mathbb{Z}^n} (1 + 4\pi^2|\xi|^2)^{-\frac{\tau}{2}}.$$

If additionally $r \leq \frac{2}{3}$, then

$$\mathrm{Tr}(\alpha(I - \Delta)^{-\frac{\tau}{2}}) = \int_{\mathbb{T}^n} \alpha(x) dx \cdot \sum_{\xi \in \mathbb{Z}^n} (1 + 4\pi^2|\xi|^2)^{-\frac{\tau}{2}} = \sum_{j=1}^{\infty} \lambda_j,$$

where λ_j ($j = 1, 2, \dots$) are the eigenvalues of $\alpha(I - \Delta)^{-\frac{\tau}{2}}$ on $L^{p(\cdot)}(\mathbb{T}^n)$ with multiplicities taken into account.

Acknowledgements

The authors express their gratitude to the anonymous referee who has pointed out several corrections and valuable suggestions leading to a substantial improvement of the manuscript.

REFERENCES

- [ACPP05] G. Alberti, M. Csörnyei, A. Pełczyński, and D. Preiss. BV has the bounded approximation property. *J. Geom. Anal.*, 15(1):1–7, 2005.
- [CUF13] D. Cruz-Uribe and A. Fiorenza. *Variable Lebesgue Spaces: Foundations and Harmonic Analysis*. Birkhäuser, Basel, 2013.
- [CUFRW14] D. Cruz-Uribe, A. Fiorenza, M. Ruzhansky, and J. Wirth. *Variable Lebesgue Spaces and Hyperbolic Systems, Advanced Courses in Mathematics - CRM Barcelona*. Birkhäuser, Basel, 2014.
- [DF93] A. Defant and K. Floret. *Tensor norms and operator ideals*, volume 176 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1993.

- [DFS08] J. Diestel, J. H. Fourie, and J. Swart. *The metric theory of tensor products. Grothendieck's résumé revisited*. American Mathematical Society, Providence, RI, 2008.
- [DHHR11] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička. *Lebesgue and Sobolev spaces with variable exponents, Lecture Notes in Mathematics*. Springer-Verlag, Heidelberg, 2011.
- [DR14a] J. Delgado and M. Ruzhansky. Fourier multipliers, symbols and nuclearity on compact manifolds. *to appear in J. Anal. Math. arXiv:1404.6479v2*, 2014.
- [DR14b] J. Delgado and M. Ruzhansky. Kernel and symbol criteria for Schatten classes and r -nuclearity on compact manifolds. *C. R. Acad. Sci. Paris, Ser. I* 352:779–784, 2014.
- [DR14c] J. Delgado and M. Ruzhansky. L_p -nuclearity, traces, and Grothendieck-Lidskii formula on compact Lie groups. *J. Math. Pures. Appl.*, 102:153–172, 2014.
- [DR14d] J. Delgado and M. Ruzhansky. Schatten classes on compact manifolds: Kernel conditions. *J. Funct. Anal.*, 267:772–798, 2014.
- [DRT] J. Delgado, M. Ruzhansky, and N. Tokmagambetov. Schatten classes and nuclearity of boundary value problems. *J. Math. Pures. Appl.*, <http://dx.doi.org/10.1016/j.matpur.2016.10.005>.
- [DRW16a] J. Delgado, M. Ruzhansky, and B. Wang. Approximation property and nuclearity on mixed-norm L^p , modulation and Wiener Amalgam spaces. *J. Lond. Math. Soc.*, 94(2):391–408, 2016.
- [DRW16b] J. Delgado, M. Ruzhansky, and B. Wang. Grothendieck–Lidskii trace formula for mixed-norm and variable Lebesgue spaces. *J. Spectr. Theory*, 6(4):781–791, 2016.
- [Enf73] P. Enflo. A counterexample to the approximation problem in Banach spaces. *Acta Math.*, 130:309–317, 1973.
- [FJ73] T. Figiel and W. B. Johnson. The approximation property does not imply the bounded approximation property. *Proc. Amer. Math. Soc.*, 41:197–200, 1973.
- [FJP11] T. Figiel, W. B. Johnson, and A. Pełczyński. Some approximation properties of Banach spaces and Banach lattices. *Israel J. Math.*, 183:199–231, 2011.
- [Gro55] A. Grothendieck. Produits tensoriels topologiques et espaces nucléaires. *Mem. Amer. Math. Soc.*, 1955(16):140, 1955.
- [Hud76a] H. Hudzik. A generalization of Sobolev spaces. I. *Funct. Approximatio Comment. Math.*, (2):67–73, 1976.
- [Hud76b] H. Hudzik. A generalization of Sobolev spaces. II. *Funct. Approximatio Comment. Math.*, (3):77–85, 1976.
- [Hud76c] H. Hudzik. On generalized Orlicz-Sobolev spaces. II. *Funct. Approximatio Comment. Math.*, (4):37–51, 1976.
- [JS12] W. B. Johnson and A. Szankowski. Hereditary approximation property. *Ann. of Math. (2)*, 176(3):1987–2001, 2012.
- [KR91] O. Kováčik and J. Rákosník. On spaces $L^{p(x)}$ and $W^{k,p(x)}$. *Czechoslovak. Math. J.*, 41(4):592–618, 1991.
- [Lid59] V. B. Lidskiĭ. Non-selfadjoint operators with a trace. *Dokl. Akad. Nauk SSSR*, 125:485–487, 1959.
- [LLO10] Á. Lima, V. Lima, and E. Oja. Absolutely summing operators on $C[0, 1]$ as a tree space and the bounded approximation property. *J. Funct. Anal.*, 259(11):2886–2901, 2010.
- [LLO14] Á. Lima, V. Lima, and E. Oja. Absolutely summing operators on separable Lindenstrauss spaces as tree spaces and the bounded approximation property. *Banach J. Math. Anal.*, 8(1):190–210, 2014.
- [LO05] Á. Lima and E. Oja. The weak metric approximation property. *Math. Ann.*, 333(3):471–484, 2005.
- [Mus83] J. Musielak. *Orlicz Spaces and Modular Spaces*. Lectures Notes in Mathematics. Springer, Berlin, 1983.
- [Nak50] H. Nakano. *Modulated Semi-Ordered Linear Spaces*. Maruzen Co. Ltd., Tokyo, 1950.
- [Nak51] H. Nakano. *Topology and Linear Topological Spaces*. Maruzen Co. Ltd., Tokyo, 1951.

- [Olo72] R. Oloff. p -normierte Operatorenideale. *Beiträge Anal.*, (4):105–108, 1972. Tagungsbericht zur Ersten Tagung der WK Analysis (1970).
- [Orl31] W. Orlicz. Über konjugierte Exponentenfolgen. *Studia Math.*, (3):200–211, 1931.
- [Pie80] A. Pietsch. *Operator ideals*, volume 20 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1980. Translated from German by the author.
- [Pie87] A. Pietsch. *Eigenvalues and s -numbers*, volume 13 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1987.
- [Pie07] A. Pietsch. *History of Banach spaces and linear operators*. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [Por66] V. R. Portnov. Certain properties of the Orlicz space generated by the functions $m(x, \omega)$. *Dokl. Akad. Nauk SSSR.*, 170:1269–1272, 1966.
- [Rob14] D. Robert. Sur les traces d’opérateurs (de Grothendieck à Lidskii). (french) [The trace of operators (from Grothendieck to Lidskii)]. *Gaz. Math.*, 259(141):76–91, 2014.
- [RT07] M. Ruzhansky and V. Turunen. On the Fourier analysis of operators on the torus. In *Modern trends in pseudo-differential operators*, volume 172 of *Oper. Theory Adv. Appl.*, pages 87–105. Birkhäuser, Basel, 2007.
- [RT09] M. Ruzhansky and V. Turunen. On the toroidal quantization of periodic pseudo-differential operators. *Numer. Funct. Anal. Optim.*, 30(9-10):1098–1124, 2009.
- [RT10a] M. Ruzhansky and V. Turunen. *Pseudo-differential operators and symmetries. Background analysis and advanced topics*, volume 2 of *Pseudo-Differential Operators. Theory and Applications*. Birkhäuser Verlag, Basel, 2010.
- [RT10b] M. Ruzhansky and V. Turunen. Quantization of pseudo-differential operators on the torus. *J. Fourier Anal. Appl.*, 16(6):943–982, 2010.
- [RW14] M. Roginskaya and M. Wojciechowski. Bounded Approximation Property for Sobolev spaces on simply-connected planar domains. *arXiv:1401.7131*, 2014.
- [Rya02] R. A. Ryan. *Introduction to tensor products of Banach spaces*. Springer-Verlag London Ltd., London, 2002.
- [Sha79] I. I. Sharapudinov. The topology of the space $\mathcal{L}^{p(t)}([0, 1])$. *Math. Zametki*, 26(4):613–632, 655, 1979.
- [Sha83] I. I. Sharapudinov. *Approximation of functions in the metric of the space $\mathcal{L}^{p(t)}([a, b])$ and quadrature formulas*. In *Constructive Function Theory 81* (Varna, 1981), pp.189–193. Publ. House Bulgar. Acad. Sci. Sofia, 1983.
- [Sha86] I. I. Sharapudinov. The basis property of the Haar system in the space $\mathcal{L}^{p(t)}([0, 1])$ and the principle of localization in the mean. *Math. Sb.*, 130(2):275–283, 286, 1986.
- [Sha96] I. I. Sharapudinov. On the uniform boundedness in l^p ($p = p(x)$) of some families of convolution operators. *Math. Zametki*, 59(2):291–302, 320, 1996.
- [Sza81] A. Szankowski. $B(H)$ does not have the approximation property. *Acta Math.*, 147(1-2):89–108, 1981.
- [Tse61] I. V. Tsenov. Generalization of the problem of best approximation of a function in the space L^s . *Uch. Zap. Dagestan. Gos. Univ.*, (7):25–37, 1961.
- [Zhi82] V. V. Zhikov. Problems of convergence, duality, and averaging for a class of functionals of the calculus of variations. *Dokl. Akad. Nauk SSSR.*, 267(3):524–528, 1982.
- [Zhi92] V. V. Zhikov. Passage to the limit in nonlinear variational problems. *Mat. Sb.*, 183(8):47–84, 1992.
- [Zhi97] V. V. Zhikov. On some variational problems. *Russian J. Math. Phys.*, 5(1):105–116, 1997.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, 180 QUEEN’S GATE, LONDON SW7 2AZ, UNITED KINGDOM

E-mail address: j.delgado@imperial.ac.uk

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, 180 QUEEN’S GATE, LONDON SW7 2AZ, UNITED KINGDOM

E-mail address: m.ruzhansky@imperial.ac.uk