

The roles of proteases during developmental programmed cell death in plants

Rafael Andrade Buono^{1,2*}, Roman Hudecek^{1,2*}, Moritz K. Nowack^{1,2}✉

¹Department of Plant Biotechnology and Genetics, Ghent University, 9052 Ghent, Belgium

²VIB Center for Plant Systems Biology, 9052 Ghent, Belgium

* these authors contributed equally to the manuscript

✉ corresponding author: Moritz K. Nowack: moritz.nowack@vib.be

Email addresses of all contributing authors:

Rafael Andrade Buono: rabuo@psb.vib-ugent.be

Roman Hudecek: rohud@psb.vib-ugent.be

Moritz K. Nowack: moritz.nowack@vib.be

ORCID IDs of all contributing authors:

Rafael Andrade Buono: 0000-0002-6675-3836

Roman Hudecek: 0000-0002-2271-7104

Moritz K. Nowack: 0000-0001-8918-7577

Abstract

Proteases are among the key regulators of most forms of programmed cell death (PCD) in animals. Also in plants, many PCD processes have been associated with protease expression or activation. However, the functional evidence of the roles and actual modes of action of plant proteases in PCD remains surprisingly limited. In this review, we give an update on protease involvement in the context of developmentally regulated plant PCD. To illustrate the diversity of protease functions, we focus on several prominent developmental PCD processes, including xylem and tapetum maturation, suspensor elimination, endosperm degradation and seed coat formation, as well as plant senescence processes. Despite the substantial advance in the field, protease functions are still often only correlatively linked to developmental PCD, and the specific molecular roles of proteases in many developmental PCD processes remain to be elucidated.

Keywords:

plant, programmed cell death (PCD), developmental PCD (dPCD), protease, development, protein degradation

One sentence abstract

We review recent developments on the role of proteases in key developmental programmed cell death contexts, and discuss their modes of action.

Accepted Manuscript

Abbreviations

PCD	Programmed cell death
dPCD	Developmentally controlled PCD
ePCD	Environmentally triggered PCD
ER	Endoplasmic reticulum
GFP	Green fluorescent protein
PLCPs	Papain-like cysteine proteases
ROS	Reactive oxygen species
SI	Self-incompatibility or Self-incompatible
SI-PCD	Self-incompatibility induced programmed cell death
TUNEL	Terminal deoxynucleotidyl transferase dUTP nick end labeling
TE	Tracheary element cell in the xylem
VPEs	Vacuolar processing enzymes

Outline

1. Introduction
2. Proteases in xylem PCD
3. Proteases in tapetum PCD
4. Proteases in seed development-related PCD
 - 4.1 Embryonic suspensor
 - 4.2 Endosperm
 - 4.3 Maternal seed tissues
5. Proteases in senescence-induced PCD
 - 5.1 Leaf senescence
 - 5.2 Floral organ senescence
6. Conclusions

1. Introduction

Programmed cell death (PCD) designates different forms of genetically encoded, tightly controlled processes to eliminate no longer needed, damaged or harmful cells in a targeted fashion. There is a plethora of different PCD pathways, and many of them are vital for the development and fitness of animals and plants. In plants, PCD occurs in the course of regular development (dPCD), as well as a part of various stress responses to the biotic and abiotic environment (ePCD) (Daneva *et al.*, 2016; Huysmans *et al.*, 2017).

During plant development, a surprising number of – partly still putative – dPCD processes have been recognized in different stages of vegetative and reproductive development (Van Hautegeem *et al.*, 2015). A paradigm for dPCD in plants occurs during the development of the water-conducting xylem network, whose efficiency relies on the formation of hollow xylem cells that die and become functional in water transport (Escamez and Tuominen, 2014). A precise control of xylem cell death in early vascular plants was a key evolutionary prerequisite for the successful colonialization of terrestrial habitats. Another well-established example of dPCD is the degeneration of the tapetum layer in the floral anther. Several mutants have been characterized in which tapetum differentiation and degeneration is altered, which almost invariably leads to abortion of pollen development and thus male sterility (Wilson and Zhang, 2009). A form of dPCD also disposes of embryonic suspensor cells, which has been shown to be important for embryo development in some plant species (Smertenko and Bozhkov, 2014). After seedling germination, the organ size of the root cap in *Arabidopsis* (*Arabidopsis thaliana*) is controlled by a precisely timed dPCD event, which is important for optimal root growth and the patterning of the root system (Fendrych *et al.*, 2014; Xuan *et al.*, 2016). Finally, reproductive self-incompatibility (SI) to avoid inbreeding relies on the specific execution of a cell death program in incompatible self-pollen in some plant taxa (Eaves *et al.*, 2014) (see also article in this Special Issue by Wang *et al.*). These selected examples of dPCD processes in plants highlight the importance of PCD for successful plant growth and reproduction. Notably, PCD has to be tightly controlled in any context, as ectopic or untimely cellular death can be as detrimental to plant development as the delay or lack of PCD.

Despite substantial advances achieved over the last two decades in the discovery and characterization of plant PCD processes, we still lack a coherent mechanistic understanding of dPCD pathways. In many cases there seems to be a strong link between tissue differentiation and execution of PCD as the ultimate differentiation step (Figure 1). A number of transcription factors have been identified in the coordination of differentiation and dPCD preparation (Cubria-Radio and

Nowack, 2019; Van Durme and Nowack, 2016), emphasizing the importance of transcriptionally regulated dPCD-associated genes as a starting point for the generation of research hypotheses. In a large-scale meta-analysis of commonly regulated genes in diverse PCD contexts, we identified a core set of dPCD-associated genes (Olvera-Carrillo *et al.*, 2015). Among these, there is a substantial number encoding hydrolases (Table 1). Though transcriptional reporters based on some of these genes provide a convenient tool for dPCD research (Olvera-Carrillo *et al.*, 2015), there is still little insight into the mechanisms of action of most of the corresponding proteins. Despite substantial efforts, unveiling targets and precise functions of proteases remains a challenging task (Sueldo and van der Hoorn, 2017).

As described in more detail in other articles of this Special Issue, proteases (also called peptidases or proteinases) are enzymes that cleave target proteins, generally via hydrolysis-driven proteolysis. There are hundreds of protease-encoding genes in the genomes of multicellular organisms, for instance, there are more than 800 in *Arabidopsis* (van der Hoorn, 2008). This vast number of proteases can be grouped according to their catalytic residues as aspartic-, cysteine-, serine-, threonine-, glutamic-, asparagine-, or metalloproteases (Oda, 2012). In the MEROPS database (<http://merops.sanger.ac.uk/>, (Rawlings *et al.*, 2010)), proteases are grouped in families (based on amino-acid sequence similarity) and in clans (based on tertiary structure), with some groups comprising over 100 individual members.

Proteases likely originated in the earliest protein-producing organisms as a result of the need to recycle amino acids (Poręba *et al.*, 2013). In addition to this ancestral housekeeping function, proteases have adopted novel functions as posttranslational modifiers, thus becoming key components in the regulation of biological processes. Hydrolytic cleavage irreversibly alters target proteins; they can be activated or inactivated, stabilized or destabilized, their structure can be modified, or targets can be re-localized to different subcellular compartments (Schaller, 2004). Owing to this versatile regulatory capacity, proteases have been identified as regulators of a plethora of biological processes, including, but not restricted to, many steps of plant vegetative and reproductive development, as well as the orchestration of local and systemic defense responses (van der Hoorn, 2008).

Proteases have also been implicated in the regulation of many plant PCD processes. In fact, proteases are key regulators and executors of several cell death processes in animal systems, including apoptosis (caspase-mediated PCD) and pyroptosis (caspase-1 dependent proinflammatory PCD) (Fink and Cookson, 2005). In these contexts, the most prominent proteases are cysteine-

dependent, aspartate-specific proteases (caspases). The activation of initiator caspases occurs both in the extrinsic and the intrinsic apoptosis pathway and leads to the activation of executioner caspases. Thus, caspases form a proteolytic cascade that eventually leads to the direct or indirect cleavage of hundreds of target proteins (Poręba *et al.*, 2013). Although a few key caspase targets have been identified, it seems that the concerted cleavage of many proteins is decisive for the execution of caspase-mediated cell death (Julien and Wells, 2017). More recently, alternative caspase-independent pathways have been described. Necroptosis, for instance, does not require caspase activity to be initiated; on the contrary, caspase-8 is an important inhibitor of necroptosis (Fuchslocher Chico *et al.*, 2017). On the other hand, caspases have also been implicated in other, non-PCD-related cellular remodeling events (Julien and Wells, 2017).

Plant scientists started to investigate the role of proteases in cell death regulation over two decades ago. In several plant PCD systems, specific activity probes reported the presence of caspase-like activities, and, in some of these systems, chemical caspase inhibitors were shown to alter PCD progression (Bonneau *et al.*, 2008). The quest for caspase homologs in plants took a turn with the publication of whole genome sequences, which revealed that caspases are not conserved outside animals (Uren *et al.*, 2000), leading to the study of the structurally related metacaspases and paracaspases, grouped in the same family (C14) and clan (CD) of cysteine proteases. Metacaspases are subdivided into types I, II, and III, depending on their overall structure. Only type I and II metacaspases are found in plants. Type I metacaspases may contain an N-terminal extension, whereas type II metacaspases lack extra N-terminal motifs and generally have a longer linker region between the p10 and p20 motifs. The diversity of caspase homologs has been recently reviewed (Klemenčič and Funk, 2018). Although metacaspases became prime candidates in the search of functional caspase analogs in plants, it was finally shown that plant metacaspases lack aspartate specificity, and rather cleave after basic arginine or lysine residues (Vercammen *et al.*, 2004). Hence, although some metacaspases have been implicated in types of plant PCD (Minina *et al.*, 2017), they cannot be directly responsible for the observed caspase-like activities. Instead, several non-C14 family proteins have been shown to generate caspase-like activities in the plant PCD context, among them vacuolar processing enzymes (VPEs, (Hatsugai *et al.*, 2015), phytaspases (Galiullina *et al.*, 2015; Reichardt *et al.*, 2018), cathepsin B (Cai *et al.*, 2018), and subunits of the 26S-proteasome (Hatsugai *et al.*, 2009).

In this review, we will explore the roles of proteases that have been implicated in the regulation or progression of developmentally regulated PCD processes in plants. We will focus on dPCD occurring in the xylem, the tapetum, the embryonic suspensor, the endosperm and the seed

coat, and finally in senescing plant organs (Figure 2). Other PCD processes, including the Arabidopsis root cap turnover (Huysmans *et al.*, 2018; Kumpf and Nowack, 2015; Xuan *et al.*, 2016), lace plant (*Aponogeton madagascariensis*) leaf perforation (Dauphinee *et al.*, 2014; Lord *et al.*, 2013), or root aerenchyma formation (Takahashi *et al.*, 2015; Yamauchi *et al.*, 2016) will not be included, as there is still very little functional data available on the proteases involved in these processes. Although in many dPCD processes, proteases and protease activities have been found to be associated with different stages of cell death progression, evidence of the precise function and the specific mode of action of these proteases is rare. We attempted not only to summarize up-to-date information on dPCD proteases, but also to point out areas in which we still have to make further progress to understand dPCD regulation and execution. Of note, the roles of proteases in the context of plant immunity- and stress-related PCD processes will not be covered, and we refer to a number of excellent reviews focusing on these topics that have been recently published (Balakireva and Zamyatnin, 2018; Hou *et al.*, 2018; Thomas and van der Hoorn, 2018).

Accepted Manuscript

2. Proteases in xylem PCD

Long distance water transport in vascular plants is facilitated by the xylem. In angiosperms, the xylem is composed of an interconnected network of tracheary elements (TEs) and tracheids, as well as parenchyma and fiber cells. The differentiation of TEs and tracheids involves extensive secondary cell wall deposition and lignification. PCD and the ensuing *post-mortem* degradation of the protoplast, while maintaining the reinforced cell walls, is essential for the final function of the xylem as an efficient water transport system (Daneva *et al.*, 2016). The progression of PCD in TEs is marked by the release of vacuolar contents caused by tonoplast rupture, triggering the deterioration of the cytoplasm, the dismantling of the endomembrane system, and mitochondrial degradation (Groover *et al.*, 1997; Kuriyama, 1999; Yu *et al.*, 2002).

Xylem-expressed proteases have been identified and implicated in xylem differentiation and PCD in different plant systems (Böllhoner *et al.*, 2018; Hao *et al.*, 2008; Liu *et al.*, 2015; Obudulu *et al.*, 2016; Petzold *et al.*, 2012; Sueldo and van der Hoorn, 2017; Zhang *et al.*, 2015). In fact, xylem differentiation and PCD are tightly coordinated. VND6 (VASCULAR-RELATED NAC-DOMAIN 6) and VND7 are two central NAC transcription factors implicated in xylem differentiation in Arabidopsis and poplar (*Populus spec.*) (Nakaba *et al.*, 2015; Ohashi-Ito *et al.*, 2010; Yamaguchi *et al.*, 2010). Closely related NAC transcription factors have been implicated in xylem development of other species, including gymnosperms and non-vascular plants (Duval *et al.*, 2014; Laubscher *et al.*, 2018; Xu *et al.*, 2014), suggesting a high degree of evolutionary conservation of xylem differentiation. Although the ectopic overexpression of VND6 or VND7 leads to the expression of several genes believed to control differentiation and PCD, among them several proteases (Ohashi-Ito *et al.*, 2010; Valdivia *et al.*, 2013; Yamaguchi *et al.*, 2010; Yamaguchi *et al.*, 2011; Zhong *et al.*, 2010), only few xylem proteases have been functionally investigated in detail to date.

XCP1 (XYLEM CYSTEINE PROTEASE 1) and XCP2 have been identified as xylem-expressed papain-like cysteine proteases (PLCPs) that are produced in preparation for PCD, accumulate in the vacuolar lumen and are to a smaller extent associated with the endoplasmic reticulum (ER) and Golgi of Arabidopsis TEs (Avci *et al.*, 2008; Zhao *et al.*, 2000). Although simultaneous deletion of both gene functions does not lead to developmental disturbances or PCD impairment, transmission electron microscopy showed a delay in the PCD-associated cell clearance (Avci *et al.*, 2008). Consistent with this role in Arabidopsis, Tr-cp14, a cysteine protease with homology to XCP1 and XCP2, has been identified in white clover (*Trifolium repens*) and shown to accumulate in the ER prior to vacuolar collapse (Mulisch *et al.*, 2013). Although accumulation of XCPs seems to be part of xylem

differentiation, their activation seems to be confined to precise developmental stages. Despite the transcriptional accumulation of *PsgXCP1*, *PsgXCP2A*, and *PsgXCP2B* in poplar, ray parenchyma cells (radially arranged cells in woody tissues) remained alive for extended periods of time, even after secondary cell wall deposition (Nakaba *et al.*, 2015). Despite the fact that ray parenchyma cells might follow differentiation programs distinct from TEs, this finding suggests that XCP proteases have to be post-transcriptionally activated in order to affect *post-mortem* clearance.

In poplar, the activity of the proteasome has been linked to the caspase 3-like activity observed during TE differentiation and PCD (Han *et al.*, 2012). Although chemical inhibition of proteasome activity by clasto-lactacystin β -lactone in *Zinnia* (*Zinnia elegans*) and Arabidopsis xylogenetic cell cultures suppressed PCD, it was difficult to separate this effect from a *post-mortem* clearance effect or a potential disruption of early xylem differentiation (Han *et al.*, 2012; Woffenden *et al.*, 1998). Reinforcing the complexity of separating the potential pro-PCD effect of proteases from a role in early differentiation, chemical caspase inhibitors have been shown to affect the kinetics of TE differentiation in *Zinnia* cell cultures, but not to block PCD itself (Twumasi *et al.*, 2010).

Albeit not exerting caspase-like activity, metacaspases have been implicated in xylem PCD. In Arabidopsis, an *atmc9* mutant displays delayed xylem *post-mortem* corpse clearance, but PCD onset is not affected. Although *post-mortem* clearance is eventually achieved in triple *atmc9 xcp1 xcp2* mutants, activity profiling of PLCPs in whole-plant extracts showed overall reduction in PLCP activity in the triple mutants in a manner that cannot be explained by *atmc9* or *xcp1 xcp2* mutations alone (Bollhöner *et al.*, 2013), pointing to an interplay between the different proteolytic activities. AtMC9 has been localized to nucleocytoplasm, the apoplast, and the vacuolar lumen (Böllhoner *et al.*, 2018; Escamez *et al.*, 2016; Tsiatsiani *et al.*, 2013; Vercaemmen *et al.*, 2006) (Figure 3). Further supporting a *post-mortem* role, the acidic pH optimum for the cytoplasmic-localized AtMC9 might only be achieved after vacuolar collapse as a committing step toward PCD (Bollhöner *et al.*, 2013). Whereas the lowering of cytoplasmic pH might enhance the activity of the cytoplasmic pool of AtMC9 without necessarily changing its abundance, the apoplastic pool might be independently regulated by the abundance of apoplastic Serpin1 (Vercaemmen *et al.*, 2006). Serpin1 is a suicide protease inhibitor that has also been localized to the cytoplasm and shown to interact with other proteases such as the PLCP RESPONSIVE TO DISSECTATION 21 (RD21), likely after vacuolar collapse as RD21 accumulates in the vacuole (Lampl *et al.*, 2013; Lema Asqui *et al.*, 2018). In agreement with a non-essential role of MC9 in xylem PCD, the downregulation of putative MC9 poplar homologs, *PttMC13* and *PttMC14*, is not sufficient to alter xylem differentiation (Böllhoner *et al.*, 2018). However, the activity of *PttMC13* and *PttMC14* coincides with the processing of RD21, again suggesting a participation of metacaspase

activity in a proteolytic cascade after vacuolar collapse potentially aiding in corpse clearance (Böllhoner *et al.*, 2018). Similarly, downregulation of *AtMC9* in Arabidopsis xylogenic cell cultures fails to alter the timing or extent of TE-PCD, though in accordance with the *atmc9* mutant, *post-mortem* autolysis is affected. Interestingly, *AtMC9* downregulation causes an accumulation of autophagic bodies in TE-cell vacuoles, and is associated with an increased death rate of parenchymatic non-TE cells (Escamez *et al.*, 2016). Non-TE cells do not express *AtMC9* in these cell cultures and, in contrast to TE-cells, survive xylogenic induction. Concomitant downregulation of *AtMC9* and the autophagy-related gene *ATG2* in TE cells alleviates the ectopic cell death phenotype, suggesting that the level of autophagy in TE cells influences the survival rate of neighboring non-TE cells (Escamez *et al.*, 2016).

Although xylem PCD is one of the most intensively studied dPCD processes, and many proteases and protease activities have been linked to xylem PCD, our knowledge of roles and mechanistic details of plant protease functions in this process remains limited. For instance, although a general program of xylem differentiation including upregulation of NAC transcription factors with subsequent upregulation of protease genes has been observed for the different species studied, the subcellular localization and target identification of these proteases need better elucidation. Moreover, with most of the phenotypes of analyzed mutants being related with *post-mortem* degradation, the role of proteases in earlier TE PCD steps remains to be unraveled.

3. Proteases in tapetum PCD

Pollen development is essential for successful sexual reproduction in plants. Pollen formation takes place within the anthers and is dependent on the proper development of the tapetum, a layer of cells lining the anther locule (Daneva *et al.*, 2016). Pollen development and pollen exine (tough outer walls of pollen grains) formation depend on correct tapetum differentiation, which induces dPCD of tapetal cells at a specific stage of pollen development (Ariizumi and Toriyama, 2011; Plackett *et al.*, 2011). Prior to dPCD, sporopollenin components are exported from intact tapetum cells, whereas the pollen coat is deposited later when tapetum dPCD occurs and their cellular content is released in the locule cavity in which the pollen grains develop. The precise timing of differentiation and PCD is crucial, as precocious or delayed tapetum differentiation and PCD almost invariably result in male sterility (Bedinger, 1992; Kawanabe *et al.*, 2006; Lam *et al.*, 2000). Nevertheless, as differentiation and degeneration of the tapetum is tightly linked, it is difficult to distinguish if tapetum differentiation and early secretion or later dPCD-related defects affect pollen development in mutant plants. In this context, several protease activities have been associated with tapetum

development (Hierl *et al.*, 2012; Omidvar *et al.*, 2017; Phan *et al.*, 2012; Sheoran *et al.*, 2009; Shukla *et al.*, 2016; Solís *et al.*, 2014), but few of them have been functionally analyzed in detail.

Arabidopsis CYSTEINE ENDOPEPTIDASE 1 (*CEP1*), a KDEL-tailed cysteine protease from the PLCP family, has been identified as a key element of tapetum PCD (Zhang *et al.*, 2014). *CEP1* is expressed in the tapetum from stage 5 to 11 of anther development. *CEP1* is transported in vesicles as an inactive proenzyme to the vacuole, where maturation occurs, and the active form is released to the cytosol after vacuolar rupture. Knock-out of *CEP1* leads to abnormal and delayed tapetal PCD, and reduction in pollen fertility. Conversely, the overexpression of *CEP1* under the 35S promoter causes premature tapetal PCD and, as in the *cep1* mutant, a reduction in the number of viable pollen grains (Zhang *et al.*, 2014). Interestingly, transcriptome analyses of *cep1* mutant flower buds revealed the upregulation of hundreds of genes in comparison to the wild type already in early stages of anther development before tapetum PCD (Zhang *et al.*, 2014), suggesting an additional role for *CEP1* in early tapetum differentiation.

Another cysteine protease, *AtCP51*, has been identified as being expressed during anther development (Yang *et al.*, 2014). Its expression was observed in different parts of the anther with strong promoter activity detected in tapetal cells and microspores. Contrarily to *CEP1*, *AtCP51* seems to have a pro-survival function, because silencing of *AtCP51* by RNAi results in male sterility as a result of early tapetum degradation. Homologs of *AtCP51* have been found in various plant species such as *OsCP1* in rice (*Oryza sativa*) and *NtCP56* in tobacco (*Nicotiana tabacum*) (Yang *et al.*, 2014). Analysis of promoter activity suggested *OsCP1* expression in late anther development, including pollen and tapetum. The *oscp1* mutant shows overall delay in plant growth, and abnormal pollen development with reduced male fertility (Lee *et al.*, 2004). However, whether *OsCP1* controls pollen viability directly or via tapetum cell death has not been established. *OsCP1* together with *Osc6*, encoding a putative protease inhibitor, have been shown to be directly regulated by the tapetum-specific transcription factor *TDR* (*Tapetum Degeneration Retardation*). Knock-out of *TDR* results in a delayed tapetum degeneration in rice, characterized by lower *OsCP1* and *Osc6* levels (Li *et al.*, 2006), confirming the link between *OsCP1* expression and tapetal PCD.

Next to cysteine proteases, aspartic proteases have been implicated in PCD processes throughout plant development, including tapetum PCD (Gao *et al.*, 2017a; Gao *et al.*, 2017b; Ge *et al.*, 2005). Two aspartic proteases in rice, *OsAP25* and *OsAP37*, have been found to be under direct transcriptional control of ETERNAL TAPETUM1 (*EAT1*) (Niu *et al.*, 2013). *EAT1* is a tapetum-expressed basic helix-loop-helix transcription factor and *EAT1* knock-out leads to a delay in PCD, causing male

sterility. Both *OsAP25* and *OsAP37* are strongly downregulated in *eat1* mutants, and transient expression of both proteases causes ectopic cell death in *Nicotiana benthamiana* leaves, which was blocked in presence of the aspartic protease inhibitor pepstatin A. These results suggest that the protease activities of *OsAP25* and *OsAP37* are necessary and sufficient to drive cells into PCD. (Niu *et al.*, 2013)

In contrast to *OsAP25* and *OsAP37*, the Arabidopsis aspartic protease UNDEAD is a negative regulator of tapetum PCD. *UNDEAD* expression is controlled by the tapetum transcription factor AtMYB80, a key regulator of tapetum differentiation and PCD (Phan *et al.*, 2011). Promoter activity analysis showed expression in the tapetum throughout the development of anthers, but mainly in young floral stages. The presence of a putative mitochondrial targeting signal suggests that UNDEAD might be localized to, and act in, mitochondria (Figure 3). RNAi-mediated knock-down of *UNDEAD* mimics the *myb80* phenotype of premature tapetal degeneration, characterized by an early onset of chromatin degradation in tapetum cells (Phan *et al.*, 2011). These results establish UNDEAD as a pro-survival protease that fine-tunes the timing of tapetum PCD downstream of MYB80. It is tempting to speculate that UNDEAD targets and inactivates mitochondria-localized pro-PCD factors, whose premature accumulation leads to early cell death in *myb80* and *undead* mutants. Future studies with *UNDEAD* overexpression might help to elucidate its pro-survival role downstream of MYB80.

4. Proteases in seed development-related PCD

The plant seed is a complex organ consisting of both zygotic fertilization products (embryo and endosperm) as well as maternally derived tissues (seed integuments, nucellus). While the embryo develops to carry on life to a new sporophytic generation, all other seed tissues die at specific stages during seed development (Daneva *et al.*, 2016; López-Fernández and Maldonado, 2015). In the following section, we will review the role of proteases in cell death processes occurring in the different seed compartments during seed development and germination.

4.1 Proteases in embryonic suspensor PCD

The embryonic suspensor is formed after the asymmetric division of the zygote. The terminal cell gives rise to the embryo proper, the basal cell to the suspensor cells. The suspensor promotes early embryo growth and undergoes a form of dPCD when the embryo enters the maturation phase (Kawashima and Goldberg, 2010; López-Fernández and Maldonado, 2015; Smertenko and Bozhkov, 2014). Although suspensor dPCD has only been investigated and described in few species, it is clear

that correct PCD progression in suspensor cells can be critical for embryo development (Daneva *et al.*, 2016).

One of the major model systems to study suspensor dPCD is the suspensor of somatically generated embryos of Norway spruce (*Picea abies*). In this *in-vitro* system, the large spruce suspensor is easily accessible for biochemical, pharmacological, and genetic approaches (Filonova *et al.*, 2008). Caspase-like VEIDase activity has been detected specifically in embryos at the onset of suspensor PCD, and its inhibition has been found to disturb embryo development by interfering with suspensor differentiation (Bozhkov *et al.*, 2004). In this context, a type II metacaspase, *mclI-Pa*, has been identified as being expressed in suspensor cells that are committed to PCD (Suarez *et al.*, 2004). While silencing of *mclI-Pa* leads to a reduced caspase-like activity on a VEID-based substrate (Suarez *et al.*, 2004), it has been shown that *mclI-Pa*, like other metacaspases, does not possess an intrinsic caspase activity (Bozhkov *et al.*, 2005). Nonetheless, *mclI-Pa* has been described to localize to the nucleus in dying cells (Figure 3), recombinant *mclI-Pa* is able to promote the degradation of nuclei in suspensor cell extracts, and its silencing reduces the frequency of DNA degradation as seen by TUNEL staining (Bozhkov *et al.*, 2005; Suarez *et al.*, 2004). A recent RNA-sequencing-based transcriptomics analysis revealed that several dPCD-associated genes are expressed in suspensor cells of Norway spruce. Those include the transcription factors XND1 and ANAC075, as well as putative spruce homologs of MC9, CEP1, and cathepsin B-like genes (Reza *et al.*, 2018).

An interplay between *mclI-pa* and autophagy during the development of suspensor cells has been observed in Norway spruce, pointing to a regulatory role for protease activities. The downregulation of autophagy has been shown to phenocopy embryo and suspensor development defects and leads to aberrant suspensor cell differentiation with divergent cell death (Minina *et al.*, 2013). Moreover, the downregulation of *mclI-Pa* leads to reduced autophagy, potentially reducing the cytoprotective activity of the latter in the rapidly differentiating suspensor cells. Although the mechanisms for the action of *mclI-Pa* in autophagy are not yet clear, a role for this metacaspase in the inactivation of an autophagy repressor has been suggested by Minina and colleagues (2014).

In tobacco, the differentiation and final PCD of suspensor cells have been linked to the opposing activity of NtCP14, a cathepsin H-like PLCP, and the interacting cystatin protease inhibitor NtCYS (Zhao *et al.*, 2013). While upregulation of *NtCP14* is sufficient to induce embryonic cell death and abortion, both overexpression of *NtCYS* or silencing of *NtCP14* leads to a delay in the onset of suspensor PCD. However, downregulation of *NtCYS* also leads to increased caspase 1-, 3-, and 6-like activities that do not seem to be caused by NtCP14, suggesting other targets or indirect effects (Zhao

et al., 2013). NtCYS has also been shown to localize, together with NtCP14, to the ER (Zhao *et al.*, 2013), suggesting either a subcellular confinement of its action to the ER or the existence of roles that are only fulfilled after the release of ER content (Figure 3). Nine other cystatins have been identified in tobacco and their differential localizations among the cytoplasm, ER, and vacuolar lumen can be a hint at different targets and processes (Zhao *et al.*, 2014). Interestingly, the control of suspensor PCD by *NtCYS* in tobacco interspecific hybrids is controlled in a parent-of-origin dependent fashion. The levels of the maternally, but not the paternally, inherited allele are controlled to ensure the correct timing of suspensor PCD (Luo *et al.*, 2016).

In contrast to the spruce tobacco suspensors, little is known about the regulation of dPCD in the Arabidopsis 7- to 9-cell suspensor (Peng and Sun, 2018), and even less about the role of proteases in this process. *CEP1* is expressed in suspensor cells prior to PCD execution where it localizes to subdomains of the ER (Zhou *et al.*, 2016). This localization prompts the hypothesis that CEP1 is stored in these structures until triggering of PCD leads to its release, unleashing its proteolytic activity (Figure 3). However, its actual role in the dying suspensor remains uncertain, as no suspensor phenotype has been identified in the respective mutants (Zhou *et al.*, 2016). KISS OF DEATH (KOD), a PCD-inducing peptide expressed in Arabidopsis suspensor cells, has been shown to be involved in suspensor PCD. Loss-of-function *kod* mutants have a reduced proportion of suspensor cells initiating PCD. In agreement with a pro-PCD role, ectopic overexpression of *KOD* in both tobacco leaves and Arabidopsis seedlings is sufficient to induce cell death. Interestingly, the expression of *KOD* leads to increased caspase 3-like activity, and co-expression of the pan-caspase inhibitor *p35* with *KOD* causes a reduction of cell death (Blanvillain *et al.*, 2011). These data suggest that the enhancement of caspase-like activity is crucial for KOD-induced death, though the mechanism of peptide signaling in the regulation of proteolytic activity is yet poorly understood.

Further suggesting a role for proteases in suspensor development and its final PCD, caspase-like activities have also been observed in the large suspensor of runner bean (*Phaseolus coccineus*) (Lombardi *et al.*, 2007). Altogether, an enhancement of proteolytic activity during suspensor development culminating in PCD seems to be a general phenomenon affecting embryo development.

4.2 Proteases in endosperm PCD

Next to the embryo, the endosperm is the second fertilization product generated by the double fertilization process in seed plants. In contrast to the embryo, however, the endosperm does not survive past the seed stage. Nevertheless, the endosperm has important functions as hybridization barrier and as nutritive tissue for the embryo during seed development and

germination (Becraft and Gutierrez-Marcos, 2012; Lafon-Placette and Köhler, 2016). Because of its role as nutrient storage in mature cereal seeds, several cereal species have been domesticated over the past millennia and now form the most important source of carbohydrate nutrition for humankind.

Several instances of dPCD terminate the life of different endosperm tissues at specific stages of seed development (Costa *et al.*, 2004; Daneva *et al.*, 2016; López-Fernández and Maldonado, 2015). In many plant species, the endosperm adjacent to the developing embryo will disappear, providing space for the growing embryo (Sabelli and Larkins, 2009). In cereals, most of the nutrients for seedling germination are stored in the starchy endosperm, which undergoes PCD after seed filling is completed (Sabelli and Larkins, 2009). In contrast to lytic endosperm death, starchy endosperm death preserves the intact starch-filled cell corpses until seed germination. Unlike the central endosperm, the peripheral aleurone layer of the endosperm shows similarities to the embryo in that both survive desiccation during seed maturation (Becraft and Gutierrez-Marcos, 2012). In cereals, the aleurone organizes nutrient remobilization during germination by secreting lytic enzymes into the dead mass of the starchy endosperm (Domínguez and Cejudo, 2014). When this function has been fulfilled, the aleurone as last surviving endosperm tissue will in turn undergo cell death (Bethke *et al.*, 2007; Sabelli and Larkins, 2009). Though many protease genes and specific protease activities have been found to be expressed in association with endosperm degeneration (Domínguez and Cejudo, 1999; Domínguez *et al.*, 2002; Rocha *et al.*, 2013; Sreenivasulu *et al.*, 2006; Szewińska *et al.*, 2016; Tran *et al.*, 2014), our knowledge about functional aspects of these proteases is still very restricted.

During lytic endosperm breakdown in the embryo-surrounding region of Arabidopsis, the bHLH transcription factor ZHOUP1 (ZOU) appears to have a central role, as *zou* mutant endosperm persists, severely impeding embryo growth (Waters *et al.*, 2013). The action of ZOU appears to result in a weakening of endosperm cell walls, allowing the growing embryo to exert mechanical pressure on the endosperm cells (Fourquin *et al.*, 2016), causing either a passive disruption, or a mechanically triggered active PCD process in the embryo-surrounding endosperm. The expression of several proteases depends on ZOU, including the PCD-associated PUTATIVE ASPARTIC PROTEINASE A3 (PASPA3) and the subtilisin-like protease ABNORMAL LEAF SHAPE1 (ALE1) (Fourquin *et al.*, 2016; Yang *et al.*, 2008). Although the functional role of PASPA3 is still unknown, ALE1 is not involved in endosperm breakdown, but in the epidermal surface formation of the developing embryo (Yang *et al.*, 2008). Recently, putative *ZOU* orthologs have been described in other species in association with endosperm development (Dou *et al.*, 2018; Feng *et al.*, 2018; Zhang *et al.*, 2017b), but whether the role of ZOU in endosperm breakdown is conserved in all plant clades is unclear.

During starchy endosperm PCD in cereal seeds, the pattern of cell death progression varies (Becraft and Gutierrez-Marcos, 2012). In wheat, a rather stochastic cell death pattern has been observed (Young and Gallie, 1999), whereas maize (*Zea mays*) and rice endosperm cell death is initiated in the central part and progresses outward (Kobayashi *et al.*, 2013; Young *et al.*, 1997). In several cases, caspase-like activities have been detected in association with different instances of starchy endosperm cell death. For example, an increase in several caspase-like activities has been detected in the maturing endosperm of barley (*Hordeum vulgare*) and found to coincide with the expression of protease-encoding genes such as *HvVPE1* and *HvPhS1*, and with nuclear DNA degradation in the starchy endosperm (Tran *et al.*, 2014). Previously, VEIDase (caspase 6-like) activity has been detected, though preferentially in young, rapidly developing tissues of the barley caryopsis, not corresponding to the onset of DNA degradation (Borén *et al.*, 2006). Also in the starchy endosperm of rice, VEIDase activity has been detected prior to the onset of endosperm degradation (Kobayashi *et al.*, 2013). In the later stage of starchy endosperm development, several protease-encoding genes are upregulated. Endosperm-expressed proteases include metalloproteases and subunits of the 20S/26S proteasome complex. Interestingly, serine and aspartic protease transcripts that are present in pericarp tissues undergoing cell death are not expressed in the starchy endosperm (Sreenivasulu *et al.*, 2006), suggesting that the preservation of starchy endosperm corpses might depend on the lack of protease activity. Specific protease activity might be further regulated by protease inhibitors that are expressed in the endosperm, but not in the degenerating pericarp (Sreenivasulu *et al.*, 2006). Despite these expression data, to date there is still little knowledge about the role of proteases in the direct regulation of cell death in the starchy endosperm in cereals.

In cereals and other taxa, the aleurone endosperm layer is the only endosperm tissue that survives seed development. In developing seeds of wheat (*Triticum aestivum*), both the embryo and the aleurone express a cystatin protease inhibitor named WC5 (Wheat Cystatin 5) (Corre-Menguy *et al.*, 2002), possibly to protect these tissues from death-associated protease activities. Nevertheless, the aleurone finally dies during germination. Aleurone PCD is a well-established process, characterized by an accumulation of reactive oxygen species and the occurrence of internucleosomal DNA fragmentation (Domínguez and Cejudo, 2014). Before dying, the aleurone secretes large quantities of enzymes, including proteases, to mobilize the storage starch and proteins in the dead bulk of the starchy endosperm (Domínguez and Cejudo, 1999; Martinez *et al.*, 2009). Recently, the N-end rule pathway controlling targeted proteolysis has been implicated in the storage reserve mobilization process in *Arabidopsis* (Zhang *et al.*, 2018). However, proteases involved in aleurone

PCD remain little investigated. In dying barley aleurone cells, a spectrum of nuclease and protease activities has been described (Fath *et al.*, 2000). The aleurone-expressed wheat carboxypeptidase CPIII has also been found to be associated with xylem TE differentiation and PCD in wheat seedlings, suggesting a common PCD-relevant function in both tissues (Domínguez *et al.*, 2002). However, whether CPIII is functionally involved in xylem cell death and might have other roles than the mobilization of endosperm proteins during germination, remains to be investigated.

In other plant taxa, for instance, castor bean (*Ricinus communis*), the bulk of the endosperm undergoes cell death only during germination (Schmid *et al.*, 1999). In castor bean, endopeptidases with a C-terminal KDEL retention motif have been identified in specific ER-derived compartments, dubbed ricinosomes (Schmid *et al.*, 1998; Schmid *et al.*, 2001). Ricinosomes accumulate the KDEL-proteases and release them during PCD into the cytoplasm, where they become activated by cleavage of a pro-peptide (Schmid *et al.*, 1999). Similar KDEL-proteases, CEP1, CEP2, and CEP3, have also been identified in Arabidopsis as being expressed in association with dPCD (Hierl *et al.*, 2014; Zhou *et al.*, 2016) and involved in pathogen defense (Höwing *et al.*, 2017; Höwing *et al.*, 2014), though their precise mode of action remains to be elucidated. CEP2 was found to accumulate in ricinosome-like vesicles and to be activated by a pH-dependent proteolytic step (Hierl *et al.*, 2014). Similarly, in the endosperm of germinating tomato (*Solanum lycopersicum*) seeds, ricinosome-like structures and KDEL-tailed cysteine protease activities have been detected (Trobacher *et al.*, 2013).

Although there is ample evidence of protease expression and activity during the different cell death and degradation processes that occur in the endosperm, it remains challenging to define the precise mechanism of protease action in control of the cell death process.

4.3 Proteases in seed integument and nucellus PCD

The seed integuments are a multi-layered structure of maternal tissue that surrounds the developing embryo and endosperm in seed plants. During seed development and maturation, the different layers follow individual differentiation programs (Beeckman *et al.*, 2000). All cell layers eventually undergo dPCD, leading to the formation of the seed coat or testa, a tough protective layer surrounding the mature embryo (Daneva *et al.*, 2016). Further to its protective role, the seed coat stores an abundance of different proteins, including proteases, that might fulfil important functions during seed dormancy, protection, and germination (Raviv *et al.*, 2017).

Accumulation and upregulation of proteases concomitant with the development of the seed coat and PCD have been reported for different species. Several proteases are upregulated during

seed coat formation in the oil-producing tree *Jatropha curcas*, including cysteine and serine proteases, and subtilases (Rocha *et al.*, 2013; Soares *et al.*, 2017). In tomato, five *VPE* genes, named *SIVPE1* through *SIVPE5*, have been identified. Based on phylogenetic and expression analysis, *SIVPE1* and *SIVPE2* are expressed in the seed coat, similarly to δVPE in Arabidopsis. However, even with the downregulation of all five *SIVPEs* by RNAi, the seed coat is able to differentiate normally (Ariizumi *et al.*, 2011). In rapeseed (*Brassica napus*), BnCysP1, a putative cysteine protease that lacks a KDEL motif and is structurally related to the leaf senescence-associated genes BnSAG-12-1 and BnSAG-12-1, has been identified to be expressed only in the inner integument prior to cell death (Obermeier *et al.*, 2009; Wan *et al.*, 2002).

In Arabidopsis, the seed coat develops from five integument layers that differentiate and finally undergo cell death in a specific succession (Haughn and Chaudhury, 2005). The Arabidopsis genome codes for four different *VPE* genes, with δVPE being specifically expressed in the inner integument (Nakaune *et al.*, 2005). While the other three Arabidopsis *VPEs* (αVPE , βVPE , and γVPE) have been shown to localize to vacuoles, δVPE was found in aggregates between the cell wall and plasma membrane of shrinking integument cells (Figure 3). In the $\delta vpe-1$ mutant, PCD characterized by nuclear degradation and cellular breakdown is delayed. However, mature $\delta vpe-1$ seeds show a similar appearance, dormancy, and germination rate as wild-type seeds. Using a YVAD-based substrate, δVPE has been shown to have caspase 1-like activity (Nakaune *et al.*, 2005). In the Arabidopsis seed coat, loss of function of the also inner integument-accumulated PDI5 (Protein Disulfide Isomerase 5), a Cys protease inhibitor, causes premature PCD of the inner seed coat layer and loss of embryo viability (Andème Ondzighi *et al.*, 2008). The *pdi5* mutant phenotype suggests that the regulation of proteolytic activity during seed coat development is crucial for proper seed set. PDI5 has been shown to interact with the Cys proteases RD21, CP43, and CP19 in integument cells. Both RD21 and PDI5 traffic together through the endomembrane system towards the vacuole, where these accumulate. Accumulation of PDI5 has been observed in flowers, stems, leaves, siliques, and immature seeds (Andème Ondzighi *et al.*, 2008). The reduction of PDI5 levels prior to the onset of PCD might indicate that PDI5 has a pro-survival role by maintaining the suppression of proteolytic activity to avoid premature seed coat PCD.

The nucellus is a tissue of maternal origin that gives rise to and supports the early female gametophyte and developing seed, and in many species degenerates through PCD (Lu and Magnani, 2018). In aubergine (*Solanum melongena*), a putative cysteine protease similar to tobacco's CYP-8 has been identified and is expressed in nucellar cells as well as in several other tissue types that undergo PCD (Xu and Chye, 1999). In castor bean, the expression of CysEP, a cysteine protease found

in ricinosomes in senescing tissues, coincides with nucellar PCD (Greenwood *et al.*, 2005). In cardoon (*Cynara cardunculus*), aspartic proteases named cardosins accumulate in flowers. These are traditionally used for cheese manufacture in the Iberian Peninsula. The nucellus of cardoon has been shown by TUNEL staining to undergo PCD and also to accumulate cardosin B (Figueiredo *et al.*, 2006). This protease localizes to the extracellular space in the stigma and the transmitting tissue in the style during floral development (Vieira *et al.*, 2001). In quinoa (*Chenopodium quinoa*), the nucellus gives rise to a layer of nutritive tissue called perisperm. It consists of dead, starch-filled cells and takes over the role of the nutritive endosperm in some angiosperms. In the forming perisperm, caspase 6-like activity has been found when starch accumulation starts and few cells already show signs of nuclear degradation (López-Fernández and Maldonado, 2013). In the degenerating nucellus of chayote (*Sechium edule*), caspase 1-, 3-, and 6-like activities have been found to be upregulated during cell death (Lombardi *et al.*, 2007). The nucellus has also been shown to produce nitric oxide and hydrogen peroxide during its development, and exogenous applications of these compounds enhances the observed caspase-like activity and cell death (Lombardi *et al.*, 2010).

In barley, seven *VPE* genes, named *HvVPE1* through *HvVPE7*, have been found to be expressed in developing grains. *HvVPE2a* and *HvVPE2b*, as well as two phytaspases *HvPhS1* and *HvPhS2*, and the aspartic protease nucellin are expressed during preparation and initiation of PCD in the nucellus (Chen and Foolad, 1997; Radchuk *et al.*, 2011; Tran *et al.*, 2014). Interestingly, *HvVPE2a/nucellin* has been found to localize to the cell wall (Linnestad *et al.*, 1998), complicating the understanding of its role in *pre-mortem* intracellular events (Figure 3). In rice, suppression of the transcription factor *MADS29* leads to the downregulation of PCD-related genes in the nucellar projection, among them one encoding a Cys protease, causing defective nucellar PCD and abnormal seed development with shrunken seeds containing abnormal starch granules (Yin and Xue, 2012). These results suggest a role for transcriptionally regulated proteolytic activity in the developmentally controlled degradation of the nucellus. The investigation of PCD progression in many plant tissues, including the nucellus, is hampered by the low accessibility of the tissue. To circumvent this issue, a system using maize nucellus protoplasts has been developed to allow the rapid manipulation of the cellular environment. In this system, overexpression of *MADS29* leads to the upregulation of Cys proteases, as was observed in rice (Chen *et al.*, 2015), indicating that nucellar protoplasts might be promising to aid gene discovery and characterization in nucellar dPCD.

While the current studies showcase a significant and developmentally controlled proteolytic activity during seed coat and nucellus development, functional links between these activities and the occurring PCD are still limited. Further studies will be necessary to untangle the roles of the identified

proteases to understand if their activity is responsible for triggering or executing cell death, or whether these are responsible primarily for *post-mortem* corpse degradation activities.

5. Proteases in senescence-induced PCD

Senescence is an integrative final stage of plant development that can also be induced or accelerated by a variety of environmental stresses. Senescence can occur on the level of individual organs, but can also encompass the entire plant at the end of its life cycle. Of interest, leaf and flower senescence are tightly controlled and are crucial for the degradation of high-weight macromolecules and their remobilization to actively growing parts of the plant (Diaz-Mendoza *et al.*, 2016).

5.1 Leaf senescence

Leaf senescence is an intensely studied phenomenon in which a large number of genes have been implicated (Li *et al.*, 2017). Proteolysis is among the most important catabolic processes occurring during leaf senescence and is central in the recycling of nitrogen compounds. Accordingly, genes encoding proteases are among the most upregulated genes during leaf senescence (Guo *et al.*, 2004). Although some proteases, like SENESCENCE-ASSOCIATED GENE 12 (SAG12), represent important genetic markers for senescence, their actual functions often remain unknown, likely due to extensive genetic or functional redundancy (Pružinská *et al.*, 2017). Generally, it is assumed that protein breakdown in senescing leaves involves protease activities in different cellular compartments, maximizing the catabolism of proteins into transportable units (Diaz-Mendoza *et al.*, 2016; Schippers *et al.*, 2015).

Although often used interchangeably, leaf senescence and PCD do not refer to the same process. Whereas nutrient mobilization is an active cellular program depending on cellular viability, PCD marks the endpoint of the senescence process (Thomas, 2013). As such, senescence-induced cell death needs to be tightly coordinated with the catabolic senescence processes to optimize recuperation of nutrients (Kim *et al.*, 2016), with nuclei and mitochondria remaining active until the very late stages of senescence (Lim *et al.*, 2007). Levels of the phytohormone salicylic acid (SA) increase during senescence. *SAG12* expression and senescence-associated cell death are delayed in the SA signaling mutant *pad4*, indicating that SA has a role in the control of gene expression during developmental senescence, and that protease activity downstream of SA can have an effect on late senescence and cell death (Morris *et al.*, 2000).

Several protease-encoding genes have been shown to be differentially regulated during leaf senescence (Bhalerao *et al.*, 2003; Gregersen and Holm, 2007; Guo *et al.*, 2004; Kinoshita *et al.*, 1999; Parrott *et al.*, 2010; Roberts *et al.*, 2012). However, untangling proteases that act primarily on nutrient mobilization from those acting on the initiation and execution of senescence-induced PCD remains problematic. While the role of proteolysis in nutrient mobilization has been extensively reviewed (Diaz-Mendoza *et al.*, 2016; Roberts *et al.*, 2012; Schippers *et al.*, 2015), here we focus on the few instances in which proteases have been implicated in senescence-associated cell death.

One approach to identify PCD-regulating genes expressed during plant organ senescence is to identify genes that are also expressed in other, senescence-unrelated PCD processes. A meta-analysis of different PCD-associated transcriptional signatures revealed the upregulation of senescence-associated genes also during osmotic stresses, the hypersensitive response, and cell-differentiation induced PCD (Olvera-Carrillo *et al.*, 2015). Thus, especially proteases that are co-regulated during leaf senescence and in cell types that undergo differentiation-induced death might represent interesting candidates for death-promoting proteases during senescence. For instance, SERINE CARBOXYPEPTIDASE-LIKE48 (SCPL48), PASPA3, and some VPEs are upregulated in organ senescence as well as diverse cell types that undergo cell death as final differentiation step (Olvera-Carrillo *et al.*, 2015).

Additionally, several proteases linked with senescence-induced PCD have been reported in different protease groups. The cucumber (*Cucumis sativus*) metalloprotease Cs1-MMP is expressed in the final stage of cotyledon senescence just prior to DNA fragmentation (Delorme *et al.*, 2000). This late upregulation suggests a role in PCD regulation rather than in nutrient recycling. Additionally, the expression of the *At2-MMP* metalloprotease in *Arabidopsis* increases with organ age and senescence, and senescence-induced PCD is accelerated in *at2-mmp-1* mutants (Golldack *et al.*, 2002). Similarly, mutants of the mitochondrial metalloprotease *FtSH4* display early leaf senescence, higher autophagic activity and cell death, suggesting a pro-survival function for *FtSH4* and *At2-MMP* (Zhang *et al.*, 2017a). Naturally, the effects of these loss-of-function mutants might be pleiotropic and the influence on leaf viability might be only indirect.

Next to protease activity, also autophagy has been involved in nutrient remobilization during leaf senescence (Havé *et al.*, 2017). Interestingly, the autophagy-deficient *atg5* mutant has been found to display higher levels of protease activity, which was attributed to the proteasome, and cytosolic and vacuolar cysteine proteases (Havé *et al.*, 2018). Although these might act directly in

promoting cell death, it is possible that these activities are back-up proteolytic systems to compensate for the absence of canonical autophagy.

5.2 Floral organ senescence

Leaf and floral senescence share the upregulation of PCD- and senescence-associated genes (Trivellini *et al.*, 2016). In comparison with leaf senescence, floral senescence might be more dependent on developmental than on environmental stimuli. For example, the lifespan of the stigma is tightly controlled by NAC transcription factors, which induce PCD to terminate flower receptivity for pollen (Gao *et al.*, 2018). Also, pollination itself is an important trigger for floral organ senescence, often mediated by the phytohormone ethylene (Rogers, 2006, 2013).

Ageing floral organs display certain PCD hallmarks, including degradation of total protein content, DNA fragmentation, vacuolar rupture and protoplast shrinkage (Battelli *et al.*, 2011). The activity of serine-, cysteine-, and metalloproteases, and of KDEL-like and caspase-like proteases has been shown throughout floral senescence (Battelli *et al.*, 2014; Pak and Van Doorn, 2005; Tripathi *et al.*, 2009; Zhou *et al.*, 2016), but functional evidence for involvement in PCD is still limited. For instance, senescing lily (*Lilium longiflorum*) flowers develop caspase-like activities during late senescence, and YVADase, VEIDase, and DEVDase inhibitors strongly decrease this activity (Battelli *et al.*, 2011). However, none of the three inhibitors affect flower longevity or the progression of senescence (Battelli *et al.*, 2011). In another study on lily flowers, signs of PCD including decreasing protein content and DNA degradation have been described in mesophyll and epidermal cells (Mochizuki-Kawai *et al.*, 2015). During early senescence, mesophyll cells show an upregulation of the KDEL-tailed cysteine protease LoCYP and display signs of PCD. Only during later flower senescence, epidermal cells start to undergo PCD, with the simultaneous upregulation of the lily *SAG12* homolog, *LoSAG12* (Mochizuki-Kawai *et al.*, 2015).

6. Conclusion

Based on the enormous regulatory potential of targeted protein modification and degradation by proteases, it comes as no surprise that proteases are involved in the regulation of most biological processes, including PCD. A major question is, however, if there are plant proteases that are similarly central to plant PCD as caspases are to apoptosis and other animal cell death types, or whether proteolytic events merely accompany plant PCD without being central for its actual regulation.

In this review, we aim to provide an overview of existing data on proteases in the context of dPCD and cover a selection of relatively well-described PCD processes in which the functional involvement of proteases has been demonstrated (Figure 2). (Dauphinee *et al.*, 2014; Huysmans *et al.*, 2018; Kumpf and Nowack, 2015; Lord *et al.*, 2013; Takahashi *et al.*, 2015; Xuan *et al.*, 2016; Yamauchi *et al.*, 2016) But even in the intensively studied PCD model systems, most available data are still associative in that they describe the expression of protease-encoding genes or activity of proteases in connection with PCD processes. Only in a limited number of cases functional data exist that allow to draw causal relationships between proteases and/or protease activity and dPCD (Table 2).

The determination of the actual mode of action of PCD-associated proteases remains a key challenge. VPEs for instance have been implicated in tonoplast breakdown as a committing step in PCD (Hatsugai *et al.*, 2015), but how tonoplast disintegration is achieved remains unresolved. A substantial limitation might be presented by the genetic and functional redundancy of proteases – many plant proteases exist in large families and even higher-order mutants might fail to produce phenotypes that would allow an inference of the proteases' function. The development and application of synthetic protease inhibitors have the potential to overcome such redundancy. However, the limited and often overestimated selectivity can restrict the utility of chemical approaches (Poręba *et al.*, 2013). A way forward would consist in comprehensive approaches combining pharmacology and reverse genetics. Protein localization is key to protease function, especially regarding the sequestration of inactive zymogens and the re-localization of activated proteases (Figure 3). So far, only a few publications describe the dynamic subcellular localization of proteases, for instance by fusion proteins that take activation cleavage sites into account (Hierl *et al.*, 2014). Additionally, strategies to identify the protease targets in quantitative degradomics studies (Demir *et al.*, 2018; Huesgen and Overall, 2012) will be an important path to a functional understanding of the role proteases play in the control and execution of dPCD in plants.

Acknowledgements

We thank all members of the PCD lab at the VIB-UGENT Center for Plant Systems Biology for vivid discussions and critical feedback. We also thank Annick Bleys for help with preparing the manuscript. We gratefully acknowledge funding by the ERC StG PROCELLDEATH (Project 639234) to M.K.N. We apologize to the colleagues whose papers we were not able to cite due to space restrictions.

References

- Andème Ondzighi C, Christopher DA, Cho EJ, Chang S-C, Staehelin LA.** 2008. *Arabidopsis* protein disulfide isomerase-5 inhibits cysteine proteases during trafficking to vacuoles before programmed cell death of the endothelium in developing seeds. *Plant Cell* **20**, 2205-2220.
- Ariizumi T, Higuchi K, Arakaki S, Sano T, Asamizu E, Ezura H.** 2011. Genetic suppression analysis in novel vacuolar processing enzymes reveals their roles in controlling sugar accumulation in tomato fruits. *Journal of Experimental Botany* **62**, 2773-2786.
- Ariizumi T, Toriyama K.** 2011. Genetic regulation of sporopollenin synthesis and pollen exine development. *Annual Review of Plant Biology* **62**, 437-460.
- Avci U, Earl Petzold H, Ismail IO, Beers EP, Haigler CH.** 2008. Cysteine proteases XCP1 and XCP2 aid micro-autolysis within the intact central vacuole during xylogenesis in *Arabidopsis* roots. *Plant Journal* **56**, 303-315.
- Balakireva AV, Zamyatnin AA.** 2018. Indispensable role of proteases in plant innate immunity. *International Journal of Molecular Sciences* **19**, 629.
- Battelli R, Lombardi L, Picciarelli P, Lorenzi R, Frigerio L, Rogers HJ.** 2014. Expression and localisation of a senescence-associated KDEL-cysteine protease from *Lilium longiflorum* tepals. *Plant Science* **214**, 38-46.
- Battelli R, Lombardi L, Rogers HJ, Picciarelli P, Lorenzi R, Ceccarelli N.** 2011. Changes in ultrastructure, protease and caspase-like activities during flower senescence in *Lilium longiflorum*. *Plant Science* **180**, 716-725.
- Becraft PW, Gutierrez-Marcos J.** 2012. Endosperm development: dynamic processes and cellular innovations underlying sibling altruism. *Wiley Interdisciplinary Reviews: Developmental Biology* **1**, 579-593.
- Bedinger P.** 1992. The remarkable biology of pollen. *Plant Cell* **4**, 879-887.
- Beeckman T, De Rycke R, Viane R, Inzé D.** 2000. Histological study of seed coat development in *Arabidopsis thaliana*. *Journal of Plant Research* **113**, 139-148.
- Bethke PC, Libourel IGL, Aoyama N, Chung Y-Y, Still DW, Jones RL.** 2007. The *Arabidopsis* aleurone layer responds to nitric oxide, gibberellin, and abscisic acid and is sufficient and necessary for seed dormancy. *Plant Physiology* **143**, 1173-1188.
- Bhalerao R, Keskitalo J, Sterky F, Erlandsson R, Björkbacka H, Birve SJ, Karlsson J, Gardeström P, Gustafsson P, Lundeberg J, Jansson S.** 2003. Gene expression in autumn leaves. *Plant Physiology* **131**, 430-442.
- Blanvillain R, Young B, Cai Y-m, Hecht V, Varoquaux F, Delorme V, Lancelin J-M, Delseny M, Gallois P.** 2011. The *Arabidopsis* peptide kiss of death is an inducer of programmed cell death. *EMBO Journal* **30**, 1173-1183.
- Böllhoner B, Jokipii-Lukkari S, Bygdell J, Stael S, Adriasola M, Muñiz L, Van Breusegem F, Ezcurra I, Wingsle G, Tuominen H.** 2018. The function of two type II metacaspases in woody tissues of *Populus* trees. *New Phytologist* **217**, 1551-1565.
- Bollhöner B, Zhang B, Stael S, Denancé N, Overmyer K, Goffner D, Van Breusegem F, Tuominen H.** 2013. *Post mortem* function of AtMC9 in xylem vessel elements. *New Phytologist* **200**, 498-510.
- Bonneau L, Ge Y, Drury GE, Gallois P.** 2008. What happened to plant caspases? *Journal of Experimental Botany* **59**, 491-499.
- Borén M, Höglund AS, Bozhkov P, Jansson C.** 2006. Developmental regulation of a VEIDase caspase-like proteolytic activity in barley caryopsis. *Journal of Experimental Botany* **57**, 3747-3753.
- Bozhkov PV, Filonova LH, Suarez MF, Helmersson A, Smertenko AP, Zhivotovsky B, von Arnold S.** 2004. VEIDase is a principal caspase-like activity involved in plant programmed cell death and essential for embryonic pattern formation. *Cell Death and Differentiation* **11**, 175-182.
- Bozhkov PV, Suarez MF, Filonova LH, Daniel G, Zamyatnin AA, Jr., Rodriguez-Nieto S, Zhivotovsky B, Smertenko A.** 2005. Cysteine protease mCII-Pa executes programmed cell death during plant

embryogenesis. *Proceedings of the Institution of Mechanical Engineers Part A-Journal of Power and Energy* **102**, 14463-14468.

Cai Y-M, Yu J, Ge Y, Mironov A, Gallois P. 2018. Two proteases with caspase-3-like activity, cathepsin B and proteasome, antagonistically control ER-stress-induced programmed cell death in *Arabidopsis*. *New Phytologist* **218**, 1143-1155.

Chen F, Foolad MR. 1997. Molecular organization of a gene in barley which encodes a protein similar to aspartic protease and its specific expression in nucellar cells during degeneration. *Plant Molecular Biology* **35**, 821-831.

Chen J, Yi Q, Song Q, Gu Y, Zhang J, Hu Y, Liu H, Liu Y, Yu G, Huang Y. 2015. A highly efficient maize nucellus protoplast system for transient gene expression and studying programmed cell death-related processes. *Plant Cell Reports* **34**, 1239-1251.

Corre-Menguy F, Cejudo FJ, Mazubert C, Vidal J, Lelandais-Brière C, Torres G, Rode A, Hartmann C. 2002. Characterization of the expression of a wheat cystatin gene during caryopsis development. *Plant Molecular Biology* **50**, 687-698.

Costa LM, Gutiérrez-Marcos JF, Dickinson HG. 2004. More than a yolk: the short life and complex times of the plant endosperm. *Trends in Plant Science* **9**, 507-514.

Cubria-Radio M, Nowack MK. 2019. Transcriptional networks orchestrating programmed cell death during plant development. *Curr Top Dev Biol* **131**, 161-184.

Daneva A, Gao Z, Van Durme M, Nowack MK. 2016. Functions and regulation of programmed cell death in plant development. *Annual Review of Cell and Developmental Biology* **32**, 441-468.

Dauphinee AN, Warner TS, Gunawardena AHLAN. 2014. A comparison of induced and developmental cell death morphologies in lace plant (*Aponogeton madagascariensis*) leaves. *BMC Plant Biology* **14**, 389.

Delorme VGR, McCabe PF, Kim D-J, Leaver CJ. 2000. A matrix metalloproteinase gene is expressed at the boundary of senescence and programmed cell death in cucumber. *Plant Physiology* **123**, 917-927.

Demir F, Niedermaier S, Villamor JG, Huesgen PF. 2018. Quantitative proteomics in plant protease substrate identification. *New Phytologist* **218**, 936-943.

Díaz-Mendoza M, Velasco-Arroyo B, Santamaria ME, González-Melendi P, Martínez M, Díaz I. 2016. Plant senescence and proteolysis: two processes with one destiny. *Genetics and Molecular Biology* **39**, 329-338.

Domínguez F, Cejudo FJ. 1999. Patterns of starchy endosperm acidification and protease gene expression in wheat grains following germination. *Plant Physiology* **119**, 81-87.

Domínguez F, Cejudo FJ. 2014. Programmed cell death (PCD): an essential process of cereal seed development and germination. *Frontiers in Plant Science* **5**, 366.

Domínguez F, González M, Cejudo FJ. 2002. A germination-related gene encoding a serine carboxypeptidase is expressed during the differentiation of the vascular tissue in wheat grains and seedlings. *Planta* **215**, 727-734.

Dou M, Zhang Y, Yang S, Feng X. 2018. Identification of *ZHOUP1* orthologs in rice involved in endosperm development and cuticle formation. *Frontiers in Plant Science* **9**, 223.

Duval I, Lachance D, Giguère I, Bomal C, Morency M-J, Pelletier G, Boyle B, MacKay JJ, Séguin A. 2014. Large-scale screening of transcription factor–promoter interactions in spruce reveals a transcriptional network involved in vascular development. *Journal of Experimental Botany* **65**, 2319-2333.

Eaves DJ, Flores-Ortiz C, Haque T, Lin Z, Teng N, Franklin-Tong VE. 2014. Self-incompatibility in *Papaver*: advances in integrating the signalling network. *Biochemical Society Transactions* **42**, 370-376.

Escamez S, André D, Zhang B, Bollhöner B, Pesquet E, Tuominen H. 2016. METACASPASE9 modulates autophagy to confine cell death to the target cells during *Arabidopsis* vascular xylem differentiation. *Biology Open* **5**, 122-129.

- Escamez S, Tuominen H.** 2014. Programmes of cell death and autolysis in tracheary elements: when a suicidal cell arranges its own corpse removal. *Journal of Experimental Botany* **65**, 1313-1321.
- Fath A, Bethke P, Lonsdale J, Meza-Romero R, Jones R.** 2000. Programmed cell death in cereal aleurone. *Plant Molecular Biology* **44**, 255-266.
- Fendrych M, Van Hautegeem T, Van Durme M, Olvera-Carrillo Y, Huysmans M, Karimi M, Lippens S, Guérin CJ, Krebs M, Schumacher K, Nowack MK.** 2014. Programmed cell death controlled by ANAC033/SOMBRERO determines root cap organ size in *Arabidopsis*. *Current Biology* **24**, 931-940.
- Feng F, Qi W, Lv Y, Yan S, Xu L, Yang W, Yuan Y, Chen Y, Zhao H, Song R.** 2018. OPAQUE11 is a central hub of the regulatory network for maize endosperm development and nutrient metabolism. *Plant Cell* **30**, 375-396.
- Figueiredo R, Duarte P, Pereira S, Pissarra J.** 2006. The embryo sac of *Cynara cardunculus*: ultrastructure of the development and localisation of the aspartic proteinase cardosin B. *Sexual Plant Reproduction* **19**, 93-101.
- Filonova LH, Suárez MF, Bozhkov PV.** 2008. Detection of programmed cell death in plant embryos. *Methods in Molecular Biology* **427**, 173-179.
- Fink SL, Cookson BT.** 2005. Apoptosis, pyroptosis, and necrosis: mechanistic description of dead and dying eukaryotic cells. *Infection and Immunity* **73**, 1907-1916.
- Fourquin C, Beauzamy L, Chamot S, Creff A, Goodrich J, Boudaoud A, Ingram G.** 2016. Mechanical stress mediated by both endosperm softening and embryo growth underlies endosperm elimination in *Arabidopsis* seeds. *Development* **143**, 3300-3305.
- Fuchslocher Chico J, Saggau C, Adam D.** 2017. Proteolytic control of regulated necrosis. *Biochimica et Biophysica Acta - Molecular Cell Research* **1864**, 2147-2161.
- Galiullina RA, Kasperkiewicz P, Chichkova NV, Szalek A, Serebryakova MV, Poreba M, Drag M, Vartapetian AB.** 2015. Substrate specificity and possible heterologous targets of phytaspase, a plant cell death protease. *Journal of Biological Chemistry* **290**, 24806-24815.
- Gao H, Li R, Guo Y.** 2017a. *Arabidopsis* aspartic proteases A36 and A39 play roles in plant reproduction. *Plant Signaling & Behavior* **12**, e1304343.
- Gao H, Zhang Y, Wang W, Zhao K, Liu C, Bai L, Li R, Guo Y.** 2017b. Two membrane-anchored aspartic proteases contribute to pollen and ovule development. *Plant Physiology* **173**, 219-239.
- Gao Z, Daneva A, Salanek Y, Van Durme M, Huysmans M, Lin Z, De Winter F, Vanneste S, Karimi M, Van de Velde J, Vandepoele K, Van de Walle D, Dewettinck K, Lambrecht BN, Nowack MK.** 2018. KIRA1 and ORESARA1 terminate flower receptivity by promoting cell death in the stigma of *Arabidopsis*. *Nature Plants* **4**, 365-375.
- Ge X, Dietrich C, Matsuno M, Li G, Berg H, Xia Y.** 2005. An *Arabidopsis* aspartic protease functions as an anti-cell-death component in reproduction and embryogenesis. *EMBO Reports* **6**, 282-288.
- Golldack D, Popova OV, Dietz K-J.** 2002. Mutation of the matrix metalloproteinase At2-MMP inhibits growth and causes late flowering and early senescence in *Arabidopsis*. *Journal of Biological Chemistry* **277**, 5541-5547.
- Greenwood JS, Helm M, Gietl C.** 2005. Ricinosomes and endosperm transfer cell structure in programmed cell death of the nucellus during *Ricinus* seed development. *Proceedings of the Institution of Mechanical Engineers Part A-Journal of Power and Energy* **102**, 2238-2243.
- Gregersen PL, Holm PB.** 2007. Transcriptome analysis of senescence in the flag leaf of wheat (*Triticum aestivum* L.). *Plant Biotechnology Journal* **5**, 192-206.
- Groover A, DeWitt N, Heidel A, Jones A.** 1997. Programmed cell death of plant tracheary elements differentiating in vitro. *Protoplasma* **196**, 197-211.
- Guo Y, Cai Z, Gan S.** 2004. Transcriptome of *Arabidopsis* leaf senescence. *Plant, Cell & Environment* **27**, 521-549.
- Han J-J, Lin W, Oda Y, Cui K-M, Fukuda H, He X-Q.** 2012. The proteasome is responsible for caspase-3-like activity during xylem development. *Plant Journal* **72**, 129-141.

- Hao X, Qian J, Xu S, Song X, Zhu J.** 2008. Location of caspase 3-like protease in the development of sieve element and tracheary element of stem in *Cucurbita moschata*. *Journal of Integrative Plant Biology* **50**, 1499-1507.
- Hatsugai N, Iwasaki S, Tamura K, Kondo M, Fuji K, Ogasawara K, Nishimura M, Hara-Nishimura I.** 2009. A novel membrane fusion-mediated plant immunity against bacterial pathogens. *Genes & Development* **23**, 2496-2506.
- Hatsugai N, Yamada K, Goto-Yamada S, Hara-Nishimura I.** 2015. Vacuolar processing enzyme in plant programmed cell death. *Frontiers in Plant Science* **6**, 234.
- Haughn G, Chaudhury A.** 2005. Genetic analysis of seed coat development in *Arabidopsis*. *Trends in Plant Science* **10**, 472-477.
- Havé M, Balliau T, Cottyn-Boitte B, Déron D, Cuff G, Soulay F, Lornac A, Reichman P, Dissmeyer N, Avice J-C, Gallois P, Rajjou L, Zivy M, Masclaux-Daubresse C.** 2018. Increases in activity of proteasome and papain-like cysteine protease in *Arabidopsis* autophagy mutants: back-up compensatory effect or cell-death promoting effect? *Journal of Experimental Botany* **69**, 1369-1385.
- Havé M, Marmagne A, Chardon F, Masclaux-Daubresse C.** 2017. Nitrogen remobilization during leaf senescence: lessons from *Arabidopsis* to crops. *Journal of Experimental Botany* **68**, 2513-2529.
- Hierl G, Höwing T, Isono E, Lottspeich F, Gietl C.** 2014. Ex vivo processing for maturation of *Arabidopsis* KDEL-tailed cysteine endopeptidase 2 (AtCEP2) pro-enzyme and its storage in endoplasmic reticulum derived organelles. *Plant Molecular Biology* **84**, 605-620.
- Hierl G, Vothknecht U, Gietl C.** 2012. Programmed cell death in *Ricinus* and *Arabidopsis*: the function of KDEL cysteine peptidases in development. *Physiologia Plantarum* **145**, 103-113.
- Hou S, Jamieson P, He P.** 2018. The cloak, dagger, and shield: proteases in plant-pathogen interactions. *Biochemical Journal* **475**, 2491-2509.
- Höwing T, Dann M, Hoefle C, Hückelhoven R, Gietl C.** 2017. Involvement of *Arabidopsis thaliana* endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 (AtCEP1) in powdery mildew-induced and AtCPR5-controlled cell death. *PLoS ONE* **12**, e0183870.
- Höwing T, Huesmann C, Hoefle C, Nagel M-K, Isono E, Hückelhoven R, Gietl C.** 2014. Endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 of *Arabidopsis* (AtCEP1) is involved in pathogen defense. *Frontiers in Plant Science* **5**, 58.
- Huesgen PF, Overall CM.** 2012. N- and C-terminal degradomics: new approaches to reveal biological roles for plant proteases from substrate identification. *Physiologia Plantarum* **145**, 5-17.
- Huysmans M, Buono RA, Skorzinski N, Cubria Radio M, De Winter F, Parizot B, Mertens J, Karimi M, Fendrych M, Nowack MK.** 2018. NAC transcription factors ANAC087 and ANAC046 control distinct aspects of programmed cell death in the *Arabidopsis* columella and lateral root cap. *Plant Cell* **30**, 2197-2213.
- Huysmans M, Lema AS, Coll NS, Nowack MK.** 2017. Dying two deaths — programmed cell death regulation in development and disease. *Current Opinion in Plant Biology* **35**, 37-44.
- Julien O, Wells JA.** 2017. Caspases and their substrates. *Cell Death and Differentiation* **24**, 1380-1389.
- Kawanabe T, Ariizumi T, Kawai-Yamada M, Uchimiya H, Toriyama K.** 2006. Abolition of the tapetum suicide program ruins microsporogenesis. *Plant & Cell Physiology* **47**, 784-787.
- Kawashima T, Goldberg RB.** 2010. The suspensor: not just suspending the embryo. *Trends in Plant Science* **15**, 23-30.
- Kim J, Woo HR, Nam HG.** 2016. Toward systems understanding of leaf senescence: an integrated multi-omics perspective on leaf senescence research. *Molecular Plant* **9**, 813-825.
- Kinoshita T, Yamada K, Hiraiwa N, Kondo M, Nishimura M, Hara-Nishimura I.** 1999. Vacuolar processing enzyme is up-regulated in the lytic vacuoles of vegetative tissues during senescence and under various stressed conditions. *Plant Journal* **19**, 43-53.
- Klemenčič M, Funk C.** 2018. Structural and functional diversity of caspase homologues in non-metazoan organisms. *Protoplasma* **255**, 387-397.

- Kobayashi H, Ikeda TM, Nagata K.** 2013. Spatial and temporal progress of programmed cell death in the developing starchy endosperm of rice. *Planta* **237**, 1393-1400.
- Kumpf RP, Nowack MK.** 2015. The root cap: a short story of life and death. *Journal of Experimental Botany* **66**, 5651-5662.
- Kuriyama H.** 1999. Loss of tonoplast integrity programmed in tracheary element differentiation. *Plant Physiology* **121**, 763-774.
- Lafon-Placette C, Köhler C.** 2016. Endosperm-based postzygotic hybridization barriers: developmental mechanisms and evolutionary drivers. *Molecular Ecology* **25**, 2620-2629.
- Lam E, Fukuda H, Greenberg J.** 2000. *Programmed cell death in higher plants*. New York: Springer Science & Business Media.
- Lampf N, Alkan N, Davydov O, Fluhr R.** 2013. Set-point control of RD21 protease activity by AtSerp1 controls cell death in Arabidopsis. *Plant Journal* **74**, 498-510.
- Laubscher M, Brown K, Tonfack LB, Myburg AA, Mizrahi E, Hussey SG.** 2018. Temporal analysis of *Arabidopsis* genes activated by *Eucalyptus grandis* NAC transcription factors associated with xylem fibre and vessel development. *Scientific Reports* **8**, 10983.
- Lee S, Jung K-H, An GH, Chung Y-Y.** 2004. Isolation and characterization of a rice cysteine protease gene, *OSCP1*, using T-DNA gene-trap system. *Plant Molecular Biology* **54**, 755-765.
- Lema Asqui S, Vercammen D, Serrano I, Valls M, Rivas S, Van Breusegem F, Conlon FL, Dangl JL, Coll NS.** 2018. AtSERPIN1 is an inhibitor of the metacaspase AtMC1-mediated cell death and autocatalytic processing *in planta*. *New Phytologist* **218**, 1156-1166.
- Li N, Zhang DS, Liu HS, Yin CS, Li XX, Liang WQ, Yuan Z, Xu B, Chu HW, Wang J, Wen TQ, Huang H, Luo D, Ma H, Zhang DB.** 2006. The rice *tapetum degeneration retardation* gene is required for tapetum degradation and anther development. *Plant Cell* **18**, 2999-3014.
- Li Z, Zhao Y, Liu X, Jiang Z, Peng J, Jin J, Guo H, Luo J.** 2017. Construction of the leaf senescence database and functional assessment of senescence-associated genes. *Methods in Molecular Biology* **1533**, 315-333.
- Lim PO, Kim HJ, Nam HG.** 2007. Leaf senescence. *Annual Review of Plant Biology* **58**, 115-136.
- Linnestad C, Doan DNP, Brown RC, Lemmon BE, Meyer DJ, Jung R, Olsen O-A.** 1998. Nucellain, a barley homolog of the dicot vacuolar-processing protease, is localized in nucellar cell walls. *Plant Physiology* **118**, 1169-1180.
- Liu J, Hai G, Wang C, Cao S, Xu W, Jia Z, Yang C, Wang JP, Dai S, Cheng Y.** 2015. Comparative proteomic analysis of *Populus trichocarpa* early stem from primary to secondary growth. *Journal of Proteomics* **126**, 94-108.
- Lombardi L, Casani S, Ceccarelli N, Galleschi L, Picciarelli P, Lorenzi R.** 2007. Programmed cell death of the nucellus during *Sechium edule* Sw. seed development is associated with activation of caspase-like proteases. *Journal of Experimental Botany* **58**, 2949-2958.
- Lombardi L, Ceccarelli N, Picciarelli P, Sorce C, Lorenzi R.** 2010. Nitric oxide and hydrogen peroxide involvement during programmed cell death of *Sechium edule* nucellus. *Physiologia Plantarum* **140**, 89-102.
- López-Fernández MP, Maldonado S.** 2013. Programmed cell death during quinoa perisperm development. *Journal of Experimental Botany* **64**, 3313-3325.
- López-Fernández MP, Maldonado S.** 2015. Programmed cell death in seeds of angiosperms. *Journal of Integrative Plant Biology* **57**, 996-1002.
- Lord CEN, Dauphinee AN, Watts RL, Gunawardena AHLAN.** 2013. Unveiling interactions among mitochondria, caspase-like proteases, and the actin cytoskeleton during plant programmed cell death (PCD). *PLoS ONE* **8**, e57110.
- Lu J, Magnani E.** 2018. Seed tissue and nutrient partitioning, a case for the nucellus. *Plant Reproduction* **31**, 309-317.
- Luo A, Zhao P, Zhang L-Y, Sun M-X.** 2016. Initiation of programmed cell death in the suspensor is predominantly regulated maternally in a tobacco hybrid. *Scientific Reports* **6**, 29467.

- Martinez M, Cambra I, Carrillo L, Diaz-Mendoza M, Diaz I.** 2009. Characterization of the entire cystatin gene family in barley and their target cathepsin L-like cysteine-proteases, partners in the hordein mobilization during seed germination. *Plant Physiology* **151**, 1531-1545.
- Minina EA, Coll NS, Tuominen H, Bozhkov PV.** 2017. Metacaspases versus caspases in development and cell fate regulation. *Cell Death and Differentiation* **24**, 1314-1325.
- Minina EA, Filonova LH, Fukada K, Savenkov EI, Gogvadze V, Clapham D, Sanchez-Vera V, Suarez MF, Zhivotovsky B, Daniel G, Smertenko A, Bozhkov PV.** 2013. Autophagy and metacaspase determine the mode of cell death in plants. *Journal of Cell Biology* **203**, 917-927.
- Minina EA, Smertenko AP, Bozhkov PV.** 2014. Vacuolar cell death in plants: metacaspase releases the brakes on autophagy. *Autophagy* **10**, 928-929.
- Mochizuki-Kawai H, Niki T, Shibuya K, Ichimura K.** 2015. Programmed cell death progresses differentially in epidermal and mesophyll cells of lily petals. *PLoS ONE* **10**, e0143502.
- Morris K, MacKerness SA-H, Page T, John CF, Murphy AM, Carr JP, Buchanan-Wollaston V.** 2000. Salicylic acid has a role in regulating gene expression during leaf senescence. *Plant Journal* **23**, 677-685.
- Mulisch M, Asp T, Krupinska K, Hollmann J, Holm PB.** 2013. The Tr-cp 14 cysteine protease in white clover (*Trifolium repens*) is localized to the endoplasmic reticulum and is associated with programmed cell death during development of tracheary elements. *Protoplasma* **250**, 623-629.
- Nakaba S, Takata N, Yoshida M, Funada R.** 2015. Continuous expression of genes for xylem cysteine peptidases in long-lived ray parenchyma cells in *Populus*. *Plant Biotechnology* **32**, 21-29.
- Nakaune S, Yamada K, Kondo M, Kato T, Tabata S, Nishimura M, Hara-Nishimura I.** 2005. A vacuolar processing enzyme, δ VPE, is involved in seed coat formation at the early stage of seed development. *Plant Cell* **17**, 876-887.
- Niu N, Liang W, Yang X, Jin W, Wilson ZA, Hu J, Zhang D.** 2013. EAT1 promotes tapetal cell death by regulating aspartic proteases during male reproductive development in rice. *Nature Communications* **4**, 1445.
- Obermeier C, Hosseini B, Friedt W, Snowdon R.** 2009. Gene expression profiling via LongSAGE in a non-model plant species: a case study in seeds of *Brassica napus*. *BMC Genomics* **10**, 295.
- Obudulu O, Bygdell J, Sundberg B, Moritz T, Hvidsten TR, Trygg J, Wingsle G.** 2016. Quantitative proteomics reveals protein profiles underlying major transitions in aspen wood development. *BMC Genomics* **17**, 119.
- Oda K.** 2012. New families of carboxyl peptidases: serine-carboxyl peptidases and glutamic peptidases. *Journal of Biochemistry* **151**, 13-25.
- Ohashi-Ito K, Oda Y, Fukuda H.** 2010. *Arabidopsis* VASCULAR-RELATED NAC-DOMAIN6 directly regulates the genes that govern programmed cell death and secondary wall formation during xylem differentiation. *Plant Cell* **22**, 3461-3473.
- Olvera-Carrillo Y, Van Bel M, Van Hautegeem T, Fendrych M, Huysmans M, Simaskova M, van Durme M, Buscaill P, Rivas S, Coll NS, Coppens F, Maere S, Nowack MK.** 2015. A conserved core of programmed cell death indicator genes discriminates developmentally and environmentally induced programmed cell death in plants. *Plant Physiology* **169**, 2684-2699.
- Omidvar V, Mohorianu I, Dalmay T, Zheng Y, Fei Z, Pucci A, Mazzucato A, Večeřová V, Sedlářova M, Fellner M.** 2017. Transcriptional regulation of male-sterility in *7B-1* male-sterile tomato mutant. *PLoS ONE* **12**, e0170715.
- Pak C, Van Doorn WG.** 2005. Delay of *Iris* flower senescence by protease inhibitors. *New Phytologist* **165**, 473-480.
- Parrott DL, Martin JM, Fischer AM.** 2010. Analysis of barley (*Hordeum vulgare*) leaf senescence and protease gene expression: a family C1A cysteine protease is specifically induced under conditions characterized by high carbohydrate, but low to moderate nitrogen levels. *New Phytologist* **187**, 313-331.

- Peng X, Sun M-X.** 2018. The suspensor as a model system to study the mechanism of cell fate specification during early embryogenesis. *Plant Reproduction* **31**, 59-65.
- Petzold HE, Zhao M, Beers EP.** 2012. Expression and functions of proteases in vascular tissues. *Physiologia Plantarum* **145**, 121-129.
- Phan HA, Iacuone S, Li SF, Parish RW.** 2011. The MYB80 transcription factor is required for pollen development and the regulation of tapetal programmed cell death in *Arabidopsis thaliana*. *Plant Cell* **23**, 2209-2224.
- Phan HA, Li SF, Parish RW.** 2012. MYB80, a regulator of tapetal and pollen development, is functionally conserved in crops. *Plant Molecular Biology* **78**, 171-183.
- Plackett ARG, Thomas SG, Wilson ZA, Hedden P.** 2011. Gibberellin control of stamen development: a fertile field. *Trends in Plant Science* **16**, 568-578.
- Poręba M, Stróżyk A, Salvesen GS, Drąg M.** 2013. Caspase substrates and inhibitors. *Cold Spring Harbor Perspectives in Biology* **5**, a008680.
- Pružinská A, Shindo T, Niessen S, Kaschani F, Tóth R, Millar AH, van der Hoorn RAL.** 2017. Major Cys protease activities are not essential for senescence in individually darkened *Arabidopsis* leaves. *BMC Plant Biology* **17**, 4.
- Radchuk V, Weier D, Radchuk R, Weschke W, Weber H.** 2011. Development of maternal seed tissue in barley is mediated by regulated cell expansion and cell disintegration and coordinated with endosperm growth. *Journal of Experimental Botany* **62**, 1217-1227.
- Raviv B, Aghajanyan L, Granot G, Makover V, Frenkel O, Gutterman Y, Grafi G.** 2017. The dead seed coat functions as a long-term storage for active hydrolytic enzymes. *PLoS ONE* **12**, e0181102.
- Rawlings ND, Barrett AJ, Bateman A.** 2010. MEROPS: the peptidase database. *Nucleic Acids Research* **38**, D227-D233.
- Reichardt S, Repper D, Tuzhikov AI, Galiullina RA, Planas-Marquès M, Chichkova NV, Vartapetian AB, Stintzi A, Schaller A.** 2018. The tomato subtilase family includes several cell death-related proteinases with caspase specificity. *Scientific Reports* **8**, 10531.
- Reza SH, Delhomme N, Street NR, Ramachandran P, Dalman K, Nilsson O, Minina EA, Bozhkov PV.** 2018. Transcriptome analysis of embryonic domains in Norway spruce reveals potential regulators of suspensor cell death. *PLoS ONE* **13**, e0192945.
- Roberts IN, Caputo C, Criado MV, Funk C.** 2012. Senescence-associated proteases in plants. *Physiologia Plantarum* **145**, 130-139.
- Rocha AJ, Soares EL, Costa JH, Costa WLG, Soares AA, Nogueira FCS, Domont GB, Campos FAP.** 2013. Differential expression of cysteine peptidase genes in the inner integument and endosperm of developing seeds of *Jatropha curcas* L. (Euphorbiaceae). *Plant Science* **213**, 30-37.
- Rogers HJ.** 2006. Programmed cell death in floral organs: how and why do flowers die? *Annals of Botany* **97**, 309-315.
- Rogers HJ.** 2013. From models to ornamentals: how is flower senescence regulated? *Plant Molecular Biology* **82**, 563-574.
- Sabelli PA, Larkins BA.** 2009. The development of endosperm in grasses. *Plant Physiology* **149**, 14-26.
- Schaller A.** 2004. A cut above the rest: the regulatory function of plant proteases. *Planta* **220**, 183-197.
- Schippers JHM, Schmidt R, Wagstaff C, Jing H-C.** 2015. Living to die and dying to live: the survival strategy behind leaf senescence. *Plant Physiology* **169**, 914-930.
- Schmid M, Simpson D, Gietl C.** 1999. Programmed cell death in castor bean endosperm is associated with the accumulation and release of a cysteine endopeptidase from ricinosomes. *Proceedings of the Institution of Mechanical Engineers Part A-Journal of Power and Energy* **96**, 14159-14164.
- Schmid M, Simpson D, Kalousek F, Gietl C.** 1998. A cysteine endopeptidase with a C-terminal KDEL motif isolated from castor bean endosperm is a marker enzyme for the ricinosome, a putative lytic compartment. *Planta* **206**, 466-475.

- Schmid M, Simpson DJ, Sarioglu H, Lottspeich F, Gietl C.** 2001. The ricinosomes of senescing plant tissue bud from the endoplasmic reticulum. *Proceedings of the Institution of Mechanical Engineers Part A-Journal of Power and Energy* **98**, 5353-5358.
- Sheoran IS, Ross ARS, Olson DJH, Sawhney VK.** 2009. Differential expression of proteins in the wild type and *7B-1* male-sterile mutant anthers of tomato (*Solanum lycopersicum*): a proteomic analysis. *Journal of Proteomics* **71**, 624-636.
- Shukla P, Subhashini M, Singh NK, Ahmed I, Trishla S, Kirti PB.** 2016. Targeted expression of cystatin restores fertility in cysteine protease induced male sterile tobacco plants. *Plant Science* **246**, 52-61.
- Smertenko A, Bozhkov PV.** 2014. Somatic embryogenesis: life and death processes during apical-basal patterning. *Journal of Experimental Botany* **65**, 1343-1360.
- Soares EL, Lima MLB, Nascimento JRS, Soares AA, Coutinho IAC, Campos FAP.** 2017. Seed development of *Jatropha curcas* L. (Euphorbiaceae): integrating anatomical, ultrastructural and molecular studies. *Plant Cell Reports* **36**, 1707-1716.
- Solís M-T, Chakrabarti N, Corredor E, Cortés-Eslava J, Rodríguez-Serrano M, Biggiogera M, Risueño MC, Testillano PS.** 2014. Epigenetic changes accompany developmental programmed cell death in tapetum cells. *Plant & Cell Physiology* **55**, 16-29.
- Sreenivasulu N, Radchuk V, Strickert M, Miersch O, Weschke W, Wobus U.** 2006. Gene expression patterns reveal tissue-specific signaling networks controlling programmed cell death and ABA-regulated maturation in developing barley seeds. *Plant Journal* **47**, 310-327.
- Suarez MF, Filonova LH, Smertenko A, Savenkov EI, Clapham DH, von Arnold S, Zhivotovsky B, Bozhkov PV.** 2004. Metacaspase-dependent programmed cell death is essential for plant embryogenesis. *Current Biology* **14**, R339-R340.
- Sueldo DJ, van der Hoorn RAL.** 2017. Plant life needs cell death, but does plant cell death need Cys proteases? *FEBS Journal* **284**, 1577-1585.
- Szewińska J, Simińska J, Bielawski W.** 2016. The roles of cysteine proteases and phytocystatins in development and germination of cereal seeds. *Journal of Plant Physiology* **207**, 10-21.
- Takahashi H, Yamauchi T, Rajhi I, Nishizawa NK, Nakazono M.** 2015. Transcript profiles in cortical cells of maize primary root during ethylene-induced lysigenous aerenchyma formation under aerobic conditions. *Annals of Botany* **115**, 879-894.
- Thomas EL, van der Hoorn RAL.** 2018. Ten prominent host proteases in plant-pathogen interactions. *International Journal of Molecular Sciences* **19**, 639.
- Thomas H.** 2013. Senescence, ageing and death of the whole plant. *New Phytologist* **197**, 696-711.
- Tran V, Weier D, Radchuk R, Thiel J, Radchuk V.** 2014. Caspase-like activities accompany programmed cell death events in developing barley grains. *PLoS ONE* **9**, e109426.
- Tripathi SK, Singh AP, Sane AP, Nath P.** 2009. Transcriptional activation of a 37 kDa ethylene responsive cysteine protease gene, *RbCP1*, is associated with protein degradation during petal abscission in rose. *Journal of Experimental Botany* **60**, 2035-2044.
- Trivellini A, Cocetta G, Hunter DA, Vernieri P, Ferrante A.** 2016. Spatial and temporal transcriptome changes occurring during flower opening and senescence of the ephemeral hibiscus flower, *Hibiscus rosa-sinensis*. *Journal of Experimental Botany* **67**, 5919-5931.
- Trobacher CP, Senatore A, Holley C, Greenwood JS.** 2013. Induction of a ricinosomal-protease and programmed cell death in tomato endosperm by gibberellic acid. *Planta* **237**, 665-679.
- Tsiatsiani L, Timmerman E, De Bock P-J, Vercammen D, Stael S, van de Cotte B, Staes A, Goethals M, Beunens T, Van Damme P, Gevaert K, Van Breusegem F.** 2013. The *Arabidopsis* metacaspase9 degradome. *Plant Cell* **25**, 2831-2847.
- Twumasi P, Iakimova ET, Qian T, van Ieperen W, Schel JHN, Emons AMC, van Kooten O, Woltering EJ.** 2010. Caspase inhibitors affect the kinetics and dimensions of tracheary elements in xylogenic *Zinnia (Zinnia elegans)* cell cultures. *BMC Plant Biology* **10**, 162.

- Uren AG, O'Rourke K, Aravind L, Pisabarro MT, Seshagiri S, Koonin EV, Dixit VM.** 2000. Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Molecular Cell* **6**, 961-967.
- Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J.** 2013. Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot *Brachypodium distachyon*. *Journal of Experimental Botany* **64**, 1333-1343.
- van der Hoorn RAL.** 2008. Plant proteases: from phenotypes to molecular mechanisms. *Annual Review of Plant Biology* **59**, 191-223.
- Van Durme M, Nowack MK.** 2016. Mechanisms of developmentally controlled cell death in plants. *Current Opinion in Plant Biology* **29**, 29-37.
- Van Hautegeem T, Waters AJ, Goodrich J, Nowack MK.** 2015. Only in dying, life: programmed cell death during plant development. *Trends in Plant Science* **20**, 102-113.
- Vercammen D, Belenghi B, van de Cotte B, Beunens T, Gavigan J-A, De Rycke R, Brackenier A, Inzé D, Harris JL, Van Breusegem F.** 2006. Serpin1 of *Arabidopsis thaliana* is a suicide inhibitor for metacaspase 9. *Journal of Molecular Biology* **364**, 625-636.
- Vercammen D, van de Cotte B, De Jaeger G, Eeckhout D, Casteels P, Vandepoele K, Vandenberghe I, Van Beeumen J, Inzé D, Van Breusegem F.** 2004. Type II metacaspases Atmc4 and Atmc9 of *Arabidopsis thaliana* cleave substrates after arginine and lysine. *Journal of Biological Chemistry* **279**, 45329-45336.
- Vieira M, Pissarr J, Veríssimo P, Castanheira P, Costa Y, Pires E, Faro C.** 2001. Molecular cloning and characterization of cDNA encoding cardosin B, an aspartic proteinase accumulating extracellularly in the transmitting tissue of *Cynara cardunculus* L. *Plant Molecular Biology* **45**, 529-539.
- Wan L, Xia Q, Qiu X, Selvaraj G.** 2002. Early stages of seed development in *Brassica napus*: a seed coat-specific cysteine proteinase associated with programmed cell death of the inner integument. *Plant Journal* **30**, 1-10.
- Waters A, Creff A, Goodrich J, Ingram G.** 2013. "What we've got here is failure to communicate": Zou mutants and endosperm cell death in seed development. *Plant Signaling & Behavior* **8**, e24368.
- Wilson ZA, Zhang D-B.** 2009. From *Arabidopsis* to rice: pathways in pollen development. *Journal of Experimental Botany* **60**, 1479-1492.
- Woffenden BJ, Freeman TB, Beers EP.** 1998. Proteasome inhibitors prevent tracheary element differentiation in zinnia mesophyll cell cultures. *Plant Physiology* **118**, 419-430.
- Xu B, Ohtani M, Yamaguchi M, Toyooka K, Wakazaki M, Sato M, Kubo M, Nakano Y, Sano R, Hiwatashi Y, Murata T, Kurata T, Yoneda A, Kato K, Hasebe M, Demura T.** 2014. Contribution of NAC transcription factors to plant adaptation to land. *Science* **343**, 1505-1508.
- Xu F-X, Chye M-L.** 1999. Expression of cysteine proteinase during developmental events associated with programmed cell death in brinjal. *Plant Journal* **17**, 321-327.
- Xuan W, Band LR, Kumpf RP, Van Damme D, Parizot B, De Rop G, Opdenacker D, Möller BK, Skorzinski N, Njo MF, De Rybel B, Audenaert D, Nowack MK, Vanneste S, Beeckman T.** 2016. Cyclic programmed cell death stimulates hormone signaling and root development in *Arabidopsis*. *Science* **351**, 384-387.
- Yamaguchi M, Goué N, Igarashi H, Ohtani M, Nakano Y, Mortimer JC, Nishikubo N, Kubo M, Katayama Y, Kakegawa K, Dupree P, Demura T.** 2010. VASCULAR-RELATED NAC-DOMAIN6 and VASCULAR-RELATED NAC-DOMAIN7 effectively induce transdifferentiation into xylem vessel elements under control of an induction system. *Plant Physiology* **153**, 906-914.
- Yamaguchi M, Mitsuda N, Ohtani M, Ohme-Takagi M, Kato K, Demura T.** 2011. VASCULAR-RELATED NAC-DOMAIN 7 directly regulates the expression of a broad range of genes for xylem vessel formation. *Plant Journal* **66**, 579-590.
- Yamauchi T, Tanaka A, Mori H, Takamura I, Kato K, Nakazono M.** 2016. Ethylene-dependent aerenchyma formation in adventitious roots is regulated differently in rice and maize. *Plant, Cell & Environment* **39**, 2145-2157.

- Yang S, Johnston N, Talideh E, Mitchell S, Jeffree C, Goodrich J, Ingram G.** 2008. The endosperm-specific *ZHOUP1* gene of *Arabidopsis thaliana* regulates endosperm breakdown and embryonic epidermal development. *Development* **135**, 3501-3509.
- Yang Y, Dong C, Yu J, Shi L, Tong C, Li Z, Huang J, Liu S.** 2014. *Cysteine Protease 51 (CP51)*, an anther-specific cysteine protease gene, is essential for pollen exine formation in *Arabidopsis*. *Plant Cell, Tissue and Organ Culture* **119**, 383-397.
- Yin L-L, Xue H-W.** 2012. The *MADS29* transcription factor regulates the degradation of the nucellus and the nucellar projection during rice seed development. *Plant Cell* **24**, 1049-1065.
- Young TE, Gallie DR.** 1999. Analysis of programmed cell death in wheat endosperm reveals differences in endosperm development between cereals. *Plant Molecular Biology* **39**, 915-926.
- Young TE, Gallie DR, DeMason DA.** 1997. Ethylene-mediated programmed cell death during maize endosperm development of wild-type and *shrunk2* genotypes. *Plant Physiology* **115**, 737-751.
- Yu X-H, Perdue TD, Heimer YM, Jones AM.** 2002. Mitochondrial involvement in tracheary element programmed cell death. *Cell Death and Differentiation* **9**, 189-198.
- Zhang D, Liu D, Lv X, Wang Y, Xun Z, Liu Z, Li F, Lu H.** 2014. The cysteine protease CEP1, a key executor involved in tapetal programmed cell death, regulates pollen development in *Arabidopsis*. *Plant Cell* **26**, 2939-2961.
- Zhang H, Gannon L, Hassall KL, Deery MJ, Gibbs DJ, Holdsworth MJ, van der Hoorn RAL, Lilley KS, Theodoulou FL.** 2018. N-terminomics reveals control of *Arabidopsis* seed storage proteins and proteases by the Arg/N-end rule pathway. *New Phytologist* **218**, 1106-1126.
- Zhang S, Li C, Wang R, Chen Y, Shu S, Huang R, Zhang D, Li J, Xiao S, Yao N, Yang C.** 2017a. The *Arabidopsis* mitochondrial protease FtSH4 is involved in leaf senescence via regulation of WRKY-dependent salicylic acid accumulation and signaling. *Plant Physiology* **173**, 2294-2307.
- Zhang Y, Li X, Yang S, Feng X.** 2017b. Identification of *ZOUP1* orthologs in soybean potentially involved in endosperm breakdown and embryogenic development. *Frontiers in Plant Science* **8**, 139.
- Zhang Z, Xin W, Wang S, Zhang X, Dai H, Sun R, Frazier T, Zhang B, Wang Q.** 2015. Xylem sap in cotton contains proteins that contribute to environmental stress response and cell wall development. *Functional & Integrative Genomics* **15**, 17-26.
- Zhao C, Johnson BJ, Kositsup B, Beers EP.** 2000. Exploiting secondary growth in *Arabidopsis*. Construction of xylem and bark cDNA libraries and cloning of three xylem endopeptidases. *Plant Physiology* **123**, 1185-1196.
- Zhao P, Zhou X-m, Zhang L-y, Wang W, Ma L-g, Yang L-b, Peng X-b, Bozhkov PV, Sun M-x.** 2013. A bipartite molecular module controls cell death activation in the Basal cell lineage of plant embryos. *PLoS Biology* **11**, e1001655.
- Zhao P, Zhou X-m, Zou J, Wang W, Wang L, Peng X-b, Sun M-x.** 2014. Comprehensive analysis of cystatin family genes suggests their putative functions in sexual reproduction, embryogenesis, and seed formation. *Journal of Experimental Botany* **65**, 5093-5107.
- Zhong R, Lee C, Ye Z-H.** 2010. Global analysis of direct targets of secondary wall NAC master switches in *Arabidopsis*. *Molecular Plant* **3**, 1087-1103.
- Zhou L-Z, Howing T, Muller B, Hammes UZ, Gietl C, Dresselhaus T.** 2016. Expression analysis of KDEL-CysEPs programmed cell death markers during reproduction in *Arabidopsis*. *Plant Reproduction* **29**, 265-272.

Table 1: Proteases commonly upregulated during Arabidopsis dPCD (Olvera-Carrillo *et al.*, 2015).

AGI	Name	Annotation	Co-expressed with*	Upregulated in dPCD**
AT5G04200	AtMC9	Metacaspase 9	7 genes	b.c.
AT4G04460	PASPA3	Saposin-like aspartyl protease family protein	6 genes	12 conditions
AT3G45010	SCPL48	Serine carboxypeptidase-like 48	5 genes	12 conditions
AT4G35350	XCP1	Xylem cysteine peptidase 1	4 genes	b.c.
AT2G25940	α VPE	Alpha-vacuolar processing enzyme	3 genes	8 conditions
AT1G20850	XCP2	Xylem cysteine peptidase 2	3 genes	b.c.
AT1G23460	n.n.	Pectin lyase-like superfamily protein	3 genes	b.c.
AT1G63120	ATRBL2	RHOMBOID-like 2	3 genes	b.c.
AT5G50260	CEP1	Cysteine proteinases superfamily protein	2 genes	b.c.
AT1G01900	ATSBT1.1	Subtilase family protein	n.a.	8 conditions

* based on a Genevestigator co-expression analysis with other dPCD-associated genes

** dPCD conditions covered in meta-analysis

n.n. no name

n.a. not applicable

b.c. below cutoff value of 8 conditions

Accepted Manuscript

Table 2: A selection of proteases implicated in dPCD processes.

Tissue/Process	Name	Species	MEROPS clan/family/ID	Catalytic type	Localization	Mode of action	Reference
Xylem	XCP1	<i>At</i>	CA/C1/C01.065	Cysteine	Vacuole and to a lesser extend in cytoplasm	Post-mortem clearance	(Avci <i>et al.</i> , 2008)
	XCP2	<i>At</i>	CA/C1/C01.065	Cysteine	Vacuole and to a lesser extend in cytoplasm	Post-mortem clearance	(Avci <i>et al.</i> , 2008)
	Tr-cp14	<i>Tr</i>	CA/C1/C01.065	Cysteine	ER	n.d.	(Mulisch <i>et al.</i> , 2013)
	AtMC9	<i>At</i>	CD/C14/C14.034	Cysteine	Nucleocytoplasm, apoplast, and vacuole	Post-mortem clearance	(Vercammen <i>et al.</i> , 2006); (Escamez <i>et al.</i> , 2016); (Bollhöner <i>et al.</i> , 2013)
Tapetum	CEP1	<i>At</i>	CA/C1/A03	Cysteine	Vacuole	Pro-PCD	(Zhang <i>et al.</i> , 2014)
	AtCP51	<i>At</i>	CA/C1/A26	Cysteine	n.d.	Pro-survival	(Yang <i>et al.</i> , 2014)
	OsAP25	<i>Os</i>	AA/A1/UPW	Aspartic	n.d.	Pro-PCD	(Niu <i>et al.</i> , 2013)
	OsAP37	<i>Os</i>	AA/A1/UPW	Aspartic	n.d.	Pro-PCD	(Niu <i>et al.</i> , 2013)
	UNDEAD	<i>At</i>	AA/A1/A01.A49	Aspartic	Predicted mitochondrial	Pro-survival	(Phan <i>et al.</i> , 2011)
Suspensor	CEP1	<i>At</i>	CA/C1/C01.A03	Cysteine	Subdomains of ER	n.d.	(Zhou <i>et al.</i> , 2016)
	NtCP14	<i>Nt</i>	CA/C1/C01.A13	Cysteine	ER	Pro-PCD	(Zhao <i>et al.</i> , 2013)
	NtCYS	<i>Nt</i>	n.p.	inhibitor	ER	Pro-survival	(Zhao <i>et al.</i> , 2013)
	mclI-Pa	<i>Pa</i>	CD / C14 / C14.036	Cysteine	Nucleus	Promotes nuclei degradation	(Bozhkov <i>et al.</i> , 2005)
Endosperm	CEP2	<i>At</i>	CA/C1/C01.A01	Cysteine	Ricinosome-like vesicles	n.d.	(Hierl <i>et al.</i> , 2014)
Inner integument	RD21A	<i>At</i>	CA/C1/C01.064	Cysteine	Vacuole	n.d.	(Andème Ondzighi <i>et al.</i> , 2008)
	PDI5	<i>At</i>	n.p.	inhibitor	Vacuole	Pro-survival	(Andème Ondzighi <i>et al.</i> , 2008)
	δVPE	<i>At</i>	CD/C13/C13.A01	Cysteine	Cytosol, Apoplast	Pro-PCD	(Nakaune <i>et al.</i> , 2005)
Nucellus	CysEP	<i>Rc</i>	CA/C1/C01.010	Cysteine	Ricinosomes	n.d.	(Greenwood <i>et al.</i> , 2005)
	HvVPE2a	<i>Hv</i>	CD/C13/C13.001	Cysteine	Apoplast	n.d.	(Linnestad <i>et al.</i> , 1998)
Leaf senescence	At2-MMP	<i>At</i>	MA/M10/M10.012	Metallo	Predicted plasma membrane	Pro-survival	(Golldack <i>et al.</i> , 2002)
Floral senescence	FtSH4	<i>At</i>	MA/M41/M41.004	Metallo	Mitochondria	Pro-survival	(Zhang <i>et al.</i> , 2017a)

n.d. not determined; n.p. not present; *At* - *Arabidopsis thaliana*; *Tr* - *Trifolium repens*; *Os* - *Oryza sativa*; *Nt* - *Nicotiana tabacum*; *Pa* - *Picea abies*; *Rc* - *Ricinus communis*; *Hv* - *Hordeum vulgare*

Accepted Manuscript

Figure Legends

Figure 1. Hypothetical regulatory network of the plant dPCD process. Coordinated with cellular differentiation, dPCD competency is achieved at least partly on the transcriptional level. A range of PCD-associated genes are upregulated, and their proteins accumulate in cells that are preparing to undergo PCD. Once competent, a hypothetical trigger starts the PCD execution process. Finally, partial or complete corpse clearance is achieved by activity of degrading enzymes that are activated during PCD and complete corpse clearance post mortem. Conceivably, proteases might play roles in all of these steps, but so far only few protease functions in dPCD have been characterized in detail.

Figure 2. Proteases implicated in selected PCD processes occurring during plant development. PCD processes are an integral part of vegetative and reproductive plant development. They occur during xylem and tapetum differentiation, in various tissues during seed development and germination, and during plant organ senescence. We indicate proteases that have been implicated with aspects of these cell death processes in the individual tissues. For some of the PCD processes listed, no functional data for protease involvement exists to date. Abbreviations are: A, aleurone layer; E, endosperm; ESR, embryo-surrounding region of the endosperm; I, seed integuments; S, suspensor.

Figure 3. Selection of dPCD-associated proteases with known or predicted subcellular localizations. Many dPCD-associated proteases accumulate as inactive proenzymes or in specific subcellular compartments before they are activated. Especially the vacuole, but also the ER and ER-derived vesicles can serve as storage space for inactive proteases. Once PCD execution is triggered, the breakup of cellular compartments is thought to release these proteases into the same compartments as their targets. The specific chemical environment of subcellular compartments before and after compartment breakup can be an additional cue to activate PCD-associated proteases. Abbreviations are: A, apoplast; C, cell wall; Ch, chloroplast; ER, endoplasmic reticulum; M, mitochondrion; N, nucleus; PM, plasma membrane; R, rinosome; V, vacuole.

Fig. 1




Fig. 2




Fig. 3

