

Hindawi Publishing Corporation
Abstract and Applied Analysis
Volume 2012, Article ID 760918, 8 pages
doi:10.1155/2012/760918

Research Article

Periodic Solutions for Duffing Type p -Laplacian Equation with Multiple Constant Delays

Hong Zhang¹ and Junxia Meng²

¹ College of Mathematics and Computer Science, Hunan University of Arts and Science, Changde, Hunan 415000, China

² College of Mathematics, Physics and Information Engineering, Jiaxing University, Jiaxing, Zhejiang 314001, China

Correspondence should be addressed to Junxia Meng, mengjunxia1968@yahoo.com.cn

Received 19 November 2011; Accepted 17 January 2012

Academic Editor: Elena Braverman

Copyright © 2012 H. Zhang and J. Meng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Using inequality techniques and coincidence degree theory, new results are provided concerning the existence and uniqueness of periodic solutions for the Duffing type p -Laplacian equation with multiple constant delays of the form $(\varphi_p(x'(t)))' + Cx'(t) + g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) = e(t)$. Moreover, an example is provided to illustrate the effectiveness of the results in this paper.

1. Introduction

Referring to the work of Esmailzadeh and Nakhaie-Jazar [1], Duffing type equation is the simplest case of a vibrating system with nonlinear restoring force generator element. This is equivalent to a mechanical vibrating system with either a hard or soft spring. Thus, this equation and its modifications have been extensively and intensively studied. In particular, the existence of periodic solutions for Duffing type equations with and without delays have been discussed by various researchers (see, e.g., [2–8] and the references given therein). However, to the best of our knowledge, the existence and uniqueness of periodic solutions of Duffing type p -Laplacian equation whose delays more than two have not been sufficiently researched. Motivated by this, we shall consider the Duffing type p -Laplacian equations with multiple constant delays of the form

$$(\varphi_p(x'(t)))' + Cx'(t) + g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) = e(t), \quad (1.1)$$

where $p > 1$ and $\varphi_p : R \rightarrow R$ is given by $\varphi_p(s) = |s|^{p-2}s$ for $s \neq 0$ and $\varphi_p(0) = 0$, C and τ_k are constants, $e : R \rightarrow R$ and $g_0, g_k : R \times R \rightarrow R$ are continuous functions, τ_k and e are T -periodic, g_0 and g_k are T -periodic in the first argument, $T > 0$ and $k = 1, 2, \dots, n$. The main purpose of this paper is to establish sufficient conditions for the existence and uniqueness of T -periodic solutions of (1.1). The results of this paper are new and complement previously known results. Moreover, we give an example to illustrate the results.

2. Preliminary Results

Throughout this paper, we will denote

$$C_T^1 := \left\{ x \in C^1(R) : x \text{ is } T\text{-periodic} \right\},$$

$$|x|_k = \left(\int_0^T |x(t)|^k dt \right)^{1/k} \quad (k > 0), \quad |x|_\infty = \max_{t \in [0, T]} |x(t)|. \quad (2.1)$$

For the periodic boundary value problem

$$(\varphi_p(x'(t)))' = \tilde{f}(t, x, x'), \quad x(0) = x(T), \quad x'(0) = x'(T), \quad (2.2)$$

where $\tilde{f} \in C(R^3, R)$ is T -periodic in the first variable, we have the following lemma.

Lemma 2.1 (see [9]). *Let Ω be an open bounded set in C_T^1 , if the following conditions hold.*

(i) *For each $\lambda \in (0, 1)$ the problem*

$$(\varphi_p(x'(t)))' = \lambda \tilde{f}(t, x, x'), \quad x(0) = x(T), \quad x'(0) = x'(T), \quad (2.3)$$

has no solution on $\partial\Omega$.

(ii) *The equation*

$$F(a) := \frac{1}{T} \int_0^T \tilde{f}(t, a, 0) dt = 0 \quad (2.4)$$

has no solution on $\partial\Omega \cap R$.

(iii) *The Brouwer degree of F*

$$\deg(F, \Omega \cap R, 0) \neq 0. \quad (2.5)$$

Then, the periodic boundary value problem (2.2) has at least one T -periodic solution on $\overline{\Omega}$.

We can easily obtain the homotopic equation of (1.1) as follows:

$$(\varphi_p(x'(t)))' + \lambda C x'(t) + \lambda \left[g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) \right] = \lambda e(t), \quad \lambda \in (0, 1). \quad (2.6)$$

Lemma 2.2. Assume that the following conditions are satisfied.

(A₁) There exists a constant $d > 0$ such that

- (1) $\sum_{k=0}^n g_k(t, x_k) - e(t) < 0$ for $x_k > d$, $t \in R$, $k = 0, 1, 2, \dots, n$,
- (2) $\sum_{k=0}^n g_k(t, x_k) - e(t) > 0$ for $x_k < -d$, $t \in R$, $k = 0, 1, 2, \dots, n$.

Moreover, if $x(t)$ is a T -periodic solution of (2.6), then

$$|x|_\infty \leq d + \frac{1}{2}\sqrt{T}|x'|_2. \quad (2.7)$$

Proof. Let $x(t)$ be a T -periodic solution of (2.6). Then, integrating (2.6) over $[0, T]$, we have

$$\int_0^T \left[g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) - e(t) \right] dt = 0. \quad (2.8)$$

Using the integral mean-value theorem, it follows that there exists $t_1 \in [0, T]$ such that

$$g_0(t_1, x(t_1)) + \sum_{k=1}^n g_k(t_1, x(t_1 - \tau_k)) - e(t_1) = 0. \quad (2.9)$$

We now prove that there exists a constant $t_2 \in R$ such that

$$|x(t_2)| \leq d. \quad (2.10)$$

Indeed, suppose otherwise. Then,

$$|x(t)| > d \quad \forall t \in R. \quad (2.11)$$

Let $\tau_0 = 0$. From (A₁), (2.9), and (2.11), we see that there exist $0 \leq i, j \leq n$ such that

$$x(t_1 - \tau_i) = \max_{0 \leq k \leq n} x(t_1 - \tau_k) \geq \min_{0 \leq k \leq n} x(t_1 - \tau_k) = x(t_1 - \tau_j), \quad (2.12)$$

which, together with (2.11), implies

$$-d > x(t_1 - \tau_i) = \max_{0 \leq k \leq n} x(t_1 - \tau_k) \quad \text{or} \quad x(t_1 - \tau_j) = \min_{0 \leq k \leq n} x(t_1 - \tau_k) > d. \quad (2.13)$$

Without loss of generality, we may assume that $x(t_1 - \tau_j) > d$ (the situation is analogous for $-d > x(t_1 - \tau_i)$). Then, we have

$$x(t_1 - \tau_i) \geq x(t_1 - \tau_k) \geq x(t_1 - \tau_j) > d, \quad k = 0, 1, 2, \dots, n. \quad (2.14)$$

According to (2.14) and (A_1) , we obtain

$$0 > \sum_{k=0}^n g_k(t_1, x(t_1 - \tau_k)) - e(t_1), \quad (2.15)$$

this contradicts the fact (2.9); thus, (2.10) is true.

Let $t_2 = mT + t_0$ where $t_0 \in [0, T]$ and m is an integer. Then, by the same approach used in the proof of inequality (3.3) of [7], we have

$$|x|_\infty = \max_{t \in [t_0, t_0+T]} |x(t)| \leq \max_{t \in [t_0, t_0+T]} \left\{ d + \frac{1}{2} \left(\int_{t_0}^t |x'(s)| ds + \int_{t-T}^{t_0} |x'(s)| ds \right) \right\} \leq d + \frac{1}{2} \sqrt{T} |x'|_2. \quad (2.16)$$

This completes the proof of Lemma 2.2. \square

Lemma 2.3. *Let (A_1) holds. Assume that the following condition is satisfied.*

(A_2) *There exist nonnegative constants $\bar{b}_0, b_0, b_1, b_2, \dots, b_n$ such that*

$$\begin{aligned} \bar{b}_0 |x_1 - x_2|^2 &\leq -(g_0(t, x_1) - g_0(t, x_2))(x_1 - x_2), \\ \bar{b}_0 > b_1 + b_2 + \dots + b_n, \quad |g_k(t, x_1) - g_k(t, x_2)| &\leq b_k |x_1 - x_2|, \end{aligned} \quad (2.17)$$

for all $t, x_1, x_2 \in \mathbb{R}$, $k = 0, 1, 2, \dots, n$.

Then, (1.1) has at most one T -periodic solution.

Proof. Suppose that $x_1(t)$ and $x_2(t)$ are two T -periodic solutions of (1.1). Set $Z(t) = x_1(t) - x_2(t)$. Then, we obtain

$$\begin{aligned} &(\varphi_p(x'_1(t)) - \varphi_p(x'_2(t)))' + C(x'_1(t) - x'_2(t)) + [g_0(t, x_1(t)) - g_0(t, x_2(t))] \\ &+ \sum_{k=1}^n [g_k(t, x_1(t - \tau_k)) - g_k(t, x_2(t - \tau_k))] = 0. \end{aligned} \quad (2.18)$$

Multiplying $Z(t)$ and (2.18) and then integrating it from 0 to T , from (A_2) and Schwarz inequality, we get

$$\begin{aligned} \bar{b}_0 |Z|_2^2 &= \bar{b}_0 \int_0^T |Z(t)|^2 dt \\ &\leq - \int_0^T (x_1(t) - x_2(t)) [g_0(t, x_1(t)) - g_0(t, x_2(t))] dt \\ &= - \int_0^T (\varphi_p(x'_1(t)) - \varphi_p(x'_2(t))) (x'_1(t) - x'_2(t)) dt \\ &\quad + \sum_{k=1}^n \int_0^T [g_k(t, x_1(t - \tau_k)) - g_k(t, x_2(t - \tau_k))] Z(t) dt \end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{k=1}^n b_k \int_0^T |x_1(t - \tau_k) - x_2(t - \tau_k)| |Z(t)| dt \\
 &\leq \sum_{k=1}^n b_k \left(\int_0^T |x_1(t - \tau_k) - x_2(t - \tau_k)|^2 dt \right)^{1/2} |Z|_2 \\
 &= \sum_{k=1}^n b_k |Z|_2^2.
 \end{aligned} \tag{2.19}$$

Since $\bar{b}_0 > b_1 + b_2 + \dots + b_n$, we have

$$Z(t) \equiv 0 \quad \forall t \in R. \tag{2.20}$$

Thus, $x_1(t) \equiv x_2(t)$ for all $t \in R$. Therefore, (1.1) has at most one T -periodic solution. The proof of Lemma 2.3 is now complete. \square

3. Main Results

Theorem 3.1. *Let (A_1) and (A_2) hold. Then, (1.1) has a unique T -periodic solution in C_T^1 .*

Proof. By Lemma 2.3, it is easy to see that (1.1) has at most one T -periodic solution in C_T^1 . Thus, in order to prove Theorem 3.1, it suffices to show that (1.1) has at least one T -periodic solution in C_T^1 . To do this, we are going to apply Lemma 2.1. Firstly, we claim that the set of all possible T -periodic solutions of (2.6) in C_T^1 is bounded.

Let $x(t) \in C_T^1$ be a T -periodic solution of (2.6). Multiplying $x(t)$ and (2.6) and then integrating it from 0 to T , we have

$$-\int_0^T \varphi_p(x'(t))x'(t)dt + \lambda \int_0^T x(t) \left[g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) - e(t) \right] dt = 0. \tag{3.1}$$

Since $x(0) = x(T)$, then there exists $t_0 \in [0, T]$ such that $x'(t_0) = 0$. And since $\varphi_p(0) = 0$, we have

$$|\varphi_p(x'(t))| = \left| \int_{t_0}^t (\varphi_p(x'(s)))' ds \right| \leq \lambda \int_{t_0}^{t_0+T} \left| g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t - \tau_k)) - e(t) \right| dt, \tag{3.2}$$

where $t \in [t_0, t_0 + T]$.

In view of (3.1), (A_2) , and Schwarz inequality, we get

$$\begin{aligned}
 \bar{b}_0 |x|_2^2 &= b_0 \int_0^T |x(t)|^2 dt \\
 &\leq - \int_0^T (x(t) - 0)(g_0(t, x(t)) - g_0(t, 0)) dt
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{\lambda} \int_0^T \varphi_p(x'(t))x'(t)dt + \sum_{k=1}^n \int_0^T [g_k(t, x(t-\tau_k)) - g_k(t, 0)]x(t)dt \\
&\quad + \sum_{k=0}^n \int_0^T g_k(t, 0)x(t)dt - \int_0^T x(t)e(t)dt \\
&\leq \sum_{k=1}^n b_k \int_0^T |x(t-\tau_k)||x(t)|dt + \sum_{k=0}^n \int_0^T |g_k(t, 0)||x(t)|dt + \sqrt{T}|e|_\infty|x|_2 \\
&\leq \sum_{k=1}^n b_k \left(\int_0^T |x(t-\tau_k)|^2 dt \right)^{1/2} |x|_2 + \sqrt{T} \sum_{k=0}^n |g_k(t, 0)|_\infty |x|_2 + \sqrt{T}|e|_\infty|x|_2 \\
&= \sum_{k=1}^n b_k |x|_2^2 + \sqrt{T} \sum_{k=0}^n |g_k(t, 0)|_\infty |x|_2 + \sqrt{T}|e|_\infty|x|_2.
\end{aligned} \tag{3.3}$$

It follows that

$$|x|_2 \leq \frac{\sqrt{T} \sum_{k=0}^n |g_k(t, 0)|_\infty + \sqrt{T}|e|_\infty}{b_0 - \sum_{k=1}^n b_k} := \theta. \tag{3.4}$$

Again from (A₂) and Schwarz inequality, (3.2) and (3.4) yield

$$\begin{aligned}
|x'|_\infty^{p-1} &= \max_{t \in [t_0, t_0+T]} \{ |\varphi_p(x'(t))| \} = \max_{t \in [t_0, t_0+T]} \left\{ \left| \int_{t_0}^t (\varphi_p(x'(s)))' ds \right| \right\} \\
&\leq \int_{t_0}^{t_0+T} \left| g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t-\tau_k)) - e(t) \right| dt \\
&= \int_0^T \left| g_0(t, x(t)) + \sum_{k=1}^n g_k(t, x(t-\tau_k)) - e(t) \right| dt \\
&\leq \int_0^T |g_0(t, x(t)) - g_0(t, 0)| dt + \sum_{k=1}^n \int_0^T |g_k(t, x(t-\tau_k)) - g_k(t, 0)| dt \\
&\quad + \sum_{k=0}^n \int_0^T |g_k(t, 0)| dt + T|e|_\infty \\
&\leq b_0 \int_0^T |x(t)| dt + \sum_{k=1}^n \int_0^T b_k |x(t-\tau_k)| dt + \sum_{k=0}^n T |g_k(t, 0)|_\infty + T|e|_\infty \\
&\leq b_0 \sqrt{T}|x|_2 + \sum_{k=1}^n b_k \sqrt{T}|x|_2 + \sum_{k=0}^n T |g_k(t, 0)|_\infty + T|e|_\infty
\end{aligned}$$

$$\begin{aligned}
&\leq b_0\sqrt{T}\theta + \sum_{k=1}^n b_k\sqrt{T}\theta + \sum_{k=0}^n T|g_k(t,0)|_\infty + T|e|_\infty \\
&:= \bar{\eta},
\end{aligned} \tag{3.5}$$

which, together with (2.7), implies that there exists a positive constant $M > 1 + (\bar{\eta})^{1/(p-1)}$ such that, for all $t \in R$,

$$|x'|_\infty < M, \quad |x|_\infty \leq d + \frac{1}{2}\sqrt{T}|x'|_2 \leq d + \frac{1}{2}T|x'|_\infty < M. \tag{3.6}$$

Set

$$\Omega = \left\{ x \in C_T^1 : |x|_\infty \leq M, |x'|_\infty \leq M \right\}, \tag{3.7}$$

then we know that (2.6) has no T -periodic solution on $\partial\Omega$ as $\lambda \in (0,1)$ and when $x(t) \in \partial\Omega \cap R$, $x(t) = M$ or $x(t) = -M$, from (A_2) , we can see that

$$\begin{aligned}
&\frac{1}{T} \int_0^T \left\{ -g_0(t, M) - \sum_{k=1}^n g_k(t, M) + e(t) \right\} dt > 0, \\
&\frac{1}{T} \int_0^T \left\{ -g_0(t, -M) - \sum_{k=1}^n g_k(t, -M) + e(t) \right\} dt < 0,
\end{aligned} \tag{3.8}$$

so condition (ii) of Lemma 2.1 is also satisfied. Set

$$H(x, \mu) = \mu x - (1 - \mu) \frac{1}{T} \int_0^T \left[g_0(t, x) + \sum_{k=1}^n g_k(t, x) - e(t) \right] dt, \tag{3.9}$$

and when $x \in \partial\Omega \cap R$, $\mu \in [0,1]$, we have

$$xH(x, \mu) = \mu x^2 - (1 - \mu)x \frac{1}{T} \int_0^T \left[g_0(t, x) + \sum_{k=1}^n g_k(t, x) - e(t) \right] dt > 0. \tag{3.10}$$

Thus, $H(x, \mu)$ is a homotopic transformation and

$$\begin{aligned}
\deg\{F, \Omega \cap R, 0\} &= \deg\left\{ -\frac{1}{T} \int_0^T \left[g_0(t, x) + \sum_{k=1}^n g_k(t, x) - e(t) \right] dt, \Omega \cap R, 0 \right\} \\
&= \deg\{x, \Omega \cap R, 0\} \neq 0,
\end{aligned} \tag{3.11}$$

so condition (iii) of Lemma 2.1 is satisfied. In view of the previous Lemma 2.1, (1.1) has at least one solution with period T . This completes the proof. \square

4. Example and Remark

Example 4.1. Let $p = 4$, $g_0(t, x) = -10e^{20+\sin t}x$, $g_1(t, x) = -2e^{2+\sin t}\sin x$, and $g_2(t, x) = -3e^{3+\cos t}\cos x$ for all $t, x \in \mathbb{R}$. Then, the following Liénard type p -Laplacian equation with two constant delays

$$(\varphi_p x'(t))' + 55x'(t) + g_0(t, x(t)) + g_1(t, x(t-1)) + g_2(t, x(t-2)) = \cos t \quad (4.1)$$

has a unique 2π -periodic solution.

Proof. From (4.1), it is straight forward to check that all the conditions needed in Theorem 3.1 are satisfied. Therefore, (4.1) has at least one 2π -periodic solution. \square

Remark 4.2. Obviously, the results in [2–5] obtained on Duffing type p -Laplacian equation with single delay and without multiple delays cannot be applicable to (4.1). This implies that the results of this paper are essentially new.

Acknowledgments

The authors would like to express their sincere appreciation to the editor and anonymous referee for their valuable comments which have led to an improvement in the presentation of the paper. This work was supported by the construct program of the key discipline in Hunan province (Mechanical Design and Theory), the Scientific Research Fund of Hunan Provincial Natural Science Foundation of PR China (Grant no. 11JJ6006), the Natural Scientific Research Fund of Hunan Provincial Education Department of PR China (Grants no. 11C0916, 11C0915, 11C1186), the Natural Scientific Research Fund of Zhejiang Provincial of P.R. China (Grant no. Y6110436), and the Natural Scientific Research Fund of Zhejiang Provincial Education Department of P.R. China (Grant no. Z201122436).

References

- [1] E. Esmailzadeh and G. Nakhaie-Jazar, "Periodic solution of a Mathieu-Duffing type equation," *International Journal of Non-Linear Mechanics*, vol. 32, no. 5, pp. 905–912, 1997.
- [2] A. Sirma, C. Tunç, and S. Özlem, "Existence and uniqueness of periodic solutions for a kind of Rayleigh equation with finitely many deviating arguments," *Nonlinear Analysis*, vol. 73, no. 2, pp. 358–366, 2010.
- [3] Y. Wang, "Novel existence and uniqueness criteria for periodic solutions of a Duffing type p -Laplacian equation," *Applied Mathematics Letters*, vol. 23, no. 4, pp. 436–439, 2010.
- [4] M.-L. Tang and X.-G. Liu, "Periodic solutions for a kind of Duffing type p -Laplacian equation," *Nonlinear Analysis*, vol. 71, no. 5-6, pp. 1870–1875, 2009.
- [5] F. Gao, S. Lu, and W. Zhang, "Existence and uniqueness of periodic solutions for a p -Laplacian Duffing equation with a deviating argument," *Nonlinear Analysis*, vol. 70, no. 10, pp. 3567–3574, 2009.
- [6] Z. Wang, L. Qian, S. Lu, and J. Cao, "The existence and uniqueness of periodic solutions for a kind of Duffing-type equation with two deviating arguments," *Nonlinear Analysis*, vol. 73, no. 9, pp. 3034–3043, 2010.
- [7] H. Gao and B. Liu, "Existence and uniqueness of periodic solutions for forced Rayleigh-type equations," *Applied Mathematics and Computation*, vol. 211, no. 1, pp. 148–154, 2009.
- [8] B. Liu, "Existence and uniqueness of periodic solutions for a kind of Rayleigh equation with two deviating arguments," *Computers & Mathematics with Applications*, vol. 55, no. 9, pp. 2108–2117, 2008.
- [9] R. Manásevich and J. Mawhin, "Periodic solutions for nonlinear systems with p -Laplacian-like operators," *Journal of Differential Equations*, vol. 145, no. 2, pp. 367–393, 1998.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

