

## Original Article

# Surgical treatment of gastrointestinal stromal tumors of the duodenum: a literature review

Georgi Popivanov<sup>1</sup>, Mihail Tabakov<sup>2</sup>, George Mantese<sup>3</sup>, Roberto Cirocchi<sup>3</sup>, Irene Piccinini<sup>3</sup>, Vito D'Andrea<sup>4</sup>, Piero Covarelli<sup>3</sup>, Carlo Boselli<sup>3</sup>, Francesco Barberini<sup>3</sup>, Renata Tabola<sup>5</sup>, Ursi Pietro<sup>4</sup>, Davide Cavaliere<sup>6</sup>

<sup>1</sup>Military Medical Academy, Clinic of Endoscopic, Endocrine surgery and Coloproctology, Sofia, Bulgaria; <sup>2</sup>University Hospital Sv. Ivan Rilski, Surgical Clinic, Sofia, Bulgaria; <sup>3</sup>Department of Surgical Sciences, The University of Rome "La Sapienza", Rome, Italy; <sup>4</sup>Department of Surgical and Biomedical Sciences, University of Perugia, Italy; <sup>5</sup>Department and Clinic of Gastrointestinal and General Surgery, Wroclaw Medical University, Wroclaw, Poland; <sup>6</sup>General Surgery and Surgical Oncology, Morgagni-Pierantoni Hospital, Forlì, Italy

**Contributions:** (I) Conception and design: G Popivanov, F Barberini, G Mantese, R Cirocchi, I Piccinini, V D'Andrea, U Pietro, D Cavaliere; (II) Administrative support: G Popivanov, F Barberini, R Cirocchi, I Piccinini, V D'Andrea, U Pietro, D Cavaliere; (III) Provision of study material or patients: G Popivanov, F Barberini, G Mantese, R Cirocchi, U Pietro, D Cavaliere; (IV) Collection and assembly of data: G Popivanov, G Mantese, M Tabakov, R Cirocchi, I Piccinini, V D'Andrea, U Pietro, D Cavaliere; (V) Data analysis and interpretation: G Popivanov, M Tabakov, R Cirocchi, PR, I Piccinini, V D'Andrea; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

**Correspondence to:** George Mantese. Rua Carazinho 146 ap. 01 Porto Alegre, Brazil. Email: georgemantese@hotmail.com.

**Background:** Gastrointestinal stromal tumors (GIST) are the most frequent mesenchymal tumours in the digestive tract. The duodenal GIST (dGIST) is the rarest subtype, representing only 4–5% of all GIST, but up to 21% of the resected ones. The diagnostic and therapeutic management of dGIST may be difficult due to the rarity of this tumor, its anatomical location, and the clinical behavior that often mimic a variety of conditions; moreover, there is lack of consent for their treatment. This study has evaluated the scientific literature to provide consensus on the diagnosis of dGIST and to outline possible options for surgical treatment.

**Methods:** An extensive research has been carried out on the electronic databases MEDLINE, Scopus, EMBASE and Cochrane to identify all clinical trials that report an event or case series of dGIST.

**Results:** Eighty-six studies that met the inclusion criteria were identified with five hundred forty-nine patients with dGIST: twenty-seven patients were treated with pancreatoduodenectomy and ninety-six with only local resection (segmental/wedge resections); in four hundred twenty-six patients it is not possible identify the type of treatment performed (pancreatoduodenectomy or segmental/wedge resections).

**Conclusions:** dGISTs are a very rare subset of GISTs. They may be asymptomatic or may involve symptoms of upper GI bleeding and abdominal pain at presentation. Because of the misleading clinical presentation the differential diagnosis may be difficult. Tumours smaller than 2 cm have a low biological aggressiveness and can be followed annually by endoscopic ultrasound. The biggest ones should undergo radical surgical resection (R0). In dGIST there is no uniformly adopted surgical strategy because of the low incidence, lack of experience, and the complex anatomy of the duodenum. Therefore, individually tailored surgical approach is recommended. R0 resection with 1–2 cm clear margin is required. Lymph node dissection is not recommended due to the low incidence of lymphatic metastases. Tumor rupture should be avoided.

**Keywords:** Duodenal gastrointestinal stromal tumor (dGIST); surgery

Received: 25 February 2018; Accepted: 04 September 2018; Published: 21 September 2018.

doi: 10.21037/tgh.2018.09.04

View this article at: <http://dx.doi.org/10.21037/tgh.2018.09.04>

## Introduction

Gastrointestinal stromal tumours (GIST) are the most common type of mesenchymal tumours found in the gastrointestinal tract (1). For years, GISTs have been defined as smooth gastrointestinal muscle tumors and have been called under various names and often misdiagnosed as leiomyomas, schwannomas and sarcomas (2). Currently, the distinctive feature of GIST is the mutation in the *c-kit* protooncogene leading to gain-of-function and subsequent cell proliferation (3-5). GISTs are rare, with relative annual incidence of 14.5 per million and prevalence of 129 per million (6). GIST may occur anywhere in the digestive tract, but are more frequently located in the stomach (60–70%) and midgut (25%) and less often in colon and rectum (5–10%) (7). Extravisceral GIST occurs in less than 10% of patients, most frequently in mesentery, intrabdominal, pelvis and retroperitoneal space (7). Duodenal GISTs (dGIST) represent only 4–5% of all GISTs, but accounted for 6–21% of surgical resected ones (8-10). The complex anatomy of the duodeno-pancreatic region can make their diagnosis and treatment extremely challenging. Anatomical closeness to noble structures (i.e., to the head of the pancreas, kidney and biliary structures) can lead to misdiagnosis and inappropriate management (11). Additionally, several factors complicate the management of dGISTs such as the relative lack of experience, the ambiguous clinical manifestations that often mimic a wide range of clinical conditions, the anatomical complexity and the lack of consensus on treatment.


The publication of solid research findings (3,12) that characterize the pathogenesis and histology of GIST have created a discontinuity in the terminology used to describe this entity. Due to the unclear terminology of scientific literature published before 2000, it is very difficult to analyse and interpret the previous reports, but the researches published since then has significantly increased (4).

Herein, we aimed to perform a state-of-the-art review of available English literature to improve the understanding of dGIST and to outline the best options for surgical treatment.

## Methods

### Search strategy

Comprehensive research has been conducted to identify all clinical studies that report the occurrence of dGIST. The search was carried out on the electronic databases MEDLINE, Scopus, EMBASE and Cochrane Libraries. The research strategy included all studies published after



**Figure 1** Study selection process.

2000 and focused on relevant articles using the following keywords: “gist”, “duodenal”, “stromal gastrointestinal cancer”, “surgery” and associated free terms. No language restrictions have been imposed to limit results.

### Inclusion criteria

Studies were selected for consideration in this analysis provided they reported cases with dGIST. In order to select only those studies that certainly dealt with dGIST, only those studies in which the immunohistochemical analysis of the *cKIT* was reported and were positive were included.

### Study selection

The inclusion criteria were used to examine the title, abstracts and full texts of the studies obtained from the research results. Two reviewers independently examined all citations on titles and abstracts produced by the search strategy and retrieved all potentially relevant reports. Two other independent readers then screened the full text of the studies to identify the entries that met the inclusion criteria. Any disagreements have been resolved by consensus. Differences in data extraction have been resolved by consensus, with references to the original article.

## Results

The initial search produced 185 potentially relevant articles (Figure 1, Tables 1,2,3,4,5) that were suitable for subsequent screening. The titles and abstracts of these articles were screened to determine the relevance and eligibility for inclusion—53 papers were excluded for different reasons:

**Table 1** Characteristics of included studies: pancreaticoduodenectomy

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patient (age/gender)	Duodenal portion location	Symptoms and signs at admission
Takahashi 2001 (13)	Japan	1	77; F	Second	Mild anemia
Uchida 2004 (14)	Japan	1	53; F	Extramural	Asymptomatic
Hompes 2004 (15)	Belgium	2	32; F	Second	Right superior abdominal discomfort
			72; F	Second	Hypovolemic shock, melena, pallor, weakness, fatigue, anemia
Kim 2004 (16)	Korea	1	37; F	Second	Weakness, nausea, melena, anemia
Sakakima 2004 (17)	Japan	1	46; F	Second	Epigastralgia, melena, anemia
Sakakura 2006 (18)	Japan	1	37; M	Second	Abdominal discomfort
Ludvigsen 2007 (19)	Denmark	1	41; M	Second	Anemia
Fiore 2009 (20)	Italy	3	49; F	NR	NR
			66; M	NR	NR
			71; M	NR	NR
Hecker 2010 (21)	Germany	1	58	Second	Recurrent episodes of upper abdominal pain, anemia, weight loss, fatigue and fever attacks, upper gastrointestinal bleeding after neoadjuvant imatinib
Frampton 2010 (22)	UK	1	37; F	Second	Right upper quadrant pain, sweating, nausea, hot flushes, palpitations
Wall 2010 (23)	–	1	36; F	–	Diarrhea, bloating, weight loss, lethargy, collapse, melena, hypotension, tachycardia
Machado 2011 (24)	Oman	1	58; M	Second	Melena, weight loss
Morcos 2011 (25)	Jordan	1	38; F	Second	Epigastric discomfort, abdominal pain, anemia
Singh 2012 (26)	India	1	30; M	Second	Mass in the right upper abdomen, abdominal pain
Koontz 2012 (27)	USA	1	36; F	Third	Upper gastrointestinal bleeding, abdominal fullness, early satiety, fatigue, anemia, syncope
Blandamura 2014 (28)	Italy	1	72; M	Third	Increasing vomiting, weight loss
Bormann 2014 (29)	Germany	1	64; F	Second	Slowly increasing pain in right upper abdomen
Kobayashi 2014 (30)	Japan	1	36; M	Second	Loss of consciousness
Parisi 2014 (31)	Italy	1	68	Second	Fatigue, anemia, positive fecal occult blood
Bhambare 2015 (32)	India	1	38; M	Second	Large abdominal lump, abdominal pain
Okasha 2015 (33)	Egypt	1	42; F	Second	Abdominal discomfort, melena, hematemesis
Niikura 2016 (34)	Japan	1	75; M	First	Anemia, melena, hemorrhagic shock
Yamamoto 2016 (35)	Japan	1	81; F	Second	Diarrhea
Tornambe 2017 (36)	Italy	1	69; M	Second	Melena, asthenia, dizziness, severe anemia

NR, information not reported

**Table 2** Characteristics of included studies: local resection

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patient (age; gender)	Duodenal portion location	Symptoms and signs at admission
Sawaki 2003 (37)	Japan	1	54; F	First	Asymptomatic
Sakamoto 2003 (38)	Japan	1	31; F	Third	NR
Kurihara 2005 (39)	Japan	1	64; M	Second	Tarry stools, dizziness, and severe anemia
Goh 2005 (40)	Singapore	2	72; F 69; M	Second Second and third	Epigastralgia, melena, anemia Abdominal discomfort
Cavallini 2005 (41)	Italy	1	66; F	Second	Anemia, melena
Vu 2005 (42)	Japan	1	43; M	Third	NR
Towu 2006 (43)	UK	1	7; M	Second and third	Severe upper gastrointestinal bleeding
Kwon 2007 (44)	Korea	1	49; M	Second and third	Asymptomatic
Graham 2007 (45)	UK	1	57; M	Fourth	Melena, anemia
Gupta 2007 (46)	USA	1	63; M	Second and third	Gastrointestinal bleeding
Mohiuddin 2007 (47)	UK	1	56; M	Second	Melena, anemia
Fernández Salazar 2007 (48)	Spain	1	44; M	NR	GI bleeding
Mennigen 2008 (49)	Germany	1	29; M	Third	Acute upper gastrointestinal bleeding
Takahashi 2009 (50)	Japan	1	57; F	First	Bloody stools
Takeuchi 2009 (51)	Japan	1	55; M	Third	Asymptomatic
Seçkin 2009 (52)	Turkey	1	56; M	Third	Melena, anemia
Hirashima 2009 (53)	Japan	1	68; F	NR	Asymptomatic
Mehta 2011 (54)	USA	1	33; F	Third	Melena, dizziness, anemia, hypotension, tachycardia
Chung 2011 (55)	Korea	2	65; M 49; F	Third Fourth	Abdominal pain Upper gastrointestinal bleeding, melena
Cameron 2011 (56)	Germany	1	62; M	Second	NR
Kato 2011 (57)	Japan	1	60; M	Third	NR
Chen 2012 (58)	Australia	1	52; M	NR	Anemia
El-Gendi 2012 (59)	Egypt	12	60; F, n=5, M, n=7	First, n=3 Second n=1 Third, n=3 Fourth, n=1	Melena, anemia
Acar 2013 (60)	Turkey	1	65; F	Third	Abdominal pain
Mouaqit 2013 (61)	Morocco	1	65; F	Second and third	Abdominal pain, anemia
Mokhtare 2013 (62)	Iran	1	2; M	Third	Acute upper gastrointestinal bleeding

**Table 2** (continued)

Table 2 (continued)

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patient (age; gender)	Duodenal portion location	Symptoms and signs at admission
Shaw 2013 (63)	UK	1	61; M	Third and fourth	Hypovolemic shock, upper gastrointestinal bleeding, melaena, hematemesis, temporary loss of consciousness
Ueda 2014 (64)	Japan	1	72; F	Third	Asymptomatic
Hankiewicz-Ziołkowska 2014 (65)	Poland	1	65; F	First	Asymptomatic
Manxhuka-Kerliu 2014 (66)	Kosovo	1	30	Fourth-jejunum	Abdominal pain, nausea, vomiting
Fukuyama 2014 (67)	Japan	1	69; M	Second	Anemia, melena
Borgaonkar 2015 (68)	India	2	52; F	NR	NR
			23; M	NR	NR
Jones 2015 (69)	USA	2	40; F	NR	Epigastric pain, recurrent gastrointestinal bleeding
			29; F	Second	Asymptomatic
Mrak 2015 (70)	Austria	1	68; F	Upper portions	Abdominal pain, melena, hemodynamic instability
Kumar 2015 (71)	India	1	55; F	Second and Third	Epigastric pain
Chung 2015 (72)	Korea	21	59; F, n=9, M, n=12	First, n=7	Incidental finding, abdominal pain, bleeding
				Second, n=5	
				Third, n=5	
				Fourth, n=4	
Graziosi 2015 (73)	Italy	1	51; F	Second	Anemia, melena
Boselli 2016 (74)	Italy	3	75; F	Fourth	Melena, anemia
			82; F	Passage between second and third	Melena, syncope, anemia, hemodynamic instability
			76; M	Passage between second and third	Melena, syncope, anemia, hemodynamic instability
Caruso 2016 (75)	Italy	1	73	Fourth, duodenojejunal junction	Anemia, melena
Jones 2016 (76)	USA	1	71; F	Second	Recurrent GI bleeding
Turculeț 2016 (77)	Romania	1	48; M	Third and Fourth	Melena, asthenia, dizziness, severe anemia
Huo 2016 (78)	China	1	58; F	First	Epigastric abdominal discomfort, diarrhea, recurrent vomiting
Mori 2016 (79)	Japan	1	74; M	Third	NR
Valli 2016 (80)	Switzerland	1	19; F	Second	Fainting, anemia, melena

Table 2 (continued)

Table 2 (continued)

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patient (age; gender)	Duodenal portion location	Symptoms and signs at admission
Crocetti 2016 (81)	Italy	9	70; F, n=9, M, n=12	NR	Pain, bleeding, dyspepsia
Thillai 2017 (82)	India	1	50; F	Second and third	Abdominal pain
Elston 2017 (83)	Australia	1	29; M	Second	Acute upper gastrointestinal hemorrhage, melena
Vasile 2017 (84)	Romania	1	59; F	First and second	Abdominal pain, upper gastrointestinal bleeding (melena), faintness, anemia
Zioni 2017 (85)	Israel	1	68; M	Second and third	Upper gastrointestinal bleeding, general weakness, melena, anemia
Perfetti 2017 (86)	Italy	1	33; M	Second	Gastrointestinal bleeding, anemia
Hakozaki 2017 (87)	Japan	1	70; F	First	Positive fecal occult blood test

NR, information not reported.

Table 3 Characteristics of included studies: PD and local resections

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patients (age; gender)	Duodenal portion location	Symptoms and sign at admission
Miettinen 2003 (9)	USA	156	56; F, n=71, M, n=85	First, n=10 Second, n=42 Third, n=17 Second-third, n=7 Fourth, n=11	Anemia, melena, obstruction, acute abdomen
Relles 2009 (88)	USA	2	43; M 31; M	Second Ligament of Treitz	Melena, fatigue, shortness of breath Solid food dysphagia, abdominal discomfort
Tien 2010 (89)	Taiwan	25	62; F, n=9, M, n=16	3 first 13 second 5 third 4 fourth	Bleeding, pain, mass
Miki 2010 (90)	Japan	6	64; F 70; F 67; F 39; F 65; M 75; F	Second Fourth First First Second Second	Melena Abdominal mass Asymptomatic Melena Anemia Anemia
Yagishita 2011 (91)	Japan	4	62; F 69; M 76; M 72; M	Second First Second First	NR NR Anemia NR

Table 3 (continued)

Table 3 (continued)

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patients (age; gender)	Duodenal portion location	Symptoms and sign at admission
Liang 2013 (92)	Japan	28	28; F	Second	Melena
			48; F	Second	Asymptomatic
			60; M	Second	Melena
			70; M	Second	Abdominal pain
			71; M	Third	Asymptomatic
			76; F	Second	Melena
			42; F	Second	Melena
			74; M	First	Melena
			53; F	First	Melena
			47; F	Second	Melena
			55; F	Fourth	Melena
			51; M	Second	Melena
			50; M	Second	Hematemesis
			69; F	Third	Abdominal pain
			65; M	Third	Melena
			63; M	Second	Acute abdomen
			44; F	Second	Hematemesis
			57; F	Third	Discomfort
			20; F	Second	Melena
			52; M	Fourth	Abdominal pain
			53; F	Second	Abdominal pain
			71; F	Second	Early satiety
			53; M	First	Early satiety
			50; F	First	Melena
			46; F	Second	Abdominal pain
			55; M	Second	Melena
			51; M	Third	Asymptomatic
			46; F	Second	Melena
Hoepfner 2013 (93)	Germany	9	51; M	Second	Bleeding
			63; M	Third	Bleeding
			52; M	First	Bleeding
			62; F	First	Jaundice
			58; M	First	Abdominal pain
			69; F	Second	Incidental finding
			43; F	Fourth	Bleeding
			75; M	Third	Bleeding
49; F	Second	Bleeding			

Table 3 (continued)

Table 3 (continued)

Study (year of publication)	Nation	Number of patients with duodenal GIST	Patients (age; gender)	Duodenal portion location	Symptoms and sign at admission
Yang 2013 (94)	China	22	58; F, n=7, M, n=15	First, n=3 Second, n=14 Third-fourth, n=5	Asymptomatic, abdominal discomfort/pain, melena, weight loss, fatigue, abdominal distension, anorexia, back pain, hematemesis, jaundice, anemia, palpable abdominal mass
Zhou 2013 (95)	China	48	53; F, n=20, M, n=28	First, n=11 Second, n=17 Third, n=6 Fourth, n=2 First-second, n=2 Second-third, n=4	Bleeding, abdominal pain, abdominal discomfort, jaundice
Duffaud 2014 (96)	France	114	57; F, n=55, M, n=59	First, n=8 Second, n=38 Third, n=27 Fourth, n=15	Pain, GI bleeding, anemia, asymptomatic
Ucar 2015 (97)	Turkey	2	65; M 60; M	First Second	Upper gastrointestinal bleeding Ileus, acute abdomen, hemorrhagic pancreatitis
Lv 2017 (98)	China	10	44; F 50; M 43; M 50; F 60; M 44; M 50; F 44; M 65; M 64; F	Third Second Second Second and third Second Second Third Second and third Second Third	Abdominal pain Asymptomatic Melena Abdominal pain Abdominal discomfort Abdominal discomfort Melena Melena Melena Abdominal pain

NR, information not reported.

Table 4 Characteristics of included studies: PD and local resections

Study (year of publication)	Histologic characteristics					
	c-kit	DOG1	CD34	Smooth muscle actin	S100	Desmin
Miettinen 2003 (9)	109	NR	49	38	19	No
Relles 2009(88)	Yes	NR	Yes	NR	No	NR
	Yes	No	Yes	No	No	No
Tien 2010 (89)	25	NR	5	5	4	No

Table 4 (continued)



Table 4 (continued)

Study (year of publication)	Histologic characteristics					
	c-kit	DOG1	CD34	Smooth muscle actin	S100	Desmin
Miki 2010 (90)	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
Yagishita 2011 (91)	Yes	No	Yes	No	No	NR
	Yes	No	Yes	No	No	NR
	Yes	No	Yes	No	No	NR
	Yes	No	No	No	No	NR
Liang 2013 (92)	Yes	NR	No	Yes	No	No
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	No	Yes	No	NR
	Yes	NR	Yes	Yes	Yes	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	No	Yes	No
	No	NR	Yes	No	No	NR
	Yes	NR	Yes	Yes	Yes	NR
	Yes	NR	No	Yes	Yes	NR
	Yes	NR	Yes	No	Yes	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	No	Yes	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	Yes	No	NR
	Yes	NR	Yes	No	Yes	NR
	Yes	NR	Yes	No	No	NR
	Yes	NR	No	Yes	No	NR
	Yes	NR	No	Yes	No	NR
Yes	NR	Yes	No	Yes	NR	
Yes	NR	Yes	Yes	Yes	NR	

Table 4 (continued)

Table 4 (continued)

Study (year of publication)	Histologic characteristics					
	c-kit	DOG1	CD34	Smooth muscle actin	S100	Desmin
Hoepfner 2013 (93)	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
Yang 2013 (94)	18	NR	17	10	8	No
Zhou 2013 (95)	47	NR	32	NR	5	6
Duffaud 2014 (96)	105	NR	58	NR	NR	NR
Ucar 2015 (97)	Yes	Yes	Yes	NR	NR	NR
	Yes	Yes	Yes	NR	NR	NR
Lv 2017 (98)	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR

NR, information not reported.

Table 5 Characteristics of included studies: PD and local resections

Study (year of publication)	Tumor extension out duodenum	Type of treatment	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Survival (years; months)
Miettinen 2003 (9)	Duodenum	15 enucleation; 48 segmental resection; 21 wedge resection; 21 pancreaticoduodenectomy; 51 NR	NR	NR	NR
Relles 2009 (88)	Duodenum	Pancreaticoduodenectomy	NR	NR	NR
	Duodenum	Local resection	NR	NR	NR
Tien 2010 (89)	Duodenum	9 pancreaticoduodenectomy; 16 local resection	NR	NR	NR

Table 5 (continued)

Table 5 (continued)

Study (year of publication)	Tumor extension out duodenum	Type of treatment	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Survival (years; months)
Miki 2010 (90)	Duodenum	Duodenectomy	NR	No	NR
		Local resection	NR	Yes	NR
		Local resection	NR	No	NR
		Local resection	NR	No	NR
		Local resection	NR	No	NR
		Pancreaticoduodenectomy	NR	No	NR
Yagishita 2011 (91)	Duodenum	Partial duodenectomy	NR	NR	NR
	Duodenum	No treatment	NR	NR	NR
	Duodenum	No treatment	NR	NR	NR
	Duodenum, pancreas	Pancreaticoduodenectomy	NR	NR	NR
Liang 2013 (92)	Duodenum	Segmental resection	No	No	164 m
		Wedge resection		No	146 m
		Pancreaticoduodenectomy		No	61 m
		Pancreaticoduodenectomy		No	35 m
		Segmental resection		No	23 m
		Segmental resection		No	25 m
		Pancreaticoduodenectomy		No	73 m
		Wedge resection		No	61 m
		Segmental resection		No	116 m
		Pancreaticoduodenectomy		No	111 m
		Segmental resection		No	55 m
		Pancreaticoduodenectomy		No	23 m
		Wedge resection		No	102 m
		Pancreaticoduodenectomy		No	NR
		Segmental resection		No	68 m
		Pancreaticoduodenectomy		No	49 m
		Pancreaticoduodenectomy		No	33 m
		Segmental resection		No	57 m
		Segmental resection		No	86 m
		Segmental resection		No	33 m
		Wedge resection		No	40 m
		Segmental resection		No	59 m
Wedge resection		No	75 m		
Segmental resection		No	73 m		
Segmental resection		Yes	71 m		
Pancreaticoduodenectomy		No	70 m		
Segmental resection		Yes	63 m		
Pancreaticoduodenectomy		No	61 m		

Table 5 (continued)

Table 5 (continued)

Study (year of publication)	Tumor extension out duodenum	Type of treatment	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Survival (years; months)
Hoepfner 2013 (93)	Duodenum	Open wedge resection	No	No	6 m
		Open wedge resection	No	Yes	7 m
		Laparoscopic wedge resection	No	No	13 m
		Pancreaticoduodenectomy	No	No	Died at 3 m
		Open wedge resection	No	No	90 m
		Open wedge resection	No	No	111 m
		Segmental resection	No	No	92 m
		Open wedge resection	Yes	No	Died at 37 m
		Tumor resection	Yes	Yes	39 m
Yang 2013 (94)	Duodenum	9 pancreaticoduodenectomy; 13 local resection	No	5	NR
Zhou 2013 (95)	Duodenum	34 local resection; 14 pancreaticoduodenectomy	NR	NR	NR
Duffaud 2014 (96)	Duodenum	82 local resection; 23 pancreaticoduodenectomy	11	20	NR
Ucar 2015 (97)	Duodenum	Local resection	NR	NR	NR
		Pancreaticoduodenectomy	NR	NR	NR
Lv 2017 (98)	Duodenum	Segmental resection	Yes	NR	36 m
		Tumor resection	Yes	NR	50 m
		Radical resection	Yes	NR	44 m
		Segmental resection	Yes	NR	47 m
		Tumor resection	Yes	NR	17 m
		Tumor resection	Yes	NR	36 m
		Segmental resection	Yes	NR	17 m
		Radical resection	Yes	NR	16 m
		Segmental resection	Yes	NR	16 m
		Pancreaticoduodenectomy	Yes	NR	41 m

NR, information not reported.

non-duodenal GISTs, no surgery type reported, and no patient data reported leaving 132 studies for inclusion in the present review. Of them, 46 studies did not report the data about *cKIT* and were excluded thus leaving 86 studies included in the analysis. The patients excluded for non-duodenal GIST were 7,142, those excluded for no surgery type reported were 1,039 and 1,002 patients were excluded because of no *cKIT* reported. The characteristics of the included studies have been summarized in *Tables 1,2,3,4,5*. All studies were published between 2001 and 2017. Eighty-two papers with 539 cases reported data on the signs and symptoms at the time of hospital admission, whereas in four studies there were no data. Upon admission, the most

common symptoms were GI bleeding and melena, followed by abdominal pain and discomfort. Symptoms of fatigue, fainting, vomiting, and ankle edema were also reported but occurred far less frequently. In the majority of cases dGISTs were located in the second portion of the duodenum. Several cases reported dGIST localization in both the second and third portions of the duodenum.

The included studies summoned a total of 549 patients with dGIST—27 patients treated with pancreatoduodenectomy (PD) (*Table 1*), 96 patients treated with local resection (*Table 2*), and 426 patients treated with either PD or segmental/wedge resections (*Tables 3,4,5*).

Among patients treated with PD (*Tables 1,6*), the majority

**Table 6** Characteristics of included studies: pancreaticoduodenectomy

Study (year of publication)	Local tumor extension	Surgical access	Neoadjuvant treatment with imatinib	Postoperative treatment with imatinib	Multivisceral resection	Survival (years, months)
Takahashi 2001 (13)	Duodenum, pancreatic head, gall bladder, gastric antrum, regional lymph nodes	Open	NR	NR	Gastric antrum	NR
Uchida 2004 (14)	Duodenum	Open	No	No	No	NR
Hompes 2004 (15)	Duodenum, pancreas, mesocolon	Open	No	No	Ascending colon	NR
Kim 2004 (16)	Duodenum	Open	No	No	No	5 m
Sakakima 2004 (17)	Duodenum	Open	NR	NR	No	NR
Sakakura 2006 (18)	Duodenum	Open	No	Yes	No	24 m
Ludvigsen 2007 (19)	Duodenum, hilum, liver, right kidney, right hemicolon, right hemipancreas, celiac trunk, inferior vena cava, right renal vein, right renal pelvis	Open	Yes	Yes	Right kidney, adrenal gland	NR
Fiore 2009 (20)	Duodenum	Open	Yes	NR	No	NR
			Yes	NR	No	NR
			Yes	NR	No	NR
Hecker 2010 (21)	Duodenum, pancreas, right flexure of the colon	Open	Yes	Yes	Right hemicolon	NR
Frampton 2010 (22)	Duodenum	Open	NR	NR	No	NR
Wall 2010 (23)	Duodenum	Open	No	No	No	NR
Machado 2011 (24)	Duodenum, right hemicolon	Open	No	Yes	Right hemicolon	2 m
Morcos 2011 (25)	Duodenum	Open	No	No	No	34 m
Singh 2012 (26)	Duodenum	Open	No	Yes	No	NR
Koontz 2012 (27)	Duodenum	Open	Yes	Yes	No	NR
Blandamura 2014 (28)	Duodenum	Open	No	No	No	NR
Bormann 2014 (29)	Duodenum, pancreas	Open	No	No	No	NR
Kobayashi 2014 (30)	Duodenum	Open	No	No	No	18 m
Bhambare 2015 (32)	Duodenum	Open	No	Yes	No	NR
Okasha 2015 (33)	Duodenum	Open	No	No	No	NR
Niikura 2016 (34)	Duodenum	Open	No	Yes	No	6 y
Yamamoto 2016 (35)	Duodenum	Open	No	No	No	NR
Tornambe 2017 (36)	Duodenum	Open	No	No	No	12 m

NR, information not reported.

were females (n=14, 52%; males: n=11, 41%; not reported: n=2, 7%), with a median age of 50 years (Table 1). Most patients presented with symptoms of anemia (n=11, 40%), melena (n=8, 27%), fatigue (n=4, 15%), and weight loss (n=4, 15%). However, in 3 patients the information was not reported. For most patients, the dGIST were located in the second portion of the duodenum (first: n=1, 4%; second: n=19, 70%; third: n=1, 4%; not reported: n=5, 18%) (Table 1). Four studies (13,19,21,25) reported dGIST tumors that extended beyond the duodenum. All PDs were performed by open procedure. While some patients received imatinib as a neoadjuvant treatment (n=7, 26%), a greater proportion received the treatment post-operatively (n=9, 33%) (Table 6). For most patients, no long term survival data were reported (n=21, 75%). Notably, the PD studies reported histological information on KIT, DOG1, or CD34.

Among patients treated with local resection (Tables 2,7) there was a slight preponderance of males (n=51, 53%; females: n=45, 47%) with a median age of 57.3 years (Table 2). Upon admission, patients most commonly presented with nausea (n=1, 1%), abdominal pain (n=8, 8%), undetermined pain (n=18, 19%), melena (n=20, 21%), dizziness (n=3, 3%), and/or GI bleeding (n=14, 16%). No data on presenting signs and symptoms was available for 7 patients (Table 2).

Within this group, most dGIST were located in the second and third portions of the duodenum (first: n=16, 17%; second: n=29, 30%, third: n=32, 33%; fourth: n=12, 12%; not reported: n=7, 7%), with tumors extending beyond the duodenum in four cases. For all local resections reported, surgical access was predominantly achieved by either open (n=89, 92%) or laparoscopic procedure (n=5, 5%). Patients treated with a local resection received less neoadjuvant (n=13, 13%) and post-operative (n=22, 23%) treatment with imatinib (Table 8), as compared to patients treated with a pancreaticoduodenectomy (Table 6), and four patients underwent multiple visceral resections.

Of the patients whose histological or molecular data were provided (Table 7), most tested positive for *cKIT* (n=85, 46%), CD34 (n=48, 26%) and DOG1 (n=20, 11%). A minority of patients tested positive for either smooth muscle actin (n=5, 3%), S100 (n=7, 4%) or desmin (n=1, 1%).

Four hundred twenty-six patients were treated with either PD or segmental/wedge resections. Within this group, there was preponderance of males (n=218, 51%) *vs.* females (n=183, 43%), not reported (n=25, 6%). At the time of admission, patients frequently presented with symptoms of bleeding (n=137, 32%), abdominal pain or discomfort (n=258, 60%) and melena (n=235, 55%). Five patients were

asymptomatic (Tables 3,4,5). dGISTs were predominantly located in the second and third portions of the duodenum in this group (first: n=47, 11%; second: n=164, 38%; third: n=83, 19%; fourth: n=41, 9%). In many cases the tumors extended into multiple portions of the duodenum simultaneously. All of these studies reported data on *cKIT* and several studies reported data on CD34 and DOG1.

Within this cohort the most common type of surgical treatment was local resection (n=151, 35%), followed by PD (n=92, 21%) and segmental (n=97, 15%) or wedge resections (n=32, 7%). No data on the type of surgical treatment were available for 51 patients. Within this group of patients, both neoadjuvant and post-operative treatment with imatinib were rarely applied (neoadjuvant treatment: n=23, 5%; adjuvant treatment: n=30, 7%). Most patients received no imatinib treatment prior to surgery (n=160, 38%) or post-operatively (n=149, 35%).

### Surgical treatment

There are three main surgical options: pancreatoduodenectomy (PD) (10,24), wedge resection (40,60,63,97) and segmental resection. PD is indicated in the cases with involvement of major duodenal papilla, pancreas or pancreatic duodenal wall and is required in 20–40%. Based on the literature, the following options for surgical reconstruction according to the localization and extent of duodenal resection are available:

First and second duodenal portions:

- ❖ wedge resection with primary transverse closure of duodenum (39) or retrocolic Roux-en-Y loop to cover a large defect (97);
- ❖ segmental resection, closure of distal stump and duodenojejunostomy by retrocolic Roux-en-Y loop (99);
- ❖ segmental resection with antrectomy with side-to-side posterior or Roux-en-Y gastrojejunostomy (55,59);
- ❖ subtotal resection with side-to-side duodenojejunostomy (100);
- ❖ sleeve resection with gastroduodenostomy (the case with situs inversus totalis) (82);
- ❖ ampullectomy with sphincteroplasty (55).

Third and fourth duodenal portions:

- ❖ segmental resection with closure of distal stump and side-to-side (93) or end-to-side/end-to-end duodenojejunostomy (62,92);
- ❖ segmental resection, end-to-end anastomosis, pylorus closure and gastroenterostomy with/without feeding jejunostomy (77);
- ❖ segmental resection, closure of distal stump near leg.

**Table 7** Characteristics of included studies: local resection

Study (year of publication)	Histologic characteristics					
	C-kit/CD117	DOG1	CD34/PDGFR-alfa	Smooth muscle actin	S100	Desmin
Sawaki 2003 (37)	Yes	NR	Yes	No	No	No
Sakamoto 2003 (38)	Yes	NR	NR	NR	NR	NR
Kurihara 2005 (39)	Yes	NR	Yes	Yes	No	No
Goh 2005 (40)	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
Cavallini 2005 (41)	Yes	NR	NR	No	Yes	No
Vu 2005 (42)	Yes	NR	No	Yes	No	No
Towu 2006 (43)	Yes	NR	No	No	NR	NR
Kwon 2007 (44)	Yes	NR	Yes	No	NR	NR
Graham 2007 (45)	Yes	NR	Yes	NR	NR	NR
Gupta 2007 (46)	Yes	NR	NR	No	No	NR
Mohiuddin 2007 (47)	Yes	NR	NR	NR	NR	NR
Fernández Salazar 2007 (48)	Yes	NR	NR	NR	Yes	NR
Menningen 2008 (49)	Yes	NR	Yes	No	No	NR
Takahashi 2009 (50)	Yes	No	Yes	No	Yes	No
Takeuchi 2009 (51)	Yes	NR	Yes	No	No	NR
Seçkin 2009 (52)	Yes	NR	Yes	NR	Yes	NR
Hirashima 2009 (53)	Yes	NR	Yes	NR	No	No
Mehta 2011 (54)	Yes	No	Yes	Yes	No	No
Chung 2011 (55)	Yes	NR	NR	NR	NR	NR
	Yes	NR	NR	NR	NR	NR
Cameron 2011 (56)	Yes	NR	NR	NR	NR	NR
Kato 2011 (57)	Yes	NR	NR	NR	NR	NR
Chen 2012 (58)	Yes	NR	NR	NR	NR	NR
El-Gendi 2012 (59)	12	NR	8	NR	2	No
Acar 2013 (60)	Yes	NR	NR	NR	Yes	NR
Mouaqit 2013 (61)	Yes	NR	Yes	NR	NR	NR
Mokhtare 2013 (62)	Yes	Yes	Yes	NR	NR	NR
Shaw 2013 (63)	Yes	NR	Yes	NR	NR	NR
Ueda 2014 (64)	Yes	NR	Yes	NR	NR	NR
Hankiewics-Ziolkowska 2014 (65)	Yes	NR	Yes	NR	NR	NR
Manxhuka-Kerliu 2014 (66)	Yes	NR	Yes	Yes	No	No

**Table 7** (continued)

Table 7 (continued)

Study (year of publication)	Histologic characteristics					
	C-kit/CD117	DOG1	CD34/PDGFR-alfa	Smooth muscle actin	S100	Desmin
Fukuyama 2014 (67)	Yes	NR	NR	NR	NR	NR
Mrak 2015 (70)	Yes	No	No	No	No	No
Kumar 2015 (71)	Yes	NR	Yes	NR	NR	NR
Chung 2015 (72)	21	NR	14	NR	10	No
Graziosi 2015 (73)	Yes	Yes	NR	NR	NR	NR
Boselli 2016 (74)	Yes	Yes	Yes	No	No	No
	Yes	Yes	No	No	No	No
	No	No	No	No	No	Yes
Caruso 2016 (75)	Yes	No	Yes	No	No	No
Jones 2016 (76)	Yes	NR	Yes	No	No	No
Turculeț 2016 (77)	Yes	Yes	No	Yes	No	NR
Huo 2016 (78)	Yes	No	Yes	No	No	No
Mori 2016 (79)	Yes	Yes	No	No	No	No
Valli 2016 (80)	Yes	Yes	Yes	No	No	No
Crocetti 2016 (81)	Yes	Yes	NR	NR	NR	NR
Thillai 2017 (82)	Yes	No	Yes	No	No	No
Elston 2017 (83)	Yes	Yes	No	No	No	No
Vasile 2017 (84)	Yes	NR	Yes	No	No	NR
Zioni 2017 (85)	Yes	NR	NR	NR	NR	NR
Perfetti 2017 (86)	Yes	Yes	Yes	No	No	No
Hakozaki 2017 (87)	Yes	NR	Yes	NR	No	NR

Table 8 Characteristics of included studies: local resection

Study (year of publication)	Tumor extension out duodenum	Type of surgical access	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Multiple visceral resection	Survival (years; months)
Sawaki 2003 (37)	Duodenum	Open	No	Yes	No	NR
Sakamoto 2003 (38)	Duodenum	Open	No	No	No	NR
Kurihara 2005 (39)	Duodenum	Open	No	No	No	NR
Goh 2005 (40)	Duodenum	Open	No	No	No	NR
	Duodenum, transverse mesocolon, small bowel mesentery	Open	No	No	Small bowel resection and transverse colectomy	NR
Cavallini 2005 (41)	Duodenum	Open	No	No	No	4 y

Table 8 (continued)



Table 8 (continued)

Study (year of publication)	Tumor extension out duodenum	Type of surgical access	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Multiple visceral resection	Survival (years; months)
Vu 2005 (42)	Duodenum	Open	NR	NR	NR	NR
Towu 2006 (43)	Duodenum	Open	No	No	No	NR
Kwon 2007 (44)	Duodenum	Open	No	No	No	NR
Graham 2007 (45)	Duodenum	NR	No	Yes	No	NR
Gupta 2007 (46)	Duodenum	Open	No	No	No	NR
Mohiuddin 2007 (47)	Duodenum	Open	No	Yes	No	42 m
Fernández Salazar 2007 (48)	Duodenum	Open	NR	NR	No	NR
Mennigen 2008 (49)	Duodenum	Open	No	No	No	NR
Takahashi 2009 (50)	Duodenum	NR	No	No	No	NR
Takeuchi 2009 (51)	Duodenum	Open	No	No	No	24 m
Seçkin 2009 (52)	Duodenum	Open	NR	NR	No	NR
Hirashima 2009 (53)	Duodenum	Open	No	No	No	NR
Mehta2011 (54)	Duodenum	NR	No	No	No	NR
Chung 2011 (55)	Duodenum	Open	No	No	No	42 m
		Open	No	No	No	13 m
Cameron 2011 (56)	Duodenum	Open	No	Yes	No	10 y
Kato 2011 (57)	Duodenum	Laparoscopic- endoscopic	NR	NR	No	NR
Chen 2012 (58)	Duodenum	Open	NR	NR	No	NR
El-Gendi 2012 (59)	Duodenum	Open	No	10	No	45 m
Acar 2013 (60)	Duodenum	Open	No	No	No	NR
Mouaqit 2013 (61)	Duodenum	Open	No	Yes	No	24 m
Mokhtare 2013 (62)	Duodenum	Open	No	No	No	NR
Shaw 2013 (63)	Duodenum	Open	NR	NR	No	NR
Ueda 2014 (64)	Duodenum	Laparoscopic	No	No	No	NR
Hankiewics-Ziolkowska 2014 (65)	Duodenum	Open	No	No	No	NR
Manxhuka-Kerliu 2014 (66)	Duodenum, jejunum	Open	No	No	No	NR
Fukuyama 2014 (67)	Duodenum	Open	Yes	Yes	No	12 m
Borgaonkar 2015 (68)	Duodenum	Open	NR	2	No	NR
Jones 2015 (69)	Duodenum	Open	Yes	Yes	No	NR
Mrak 2015 (70)	Duodenum	Open	No	Yes	No	NR
Kumar 2015 (71)	Duodenum	Open	No	No	No	6 m

Table 8 (continued)

Table 8 (continued)

Study (year of publication)	Tumor extension out duodenum	Type of surgical access	Neoadjuvant treatment with Imatinib	Postoperative treatment with Imatinib	Multiple visceral resection	Survival (years; months)
Chung 2015 (72)	Duodenum	19 open, 2 laparoscopic	No	No	No	NR
Graziosi 2015 (73)	Duodenum	Open	No	Yes	No	NR
Boselli 2016 (74)	Duodenum	Open	No	No	No	NR
	Duodenum	Open	No	No	No	NR
	Duodenum	Open	No	No	No	NR
Caruso 2016 (75)	Duodenum, jejunum	Open	No	No	No	NR
Jones 2016 (76)	Duodenum	Open	No	No	No	NR
Turculeț 2016 (77)	Duodenum	Open	No	No	No	NR
Huo 2016 (78)	Duodenum	Open	No	No	No	12 m
Mori 2016 (79)	Duodenum	Open	No	No	No	NR
Valli 2016 (80)	Duodenum	Open	Yes	No	No	NR
Crocetti 2016 (81)	NR	Open	Yes	No	No	NR
Thillai 2017 (82)	Duodenum, adherent to ascending colon and Gerota's fascia	Open	No	Yes	No	NR
Elston 2017 (83)	Duodenum	Open	No	No	No	NR
Vasile 2017 (84)	Duodenum	Open	No	Yes	Gastric antrum, distal common bile duct, pancreatic head	24 m
Zioni 2017 (85)	Duodenum	Laparoscopic	No	No	No	1,5 y
Perfetti 2017 (86)	Duodenum	Open	Yes	Yes	No	1,5 y
Hakozaki 2017 (87)	Duodenum	Open	No	No	No	NR

NR, information not reported.

Treitz and end-to-side duodenojejunosomy with first jejunal loop (99);

- ❖ segmental resection with end-to-side or end-to-end duodenojejunosomy to the left of the superior mesenteric vessels (66,72,75) segmental resection, closure of distal stump on the right of superior mesenteric vessels, papilloplasty, end-to-side duodenojejunosomy, transcystic tube and transjejunal tubes draining the main pancreatic duct and decompressing duodenum (38);
- ❖ segmental resection of third and fourth part of duodenum, closure of second portion stump and side-to-end duodenojejunosomy (49);

## Discussion

Duodenal GIST is typically seen in individuals between the ages of 60–70 years old. There were few papers reporting an occurrence of dGIST in children and teenagers, all of which manifested with severe bleeding (80,97,101). The median age of the patients with dGIST is 56 years with a slight preponderance of males (54% *vs.* 46%) (1,5).

### Risk factors

Most cases are sporadic, whereas familial clustering is reported in only 1.5–4% of the cases (69,102). Familial

GIST is an autosomal dominant condition caused by germline mutations of *cKIT* or *PDGFRA* and manifests at earlier age. GIST may be part of Carney's triad (gastric GIST, paraganglioma, and pulmonary chondroma) or Carney-Stathakis syndrome, which are caused by germline mutations in *SDH* (89). It is a well-known fact that von Recklinghausen neurofibromatosis type 1 (NF 1) is associated with an increased risk for GIST. GISTs in NF 1 occur in 6–7%, affect duodenum in 22–31%, and tend to be smaller, multiple and with lower mitotic count and occur at younger age (89,103,104). Hakoziaki *et al.* described a case of dGIST and rectal cancer in a patient with NF 1 (87). Malignant carcinoid of ampulla Vateri and bilateral feochromocytoma have been also described with low responses to imatinib (105). Few papers reported extremely rare combination with pancreatic neuroendocrine neoplasia (64) and somatostatinoma in NF 1 (35) and Brunner's gland cysts (78).

### Localization

Most dGISTs develop in the second (59–63%) and third portion (22%) of duodenum (55,106), whereas the first and fourth are less frequently affected. GIST of ampulla Vateri is extremely rare and according to the review of Kobayashi *et al.* only 12 cases have been described till 2014 (30). In all cases pain, jaundice and melena were the most frequent symptoms. Despite its rarity, this localization requires meticulous differential diagnosis (neuroendocrine tumors, carcinoma, paraganglioma) because usually the radical treatment requires PD (107).

### Histopathology and molecular characteristics

Histologically, dGISTs do not differ from other GIST localizations. The most frequently reported pattern was spindle cell (67%), but epithelioid (11%), pleomorphic, mixed (22%), hemangioma-like or hemangiopericytoma-like patterns can also be seen (9,13,55,93,99,101,108-114). Immunohistochemistry staining of the specimens revealed the following distribution of the markers: CD117 (*c-kit*) (92–100%), and less frequently CD34 (54–70%), smooth muscle actin (20–30%), S-100 protein (10–20%), and neurofilament 68 (14%), DOG1 protein (6%) (9,12,65,94,115). Lack of DOG-1 expression was associated with poor prognosis in a recent study with 332 patients (93). The differential diagnosis includes fibromatosis (110), schwannomas, leiomyomas,

inflammatory fibroid tumors, solitary fibrous tumor, mesenteric sclerosing fibrotic lesions, sarcomas, metastasis from malignant melanoma, glomus tumors, paragangliomas, ectopic pancreas (65,92,98).

At the molecular level, approximately 96% of GISTs have *cKIT* (CD117) mutation, typically in exon 11, but mutations in exons 9, 13, and 17 may also occur (101). Exon 11 mutations carry a better prognosis and respond well to standard dose (400 mg/dayly) of imatinib (116). Some authors reported worse outcomes in exon 9 mutation (102). Approximately 8% of the cases have *PDGFRA* mutation, which is mutually exclusive with *cKIT* but is associated with longer relapse-free interval (43,59,116). Some *PDGFRA* are resistant to imatinib (65). The so-called wild type is observed in 10–15% of the cases and in 90% of children. It is caused by germline mutations in *BRAF* V600E, *RAS* family or succinate dehydrogenase subunits (*SDH* A-D). GISTs in NF1 are usually wild type with possible implication of *RAS*-*MAPK* pathway (104,117-119). This subset of GISTs is relatively resistant to the treatment with imatinib. Further large studies are needed to elucidate the individual prognosis according to the mutational status.

### Clinical manifestation

Approximately 70% of the cases manifest with symptoms, while 21% are found incidentally and 10% on autopsy (120). Our analysis revealed that in contrast to the other localizations the most frequent manifestation of dGIST is the upper gastrointestinal bleeding, which is in accordance with the literature (21,74,121). Constant/intermittent dull pain/discomfort or abdominal palpable mass are less frequent initial symptoms (10,32,71). Rarely, however, dGISTs may have various initial presentations leading to misdiagnosis. Millonig *et al.* reported extrahepatic cholestasis and Takotsudo cardiomyopathy in a patient with undiagnosed NF1 (122). The extremely rare dGIST of ampulla Vateri can also manifest with jaundice (30). In certain cases, extramural growth may mimic pancreatic head tumor (26,44,84,97), large duodenal cyst (61), bleeding or uncomplicated duodenal diverticula (33, 46). Wall *et al.* reported a rare case with duodenal-jejunal intussusception (23). Lin *et al.* described two cases with bleeding small dGISTs (<2 cm) initially misdiagnosed as hemobilia (123). A case mimicking refractory peptic ulcer treated for 8 years was also described (76), even hypercalcemia due to elevated serum calcitriol in metastatic disease as well (124).

### **Risk stratification**

The recurrence risk and survival outcomes for dGIST are difficult to be determined based solely on the histopathological characteristics. Multiple factors have been proposed as predictive of survival outcomes, including tumor location and size, mitotic rate, kinase mutational status, and incidence of tumor rupture. However, tumor size and mitotic rate are the two most widely accepted risk indicators (125,126).

The Fletcher's and subsequently the modified National Institute of Health classification are the most popular risk classifications (95). Miettinen *et al.*, based on the follow-up of 140 cases with dGIST, proposed a distinct risk stratification of dGIST in which group 1 (<2 cm and <5 mitoses) is considered benign, whereas group 6 (>5 cm and >5 mitoses) carries an extremely poor prognosis with 86% mortality (18/21) within 21 months (9). The overall mortality in their series was 34%. This classification was externally validated by the French Sarcoma group in a series with 114 patients (119). The overall doubling time for dGIST is 17 months in comparison to leiomyoma (231 months). According to the risk group the doubling time is 24, 17 and 4 months in low, intermediate and high risk dGISTs (115).

Generally, there are contradictory results in the literature regarding the prognosis of dGIST in comparison to the other localizations. Certain authors reported worse outcome in comparison to small bowel and gastric GISTs (43), whereas others reported similar prognosis with small bowel localization. The more recent, population-based study of Guller *et al.* reported similar survival in gastric, duodenal and small bowel GISTs in contrast to colonic and extraviseral localizations (127). This finding was corroborated after subgroup analysis of two periods (1998–2004 *vs.* 2005–2011) thus eliminating a possible confounding factor associated with the implementation of imatinib therapy.

### **Diagnosis**

Although it may be straightforward in most cases, certain considerations should be kept in mind due to their occasionally misleading manifestation (128). Differential diagnosis includes adenocarcinoma, endocrine tumors, benign tumors and rare entities such as intrabdominal fibromatosis (110), ectopic pancreas (129), and Brunner's gland cysts (78).

Because most of dGISTs present with acute bleeding or chronic anemia endoscopic evaluation of upper gastrointestinal tract should be the first step (101). It allows for biopsy and can be also therapeutic. In case of failure, transarterial embolization is a method of choice, either as definitive hemostasis or as a bridging procedure before surgical intervention (39). Occasionally, endoscopy may be misleading, especially in small intramural lesions without mucosal involvement (ulceration or central depression) or located near the papilla Vateri. In such cases of diagnostic uncertainty, endoscopic ultrasound (EUS) is an invaluable modality. Usually dGISTs appears as hypoechoic and well-vascularized lesion. Two small series reported a significant correlation between the presence of intratumoral vessels on contrast-enhanced or color Doppler EUS and the malignant potential of dGISTs (130,131). EUS provides a precise evaluation of the size, border, layer of origin, echogenicity and heterogeneity of the lesions thus facilitating the differential diagnosis (lipomas, hemangiomas, ectopic pancreas, and cysts) and decision-making process (112). Additionally, EUS allows also fine needle or trucut biopsy with 100% specificity and 84% sensitivity (92,112,131,132). Percutaneous biopsy should be avoided when possible because of the risk for tumor spillage and dissemination.

Abdominal US is useful screening tool in the cases with dull pain in upper abdomen, but computed tomography (CT) and magnetic resonance imaging (MRI) are mandatory to make an exact staging and preoperative planning of surgery (107,113). On CT dGISTs appear as well-defined, heterogeneously enhanced, hypervascular mass with prominent feeding arteries and intra- or extramural growth (107,114). In contrast, the ectopic pancreas has intraluminal growth and ill-defined border with enhancement of the overlying mucosa (129). The periampullary pancreatic cancer is usually hypodense on arterial phase with concomitant pancreatic duct dilatation, periampullary neuroendocrine tumors reveal hypervascular enhancement, calcifications, lack of ductal obstruction, central necrosis and cystic degeneration, whereas solid periampullary tumors demonstrate heterogeneous hypoenhancement in both phases (114). Fluorodeoxyglucose positron emission tomography (FDG-PET) is not routine tool but can be useful to monitor the effect from imatinib treatment and follow-up (133,134).

### **Surgical treatment**

The ESMO guideline suggests that tumors under 2 cm has low aggressive behaviour and therefore could be followed

annually by endoscopic ultrasound, “although an evidence-based optimal surveillance policy is lacking” (109). In contrast to the other localizations, in dGIST there is no uniformly adopted surgical strategy because of the low incidence, lack of enough experience, and the complex anatomy of the duodenum. Therefore, individually tailored surgical approach is recommended. In fact, there are three main surgical options: pancreatoduodenectomy (PD), wedge resection and segmental resection.

PD is indicated in the cases with involvement of major duodenal papilla, pancreas or pancreatic duodenal wall and is required in 20–40% (101,135). In some cases, it can be performed successfully in emergency setting due to life-threatening bleeding (10,24,111) even with second-stage pancreatojejunostomy (17). Tien *et al.* found that size >5 cm and preoperative diagnosis of dGIST were more frequently treated by PD (89). In another series, the patients with PD had larger tumor size and higher mitotic count (118). A recent meta-analysis of seven comparative studies confirmed categorically the above-mentioned findings (89).

Imatinib mesylate have played a key role as a neoadjuvant therapy in the management of GISTs (21,122). In locally advanced disease neoadjuvant imatinib may downstage the tumor to allow R0 resection or even an organ preserving intervention (11,122,136–138) despite the risk of bleeding (111). Ludvigsen *et al.* reported a successful PD en-block with right kidney and suprarenal gland (19). Fukuyama *et al.* reported even avoidance of PD after downstaging neoadjuvant therapy with imatinib (67). Recently, successful organ-preserving duodenectomies after neoadjuvant therapy in 9 of 10 cases was reported (92) similarly to other authors (82).

PD is, however, burdened by significant morbidity and longer hospital stay in comparison to local resection, probably due to the “soft” pancreas and small caliber of pancreatic duct (116,118,119). Moreover, several studies reported no significant difference in recurrence rate and disease specific survival between limited resection and pancreatoduodenectomy (55,97,101,104), which is categorically supported from the meta-analysis of Chok *et al.* (122) and the experience of the French Sarcoma Group (119).

R0 resection with 1–2 cm clear margin is sufficient treatment and lymph node dissection is not recommended due to the low incidence of lymphatic metastases (102). There are several options for local resection and surgical reconstruction according to the localization of the tumour. The rupture of GIST during the surgery should be avoided because is associated with nearly 100% risk for recurrence (125). Therefore, the good knowledge of anatomy, gentle handling

of the tissues, and careful dissection of duodenal wall from the inferior border of pancreas, meticulous hemostasis and knowing of the possible options for duodenal reconstruction are mandatory for successful outcomes.

Laparoscopic resection is feasible and safe with reported subtotal resection with side-to-side duodenojejunostomy (100) and wedge resection (85,139). Tanaka *et al.* reported eighth cases successful segmental resections of duodenojejunal junction with side-to-side anastomosis comparable with 11 open procedures (140). Some authors reported laparoscopically assisted endoscopic submucosal resection of 20 mm GIST located in the third portion (57).

There are six reported cases of dGIST localized in first and second portion removed by robotic surgery: wedge resection (140–142) and segmental resection with side-to-side duodenojejunostomy (143).

Although in most cases wedge and segmental resections could be relatively straightforward, severe complications may occur such as acute pancreatitis, pancreatic fistula, significant blood loss and anastomotic stenosis (117,118). Delayed gastric emptying treating with gastrojejunostomy and the fearsome anastomotic failure necessitated secondary PD (101) and injury of the mesenteric root managed by total enterectomy have been described in the literature (132). The pitfalls in segmental resections are associated with the superior mesenteric vessels lying on the third portion, adjacent pancreas and the common blood supply, necessity to preserve ampulla Vateri (117,107).

## Conclusions

dGIST is a very rare entity. It may be asymptomatic or may involve symptoms of upper GI bleeding and abdominal pain at presentation. Because of the misleading clinical presentation the differential diagnosis may be difficult. Gastrointestinal endoscopy is the most common initial diagnostic procedure while abdominal and thoracic CT scan are mandatory for accurate oncologic staging and surgical planning.

Tumours smaller than 2 cm have a low biological aggressiveness and can be followed annually by endoscopic ultrasound; the biggest tumors should undergo radical surgery (R0) (144). In contrast to the other localizations, dGIST have no uniformly adopted surgical strategy because of the low incidence, lack of experience, and complex anatomy of the duodeno-pancreatic region. Therefore, individually tailored surgical approach is recommended. R0 resection with 1–2 cm clear margin is sufficient treatment

and lymph node dissection is not recommended due to the low incidence of lymphatic metastases. Tumor rupture should be avoided.

### Acknowledgements

None.

### Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

*Ethical Statement:* The review of literature does not need evaluation by Ethical committee and by Institutional/Regional/National Ethics committees/boards.

### References

1. Yang WL, Yu JR, Wu YJ, et al. Duodenal gastrointestinal stromal tumor: clinical, pathologic, immunohistochemical characteristics, and surgical prognosis. *J Surg Oncol* 2009;100:606-10.
2. Cirocchi R, Patelli S, Castellani E, et al. Right hemicolectomy plus pancreaticoduodenectomy vs partial duodenectomy in treatment of locally advanced right colon cancer invading pancreas and/or only duodenum. *Surg Oncol* 2014;23:92-8.
3. Hirota S, Isozaki K, Moriyama Y, et al. Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. *Science* 1998;279:577-80.
4. Biasco G, Velo D, Angriman I, et al. Gastrointestinal stromal tumors: report of an audit and review of the literature. *Eur J Cancer Prev* 2009;18:106-16.
5. Miettinen M, Lasota J. Gastrointestinal stromal tumors (GISTs): definition, occurrence, pathology, differential diagnosis and molecular genetics. *Pol J Pathol* 2003;54:3-24.
6. Nilsson B, Bummig P, Meis-Kindblom JM, et al. Gastrointestinal stromal tumors: the incidence, prevalence, clinical course, and prognostication in the pre-Imatinib mesylate era – a population-based study in western Sweden. *Cancer* 2005;103:821-9.
7. Sawaki A. Rare gastrointestinal stromal tumors (GIST): omentum and retroperitoneum. *Transl Gastroenterol Hepatol* 2017;2:116.
8. Agaimy A, Wunsch PH. Gastrointestinal stromal tumors: a regular origin in the muscularis propria, but an extremely diverse gross presentation. *Langenbecks Arch Surg* 2006;391:322-9.
9. Miettinen M, Kopczynski J, Makhlof HR, et al. Gastrointestinal stromal tumors, intramural leiomyomas and leiomyosarcomas in the duodenum. *Am J Surg Pathol* 2003;27:625-41.
10. Winfield RD, Hochwald SN, Vogel SB, et al. Presentation and management of gastrointestinal stromal tumors of the duodenum. *Am Surg* 2006;72:719-22; discussion 722-3.
11. Johnston FM, Kneuert PJ, Cameron JL, et al. Presentation and management of gastrointestinal stromal tumors of the duodenum: A multi-institutional analysis. *Ann Surg Oncol* 2012;19: 3351-60.
12. Kindblom LG, Remotti HE, Aldenborg F, et al. Gastrointestinal pacemaker cell tumor (GIPACT): gastrointestinal stromal tumors show phenotypic characteristics of the interstitial cells of Cajal. *Am J Pathol* 1998;152:1259-69.
13. Takahashi Y, Noguchi T, Takeno S, et al. Gastrointestinal stromal tumor of the duodenal ampulla: report of a case. *Surg Today* 2001;31:722-6.
14. Uchida H, Sasaki A, Iwaki K, et al. An extramural gastrointestinal stromal tumor of the duodenum mimicking a pancreatic head tumor. *J Hepatobiliary Pancreat Surg* 2005;12:324-7.
15. Hompes D, Topal B, Ectors N, et al. Gastro-intestinal stromal tumour of the duodenum: extreme presentation in two cases. *Acta Chir Belg* 2004;104:110-3.
16. Kim SH, Kim JH, Baik GH, et al. Malignant gastrointestinal stromal tumor of the ampulla of Vater: a case report. *Korean J Gastroenterol* 2004;43:66-70.
17. Sakakima Y, Inque S, Fuji T, et al. Emergency pylorus-preserving pancreatoduodenectomy followed by second-stage pancreatojejunostomy for a gastrointestinal stromal tumor of the duodenum with an intratumoral gas figure: report of a case. *Surg Today* 2004;34:701-5.
18. Sakakura C, Hagiwara A, Soga K, et al. Long-term survival of a case with multiple liver metastases from duodenal gastrointestinal stromal tumor drastically reduced by the treatment with imatinib and hepatectomy. *World J Gastroenterol* 2006;12:2793-7.
19. Ludvigsen L, Toxverd A, Mahdi B, et al. Successful resection of an advanced duodenal gastrointestinal stromal tumor after down-staging with Imatinib: report of a case. *Surg Today* 2007;37:1105-9.
20. Fiore M, Palassini E, Fumagalli E, et al. Preoperative imatinib mesylate for unresectable or locally advanced primary gastrointestinal stromal tumors (GIST). *Eur J Surg Oncol* 2009;35:739-45.

21. Hecker A, Hecker B, Bassaly B, et al. Dramatic regression and bleeding of a duodenal GIST during preoperative imatinib therapy: case report and review. *World J Surg Oncol* 2010;8:47.
22. Frampton AE, Bong JJ, Kyriakides C, et al. En bloc resection of the pancreatic head and second part of duodenum for a duodenal gastrointestinal stromal tumor: a multi-media report. *JOP* 2010;11:396-400.
23. Wall ML, Ghallab MA, Farmer M, et al. Gastrointestinal stromal tumor presenting with duodenal-jejunal intussusception: a case report. *Ann R Coll Surg Engl* 2010;92:e32-4.
24. Machado NO, Chopra PJ, Al-Haddabi IH, et al. Large duodenal gastrointestinal stromal tumor presenting with acute bleeding managed by a Whipple resection. A review of surgical options and the prognostic indicators of outcome. *JOP* 2011;12:194-9.
25. Morcos B, Al-Ahmad F. A large gastrointestinal stromal tumor of the duodenum: a case report. *J Med Case Rep* 2011;5:457.
26. Singh S, Paul S, Khandelwal P, et al. Duodenal GIST presenting as large pancreatic head mass: an uncommon presentation. *JOP* 2012;13:696-9.
27. Koontz MZ, Visser BM, Kunz PL. Neoadjuvant imatinib for borderline resectable GIST. *J Natl Compr Canc Netw* 2012;10:1477-82.
28. Blandamura S, Alessandrini L, Bertorelle R, et al. Multiple sporadic gastrointestinal stromal tumors concomitant with ampullary adenocarcinoma: a case report with KIT and PDGFRA mutational analysis and miR-221/222 expression profile. *Pathol Res Pract* 2014;210:392-6.
29. Bormann F, Wild W, Aksoy H, et al. A Pancreatic Head Tumor Arising as a Duodenal GIST: A Case Report and Review of the Literature. *Case Rep Med* 2014;2014:420295.
30. Kobayashi M, Hirata N, Nakaji S, et al. Gastrointestinal stromal tumor of the ampulla of Vater. *World J Gastroenterol* 2014;20:4817-21.
31. Parisi A, Desiderio J, Trastulli S, et al. Robotic pancreaticoduodenectomy in a case of duodenal gastrointestinal stromal tumor. *World J Surg Oncol* 2014;12:372.
32. Bhambare MR, Pandya JS, Waghmare SB, et al. Gastrointestinal stromal tumor presenting as palpable mass: a rare entity. *World J Gastrointest Surg* 2015;7:98-101.
33. Okasha HH, Amin KM, Al-Shazli M, et al. A duodenal gastrointestinal stromal tumor with a large central area of fluid and gas due to fistulization into the duodenal lumen, mimicking a large duodenal diverticulum. *Endosc. Ultrasound* 2015;4:253-6.
34. Niikura R, Serizawa T, Yamada A, et al. Safety of Regular-Dose Imatinib Therapy in Patients with Gastrointestinal Stromal Tumors Undergoing Dialysis. *Case Rep Gastroenterol* 2016;10:17-23.
35. Yamamoto R, Kato S, Maru T, et al. The coexistence of somatostatinoma and gastrointestinal stromal tumor in the duodenum of a patient with von Recklinghausen's disease. *Intern Med* 2016;55:617-22.
36. Tornambe A, Tornambe G. Duodenal gastrointestinal stromal tumor. A case report. *Ann Ital Chir.* 2017;6.
37. Sawaki A, Ohashi K, Yamao K, et al. Effect of a tyrosine kinase inhibitor STI571 in a patient with hepatic metastases from a duodenal gastrointestinal stromal tumor. *J Gastroenterol* 2003;38:690-4.
38. Sakamoto Y, Yamamoto J, Takahashi H, et al. Segmental resection of the third portion of the duodenum for a gastrointestinal stromal tumor: a case report. *Jpn J Clin Oncol* 2003;33:364-6.
39. Kurihara N, Kikuchi K, Tanabe M, et al. Partial resection of the second portion of the duodenum for gastrointestinal stromal tumor after effective transarterial embolization. *Int J Clin Oncol* 2005;10:433-7.
40. Goh BK, Chow PKH, Ong H, et al. Gastrointestinal stromal tumor involving the second and third portion of the duodenum: treatment by partial duodenectomy and Roux-en-Y duodenectomy. *J Surg Oncol* 2005;91:273-5.
41. Cavallini M, Cecera A, Ciardi A, et al. Small periampullary duodenal gastrointestinal stromal tumor treated by local excision: report of a case. *Tumori* 2005;91:264-6.
42. Vu HA, Xinh PT, Kikushima M, et al. A recurrent duodenal gastrointestinal stromal tumor with a frameshift mutation resulting in a stop codon in KIT exon 13. *Genes Chromosomes Cancer* 2005;42:179-83.
43. Towu E, Stanton M. Gastrointestinal stromal tumor presenting with severe bleeding: a review of the molecular biology. *Pediatr Surg Int* 2006;22:462-4.
44. Kwon SH, Cha HJ, Jung SW, et al. A gastrointestinal stromal tumor of the duodenum masquerading as a pancreatic head tumor. *World J Gastroenterol* 2007;13:3396-9.
45. Graham J, Debiec-Rychter M, Corless CL, et al. Imatinib in the management of multiple gastrointestinal stromal tumors associated with a germline KIT K642E mutation. *Arch Pathol Lab Med* 2007;131:1393-6.
46. Gupta N, Schrimmer BD, Mishra R, et al. Malignant GIST masquerading as a bleeding duodenal diverticulum.

- Endoscopy 2007;39:E142-3.
47. Mohiuddin K, Nizami S, Munir A, et al. Metastatic duodenal GIST: role of surgery combined with imatinib mesylate. *Int Semin Surg Oncol* 2007;4:9.
  48. Fernández Salazar LI, Alvarez Gago T, Sanz Rubiales A, et al. Gastrointestinal stromal tumors (GISTs): clinical aspects. *Rev Esp Enferm Dig* 2007;99:19-24.
  49. Mennigen R, Wolters HH, Schule B, et al. Segmental resection of the duodenum for gastrointestinal stromal tumor (GIST). *World J Surg Oncol* 2008;6:105.
  50. Takahashi Y, Shimizu S, Sakurai S, et al. Gastrointestinal stromal tumor in the duodenum exhibiting hemangiopericytoma-like histological pattern. *Pathol Int* 2009;59:98-101.
  51. Takeuchi H, Matsumoto T, Kusumoto T, et al. Duodenal Gastrointestinal Stromal Tumor Treated by Wedge Resection in a Patient with Neurofibromatosis Type 1: Report of a Case and Review of the Japanese Literature. *Case Rep Gastroenterol* 2009;3:343-9.
  52. Seçkin Y, Koyuncu A, Yönel O, et al. Duodenal stromal tumor: a rare and overlooked cause of massive gastrointestinal bleeding. *Turk J Gastroenterol* 2009;20:150-1.
  53. Hirashima K, Takamori H, Hirota M, et al. Multiple gastrointestinal tumors in neurofibromatosis type 1: report of a case. *Surg Today* 2009;39:979-83.
  54. Mehta C, Gumaste VV, Leytin A, et al. An unusual cause of upper gastrointestinal bleeding: duodenal GIST. A case report and literature review. *Acta Gastroenterol Belg* 2011;74:347-51.
  55. Chung JC, Kim HC, Chu CW. Segmental resection with duodenojejunostomy of gastrointestinal stromal tumor involving the duodenum. *J Kor Surg Soc* 2011;80:12-6.
  56. Cameron S, Schaefer IM, Schwoerer H, et al. Ten Years of Treatment with 400 mg Imatinib per Day in a Case of Advanced Gastrointestinal Stromal Tumor. *Case Rep Oncol* 2011;4:505-11.
  57. Kato M, Nakajima K, Nishida T, et al. Local resection by combined laparoendoscopic surgery for duodenal gastrointestinal tumor. *Diagn Ther Endosc* 2011;2011:645609.
  58. Chen J, Gundara JS, Haddad R, et al. Clinicopathological and molecular aspects of foregut gastrointestinal tumors. *ANZ J Surg* 2014;84:52-8.
  59. El-Gendi A, El-Gendi S, El-Gendi M. Feasibility and oncological outcomes of limited duodenal resection in patients with primary nonmetastatic duodenal GIST. *J Gastrointest Surg* 2012;16:2197-202.
  60. Acar F, Sahin M, Ugras S, et al. A gastrointestinal stromal tumor of the third portion of the duodenum treated by wedge resection: a case report. *World J Gastrointest Surg* 2013;5:332-6.
  61. Mouaqit O, Chbani L, Maazaz K, et al. A large gastrointestinal stromal tumor of the duodenum treated by partial duodenectomy with Roux-en-Y duodenojejunostomy: a case report. *J Med Case Rep* 2013;7:184.
  62. Mokhtare M, Taghvaei T, Fakheri HT. Acute bleeding in duodenal gastrointestinal stromal tumor. *Middle East J Dig Dis* 2013;5:47-51.
  63. Shaw A, Jeffery J, Dias L, et al. Duodenal wedge resection for large gastrointestinal stromal tumor presenting with life-threatening haemorrhage. *Case Rep Gastrointest Med* 2013;2013:562642.
  64. Ueda K, Hijioka M, Lee L, et al. A synchronous neuroendocrine tumor and duodenal gastrointestinal stromal tumor. *Intern Med* 2014;53:2483-8.
  65. Hankiewicz-Ziołkowska K, Soboń M, Szyłberg T, et al. Duodenal bulb tumour of unknown origin. *Prz Gastroenterol* 2014;9:365-70.
  66. Manxhuka-Kerliu S, Sahatciu-Meka V, Kerliu I, et al. Small intestine gastrointestinal stromal tumor in a young adult woman: a case report and review of the literature. *J Med Case Rep* 2014;8:321.
  67. Fukuyama K, Fujikawa T, Kuramitsu S, et al. Successful treatment of bleeding large duodenal gastrointestinal stromal tumor in a patient under dual antiplatelet therapy after recent drug-eluting coronary stent implantation. *BMJ Case Rep* 2014;2014:bcr2014204462.
  68. Borgaonkar V, Deshpande S, Borgaonkar V, et al. Gastrointestinal Stromal Tumor-Single-Center Experience with Review of the Literature. *Indian J Surg* 2015;77:678-81.
  69. Jones DH, Caracciolo JT, Hodul PJ, et al. Familial gastrointestinal stromal tumor syndrome: report of 2 cases with KIT Exon 11 mutation. *Cancer Control* 2015;22:102-8.
  70. Mrak K, Liegl-Atzwanger B, Haybaeck J, et al. Surgical Management of Duodenal Gastrointestinal Stromal Tumors: A Case Report. *Anticancer Res* 2015;35:6321-4.
  71. Kumar A, Jakhmola CK, Chauhan SS, et al. Atypical presentation of gastrointestinal stromal tumor masquerading as a large duodenal cyst: a case report. *Int J Surg Case Rep* 2015;9:123-6.
  72. Chung JC, Kim HC, Hur SM. Limited resections for duodenal gastrointestinal stromal tumors and their oncologic outcomes. *Surg Today* 2016;46:110-6.
  73. Graziosi L, Marino E, Ludovini V, et al. Unique case of



- sporadic multiple gastro intestinal stromal tumour. *Int J Surg Case Rep* 2015;9:98-100.
74. Boselli C, Cirocchi R, Gemini A, et al. Urgency surgical treatment for duodenal GISTs: analysis of aged patients and review of the literature. *Aging Clin Exp Res* 2017;29:1-6.
  75. Caruso F, Nencioni M, Zefelippo A, et al. Is duodenojejunal anastomosis to the left of the superior mesenteric vessels a feasible option for tumors of the angle of Treitz. *Tumori* 2016;102:71-3.
  76. Jones JD, Oh S, Clark C, Pawa R. A Bleeding Duodenal GIST Masquerading as Refractory Peptic Ulcer Disease. *ACG Case Rep J* 2016;3:e189.
  77. Turculeț C, Ene D, Georgescu TF, et al. A rare case of upper digestive hemorrhage due to bleeding duodenal tumor. *Chirurgia (Bucur)* 2016;111:505-8.
  78. Huo X, Wei J, Liu X, et al. Brunner's gland cyst in combination with gastrointestinal stromal tumor: a case report. *Oncology Letters* 2016;11:3409-12.
  79. Mori N, Ichikawa T, Hashimoto J, et al. Cholangiolocellular Carcinoma of the Liver Exhibiting High F-18 FDG Uptake. *Tokai. J Exp Clin Med* 2016;41:60-4.
  80. Valli PV, Valli C, Pfammatter T, et al. Life-threatening bleeding of a duodenal gastrointestinal stromal tumor in a teenager: a rare case report. *Endosc Int Open* 2016;4:E1244-6.
  81. Crocetti D, Sapienza P, Cisano C, et al. Pancreas preserving surgery for duodenal gastrointestinal stromal tumor removal. *Minerva Chirurgica* 2016;71:281-5.
  82. Thillai M, Alexander N, Paramasivam S, et al. Malignant duodenal GIST in a patient with situs inversus totalis – a rare association and brief review of literature. *J Gastrointest Cancer* 2017. [Epub ahead of print].
  83. Elston MS, Sehgal S, Dray M, et al. A Duodenal SDH-Deficient Gastrointestinal Stromal Tumor in a Patient With a Germline SDHB Mutation. *J Clin Endocrinol Metab* 2017;102:1447-50.
  84. Vasile D, Iancu G, Iancu RC, et al. Duodenal gastrointestinal stromal tumor presenting as pancreatic head mass – a case report. *Rom J Morphol Embryol* 2017;58:255-9.
  85. Zioni T, Dizengof V, Kirshtein B. Laparoscopic resection of duodenal gastrointestinal stromal tumour. *J Minim Access Surg* 2017;13:157-60.
  86. Perfetti V, Laurini E, Aulić S, et al. Molecular and functional characterization of a new 3' end KIT juxtamembrane deletion in a duodenal GIST treated with neoadjuvant Imatinib. *Oncotarget* 2017;8:56158-67.
  87. Hakozaiki Y, Sameshima S, Tatsuoka T, et al. Rectal carcinoma and multiple gastrointestinal tumors (GIST) of the small intestine in a patient with neurofibromatosis type 1: a case report. *World J Surg Oncol* 2017;15:160.
  88. Relles D, Baek J, Witkiewicz A, et al. Periampullary and duodenal neoplasms in neurofibromatosis type 1: two cases and an updated 20-year review of the literature yielding 76 cases. *J Gastrointest Surg* 2010;14:1052-61.
  89. Tien YW, Lee CY, Huang CC, et al. Surgery for gastrointestinal stromal tumors of the duodenum. *Ann Surg Oncol* 2010;17:109-14.
  90. Miki Y, Kurokawa Y, Hirao M, et al. Survival analysis of patients with duodenal gastrointestinal stromal tumors. *J Clin Gastroenterol* 2010;44:97-101.
  91. Yagishita A, Matsubayashi H, Kakushima N, et al. Gastrointestinal tumors of the duodenum: report of four cases. *Clin J Gastroenterol* 2011;4:162-6.
  92. Liang X, Yu H, Zhu LH, et al. Gastrointestinal stromal tumors of the duodenum: surgical management and survival results. *World J Gastroenterol* 2013;19:6000-10.
  93. Hoepfner J, Kuleman B, Marjanovic G, et al. Limited resection for duodenal gastrointestinal stromal tumors: surgical management and clinical outcome. *World J Gastrointest Surg* 2013;5:16-21.
  94. Yang F, Jin C, Du Z, et al. Duodenal gastrointestinal stromal tumor: clinicopathological characteristics, surgical outcomes, long term survival and predictors for adverse outcomes. *Am J Surg* 2013;206:360-7.
  95. Zhou B, Zhang M, Wu J, et al. Pancreatoduodenectomy versus local resections in the treatment of gastrointestinal tumors of the duodenum. *World J Surg Oncol* 2013;11:196.
  96. Duffaud F, Meeus P, Bachet JB, et al. Conservative vs. duodenopancreatectomy in primary duodenal gastrointestinal stromal tumors (GIST): a retrospective review of 114 patients from the French Sarcoma Group (FSG). *Eur J Surg Oncol* 2014;40:1369-75.
  97. Uçar AD, Oymaci E, Carti EB, et al. Characteristics of emergency gastrointestinal stromal tumor (GIST). *Hepatogastroenterology* 2015;62:635-40.
  98. Lv A, Qian H, Qiu H, et al. Organ-preserving surgery for locally advanced duodenal gastrointestinal tumor after neoadjuvant treatment. *Biosci Trends* 2017;11:483-9.
  99. Beham A, Schaefer IM, Cameron S, et al. Duodenal GIST: a single center experience. *Int J Colorectal Dis* 2013;28:581-90.
  100. Corcione F, Pirozzi F, Sciuto A, et al. Laparoscopic

- pancreas-preserving subtotal duodenectomy for gastrointestinal tumor. *Minim Invasive Ther Allied Technol* 2013;22:187-90.
101. Sugase T, Takahashi T, Nakajima K, et al. Clinicopathological characteristics, surgery and survival outcomes of patients with duodenal gastrointestinal stromal tumors. *Digestion* 2016;94:30-6.
  102. Joo MK, Park JJ, Kim H, et al. Endoscopic versus surgical resection of GI stromal tumors in the upper GI tract. *Gastrointest Endosc* 2016;83:318-26.
  103. Salvi PF, Lorenzon L, Caterino S, et al. Gastrointestinal stromal tumor associated with neurofibromatosis 1: a single center experience and systematic review of the literature including 252 cases. *Int J Surg Oncol* 2013;2013:398570.
  104. Sapalidis K, Panteli N, Strati TM, et al. Surgical management of gastrointestinal stromal tumors: a single centre's experience. *Hippokratia* 2015;19:73-5.
  105. Millonig G, Giesel FL, Reimann FM, et al. Gastrointestinal stromal tumor in a patient with undiagnosed neurofibromatosis type 1: an uncommon cause of extrahepatic cholestasis. *Z Gastroenterol* 2010;48:479-81.
  106. Shen C, Chen H, Yin Y, et al. Duodenal gastrointestinal stromal tumors: clinicopathological characteristics, surgery, and long-term outcome. *BMC Surg* 2015;15:98.
  107. Cirocchi R, Kelly MD, Griffiths EA, et al. A systematic review of the management and outcome of ERCP related duodenal perforations using a standardized classification system. *Surgeon* 2017;15:379-87.
  108. Chung JC, Chu CW, Cho GS, et al. Management and outcome of gastrointestinal stromal tumors of duodenum. *J Gastrointest Surg* 2010;14:880-3.
  109. Buchs NC, Bucher P, Gervaz P, et al. Segmental duodenectomy for gastrointestinal stromal tumor of the duodenum. *World J Gastroenterol* 2010;16:2788-92.
  110. Kamath AS, Sarr MG, Nagorney DM, et al. Gastrointestinal stromal tumor of the duodenum: single center institutional experience. *HPB* 2012;14:774-6.
  111. Uehara K, Hasegawa H, Ogiso S, et al. Gastrointestinal stromal tumor of the duodenum: diagnosis and treatment. *Geka* 2001;63:1058-61.
  112. Săftoiu A, Vilmann P, Hassan H. Utility of colour doppler endoscopic ultrasound evaluation and guided therapy of submucosal tumours of the upper gastrointestinal tract. *Ultraschall Med* 2005;26:487-95.
  113. DeWitt J, Emerson RE, Sherman S, et al. Endoscopic ultrasound-guided Trucut biopsy of gastrointestinal mesenchymal tumor. *Surg Endosc* 2011;25:2192-202.
  114. Matsui T, Mitsui H, Sekigawa K, et al. A case of a duodenal gastrointestinal stromal tumor diagnosed with the aid of diffusion-weighted magnetic resonance imaging. *Clin J Gastroenterol* 2009;2:384-7.
  115. Lee SY, Goh BKP, Sadot E, et al. Surgical strategy and outcomes in duodenal gastrointestinal stromal tumor. *Ann Surg Oncol* 2017;24:202-10.
  116. Crown A, Biehl TR, Rocha FG. Local resection for duodenal gastrointestinal stromal tumors. *Am J Surg* 2016;211:867-70.
  117. García-Molina FJ, Mateo-Vallejo F, Franco-Osorio J, et al. Surgical approach for tumors of the third and fourth part of the duodenum. Distal pancreas-preserving duodenectomy. *Int J Surg* 2015;18:143-8.
  118. Lupașcu C, Andronic D, Moldovanu R, et al. Treatment of gastrointestinal stromal tumors--initial experience. *Chirurgia (Bucur)* 2010;105:657-62.
  119. Colombo C, Ronellfisch U, Yuxin Z, et al. Clinical, pathological and surgical characteristics of duodenal gastrointestinal tumor and their influence on survival: a multi-center study. *Ann Surg Oncol* 2012;19:3361-7.
  120. Rabin I, Chikman B, Lavy R, et al. Gastrointestinal stromal tumors: a 19 year experience. *Isr Med Assoc J* 2009;11:98-102.
  121. Sorour MA, Kassem MI, El-Hamid A, et al. Gastrointestinal stromal tumors (GIST) related emergencies. *Int J Surg* 2014;12:269-80.
  122. Chok AY, Koh YX, Ow MY, et al. A systematic review and meta-analysis comparing pancreatoduodenectomy versus limited resection for duodenal gastrointestinal stromal tumors. *Ann Surg Oncol* 2014;21:3429-38.
  123. Lin C, Chang Y, Zhang Y, et al. Small duodenal gastrointestinal stromal tumors presenting with acute bleeding misdiagnosed as hemobilia: two case reports. *Oncol Lett* 2012;4:1069-71.
  124. Hygum K, Wulff CN, Harslof T, et al. Hypercalcemia in metastatic GIST caused by systemic elevated calcitriol: a case report and review of the literature. *BMC Cancer* 2015;15:788.
  125. Goh BK, Chow PK, Kesavan S, et al. Outcome after surgical treatment of suspected gastrointestinal tumors involving duodenum. *J Surg Oncol* 2008;97:388-91.
  126. Hohenberger P, Ronellenfisch U, Oladeji O, et al. pattern of recurrence in patients with ruptured primary gastrointestinal stromal tumour. *Br J Surg* 2010;97:1854-9.
  127. Guller U, Tarantino I, Cerny T, et al. Revisiting a dogma: similar survival of patients with small bowel and gastric GIST. A population-based propensity score SEER analysis.

- Gastric Cancer 2017;20:49-60.
128. Chiarugi M, Galatioto C, Lippolis P, et al. Gastrointestinal stromal tumor of the duodenum in childhood: a rare case report. *BMC Cancer* 2007;7:79.
  129. Kim JY, Lee JM, Kim KW, et al. Ectopic pancreas: CT findings with emphasis on differentiation from small gastrointestinal stromal tumor and leiomyoma. *Radiology* 2009;252:92-100.
  130. Yamashita Y, Kato J, Ueda K, et al. Contrast-enhanced endoscopic ultrasonography can predict a higher malignant potential of gastrointestinal stromal tumors by visualizing large newly formed vessels. *J Clin Ultrasound* 2015;43:89-97.
  131. Hoda KM, Rodriguez SA, Faigel DO. EUS-guided sampling of suspected GI stromal tumors. *Gastrointest Endosc* 2009;69:1218-23.
  132. Yamashita S, Sakamoto Y, Saiura A, et al. Pancreas-sparing duodenectomy for gastrointestinal stromal tumor. *Am J Surg* 2014;207:578-83.
  133. Jang SK, Kim JH, Joo I, et al. Differential diagnosis of pancreatic cancer from other solid tumours arising from the periampullary area on MDCT. *Eur Radiol* 2015;25:2880-8.
  134. Goldstein D, Tan BS, Rossleigh M, et al. Gastrointestinal stromal tumors: correlation of 18F-FDG Gamma camera-based coincidence positron emission tomography with CT for assessment of treatment response – an AGITG study. *Oncology* 2005;69:326-32.
  135. ESMO/European Sarcoma Network Working Group. Gastrointestinal stromal tumors: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014;25 Suppl 3:iii21-6.
  136. Cirocchi R, Farinella E, La Mura F, et al. Efficacy of surgery and imatinib mesylate in the treatment of advanced gastrointestinal stromal tumor: a systematic review. *Tumori* 2010;96:392-9.
  137. Gold JS, van der Zwan SM, Gonen M, et al. Outcome of metastatic GIST in the era before tyrosine kinase inhibitors. *Ann Surg Oncol* 2007;14:134-42.
  138. Bourgouin S, Hornez E, Guiramand J, et al. Duodenal gastrointestinal stromal tumors (GISTs): arguments for conservative surgery. *J Gastrointest Surg* 2013;17:482-7.
  139. Abe N, Hashimoto Y, Takeuchi H, et al. Laparoscopy-assisted full-thickness resection of the duodenum for patients with gastrointestinal stromal tumor with ulceration. *Asian J Endosc Surg* 2017;10:388-93.
  140. Xie YB, Liu H, Cui L, et al. Tumors of the angle of Treitz: a single-center experience. *World J Gastroenterol* 2014;20:3628-34.
  141. Tanaka E, Kim M, Lim JS, et al. Usefulness of laparoscopic side-to-side duodenojejunostomy for gastrointestinal stromal tumors located at the duodenojejunal junction. *J Gastrointest Surg* 2015;19:313-8.
  142. Noshiro H, Nomura A, Akashi M, et al. Pure robotic surgery for intraluminally growing gastrointestinal stromal tumors around the esophagogastric junction or pyloric ring. *Hepatogastroenterology* 2015;62:629-34.
  143. Downs-Canner S, Van de Vliet WJ, Tholen SJJ, et al. Robotic surgery for benign duodenal tumors. *J Gastrointest Surg* 2015;19:306-12.
  144. Huscher CG, Mingoli A, Sgarzini G, et al. Transoral extraction of a laparoscopically resected large gastric GIST. *J Laparoendosc Adv Surg Tech A* 2013;23:707-9.

doi: 10.21037/tgh.2018.09.04

**Cite this article as:** Popivanov G, Tabakov M, Mantese G, Cirocchi R, Piccinini I, D'Andrea V, Covarelli P, Boselli C, Barberini F, Tabola R, Pietro U, Cavaliere D. Surgical treatment of gastrointestinal stromal tumors of the duodenum: a literature review. *Transl Gastroenterol Hepatol* 2018;3:71.