



DISTANCE k -DOMINATION IN SOME CYCLE RELATED GRAPHS

S. K. VAIDYA AND N. J. KOTHARI

Received 12 March, 2014

Abstract. In this paper we determine distance k -domination number of graph obtained by duplication of vertices altogether by edges in cycle C_n , splitting graph of cycle C_n as well as graph obtained by duplication of edges altogether by vertices in cycle C_n .

2010 *Mathematics Subject Classification:* 05C69; 05C12; 05C38; 05C76

Keywords: dominating set, domination number, distance domination, splitting graph

1. INTRODUCTION

Graph $G = (V(G), E(G))$, we mean simple, finite, connected and undirected graph. The open neighbourhood $N(v)$ of $v \in V(G)$ is the set of all vertices adjacent to v . That is, $N(v) = \{u \in V(G)/uv \in E(G)\}$. The closed neighbourhood of $v \in V(G)$ is the set $N[v] = N(v) \cup \{v\}$. The distance $d(u, v)$ between two vertices u and v is the length of the shortest uv -path in G , if exists, otherwise $d(u, v) = \infty$. The open k -neighbourhood $N_k(v)$ of a vertex $v \in V(G)$ is the set of all vertices of G which are different from v and at a distance at most k from v in G . That is $N_k(v) = \{u \in V(G)/d(u, v) \leq k\}$. The closed k -neighbourhood set is defined as $N_k[v] = N_k(v) \cup \{v\}$. It is obvious that $N(v) = N_1(v)$. A set $D \subseteq V(G)$ is called a dominating set if every vertex in $V(G) - D$ is adjacent to at least one vertex in D . For terminology and notation not defined here we follow West [13] and Haynes *et al.* [3]. The concept of distance dominating set was initiated by Slater [7] with special reference to communication network, while the term distance k -dominating set was given by Henning *et al.* [5]. For an integer $k \geq 1$, $D \subseteq V(G)$ is a distance k -dominating set of G , if every vertex in $V(G) - D$ is within the distance k from some vertex $v \in D$. That is, $N_k[D] = V(G)$. The minimum cardinality among all the distance k -dominating sets of G is called the distance k -domination number of G and it is denoted by $\gamma_k(G)$. It is obvious that $\gamma(G) = \gamma_1(G)$. A distance k -dominating set of cardinality $\gamma_k(G)$ is called a γ_k -set. Many reserchers have explored the concept of distance k -domination in graphs. The distance domination number for cartesian products of two paths has been investigated by Klobucar [6]

while the distance domination in the context of spanning tree of the graph is discussed by Griggs and Hutchinson [2]. The bounds on the distance two-domination number and the classes of graphs attaining these bounds are reported in Sridharan *et al.* [8]. Tian and Xu [9] have established upper bound for distance k -domination for connected graph G and show that $\gamma_k(G) \leq \left\lfloor \frac{n-\Delta+k-1}{k} \right\rfloor$. The same authors in [10] have studied average distance and distance domination number and established an upper bound of average distance in terms of distance domination. Fischermann and Volkmann [1] have characterized the graphs whose distance n -domination number is equal to half of their number of vertices, when the diameter is greater or equal to $2n-1$. Vaidya and Kothari [12] have investigated distance k -domination number of total graph, shadow graph and middle graph of path P_n . The same authors in [11] have investigated distance k -domination number for the graphs obtained by graph operations on some standard graphs. For more bibliographic references on distance k -domination, the readers are advised to refer a survey article by Henning [4].

2. RESULTS

Proposition 1 ([4]). *Let $k \geq 1$ and D be a distance k -dominating set of a graph G . Then D is a minimal distance k -dominating set of G if and only if each $d \in D$ has at least one of the following two properties hold.*

- (1) *There exist a vertex $v \in V(G) - D$ such that $N_k(v) \cap D = \{d\}$.*
- (2) *The vertex d is at distance at least $k+1$ from every other vertex d' of D in G .*

Definition 1. Duplication of a vertex v by a new edge $e = v'v''$ of graph G produces a new graph G' such that $N(v') \cap N(v'') = \{v\}$.

Theorem 1. *If G is a graph obtained by duplication of vertices altogether by edges in cycle C_n ($n \leq 2k-1$) then $\gamma_k(G) = 1$.*

Proof. Let G be a graph obtained by duplication of vertices v_1, v_2, \dots, v_n by edges $u_{2i-1}u_{2i}$ ($1 \leq i \leq n$) in cycle C_n . Then $D = \left\{ v \left\lfloor \frac{n}{2} \right\rfloor \right\}$ is distance k -dominating set of G as $n \leq 2k-1$. Hence $\gamma_k(G) = 1$. \square

Theorem 2. *If G is a graph obtained by duplication of vertices altogether by edges in cycle C_n ($n > 2k-1$) then*

$$\gamma_k(G) = \begin{cases} \left\lfloor \frac{n}{2k-1} \right\rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k-2 \pmod{2k-1} \\ \frac{n}{2k-1} & \text{for } n \equiv 0 \pmod{2k-1} \end{cases}$$

Proof. Let G be a graph obtained by duplication of vertices v_1, v_2, \dots, v_n by edges $u_{2i-1}u_{2i}$ ($1 \leq i \leq n$) in cycle C_n . One can observe that v_i 's dominate more vertices than u_i 's. Here vertices from $v_{n-(k-1)}$ to v_{k+1} , $u_{2n-(2k-3)}$ to u_{2n} and u_1

to u_{2k} are dominated by a vertex v_1 at a distance k . Also $d(v_1, u_{2k}) = k$, and $d(u_{2n-(2k-3)}, v_1) = k$. Hence $2k - 1$ consecutive vertices from u_i 's dominated by only one vertex v_1 . Hence

$$\gamma_k(G) \geq \left\lfloor \frac{n}{2k-1} \right\rfloor \tag{2.1}$$

Now depending upon the number of vertices of C_n , consider the following subsets, For $n \equiv 1, 2, \dots, 2k - 2 \pmod{2k - 1}$

$$D = \left\{ v_{1+(2k-1)j/0} \leq j \leq \left\lfloor \frac{n}{2k-1} \right\rfloor \right\}, |D| = \left\lfloor \frac{n}{2k-1} \right\rfloor + 1,$$

for $n \equiv 0 \pmod{2k - 1}$

$$D = \left\{ v_{1+(2k-1)j/0} \leq j < \frac{n}{2k-1} \right\}, |D| = \frac{n}{2k-1}.$$

We claim that each D is a distance k -dominating set as

For $j \neq 0$,

$$d(v_{1+(2k-1)j}, v_{i+(2k-1)j}) \leq k, \text{ where } -k + 1 \leq i \leq k + 1, i \neq 1$$

$$d(v_{1+(2k-1)j}, u_{i+(2k-1)j}) \leq k, \text{ where } (3 - 2k) + (2k - 1)j \leq i \leq 2k + (2k - 1)j$$

and for $j = 0$,

$$d(v_1, v_i) \leq k, \text{ where } n - k + 1 \leq i \leq n, 2 \leq i \leq k + 1$$

$$d(v_1, u_i) \leq k, \text{ where } 1 \leq i \leq 2k, 2n - (2k - 3) \leq i \leq 2n$$

Therefore for $j \neq 0$,

$$\begin{aligned} N_k(v_{1+(2k-1)j}) = & \{v_{2+(2k-1)j}, v_{3+(2k-1)j}, \dots, v_{(k+1)+(2k-1)j}, \dots, v_{(2k-1)j}, \\ & v_{(2k-1)j-1}, \dots, v_{(2k-1)j-(k-1)}, u_{1+2(2k-1)j}, u_{2(1+(2k-1)j)}, \\ & \dots, u_{2k+2((2k-1)j)}, u_{(3-2k)+2((2k-1)j)}, u_{(4-2k)+2((2k-1)j)}, \\ & \dots, u_{2(2k-1)j}\}. \end{aligned}$$

While for $j = 0$,

$$\begin{aligned} N_k(v_1) = & \{v_2, v_3, \dots, v_{k+1}, v_{n-(k-1)}, v_{n-k}, \dots, v_n, u_1, u_2, \dots, u_{2k}, u_{2n}, \\ & u_{2n-1}, \dots, u_{2n-(2k-3)}\}. \end{aligned}$$

Then $N_k[D] = N_k[v_1] \cup N_k[v_{1+(2k-1)j}] = V(G)$.

For some $j = j_1, v_{1+(2k-1)j_1} \in D$ and for some $j = j_2, v_{1+(2k-1)j_2} \in D$,

$$d(v_{1+(2k-1)j_1}, v_{1+(2k-1)j_2}) = (j_2 - j_1)(2k - 1) \geq k + 1.$$

Which implies that every vertex d of D is at a distance $k + 1$ apart from every other vertex of D in G . Thus by Proposition 1 above defined D is a minimal distance

k -dominating set of G and by expression 2.1 it is also of minimum cardinality for $n > 2k - 1$. Hence

$$\gamma_k(G) = \begin{cases} \left\lfloor \frac{n}{2k-1} \right\rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k-2 \pmod{2k-1} \\ \frac{n}{2k-1} & \text{for } n \equiv 0 \pmod{2k-1} \end{cases}$$

□

Illustration 1. Distance 3-dominating set in graph G obtained by duplication of vertices in cycle C_{22} altogether by edges is shown by solid vertices in FIGURE 1.




FIGURE 1.

Definition 2. For a graph G , the splitting graph $S'(G)$ of graph G is obtained by adding a new vertex v' corresponding to each vertex v of G such that $N(v) = N(v')$.

Theorem 3. If $n \leq 2k + 1, k \neq 1$, then $\gamma_k(S'(C_n)) = 1$.

Proof. Let v_1, v_2, \dots, v_n be the vertices of cycle C_n and u_1, u_2, \dots, u_n be the vertices corresponding to v_1, v_2, \dots, v_n which are added to obtain $S'(C_n)$. Then $D = \left\{ v_{\left\lfloor \frac{n}{2} \right\rfloor} \right\}$ is distance k -dominating set of $S'(C_n)$ as $n \leq 2k + 1$. Hence $\gamma_k(S'(C_n)) = 1$. □

Theorem 4. If $n > 2k + 1$ then

$$\gamma_k(S'(C_n)) = \begin{cases} \left\lfloor \frac{n}{2k+1} \right\rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k \pmod{2k+1} \\ \frac{n}{2k+1} & \text{for } n \equiv 0 \pmod{2k+1} \end{cases}$$

Proof. Let v_1, v_2, \dots, v_n be the vertices of cycle C_n and u_1, u_2, \dots, u_n be the vertices corresponding to v_1, v_2, \dots, v_n which are added to obtain $S'(C_n)$. One can observe that v_i 's dominate more vertices than u_i 's at a distance k . In graph $S'(C_n)$

vertices $v_{n-(k-1)}$ to v_{k+1} and $u_{n-(k-1)}$ to u_{n+k} are dominated by a vertex v_1 at a distance k . Also $d(v_{n-(k-1)}, v_{k+1}) = 2k + 1$, and $d(u_{n-(k-1)}, u_{k+1}) = 2k + 1$. Hence $2k + 1$ consecutive vertices from v_i 's and $(2k + 1)$ consecutive vertices from u_i 's dominated by only one vertex at a distance k . Which implies that

$$\gamma_k (S'(C_n)) \geq \left\lfloor \frac{n}{2k + 1} \right\rfloor \tag{2.2}$$

Now depending upon the number of vertices of C_n , consider the following subsets, For $n \equiv 1, 2, \dots, 2k \pmod{2k + 1}$

$$D = \left\{ v_{1+(2k+1)j} / 0 \leq j \leq \left\lfloor \frac{n}{2k + 1} \right\rfloor \right\}, |D| = \left\lfloor \frac{n}{2k + 1} \right\rfloor + 1,$$

for $n \equiv 0 \pmod{2k + 1}$

$$D = \left\{ v_{1+(2k+1)j} / 0 \leq j < \frac{n}{2k + 1} \right\}, |D| = \frac{n}{2k + 1}.$$

We claim that each D is a distance k -dominating set as for $j \neq 0$,

$$d(v_{1+(2k+1)j}, v_{i+(2k+1)j}) \leq k, \text{ where } -k + 1 \leq i \leq k + 1, i \neq 1$$

$$d(v_{1+(2k+1)j}, u_{i+(2k+1)j}) \leq k, \text{ where } -k + 1 \leq i \leq k + 1$$

while for $j = 0$

$$d(v_1, v_i) \leq k, \text{ where } n - k + 1 \leq i \leq n, 1 \leq i \leq k + 1$$

$$d(v_1, u_i) \leq k, \text{ where } 1 \leq i \leq k + 1, n - k + 1 \leq i \leq n$$

Therefore $j \neq 0$,

$$N_k(v_{1+(2k+1)j}) = \{v_{2+(2k+1)j}, v_{3+(2k+1)j}, \dots, v_{(k+1)+(2k+1)j}, v_{(2k+1)j}, v_{(2k+1)j-1}, \dots, v_{(2k+1)j-(k-1)}, u_{1+(2k+1)j}, u_{2+(2k+1)j}, \dots, u_{(k+1)+(2k+1)j}, u_{(2k+1)j}, u_{(2k+1)j-1}, \dots, u_{(2k+1)j-(k-1)}\}.$$

While for $j = 0$,

$$N_k(v_1) = \{v_2, v_3, \dots, v_{k+1}, v_n, v_{n-1}, v_{n-(k+1)}, u_1, u_2, \dots, u_{k+1}, u_{n-k+1}, u_{n-k+2}, \dots, u_n\}.$$

Then $N_k[D] = N_k[v_1] \cup N_k[v_{1+(2k+1)j}] = V(S'(C_n))$.

For some $j = j_1, v_{1+(2k+1)j_1} \in D$ and for some $j = j_2, v_{1+(2k+1)j_2} \in D$,

$$d(v_{1+(2k+1)j_1}, v_{1+(2k+1)j_2}) = (j_2 - j_1)(2k + 1) \geq k + 1$$

This implies that every vertex d of D is at a distance $k + 1$ apart from every other vertex of D in $S'(C_n)$. Thus by Proposition 1 above defined D is a minimal distance

k -dominating set of G and by expression 2.2 it is also of minimum cardinality for $n > 2k - 1$. Hence

$$\gamma_k(S'(C_n)) = \begin{cases} \left\lfloor \frac{n}{2k+1} \right\rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k \pmod{2k+1} \\ \frac{n}{2k+1} & \text{for } n \equiv 0 \pmod{2k+1} \end{cases}$$

□

Illustration 2. Distance 4-dominating set in $S'(C_{20})$ is shown by solid vertices in FIGURE 2.




FIGURE 2.

Definition 3. Duplication of an edge $e = uv$ by a new vertex w in a graph G produces a new graph G' such that $N(w) = \{u, v\}$.

Theorem 5. If G is a graph obtained by duplication of edges altogether by vertices in cycle C_n ($n \leq 2k$) then $\gamma_k(G) = 1$.

Proof. Let G be a graph obtained by duplication of edges $v_i v_{i+1}$ altogether by vertices u_i , ($1 \leq i < n$) in cycle C_n . Then $\left\{ v_{\lfloor \frac{n}{2} \rfloor} \right\}$ is distance k -dominating set of G as $n \leq 2k$. Hence $\gamma_k(G) = 1$. □

Theorem 6. *If G is a graph obtained by duplication of edges altogether by vertices in cycle C_n ($n > 2k$) then*

$$\gamma_k(G) = \begin{cases} \lfloor \frac{n}{2k} \rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k - 1 \pmod{2k} \\ \frac{n}{2k} & \text{for } n \equiv 0 \pmod{2k} \end{cases}$$

Proof. Let G be a graph obtained by duplication of edges $v_i v_{i+1}$ altogether by vertices u_i ($1 \leq i < n$) in cycle C_n . One can observe that v_i 's dominate more vertices than u_i 's at a distance k . In graph G vertices v_{n-k} to v_{k+1} and $u_{n-(k-1)}$ to u_k are dominated by a vertex v_1 at a distance k . Also $d(u_{n-(k-1)}, v_1) = k$ and $d(v_1, u_k) = k$. Hence $2k$ consecutive vertices of u_i 's are dominated by only one vertex at a distance k . Hence

$$\gamma_k(G) \geq \lfloor \frac{n}{2k} \rfloor \tag{2.3}$$

Now depending upon the number of vertices of C_n , consider the following subsets. For $n \equiv 1, 2, \dots, 2k - 1 \pmod{2k}$

$$D = \left\{ v_{1+(2k)j} / 0 \leq j \leq \lfloor \frac{n}{2k} \rfloor \right\}, |D| = \lfloor \frac{n}{2k} \rfloor + 1,$$

for $n \equiv 0 \pmod{2k}$

$$D = \left\{ v_{1+(2k)j} / 0 \leq j < \frac{n}{2k} \right\}, |D| = \frac{n}{2k}.$$

Now we claim that each D is a distance k -dominating set as for $j \neq 0$,

$$d(v_{1+2kj}, v_{i+2kj}) \leq k, \text{ where } -k + 1 \leq i \leq k + 1, i \neq 1,$$

$$d(v_{1+2kj}, u_{i+2kj}) \leq k, \text{ where } -k + 1 \leq i \leq k$$

while for $j = 0$,

$$d(v_1, v_i) \leq k, \text{ where } n - k + 1 \leq i \leq n, 2 \leq i \leq k + 1$$

$$d(v_1, u_i) \leq k, \text{ where } n - k + 1 \leq i \leq n, 1 \leq i \leq k$$

Therefore for $j \neq 0$,

$$N_k(v_{1+2kj}) = \{v_{2+2kj}, v_{3+2kj}, \dots, v_{(k+1)+2kj}, v_{2kj}, v_{2kj-1}, \dots, v_{2kj-(k-1)}, u_{1+2kj}, u_{2+2kj}, \dots, u_{k+2kj}, u_{2kj}, u_{2kj-1}, \dots, u_{2kj-(k-1)}\}.$$

While for $j = 0$,

$$N_k(v_1) = \{v_2, v_3, \dots, v_{k+1}, v_{n-k+1}, v_{n-k+2}, \dots, v_n, u_1, u_2, \dots, u_k, u_{n-k+1}, u_{n-k+2}, \dots, u_n\}.$$

This implies that $N_k[D] = N_k[v_1] \cup N_k[v_{1+(2k)j}] = V(G)$.

For some $j = j_1, v_{1+(2k)j_1} \in D$ and for some $j = j_2, v_{1+(2k)j_2} \in D$,

$$d(v_{1+(2k)j_1}, v_{1+(2k)j_2}) = (j_2 - j_1)2k \geq k + 1$$

Which implies that every vertex d of D is at a distance $k + 1$ apart from every other vertex of D in G . Thus by Proposition 1 above defined D is a minimal distance k -dominating set of G and by expression 2.3 it is also of minimum cardinality for $n > 2k$. Hence

$$\gamma_k(G) = \begin{cases} \left\lfloor \frac{n}{2k} \right\rfloor + 1, & \text{for } n \equiv 1, 2, \dots, 2k - 1 \pmod{2k} \\ \frac{n}{2k} & \text{for } n \equiv 0 \pmod{2k} \end{cases}$$

□

Illustration 3. Distance 2-dominating set in graph G obtained by duplication of edges in C_{18} altogether by vertices is shown by solid vertices in FIGURE 3.




FIGURE 3.

REFERENCES

[1] M. Fischermann and L. Volkmann, “Graphs having distance- n domination number half their order.” *Discrete Appl. Math.*, vol. 120, no. 1-3, pp. 97–107, 2002, doi: [10.1016/S0166-218X\(01\)00284-0](https://doi.org/10.1016/S0166-218X(01)00284-0).

[2] J. R. Griggs and J. P. Hutchinson, “On the r -domination number of a graph.” *Discrete Math.*, vol. 101, no. 1-3, pp. 65–72, 1992, doi: [10.1016/0012-365X\(92\)90591-3](https://doi.org/10.1016/0012-365X(92)90591-3).

[3] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Fundamentals of Domination in Graphs*. Marcel Dekker, New York, 1998.

- [4] M. A. Henning, "Distance domination in graphs," in *Domination in Graphs: Advanced Topics*, T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, Eds. Marcel Dekker, New York, 1998, pp. 321–349.
- [5] M. Henning, O. R. Oellermann, and H. C. Swart, "Bounds on distance domination parameters." *J. Comb. Inf. Syst. Sci.*, vol. 16, no. 1, pp. 11–18, 1991.
- [6] A. Klobučar, "On the k -dominating number of Cartesian products of two paths." *Math. Slovaca*, vol. 55, no. 2, pp. 141–154, 2005.
- [7] P. J. Slater, "R-domination in graphs." *J. Assoc. Comput. Mach.*, vol. 23, pp. 446–450, 1976, doi: [10.1145/321958.321964](https://doi.org/10.1145/321958.321964).
- [8] N. Sridharan, V. Subramanian, and M. Elias, "Bounds on the distance two-domination number of a graph." *Graphs Comb.*, vol. 18, no. 3, pp. 667–675, 2002, doi: [10.1007/s003730200050](https://doi.org/10.1007/s003730200050).
- [9] F. Tian and J.-M. Xu, "A note on distance domination numbers of graphs." *Australas. J. Comb.*, vol. 43, pp. 181–190, 2009.
- [10] F. Tian and J.-M. Xu, "Average distances and distance domination numbers." *Discrete Appl. Math.*, vol. 157, no. 5, pp. 1113–1127, 2009, doi: [10.1016/j.dam.2008.03.024](https://doi.org/10.1016/j.dam.2008.03.024).
- [11] S. K. Vaidya and N. J. Kothari, "Distance k -domination of some path related graphs." *International Journal of Mathematics and Soft Computing*, vol. 4, no. 1, pp. 1–5, 2014.
- [12] S. K. Vaidya and N. J. Kothari, "Some new results on distance k -domination in graphs." *Int. J. Comb.*, vol. 2013, p. 4, 2013, doi: [10.1155/2013/795401](https://doi.org/10.1155/2013/795401).
- [13] D. B. West, *Introduction to graph theory*, 2nd ed. Prentice-Hall, New Delhi, India, 2003.

Authors' addresses

S. K. Vaidya

Saurashtra University, Department of Mathematics, Rajkot - 360005, Gujarat, INDIA

E-mail address: samirkvaidya@yahoo.co.in

N. J. Kothari

L. E. College (Diploma), General Department, Morbi-363642, Gujarat, INDIA

E-mail address: nirang.kothari@yahoo.com