

Editorial Manager(tm) for Neurocritical Care
Manuscript Draft

Manuscript Number:

Title: The prognostic value of brain extracellular fluid nitric oxide metabolite after traumatic brain injury

Article Type: Original Research

Keywords: traumatic brain injury, neurocritical care, neuromonitoring, microdialysis, biomarker

Corresponding Author: Martin Miles Tisdall, MD

Corresponding Author's Institution: The National Hospital for Neurology and Neurosurgery

First Author: Martin Miles Tisdall, MD

Order of Authors: Martin Miles Tisdall, MD;Konrad Redjak;Neil Kitchen;Martin Smith;Axel Petzold

1
2
3
4
5
6
7
8
9

The prognostic value of brain extracellular fluid nitric oxide metabolite after traumatic brain injury

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

Authors: ^{1,2}Martin M. Tisdall FRCS(SN) MD; ³Konrad Rejdak, PhD; ²Neil D.

Kitchen FRCS(SN) MD; ¹Martin Smith MBBS FRCA; ^{1,4}Axel Petzold MD PhD

Departments of ¹Neuroanaesthesia and Neurocritical Care, and ²Neurosurgery,
The National Hospital for Neurology and Neurosurgery, Queen Square, London,
UK, WC1N 3BG.

³ Department of Neurology, Medical University of Lublin, Lublin, Poland

⁴ VUmc, Department of Neurology, Amsterdam, NL.

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Address requests for reprints to: Axel Petzold, UCL Institute of Neurology, Box
76, Queen Square, London, WC1N 3BG, a.petzold@ion.ucl.ac.uk

MT is supported by a Wellcome Clinical Research Training Fellowship, Grant No
075608.

*Keywords: Traumatic Brain Injury, Microdialysis, Nitric Oxide, Critical Care,
Patient Monitoring*

1
2
3
4 **Abstract**
5

6 *Introduction:* Nitric oxide (NO) is a compound with both protective and damaging
7 effects on neurons. Quantification of NO metabolites in humans is limited by
8 sample contamination with blood. *In vivo* cerebral microdialysis may offer an
9 alternative approach as sampling of extracellular fluid (ECF) adjacent to neurons
10 becomes possible. We investigate the prognostic value of brain ECF NO
11 metabolites in patients with traumatic brain injury (TBI).
12
13
14
15
16
17
18
19
20

21 *Methods:* A prospective case cohort of 195 ECF samples collected from 11
22 cases over four days following TBI were collected. Nitrate and nitrite
23 concentration ([NO_x]) were quantified using a vanadium based colorimetric
24 assay.
25
26
27
28
29

30
31 *Results:* Early ECF NO_x levels (<48 hours post TBI) were significantly higher in
32 non-survivors (median 59.2 μmol/L, n=7) compared to survivors (23.3 μmol/L,
33 n=4)(p=0.04). Late (48-96 hours) ECF NO_x levels remained higher in non-
34 survivors 47.9 μmol/L compared to survivors 23.0 μmol/L but this was not of
35 significant (p=0.29). Receiver operator characteristic analysis show an optimized
36 cutoff level for ECF NO_x of 26.5 μmol/L measured <48 hours post TBI for
37 predicting non-survival (sensitivity 100%, specificity 75%).
38
39
40
41
42
43
44
45
46
47

48 *Conclusion:* Early ECF NO_x levels are of prognostic value after TBI. ECF NO_x
49 may be a useful biomarker for treatment trials targeted at nitric oxide metabolism.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **INTRODUCTION**
8

9 Nitric oxide (NO) is a free radical involved in diverse physiological and
10 pathophysiological processes in the central nervous system. After traumatic brain
11 injury (TBI), NO may have both protective and damaging effects(1) and this
12 relates to NO's role in cerebral blood flow regulation(2), its reaction with
13 superoxide to form peroxynitrite, and its inhibition of cytochrome c oxidase(3).
14 TBI remains the major cause of death in young people and survival of TBI is
15 associated with significant morbidity. Improved prognostication after TBI might
16 facilitate patient selection for aggressive management strategies, which may
17 improve mortality but carry the risk of increased morbidity if patient selection is
18 suboptimal.
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Due to its short half life, it is difficult to measure nitric oxide concentration
34 directly, but measurement of the concentration of downstream NO metabolites,
35 nitrate and nitrite (NOx), is an accepted method of estimating endogenous NO
36 production(4). NOx concentration ([NOx]) in the cerebrospinal fluid (CSF) of
37 patients after TBI has been reported (5,6) but the presence of subarachnoid
38 hemorrhage, as commonly occurs after TBI, can influence CSF [NOx]
39 measurement(4) and might reduce the usefulness of CSF [NOx] as a marker of
40 secondary injury.
41
42
43
44
45
46
47
48
49
50
51
52

53 Cerebral microdialysis is an established technique that allows focal
54 measurement of brain extracellular fluid (ECF) biochemistry(7,8). As the
55 microdialysis catheter is placed in brain tissue, it can be used to assess [NOx]
56
57
58
59
60
61
62
63
64
65

1
2
3
4 closer to NO's site of action than that measured in CSF and localise the
5
6 measurement to a particular area of interest. Microdialysate [NOx] in the brains
7
8 of rats post cortical impact injury correlates well with direct NO electrode
9
10 readings(1).
11
12
13

14 In this study we measure the concentration of nitrite and nitrate in the
15
16 brain extracellular fluid of patients with traumatic brain injury and compare it with
17
18 patient outcome.
19
20
21
22

23 **2. Materials and Methods**

24 *2.1 Patients and sample preparation*

25
26 This study was approved by the Joint Research Ethics Committee of the
27
28 National Hospital for Neurology and Neurosurgery and the Institute of Neurology
29
30 and, as all patients were unconscious at the time of the study, written assent was
31
32 obtained from their personal representatives.
33
34
35
36
37

38 We studied eleven adults with TBI undergoing protocolised management
39
40 to control cerebral perfusion pressure (CPP) and intracranial pressure (ICP) on
41
42 the neurocritical care unit. The patients were all male with median age 26.5 years
43
44 (range 17-68). As part of routine clinical monitoring of brain ECF biochemistry
45
46 each patient had a commercially available microdialysis catheter (CMA 70,
47
48 CMA/Microdialysis, Solna, Sweden) with a 10 mm dialysis membrane length and
49
50 20 kDa molecular weight cut-off inserted into their brain tissue through a skull
51
52 bolt (Technicam Ltd, Newton Abbott, UK). Probe positioning followed the
53
54 recommendations of the consensus meeting on microdialysis in neurointensive
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 care(9). The probe was perfused with artificial CSF (Perfusion Fluid CNS, CMA),
5
6 using a CMA 106 pump, at a rate of 0.3 μ L/min.
7
8

9 Systemic data including CCP and ICP were collected electronically. After
10 routine measurement of standard microdialysis variables, including
11 lactate:pyruvate ratio (LPR), using the CMA 600 analyser (CMA/Microdialysis,
12 Solna, Sweden), the microdialysate samples were stored at -80 °C prior to NOx
13
14
15
16
17
18
19 assay.
20

21 Microdialysate sampling was continued until invasive intracranial
22 monitoring was no longer clinically indicated and at that stage the microdialysis
23 catheter was removed. In any event only samples from the first 4 days post injury
24 were included in this analysis. Peak CSF [NOx] occurs within the first 48 hours
25
26
27
28
29
30
31 post TBI(5,6) and so samples for analysis were divided into those taken up to 48
32
33
34 hours, and those taken 48-96 hours, post TBI.
35

36 2.2 NOx Assay 37

38 We used a vanadium-based assay for nitrite/nitrate measurement in
39
40 biological specimens as described by Miranda *et al.*(10).
41

42 Reagents: Sulfanilamide (SULF), *N*-(1-Naphthyl) ethyl-enediamine dihydrochloride
43 (NEDD), vanadium (III) chloride (VCl_3), sodium nitrite, sodium nitrate,
44
45
46
47
48
49
50 hydrochloric acid were purchased from Sigma-Aldrich (Poole, Dorset, UK). Water
51 was purified with Milli-RO and Milli-Q systems (Millipore).
52

53 The assay was performed in a standard flat-bottomed 96-well polystyrene
54
55
56
57 microtitre plate, containing 50 μ l/well of standard or sample in duplicate.

58 Immediately before commencing the assay, Griess reagents: SULF(50 μ l) and
59
60
61
62
63
64
65

1
2
3
4 NEDD(50µl) were premixed and added to each well. Nitrite mixed with Griess
5
6 reagents forms a chromophore from the diazotization of sulfanilamide by acidic
7
8 nitrite followed by coupling with bicyclic amines such as *N*-1-(Naphthyl)
9
10 ethylenediamine. Subsequently, the vanadium solution was added in a volume of
11
12 100 µl/well, the contents mixed vigorously and the plate left to incubate for 60
13
14 minutes at 37 °C. The absorbance at 540 nm was measured to assess the total
15
16 level of nitrite and nitrate (NO_x) in the samples.
17
18
19
20

21 *2.3 Statistical Analysis*

22
23 Data were analysed using SAS software (version 9.1, SAS Institute, Cary,
24
25 NC, USA). Wilcoxon rank sum test was used to compare variables in survivors
26
27 and non-survivors. Because of a non-Gaussian distribution the median
28
29 (interquartile range) are presented. Receiver operator characteristic (ROC)
30
31 analyses are used to determine the sensitivity and specificity of threshold
32
33 variable values for non-survival. Two tailed exact p values are presented
34
35 throughout and values of less than 0.05 are considered significant.
36
37
38
39
40
41
42

43 **3. Results**

44
45 195 microdialysate samples were analysed from 11 patients.
46
47 Microdialysate [NO_x] <48 hours post TBI was significantly higher in non-survivors
48
49 (n=7, 59.2 (31.8-150.6) µmol/l compared to survivors 23.3 (15.2-32.9) µmol/l
50
51 (n=4, p=0.04). Microdialysate [NO_x] 48 to 96 hours post TBI was 23.0
52
53 (21.8-35.2) in survivors (n=4) and 47.9 (37.6-60.6) in non-survivors (n=5)
54
55 (p=0.29) (figure 1). There was no significant difference in microdialysate [NO_x]
56
57
58
59
60
61
62
63
64
65

1
2
3
4 over time for either subgroup although there was a trend toward microdialysate
5 [NOx] increasing over time in survivors and decreasing over time in non-
6
7
8 survivors. For the period <48 hours post TBI, receiver operator characteristic
9 (ROC) analysis revealed microdialysate [NOx]>26.5 µmol/l predicted non-survival
10 with sensitivity 100% and specificity 75% and the area under the curve was 0.89.
11
12 For the period 48-96 hours post TBI microdialysate [NOx]>24.1 µmol/l predicted
13 non-survival with sensitivity 100% and specificity 75% and the area under the
14 curve was 0.75 (figure 2).
15
16
17
18
19
20
21
22

23
24 There were no significant differences in ICP, CPP or microdialysate
25 lactate pyruvate ratio between survivors and non-survivors for either time period.
26
27
28
29

30 31 **4. Discussion**

32
33 This prospective study shows that it is feasible to reliably quantify NOx
34 from minute amounts of human ECF using a rapid and high throughput
35 technique. The main finding of this study is that ECF NOx samples in the initial
36 48 hours after TBI were of prognostic value. This data is in line with results from
37 CSF [NOx] in humans after TBI (5). The main advantage ECF NOx levels have
38 over CSF NOx levels is that sample contamination with systemic NOx levels
39 coming from the blood is less likely.
40
41
42
43
44
45
46
47
48
49

50
51 The potential of this measurement is suggested by the fact that we saw
52 significant differences in microdialysate [NOx] between the outcome groups in
53 the period <48 hours post TBI whilst finding no differences in other measured
54 variables. ROC analysis indicates effective discrimination between survivors and
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 non-survivors for microdialysate [NOx] <48 hours post TBI as indicated by the
5
6 large area under the ROC plot. Interestingly this overlaps with the time interval of
7
8 accelerated neuronal loss following TBI(8).
9

10
11 Our results suggest that microdialysate [NOx] might be a useful prognostic
12
13 marker after TBI and that its prognostic value may be greatest early after injury,
14
15 during the first 48 hrs in this study. This is important because if prognostication of
16
17 patient outcome is to guide clinical decision making it must be possible to do so
18
19 as soon as possible after TBI.
20
21

22
23 This study presents data from a small number of patients and the
24
25 threshold value we suggest must therefore be viewed as a preliminary result. We
26
27 present these results to demonstrate the prognostic potential of microdialysate
28
29 [NOx] and for evaluation in future studies. These microdialysate concentrations
30
31 are dependent on the relative recovery of NOx and hence the microdialysis
32
33 catheter type and perfusate flow rate and in this study we use standard
34
35 methods(7). Automatisation of the colorimetric method(10) used should be
36
37 possible and thus allow for high throughput bedside measurements of ECF NOx
38
39 levels from much larger patient cohorts.
40
41
42
43
44

45 The incidence of mortality after TBI in our cohort is high and we believe
46
47 that this is related to our selection of patients for cerebral microdialysis
48
49 monitoring. In our centre, patients in whom sedation is stopped shortly after
50
51 admission following either conservative or surgical management are not
52
53 monitored using cerebral microdialysis and this study cohort therefore represents
54
55 the more severely injured end of the spectrum of our TBI admissions.
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Microdialysate [NOx] has been previously correlated with regional cerebral
5
6 blood flow in patients with TBI(2). In the above study no significant correlation
7
8 was found between microdialysate [NOx] and neurological outcome, which is in
9
10 apparent contrast with earlier results reported by the same group where the
11
12 highest microdialysate [NOx] in the first 5 days after TBI were found in patients
13
14 surviving with moderate or severe disability as opposed to non-survivors or those
15
16 making a good recovery, with non-survivors having lower microdialysate [NOx]
17
18 than survivors(11). This seems to be at variance with our findings, perhaps
19
20 reflecting the complexity of NO's neuroprotective and neurotoxic roles, although
21
22 our results are more consistent with previous CSF studies (5).
23
24
25
26
27

28 Animal and human studies suggest that the peak in NO occurring > 6
29
30 hours post TBI is primarily related to expression of inducible nitric oxide synthase
31
32 (iNOS)(1) and, as none of our patients were studied <6 hours post TBI, the
33
34 elevated microdialysate [NOx] we observed in non-survivors may also to be due
35
36 to iNOS. Most studies suggest that NO produced by iNOS has detrimental
37
38 effects and that inhibition of iNOS activity after TBI might have benefit(1),
39
40 although this is not universally accepted. The present findings are in line with the
41
42 basic science literature.
43
44
45
46
47

48 We would like to stress that, as NO is required for a number of vital
49
50 physiological functions, any future treatment aiming to completely block NO
51
52 production or receptor binding is likely to have deleterious effects. However,
53
54 combining ECF NOx levels with a validated biomarker for neurodegeneration
55
56 (eNfH)(8) might allow determination of the *in vivo* concentration at which
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

neurotoxicity occurs and thus help establish a target treatment threshold for NOx lowering strategies.

1
2
3
4
5
6
7 **References**
8

- 9 1. Cherian L, Hlatky R, Robertson CS. Nitric oxide in traumatic brain injury.
10 Brain Pathol 2004;14:195-201.
11
12 2. Hlatky R, Goodman JC, Valadka AB, Robertson CS. Role of nitric oxide in
13 cerebral blood flow abnormalities after traumatic brain injury. J Cereb Blood
14 Flow Metab 2003;23:582-588.
15
16 3. Moncada S, Bolanos JP. Nitric oxide, cell bioenergetics and
17 neurodegeneration. J Neurochem 2006;97:1676-1689.
18
19 4. Rejdak K, Petzold A, Sharpe MA et al. Cerebrospinal fluid nitrite/nitrate
20 correlated with oxyhemoglobin and outcome in patients with subarachnoid
21 hemorrhage. J Neurol Sci 2004;219:71-76.
22
23 5. Clark RS, Kochanek PM, Obrist WD et al. Cerebrospinal fluid and plasma
24 nitrite and nitrate concentrations after head injury in humans. Crit Care Med
25 1996;24:1243-1251.
26
27 6. Uzan M, Tanriover N, Bozkus H, Gumustas K, Guzel O, Kuday C. Nitric
28 oxide (NO) metabolism in the cerebrospinal fluid of patients with severe
29 head injury. Inflammation as a possible cause of elevated NO metabolites.
30 Surg Neurol 2001;56:350-356.
31
32 7. Tisdall MM, Smith M: Cerebral microdialysis: research technique or clinical
33 tool. Br J Anaesth 2006;97:18-25.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

8. Petzold A, Tisdall MM, Girbes AR et al. In vivo monitoring of neuronal loss in traumatic brain injury: a microdialysis study. *Brain* 2011;134:464-483.
9. Bellander BM, Cantais E, Enblad P et al. Consensus meeting on microdialysis in neurointensive care. *Intensive Care Med* 2004;30:2166-2169.
10. Miranda KM, Espey MG, Wink DA. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 2001;5:62-71.
11. Hlatky R, Furuya Y, Valadka AB, Goodman JC, Robertson CS. Microdialysate nitrate/nitrite levels following severe head injury. *Acta Neurochir Suppl* 2002;81:331-333.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Figure 1: Microdialysate nitrite/nitrate concentration ([NOx]) for survivors and non-survivors of traumatic brain injury for the periods less than 48 hours, and 48-96 hours, post traumatic brain injury (ns=not significant).

Figure 2: Receiver operator characteristic analysis for microdialysate nitrite/nitrate concentration ([NOx]) as predictor of non-survival during the first 48 hours (A) and 48-96 hours (B) post traumatic brain injury.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)

