

1 **Metabolic exhaustion in infection, cancer and autoimmunity**

2 **McKinney EF and Smith KGC**

3 Department of Medicine, University of Cambridge School of Clinical Medicine, Addenbrooke's Hospital, Hills Road,
4 Cambridge CB2 0QQ, UK.

5 6 **Abstract**

7 It has become increasingly clear that changes in metabolism are not just
8 consequences of T cell activation but rather are essential drivers of that process,
9 shaping the extent and nature of differentiation and function. The process of T
10 cell exhaustion has been linked to the outcome of chronic immune responses in
11 multiple contexts including chronic infection, cancer and autoimmunity. Factors
12 regulating the development and maintenance of exhaustion are of increasing
13 interest as the target of therapeutic modulation. Recent work has shown T cell
14 immunometabolism to be integral to the control and development of T cell
15 exhaustion. Early metabolic changes are responsible for later emergence of
16 exhaustion, do not simply reflect changes secondary to chronic activation and
17 are modifiable. An increased understanding of this metabolic control promises to
18 improve our ability to modulate T cell immunity to chronic antigen stimulation
19 in multiple contexts.
20

21 **Introduction**

22 The ability to sustain T cell function despite persisting antigen stimulation
23 contributes to the clearance of both chronic infection and cancer. However,
24 where T cell immunity targets self antigen or excessively damages healthy tissue,
25 maintaining robust responses can prove harmful rather than helpful. T cell
26 exhaustion has emerged as a key mechanism through which CD8 T cells lose
27 effector capacity during persistent stimulation, curtailing their ability to cause
28 damaging immunopathology but also facilitating viral persistence or hindering
29 tumour-targeted responses. A substantial body of work using murine models of
30 chronic LCMV virus infection has identified the molecular, transcriptional and
31 functional basis of T cell exhaustion. Recent studies have implicated early
32 metabolic signalling during T cell activation in the later development of
33 exhaustion. In doing so this work has highlighted the potential for metabolic
34 signals to shape T cell differentiation, and has focussed attention on the potential
35 of immunometabolic therapies in multiple contexts.
36

37 **T cell differentiation and metabolism**

38 The process of T cell activation and differentiation involves rapid proliferation,
39 cytokine production and development of cytotoxic effector function. For this to
40 happen, cells must balance increasing energy demands with an increasing need
41 for substrates to allow growth and division. Bioenergetic needs of naïve or
42 quiescent T cells are primarily met by mitochondrial oxidative phosphorylation
43 (OXPHOS), an efficient means of generating ATP from a glucose substrate¹. On
44 activation, however, there is a switch to aerobic glycolysis² known as the
45 'Warburg effect', a comparatively inefficient process in which fewer ATP
46 molecules are generated per molecule of glucose processed³. A T cells choice of
47 aerobic glycolysis initially makes little sense as it is not only inefficient, but
48 seemingly unnecessary – cells switch to glycolysis even when abundant oxygen
49 is available. However, the trade off is that a greater number of metabolic
50 intermediates are produced for anabolic growth, including those for nucleotide

51 and amino acid synthesis⁴, maintenance of cellular redox balance
52 (NAD⁺/NADH)⁵ and acetyl-CoA production for lipid synthesis through citrate
53 processing⁶. The switch to aerobic glycolysis is also required for the
54 development of full effector function (IL-2 and IFN- γ release), as the glycolytic
55 enzyme GAPDH otherwise restricts translation of IFN- γ mRNA².
56 However, effector differentiation is not the only goal of T cell activation and the
57 switch to aerobic glycolysis is not complete². Following antigen encounter and
58 eradication an expanded, persistent population of memory T cells shows a faster,
59 more robust response to subsequent activation (**Fig. 1**). For such memory to
60 develop during a primary T cell burst, aerobic glycolysis-driven effector function
61 must be balanced with other metabolic pathways including OXPHOS⁷ and fatty
62 acid oxidation (FAO)^{8,9}. A central role for mitochondria in this balance has been
63 highlighted by marked dynamic changes in their ultrastructure: T_{mem} show
64 fusion of mitochondrial cristae into networks whereas their T_{eff} counterparts
65 undergo fission and remain expanded¹⁰. The mature memory phenotype is also
66 fuelled by an increased mitochondrial biomass¹¹ and spare respiratory capacity
67 (SRC), driven by IL-15-mediated upregulation of carnitine palmitoyltransferase
68 (CPT1a)⁷. CPT1a activity controls mitochondrial FAO and fosters the ability to
69 generate a stronger, longer burst of both aerobic glycolysis and OXPHOS on
70 restimulation¹¹. However, it should be noted that the role of FAO has been
71 demonstrated using pharmacological inhibition of CPT1a, the specificity of which
72 has been brought into question by recent genetic deletion studies¹². Much
73 complexity surrounding the role of FAO in T cell differentiation remains to be
74 unpicked.

75

76 **FIGURE 1** Balance between T_{eff}/T_{mem} and metabolic pathways associated

77

78 It is to be expected that dynamic metabolic changes must occur to keep pace with
79 the rapid changes in cellular function that accompany T cell activation¹³.
80 However, these studies have together implicated cellular metabolism – and
81 mitochondrial biomass and function in particular - as a key driver of T cell
82 phenotype, rather than simply reflecting changes in energy demand during
83 differentiation^{8,14}. With immunometabolism taking centre stage, it has now
84 become clear that the development and maintenance of T cell exhaustion are also
85 under metabolic regulation.

86

87 **Exhaustion and metabolism**

88 T cell exhaustion is associated with progressive dysfunction characterised by
89 limited proliferation, cytokine production and effector capacity that is driven by
90 persistent, high levels of antigen exposure and a relative lack of costimulatory
91 'help' signals¹⁵. Exhaustion was originally identified in the murine LCMV model
92 of chronic viral infection but has since been identified during numerous other
93 chronic infections, in the tumour microenvironment and during
94 autoinflammatory responses^{16,17}. Its development controls persistence and
95 outcome of cellular immunity to persistent antigen and has been associated with
96 disease outcome in each case^{15,17}. More recent work has demonstrated that
97 exhaustion is also associated with significant bioenergetic compromise in the
98 setting of both chronic viral infection and the tumour microenvironment¹⁸⁻²⁰.
99 The observed metabolic dysfunction precedes the development of exhaustion *per*

100 se and has been mechanistically linked to early inhibitory PD1 signals. However,
101 this is only one part of a complex cascade modulating metabolism and the
102 balance of effector function, cellular survival and exhaustion. To date, studies
103 have focussed on catabolic metabolism, mitochondrial function and on pathways
104 controlling them. Consequently, this review is similarly focussed, but this should
105 not be taken to preclude the potential for downstream synthetic pathways to
106 also play an important role.

107
108 In the LCMV model of chronic viral infection, late emergence of T cell dysfunction
109 is preceded by both transcriptional and functional changes suggesting glucose
110 deprivation, with suppression of both glycolysis and OXPHOS^{18,21}. Glucose
111 transport is limited in early exhaustion with downregulation of GLUT1-mediated
112 transport. However, alterations in glucose uptake were seen to be modest,
113 prompting a search for changes in mitochondrial mass and function. This showed
114 that mitochondria of exhausted cells (Texh) are larger, but dysfunctional.
115 Increased mitochondrial biomass was accompanied by a paradoxical reduction
116 in mitochondrial function (Oxygen consumption rate, OCR) and by increases in
117 both mitochondrial depolarisation and reactive oxygen species (ROS) production
118 relative to their effector counterparts (Teff). Ultrastructural changes in the
119 mitochondria, including elongation of cristae, were also apparent¹⁸ although it is
120 less clear how these relate to ultrastructural changes already known to
121 differentiate classic memory and effector cell populations¹⁰. It is challenging –
122 particularly in vivo - to quantify the degree of metabolic switching required to
123 induce these mitochondrial changes (as it is with associated parameters such as
124 spare respiratory capacity (SRC))⁷, and thus to understand their importance in
125 directing T cell fate.

126 CD8⁺ tumour-infiltrating lymphocytes (TIL) also exhibit an exhausted
127 phenotype^{22,23} that limits anti-tumour responses, and can be reversed by cancer
128 immunotherapy with impressive results¹⁶. Like Texh in chronic LCMV infection,
129 TIL exhaustion in multiple murine models shows evidence of glucose deprivation
130 and mitochondrial dysfunction. As in Texh in the LCMV model, reduced TIL
131 glucose uptake was matched by reduced mitochondrial function (OCR) and
132 increased depolarisation^{19,20}. However, while bioenergetic changes of Texh were
133 broadly similar in both TIL and chronic viral models, there were also differences.
134 Whereas Texh in LCMV infection showed greater mitochondrial biomass and
135 increased ROS production¹⁸, TILs showed a reduction in both¹⁹. The apparent
136 contribution of PD1 signaling to metabolic dysfunction was also different in the
137 two contexts. Early metabolic dysfunction of Texh in LCMV-cl13 infection
138 responded to either genetic deletion of PD1 (adoptive transfer of transgenic
139 *Pdcd1*^{-/-} LCMV-specific P14 T cells) or to PDL1 blockade^{18,19}. However,
140 mitochondrial biomass was not substantially altered by PD1 blockade in TILs,
141 even though the blockade resulted in effective tumour regression¹⁹. These
142 context-specific differences in metabolic features of exhaustion require further
143 investigation but some explanations can be offered based on our increasing
144 appreciation of metabolic factors controlling T cell effector function, survival and
145 exhaustion.

146
147 **Metabolic substrate availability and exhaustion: Goldilocks and the 3**
148 **substrates**

149 Emerging evidence has implicated several metabolic pathways as important
150 players in the development of T cell exhaustion. However it should be noted that,
151 as with their contribution to T cell differentiation more broadly, their respective
152 roles are actually integrated, with changes in one influencing many others.

153

154 **(FIGURE 2: metabolic substrates and T cell exhaustion)**

155

156 **Hypoxia and exhaustion**

157 Both metabolic and effector dysfunction of TIL is clearly driven by and restricted
158 to the tumour microenvironment^{19,20,22,23}. Local metabolic substrate availability
159 is an important factor shaping exhaustion and may also explain some of the
160 variation in metabolic parameters between exhaustion models (**Fig. 2a**).

161 It has long been known that the tumour microenvironment is hypoxic and that
162 this favours tumour growth²⁴. However, recent work has revealed the
163 importance of hypoxia in driving the metabolic dysfunction of exhaustion. The
164 tumour suppressor gene *Vhl* does not interact with Hif (hypoxia inducible factor)
165 subunits under hypoxic conditions, allowing Hif to translocate to the nucleus and
166 mediate transcription of target genes. Genetic deletion of *Vhl* unleashed Hif
167 activity during persistent T cell responses resulting in fatal immunopathology
168 during chronic LCMV infection but improving survival in an experimental
169 melanoma model as T cells maintained effector function instead of exhausting²⁵.
170 This work further supports the prevailing theory that exhaustion has evolved as
171 a mechanism by which chronic infection is 'tolerated' rather than risk fatal injury
172 during an attempt to clear it¹⁷. It also highlights the importance of the
173 hypoxia/*Vhl*/Hif pathway in controlling T cell effector function during persistent
174 antigen stimulation. A metabolomic study of *Vhl*-deficient murine CD8⁺ T cells
175 (with consequent high levels of Hif activity) subsequently identified S-2-
176 hydroxyglutarate (S2HG) as a key immunometabolite controlling persistent CTL
177 activity in both immunisation and tumour models²⁶. S2HG levels are increased by
178 TCR stimulation in a Hif-dependent fashion but may also be influenced by other
179 metabolic pathways including glutaminolysis promoting both effector
180 differentiation and persistence²⁶. It also appears that the contribution of hypoxic
181 *Vhl*/Hif signalling to exhaustion is distinct from the contribution of classic
182 inhibitory receptors. In the absence of *Vhl* (P14 *Vhl*^{-/-} in the LCMV-cl13 model),
183 classic inhibitory receptors such as LAG3 and PD1 were still induced during
184 chronic infection but exhaustion did not develop, resulting in lethal
185 immunopathology driven by an aggressive CD8⁺ CTL response.

186 Hypoxia is a constant presence in conditions of rapid T cell infiltration,
187 proliferation and activation whether this is in the TME, virally infected tissue or
188 in the context of autoinflammatory disease. It is clear that the T cell metabolic
189 response to this hypoxic microenvironment is an important determinant of its
190 ability to sustain an effector response and consequently of disease outcome.
191 However, the exact impact of hypoxia on T cell effector function remains unclear
192 with some studies suggesting that limiting Hif expression (by knockdown rather
193 than genetic deletion) can improve rather than attenuate effector capacity²⁷.

194

195 **Glucose and exhaustion**

196 T cell activation is characterised by aerobic glycolysis, the switch away from
197 OXPHOS despite the presence of sufficient oxygen to maintain it¹³. However, a

198 lack of glucose availability in – or a lack of uptake from - the microenvironment
199 does have important consequences for persistence of T cell effector function
200 (**Fig. 2b**). Some studies have suggested that limiting glycolysis can favour
201 OXPHOS-mediated memory formation and enhanced persistent effector
202 function²⁸, although others have demonstrated a need for ongoing glycolysis to
203 maintain effector function²⁹. This apparent discrepancy reflects a need for
204 glycolysis-driven effector function to be counterbalanced by a concurrent need
205 for memory formation, driven by alternative metabolic pathways such as Hif,
206 FAO and OXPHOS. At least some glycolysis is clearly required for initial T cell
207 activation and for the development and maintenance of proliferation and effector
208 function^{2,13}. Similarly, even in the setting of enforced constitutive glycolysis
209 (achieved through conditional deletion of *Vhl*) a long-lived memory population
210 does develop, albeit skewed towards an effector-memory phenotype²⁹. A
211 balanced contribution of both effector-associated glycolysis and memory-
212 promoting metabolism such as FAO and OXPHOS is required for optimal T cell-
213 mediated control of pathogen or tumour³⁰. Skewing the response too far in either
214 direction could hamper persistent effector function.
215 T_H cells do show an early reduction in GLUT1-mediated glucose uptake and
216 early gene expression profiles consistent with glucose deprivation^{18,19,31}. When
217 TIL or T_H cells from chronic LCMV infection were exposed to conditions of
218 glucose deprivation, they showed attenuation of both TCR and NFAT signalling
219 with associated defects in Ca²⁺ flux, akin to those characterising T cell anergy^{31,32}.
220 Altered levels of the glycolytic metabolite phosphoenolpyruvate (PEP)³¹ in T_H
221 also contribute to the observed metabolic and effector dysfunction and do not
222 simply reflect secondary changes in metabolism. Glycolysis-induced PEP
223 accumulation was identified as necessary to sustain both Ca²⁺ and NFAT
224 signalling – and hence effector function - following TCR activation, with a loss of
225 IFN- γ and CD40L expression when it was suppressed³¹.
226 This same study also reminds us that cells are continually competing for
227 metabolic substrates in their local microenvironment – and it is not just a T cells
228 ability to take up and use a substrate that determines metabolic fitness, the
229 competing ability of neighbouring cells may be just as important. Overexpression
230 of the enzyme hexokinase-2 in tumour cells increased their rate of glycolysis
231 with evidence of glucose deprivation developing in TILs as a consequence³¹. This
232 competition for substrate has been elegantly demonstrated in the TME, where
233 tumour cell-driven changes in the availability of glucose were reflected in a
234 reduction in TIL effector capacity²⁰. A reduction in tumour cell glycolysis may at
235 least in part explain the reversal of TIL exhaustion and the success of
236 ‘checkpoint’ inhibitory receptor blockade²⁰. The degree to which such
237 competition for substrate contributes to exhaustion in the context of either
238 chronic infection or autoimmune responses is less clear, although is very likely to
239 play a role.

240

241 **Amino acids, survival and effector differentiation**

242 Enhanced T cell survival in an exhausting tumour microenvironment has also
243 been shown through increased uptake and metabolism of L-Arginine, that is
244 usually specifically depleted during T cell activation³³. Increased L-Arginine
245 correlated with improved T cell survival with differentiation skewed towards the
246 replicative, CCR7⁺ central memory (T_{cm}) phenotype (**Fig. 2c**). Once again,

247 availability of the metabolic substrate controlled the response as restoring
248 depleted L-Arginine levels improved cytotoxic capacity both in vitro and in
249 vivo³³. Once again, enhanced memory differentiation was associated with
250 attenuation of effector function, in this case a reduction in IFN- γ production
251 (although, consistent with memory differentiation, equivalent levels were
252 produced on secondary restimulation)³³. Cellular uptake of other amino acids
253 (AA), including leucine³⁴ and glutamine³⁵ via the AA transporters slc7a5 and
254 ASCT2 respectively, are important for initiation, maintenance and integration of
255 TCR signals with downstream metabolic sensors mTOR. They have not as yet
256 been investigated in the context of exhaustion but their expression and AA
257 availability is likely to also play a role. The critical AA transporter CD98, a
258 downstream target of the metabolic regulator c-Myc³⁶ and a heterodimer
259 incorporating slc7a5, does show elevated expression on memory but not
260 exhausted CD8⁺ T cells during chronic infection³⁷.

261

262 Together, these data make it clear that competition for metabolic substrate
263 within the local microenvironment is a critical factor influencing the balance of
264 effector function and T cell persistence during exhaustion^{19,20,31}.

265

266 **Metabolic exhaustion in different models: the same but different**

267 The majority of studies to date consistently demonstrate metabolic dysfunction
268 preceding exhaustion in different contexts. Furthermore, the observations are
269 not restricted to murine models of disease. TIL from human cancer have been
270 shown to have a similarly dysfunctional metabolic phenotype^{19,38} while human
271 CD8⁺ T cells specific for chronic persistent (HBV)^{39,40} but not acute (influenza)⁴⁰
272 or recurrent virus infections (CMV)³⁹ have been shown to have similar
273 limitations in glucose uptake and mitochondrial biomass.

274 However, while bioenergetic changes of Texh appear broadly similar in both TIL
275 and chronic viral models, there are also differences. Whereas Texh in chronic
276 LCMV infection showed greater mitochondrial biomass and increased ROS
277 production¹⁸, TILs showed a reduction in both¹⁹. The apparent contribution of
278 PD1 signalling to metabolic dysfunction was also different in the two contexts.
279 Early metabolic dysfunction of Texh in LCMVcl13 infection responded to either
280 genetic deletion of PD1 (adoptive transfer of transgenic *Pdcd1*^{-/-} LCMV-specific
281 P14 T cells) or to PDL1 blockade^{18,19}. However, mitochondrial biomass was not
282 substantially altered by PD1 blockade in TILs, even though the blockade resulted
283 in effective tumour regression¹⁹.

284 Sampling of T cells at different stages of exhaustion (a dynamic process reflecting
285 a continuum of dysfunction⁴¹) and from microenvironments comprising different
286 substrate availabilities may explain at least some of the variance observed.

287 However, other factors may also contribute. Texh cells are known to be
288 heterogeneous, with two principal subsets defined by both their level of PD1
289 expression and reciprocal expression of the transcription factors Tbet and
290 Eomes³⁰. The Tbet^{hi}Eomes^{lo}PD1^{int} subset show greatest proliferative potential,
291 are the lineage precursors of their more cytotoxic Tbet^{lo}Eomes^{hi}PD1^{hi}
292 counterparts and are the dominant subset responding to PD1-PDL1 blockade⁴².
293 In LCMV, the Tbet^{lo}Eomes^{hi}PD1^{hi} subset was shown to have more severe
294 metabolic dysfunction consistent with its 'terminally exhausted' phenotype. The
295 distribution of these subsets amongst TIL is not currently clear^{19,38} but the lack

296 of a metabolic response to PD1 blockade¹⁹ suggests they are likely to be
297 enriched for 'terminally exhausted' cells. The observation that adoptive transfer
298 of TIL into an acute viral infection setting (VV-OVA) fails to restore their
299 function, is consistent with it¹⁹. Despite this, a bioenergetic revival following PD1
300 blockade, such as is seen in chronic infection¹⁸, may still occur in tumour models
301 and contribute to revival of tumoricidal function. However, it may be occurring
302 out with the tumour microenvironment in less terminally-exhausted T cells and
303 consequently not be visible within it.

304
305 Exhaustion is a spectrum rather than a binary state¹⁵ and both its extent and the
306 nature of its induction may differ depending on the context. TILs show an
307 exhausted phenotype only within the tumour microenvironment, remaining fully
308 functional in the periphery of tumour-bearing mice^{15,19} or in human cancer^{22,38}.
309 Indeed, TIL metabolic dysfunction could not be reproduced on co-culture of CD8⁺
310 T cells with isolated tumour cells in vitro¹⁹ indicating an important and complex
311 role of the microenvironment in its induction. By contrast, exhaustion is
312 apparent between different mice rather than between different anatomical
313 locations in the same mouse⁴³. Those receiving the clone 13 strain of LCMV
314 develop chronic infection and exhaustion, while those receiving the highly
315 related Armstrong strain have robust memory differentiation following an acute,
316 self-terminating infection.

317 It is clear that T cell bioenergetics vary markedly between different tissues
318 during infection^{18,19} and also between distinct tumour models¹⁹, highlighting the
319 importance of context for their development and maintenance. Given evidence
320 demonstrating the importance of local microenvironmental signals, substrate
321 availability and the phenotype of neighbouring cell types (both immune and
322 stromal), it is less surprising that, in different contexts, T cells come to be
323 exhausted by different paths.

324

325 **Early T cell metabolism may control exhaustion, but what controls early T** 326 **cell metabolism?**

327 Multiple lines of evidence support the assertion that metabolic pathways control
328 the development of T cell exhaustion, although exact in vivo mechanisms remain
329 to be fully elucidated. But what signals control those metabolic pathways early
330 after activation? A series of elegant studies have now identified major signalling
331 pathways controlling the balance between effector and memory differentiation.
332 More recently, these pathways have been investigated in both TIL and chronic
333 infection models of T cell exhaustion, indicating an important role there also.

334

335 During T cell activation and proliferation the intracellular serine/threonine
336 kinase mTOR (mechanistic target of rapamycin) integrates diverse
337 environmental signals including those derived from nutrient, cytokine and TCR
338 signalling pathways⁴⁴. Effector differentiation is favoured by TCR-triggered
339 PI3K/Akt activity, driving mTOR-dependent¹⁴ inactivation of the transcription
340 factor FOXO1⁴⁵. FOXO1 serves to repress effector function and promote memory
341 formation by increasing expression of the transcription factor Eomesodermin
342 and by concurrently repressing Tbet⁴⁶. In the presence of TCR signals, such
343 FOXO1 inactivation tips this balance towards Tbet-mediated effector
344 differentiation (**Fig. 3**). It is also clear that many signals feed into this complex

345 junction, including those dependent on the local availability of metabolic
346 substrates considered above, including hypoxia, glucose, immunometabolites
347 such as S2HG, PEP and amino acids such as L-arginine^{25,29,31,33}.

348

349 **FIGURE 3: metabolism integrates inhibitory and TCR signalling in** 350 **exhaustion**

351

352 **Inhibitory receptor control: PD1 and T cell metabolism**

353 It is now becoming apparent that early inhibitory receptor signalling in activated
354 T cells can modulate T cell metabolism to influence the later development of
355 exhaustion. It has long been known that T cell exhaustion is driven by high levels
356 of sustained TCR signalling^{41,47}. However recent work has shown that, despite
357 ongoing antigen exposure and TCR triggering, downstream PI3K/AKT/mTOR
358 signalling in exhaustion is not similarly sustained^{18,48} – in other words, persistent
359 antigen may drive exhaustion but this does not correspond to persistent
360 downstream TCR signals. PD1 expression is induced on antigen-stimulated cells
361 by TCR signals through NFATc⁴⁹⁻⁵¹. However, PD1 ligation is capable of
362 ‘desensitising’ the TCR signal, allowing a degree of ongoing nuclear FOXO1
363 activity that in turn sustains both further PD1 expression and ongoing TCR
364 desensitisation⁴⁸. This establishes a self-reinforcing cycle that restrains
365 development of full effector function with an associated switch away from
366 aerobic glycolytic metabolism^{48,52}.

367 These data suggest that persistent PD1 signalling actually facilitates T cell
368 survival in the face of persistent antigenic stimulus. By desensitising TCR signals,
369 PD1 may actually serve to maintain a degree of bioenergetic fitness in the face of
370 persistent antigen exposure. We have seen that exhausted T cells do show
371 evidence of metabolic dysfunction^{18,19}. However, without persistent PD1 signals
372 this could be much worse. PD1 can therefore act to facilitate an ongoing, if
373 attenuated, T cell response in the context of persistent TCR activation⁴⁸. This
374 system is complex, however – genetic deletion of PD1 can counterintuitively
375 allow the later accumulation of ‘terminally exhausted’ cells (*Pdcd1*^{-/-})⁵³. However,
376 the details of how PD1 facilitates survival with attenuated function remain
377 unclear and its effects are likely to vary with the time post stimulation (early
378 versus late, see discussion of PGC1a below).

379 This role for PD1 in T cell differentiation also makes evolutionary sense. It is
380 likely that exhaustion has evolved as a means of carefully balancing control of
381 chronic infection with the risk of causing damaging immunopathology¹⁷. During
382 acute T cell activation, a feedback loop is established whereby strong TCR signals
383 drive Akt-mediated cytotoxicity (**Fig. 3**) but also drive PD1 expression, limiting
384 that TCR signalling. Should TCR signals persist, they will be desensitised by PD1,
385 allowing preservation of bioenergetic fitness sufficient to sustain a longer-term,
386 albeit more restrained, T cell response.

387 These new insights into metabolic control of T cell exhaustion have important
388 implications for the design of therapies aimed at its modulation. This schema
389 may also initially seem at odds with observations that PD1 blockade can promote
390 restoration of cytotoxicity in both TILs¹⁶ and in chronic viral infection⁵⁴.

391 However, PD1-driven desensitisation of TCR signals and preservation of some
392 metabolic fitness serves to keep T cells in the fight – the response to persistent
393 antigen is, after all, more of a marathon than a sprint. Reversing exhaustion

394 through inhibitory receptor blockade shows much promise as a therapeutic
395 approach in chronic infection or cancer^{23,54,55}. PD1 blockade can clearly induce
396 increased cytotoxicity of exhausted CD8⁺ T cells. However, if this occurs through
397 promoting 'terminal' differentiation of more cytotoxic Tbet^{lo}Eomes^{hi}PD1^{hi} cells at
398 the expense of a more durable and metabolically-fit Tbet^{hi}Eomes^{lo}PD1^{int}
399 population^{30,53}, such therapy could effectively undercut the immune systems
400 capacity to maintain an ongoing response. The predicted result would be
401 eradication of the targeted antigen (whether tumour or virus) in some cases
402 where induced cytotoxicity was sufficient, but perhaps a 'deeper' state of
403 subsequent exhaustion where it isn't and residual metabolic fitness was lost
404 through inhibitory receptor blockade.

405

406 What is less clear at this stage is exactly how this perpetuating cycle is
407 established in the first place. While the microenvironmental signals discussed
408 above are important, costimulatory signalling is very likely to also contribute. A
409 lack of costimulatory signals during TCR triggering results in a hyporesponsive
410 state termed anergy³². Costimulatory signals, such as those received through
411 CD28 binding^{56,57} or cytokines such as IL-2 (ref ⁵⁸), are critical for upregulation of
412 glucose, AA transporters and the FAO regulator CPT1a that facilitate the
413 metabolic switch underlying T cell activation^{13,57}. A failure of such initial
414 costimulation results in comparative metabolic quiescence and anergy⁵⁹.
415 Exhaustion is distinct from anergy: T cells have received sufficient stimulatory
416 signals to activate, but they continue to receive them and become dysfunctional.
417 However, inadequate costimulation (rather than absent costimulation as in
418 anergy) also contributes to development of exhaustion^{60,61} and is likely to
419 promote the early PD1-driven cycle of bioenergetic compromise promoting later
420 exhaustion (**Fig. 3**).

421

422 **Other inhibitory receptors**

423 Just as the precise nature of metabolic dysfunction of exhausted cells appears
424 different in different contexts (see above), so the mechanism by which it
425 develops may also be distinct. A range of coinhibitory receptors are highly
426 expressed on exhausted T cells and play distinct but synergistic roles in driving
427 that phenotype⁴⁹. Mitochondrial dysfunction of TILs is apparent in the tumour
428 microenvironment of multiple mouse models^{19,20,31} and in human cancers^{19,38}.
429 However, the degree of metabolic dysfunction does not directly parallel the
430 concurrent level of inhibitory receptor expression^{19,26} in each context.
431 Exhaustion can clearly develop in the absence of specific inhibitory receptor
432 expression^{18,53} and inhibitory receptor expression can be present without the
433 development of exhaustion, for example where Vhl/Hif signals are absent²⁵.
434 Further, whereas metabolic changes are important drivers of exhaustion early
435 during TCR signalling, exhaustion can be at least partially reversed at later stages
436 with increased cytotoxicity despite no clear modulation of metabolism^{18,19}.
437 Together, these data suggest that multiple, redundant signals must contribute to
438 the development of exhaustion and its associated metabolic dysfunction. The
439 summated input from these cell surface and local environmental signals
440 contribute to the resulting degree of exhaustion.

441

442 **mTOR and metabolism in exhaustion**

443 During T cell activation mTOR acts to integrate multiple signals including from
444 metabolic, cytokine and TCR signalling pathways⁴⁴. Perhaps unsurprisingly,
445 mTOR activity has therefore been shown to have a central role in T cell
446 metabolism and, more recently, in exhaustion. It was, however, initially a
447 surprise when the immunosuppressive drug rapamycin, an inhibitor of mTOR,
448 was found to promote T cell memory differentiation and the associated switch
449 away from aerobic glycolysis towards FAO metabolism^{8,14}. It has been used for
450 some time as an immunosuppressive agent capable of limiting TCR-driven
451 activation and proliferation. However, mTOR has since been shown to directly
452 promote phosphorylation and inactivation of FOXO1, promoting effector
453 differentiation through downstream Tbet-mediated effects^{45,46,62,63}.
454 Early mTOR blockade in the LCMV-cl13 chronic viral infection model also
455 modestly improved mitochondrial fitness of exhausted cells (depolarisation and
456 size¹⁸) suggesting that chronic mTOR signals downstream of the TCR can
457 contribute to the metabolic changes of exhaustion, even if TCR signals are
458 downtuned by PD1. It has been observed that blocking mTOR actually impairs
459 TIL response to PD1 blockade, suggesting that ongoing mTOR signalling to at
460 least some degree may be needed to preserve effector function^{18,19}.

461
462 The evidence for mTOR involvement in the metabolic phenotype of exhaustion
463 highlights the complex, dynamic balance between 'terminal' effector function and
464 T cell persistence. While mTOR signalling may in part drive the cell away from
465 mitochondrial metabolism toward anaerobic glycolysis, these changes are a
466 necessary part of effector differentiation. Both are required for adequate ongoing
467 T cell responses to persistent antigen and the degree to which either path is
468 inhibited impacts on the resulting phenotype³⁰.

469
470 **FOXO signals and exhaustion: death, survival and metabolism**
471 FOXO transcription factors integrate signals from growth factors, substrate
472 availability and inflammation and in turn regulate cellular growth, differentiation
473 and metabolism⁶⁴. The balance of FOXO signalling is clearly important in
474 directing CD4⁺ regulatory T cell differentiation^{65,66}, however this balance also
475 plays an important cell-intrinsic role in CD8⁺ T cell differentiation and in
476 directing metabolic dysfunction of exhaustion. FOXOs 1 and 3 are expressed in T
477 cells⁶⁴. FOXO1 activity promotes T cell survival and persistence^{67,68} while FOXO3
478 activity limits survival through pro-apoptotic effects mediated by Bim and
479 PUMA⁶⁹⁻⁷¹. Ablating FOXO3 activity (*Foxo3*^{-/-} CD8⁺ T cells) restricts the
480 contraction phase post-stimulation allowing greater persistence of effector CD8⁺
481 T cells⁷², facilitating clearance of chronic LCMV infection⁷³ and control of HIV in
482 humans⁷⁴. By contrast, FOXO1 signalling can maintain a pluripotent stem cell
483 phenotype⁷⁵ and is required for differentiation of effector CD8⁺ T cells into a
484 functional memory subset⁷⁶ through its downstream modulation of the
485 transcription factors Tbet and Eomes⁴⁶. It is therefore likely that the balance of
486 FOXO signalling following TCR signals will determine the balance of effector fate
487 and cellular persistence during chronic antigen exposure.
488 However, the balance of FOXO-driven regulation is complex⁶⁴ and a complete
489 picture of its contribution to the metabolic dysfunction of exhaustion has yet to
490 emerge with several apparent contradictions persisting. For example, the role of
491 FOXO3 in promoting apoptosis and effector differentiation is at odds with its

492 apparent ability to maintain PGC1a expression and metabolic fitness (described
493 below). Similarly, despite its role in promoting stem cell-like persistence, FOXO1
494 also appears to be necessary for differentiation of the terminally exhausted
495 Tbet^{lo}Eomes^{hi}PD1^{hi} subset⁴⁸. These apparent discrepancies may reflect the
496 complex integration by FOXOs of multiple other pathways including hypoxic
497 signalling⁷⁷, metabolic pathways⁷⁸ and nutrient availability⁷⁹. A future challenge
498 will be to fully understand the complex role of these transcription factors in
499 regulating metabolism during exhaustion.

500

501 **PGC1a: regulation at the heart of metabolic exhaustion?**

502 The ability to sustain energy production during rapid cellular proliferation
503 requires mitochondrial biogenesis and the induction of gene expression
504 programmes regulating substrate uptake and use. This process is controlled by
505 the transcriptional coactivator PGC1a⁸⁰ through control of both nuclear
506 respiratory factors (Nrf) and mitochondrial transcription factor A (Tfam)⁸¹. It is
507 now clear that PGC1a is central to the metabolic dysfunction of exhaustion in
508 both LCMV and tumour models. PGC1a shows reduced levels of expression in
509 both TIL¹⁹ and Texh in the LCMV model¹⁸, while its overexpression limits
510 mitochondrial dysfunction^{18,19}, facilitating the emergence of polyfunctional CD8
511 T cells¹⁸ that drive tumour infiltration and regression¹⁹.

512 The FOXO family of transcription factors - both FOXO1^{19,82} and FOXO3^{19,83} - have
513 each been shown to promote PGC1a expression. Conversely, PGC1a expression is
514 repressed by TCR-driven Akt signalling⁸² which in turn phosphorylates and
515 inactivates FOXOs 1 and 3⁸¹. Therefore persistent antigen/TCR signalling may
516 serve to maintain Akt signalling and promote effector differentiation, but
517 metabolic fitness will also be curtailed as FOXO-driven PGC1a expression is
518 restricted (**Fig. 3**).

519 It remains unclear, however, exactly how complex signals from multiple
520 inhibitory receptors including PD1 are integrated to modulate PGC1a levels and
521 whether this is the only or the most important transcription factor modulating
522 early metabolic fitness. Absence of PD1 from the onset of a T cell response
523 (through genetic deletion, *Pdcd1*^{-/-}) facilitates a later expansion of exhausted
524 cells in the absence of inhibitory signals restraining cytotoxicity and terminal
525 differentiation⁵³. This demonstrates that PD1 is not essential for the
526 development of exhaustion and its expression at certain points may even limit its
527 development. Genetic ablation of PD1 also results in higher expression of
528 PGC1a¹⁸ suggesting that, if PD1 is acting to preserve a degree of early
529 bioenergetic fitness and to facilitate later survival despite ongoing stimulation, it
530 is unlikely to be doing so through PGC1a. Other data showing delayed tumour
531 progression after PD-1 blockade in the absence of metabolic changes in TIL also
532 suggest additional mechanisms²⁷. Further studies will be required to clarify the
533 complexity of these signalling events.

534

535 **Autoimmune disease, metabolism and exhaustion**

536 T cell exhaustion regulates cellular immunity during the response to persistent
537 antigenic stimulation, whether that is during chronic viral infection or in the
538 tumour microenvironment. Exhaustion has also been inversely associated with a
539 relapsing course of autoinflammatory disease, another instance of immune
540 reactivity to persistent antigen (in this case, 'self'). A transcriptional signature

541 specific for exhaustion correlates with a reduced rate of relapse in multiple
542 diseases⁶¹ and features of T cell exhaustion correlate with favourable response
543 during immunotherapy of type 1 diabetes⁸⁴. Such observations are consistent
544 with the evolution of exhaustion as a mechanism to limit immunopathology in
545 the face of persistent TCR stimulation^{17,41}. A broadly similar phenotype of
546 exhaustion is apparent in the distinct contexts of chronic viral infection and the
547 tumour microenvironment. These broad similarities are accompanied by specific
548 differences, too, likely driven by distinct signals received in each context. It is
549 likely that immunometabolism also plays a key role in control of effector,
550 persistent or exhausted T cell responses in autoimmunity.

551
552 It is clear that metabolic 'switches' are not only important for the initiation of T
553 cell activation¹³ but are also responsible for controlling an ongoing balance of
554 effector/memory differentiation and exhaustion^{18,19}. Many studies have sought
555 to directly compare measures of metabolic fitness in autoinflammatory disease,
556 typically making comparison with uninflamed, healthy control tissues and
557 sampling cells during or after active autoinflammation and immunosuppressive
558 therapy⁸⁵. In this context it is unsurprising to find altered metabolism as a
559 secondary consequence of immune activation or therapeutic modulation –
560 indeed it would be surprising not to find it. This highlights the need for
561 appropriate control for variable proliferation and immune activation in models
562 and assays of immunometabolism, such as those used during LCMV models
563 (where adoptive transfer of small numbers of transgenic, LCMV-specific P14 T
564 cells facilitates their direct comparison with host cells in an equivalent
565 context^{18,53}). Evidence to support a primary role of exhaustion-associated
566 metabolic dysfunction in autoinflammatory disease is more limited.

567
568 Several recent genetic studies have, however, implicated exhaustion-associated
569 metabolic pathways in the onset and progression of multiple autoimmune
570 phenotypes. Perhaps the most striking example is the recurrent association of a
571 variant allele in neutrophil cytosolic factor 1 (Ncf1) with onset or severity of
572 multiple autoinflammatory diseases⁸⁶. Ncf1 is a necessary for activation of
573 NADPH oxidase and consequent production of superoxide during the respiratory
574 burst. Reduction in ROS levels in the presence of a mutant allele leads to
575 increased severity and relapse in mouse models of multiple sclerosis and
576 rheumatoid arthritis^{86,87} and shows association with multiple human
577 autoimmune diseases⁸⁶. Redox signalling can be controlled by PD1 signalling in T
578 cells⁸⁸ and ROS generation has been implicated in the metabolic dysfunction of
579 exhaustion^{18,19}. However, a direct link between NCF1 function and exhaustion
580 remains to be tested.

581 The process of T cell exhaustion has thus far been associated with the
582 progression rather than onset of autoinflammatory disease^{61,84}, although
583 exhaustion-associated inhibitory receptors have also been linked to broken
584 tolerance¹⁷. A genome-wide association study of clinical outcome has identified a
585 variant allele in FOXO3 associated with disease progression (but not
586 susceptibility) in Crohn's disease, rheumatoid arthritis and malaria infection⁸⁹.
587 Alongside the association of T cell exhaustion with outcome in Crohn's disease⁶¹
588 this suggests a possible role for FOXO3-driven metabolic control of T cell
589 exhaustion in autoinflammatory disease. However, FOXO3 the Crohn's associated

590 variant allele has not as yet been functionally tested in T cells and a monocyte-
591 intrinsic signalling pathway directing TGF- β and TNF production is at least partly
592 responsible for the association⁹⁰.

593

594 **Summary**

595 During T cell activation changes in cellular metabolism are inevitably required to
596 sustain the rapid changes in cell function. Increasing evidence suggests that
597 metabolic changes do not simply react to changes in T cell differentiation but
598 drive them. It is now apparent that the integration of complex early signals from
599 a cells microenvironment is critical both for initiating activation and for
600 regulating the nature of 'downstream' differentiation. These signals direct
601 phenotypes of effector function and memory development, but also of T cell
602 exhaustion during chronic antigen exposure.

603 In both chronic infection and tumour models where exhaustion is apparent,
604 metabolic changes in T cells occurred rapidly, prior to the emergence of
605 exhaustion *per se*, and were maintained into later stages. Multiple observations
606 therefore confirm a prominent role for cellular metabolism in 'upstream' control
607 of exhaustion rather than simply reflecting changes secondary to its
608 development.

609 A wealth of evidence also indicates that signalling pathways promoting cellular
610 persistence/survival can be dissociated from those controlling effector function.
611 Signals promoting aerobic glycolysis during activation may enhance effector
612 differentiation but this may be at the expense of memory potential. Conversely,
613 the use of alternative pathways such as lipid metabolism at the expense of
614 aerobic glycolysis may do the opposite. However, both processes are required in
615 parallel to appropriately mount an effective, durable T cell response to persistent
616 antigen³⁰. As such, the metabolic balance between effector function and
617 persistence represents another example of the 'Goldilocks' principle, in which
618 effector and memory differentiation must be neither too much nor too little, but
619 'just right' in order to appropriately deal with infection without the development
620 of immunopathology. Future challenges will include quantifying the extent to
621 which each metabolic pathway contributes to T cell function, how each is
622 controlled and how we might harness that knowledge to modulate metabolism
623 and hence immunity. To achieve this, our understanding of T cell metabolism
624 during human disease will have to catch up with the advances driven by
625 informative mouse models of cancer and infection. However, modulating
626 immunometabolism shows promise as a means of controlling chronic T cell
627 responses in infection, cancer and autoinflammatory disease.

628

629

630

631 **REFERENCES**

- 632 1 MacIver, N. J., Michalek, R. D. & Rathmell, J. C. Metabolic regulation of T
633 lymphocytes. *Annu Rev Immunol* **31**, 259-283, doi:10.1146/annurev-
634 immunol-032712-095956 (2013).
- 635 2 Chang, C. H. *et al.* Posttranscriptional control of T cell effector function by
636 aerobic glycolysis. *Cell* **153**, 1239-1251, doi:10.1016/j.cell.2013.05.016
637 (2013).

638 3 Vander Heiden, M. G., Cantley, L. C. & Thompson, C. B. Understanding the
639 Warburg effect: the metabolic requirements of cell proliferation. *Science*
640 **324**, 1029-1033, doi:10.1126/science.1160809 (2009).

641 4 Pearce, E. L. & Pearce, E. J. Metabolic pathways in immune cell activation
642 and quiescence. *Immunity* **38**, 633-643,
643 doi:10.1016/j.immuni.2013.04.005 (2013).

644 5 Anastasiou, D. *et al.* Inhibition of pyruvate kinase M2 by reactive oxygen
645 species contributes to cellular antioxidant responses. *Science* **334**, 1278-
646 1283, doi:10.1126/science.1211485 (2011).

647 6 Zhao, S. *et al.* ATP-Citrate Lyase Controls a Glucose-to-Acetate Metabolic
648 Switch. *Cell reports* **17**, 1037-1052, doi:10.1016/j.celrep.2016.09.069
649 (2016).

650 7 van der Windt, G. J. *et al.* Mitochondrial respiratory capacity is a critical
651 regulator of CD8+ T cell memory development. *Immunity* **36**, 68-78,
652 doi:10.1016/j.immuni.2011.12.007 (2012).

653 8 Pearce, E. L. *et al.* Enhancing CD8 T-cell memory by modulating fatty acid
654 metabolism. *Nature* **460**, 103-107, doi:10.1038/nature08097 (2009).

655 9 O'Sullivan, D. *et al.* Memory CD8(+) T cells use cell-intrinsic lipolysis to
656 support the metabolic programming necessary for development.
657 *Immunity* **41**, 75-88, doi:10.1016/j.immuni.2014.06.005 (2014).

658 10 Buck, M. D. *et al.* Mitochondrial Dynamics Controls T Cell Fate through
659 Metabolic Programming. *Cell* **166**, 63-76, doi:10.1016/j.cell.2016.05.035
660 (2016).

661 11 van der Windt, G. J. *et al.* CD8 memory T cells have a bioenergetic
662 advantage that underlies their rapid recall ability. *Proceedings of the*
663 *National Academy of Sciences of the United States of America* **110**, 14336-
664 14341, doi:10.1073/pnas.1221740110 (2013).

665 12 Nomura, M. *et al.* Fatty acid oxidation in macrophage polarization. *Nature*
666 *immunology* **17**, 216-217, doi:10.1038/ni.3366 (2016).

667 13 Buck, M. D., O'Sullivan, D. & Pearce, E. L. T cell metabolism drives
668 immunity. *The Journal of experimental medicine* **212**, 1345-1360,
669 doi:10.1084/jem.20151159 (2015).

670 14 Araki, K. *et al.* mTOR regulates memory CD8 T-cell differentiation. *Nature*
671 **460**, 108-112, doi:10.1038/nature08155 (2009).

672 15 Wherry, E. J. T cell exhaustion. *Nature immunology* **12**, 492-499 (2011).

673 16 Pauken, K. E. & Wherry, E. J. Overcoming T cell exhaustion in infection and
674 cancer. *Trends in immunology* **36**, 265-276, doi:10.1016/j.it.2015.02.008
675 (2015).

676 17 McKinney, E. F. & Smith, K. G. T-cell exhaustion: understanding the
677 interface of chronic viral and autoinflammatory diseases. *Immunol Cell*
678 *Biol* **94**, 935-942, doi:10.1038/icb.2016.81 (2016).

679 18 Bengsch, B. *et al.* Bioenergetic Insufficiencies Due to Metabolic Alterations
680 Regulated by the Inhibitory Receptor PD-1 Are an Early Driver of CD8(+) T
681 Cell Exhaustion. *Immunity* **45**, 358-373,
682 doi:10.1016/j.immuni.2016.07.008 (2016).

683 19 Scharping, N. E. *et al.* The Tumor Microenvironment Represses T Cell
684 Mitochondrial Biogenesis to Drive Intratumoral T Cell Metabolic
685 Insufficiency and Dysfunction. *Immunity* **45**, 701-703,
686 doi:10.1016/j.immuni.2016.08.009 (2016).

- 687 20 Chang, C. H. *et al.* Metabolic Competition in the Tumor Microenvironment
688 Is a Driver of Cancer Progression. *Cell* **162**, 1229-1241,
689 doi:10.1016/j.cell.2015.08.016 (2015).
- 690 21 Wherry, E. J. *et al.* Molecular signature of CD8+ T cell exhaustion during
691 chronic viral infection. *Immunity* **27**, 670-684 (2007).
- 692 22 Baitsch, L. *et al.* Exhaustion of tumor-specific CD8(+) T cells in metastases
693 from melanoma patients. *The Journal of clinical investigation* **121**, 2350-
694 2360, doi:10.1172/JCI46102 (2011).
- 695 23 Gubin, M. M. *et al.* Checkpoint blockade cancer immunotherapy targets
696 tumour-specific mutant antigens. *Nature* **515**, 577-581,
697 doi:10.1038/nature13988 (2014).
- 698 24 Harris, A. L. Hypoxia--a key regulatory factor in tumour growth. *Nat Rev*
699 *Cancer* **2**, 38-47, doi:10.1038/nrc704 (2002).
- 700 25 Doedens, A. L. *et al.* Hypoxia-inducible factors enhance the effector
701 responses of CD8(+) T cells to persistent antigen. *Nature immunology* **14**,
702 1173-1182, doi:10.1038/ni.2714 (2013).
- 703 26 Tyrakis, P. A. *et al.* S-2-hydroxyglutarate regulates CD8+ T-lymphocyte
704 fate. *Nature* **540**, 236-241, doi:10.1038/nature20165 (2016).
- 705 27 Zhang, Y. *et al.* Enhancing CD8(+) T Cell Fatty Acid Catabolism within a
706 Metabolically Challenging Tumor Microenvironment Increases the
707 Efficacy of Melanoma Immunotherapy. *Cancer Cell* **32**, 377-391 e379,
708 doi:10.1016/j.ccell.2017.08.004 (2017).
- 709 28 Sukumar, M. *et al.* Inhibiting glycolytic metabolism enhances CD8+ T cell
710 memory and antitumor function. *The Journal of clinical investigation* **123**,
711 4479-4488, doi:10.1172/JCI69589 (2013).
- 712 29 Phan, A. T. *et al.* Constitutive Glycolytic Metabolism Supports CD8+ T Cell
713 Effector Memory Differentiation during Viral Infection. *Immunity* **45**,
714 1024-1037, doi:10.1016/j.immuni.2016.10.017 (2016).
- 715 30 Paley, M. A. *et al.* Progenitor and terminal subsets of CD8+ T cells
716 cooperate to contain chronic viral infection. *Science* **338**, 1220-1225,
717 doi:10.1126/science.1229620 (2012).
- 718 31 Ho, P. C. *et al.* Phosphoenolpyruvate Is a Metabolic Checkpoint of Anti-
719 tumor T Cell Responses. *Cell* **162**, 1217-1228,
720 doi:10.1016/j.cell.2015.08.012 (2015).
- 721 32 Schwartz, R. H. T cell anergy. *Annu Rev Immunol* **21**, 305-334 (2003).
- 722 33 Geiger, R. *et al.* L-Arginine Modulates T Cell Metabolism and Enhances
723 Survival and Anti-tumor Activity. *Cell* **167**, 829-842 e813,
724 doi:10.1016/j.cell.2016.09.031 (2016).
- 725 34 Sinclair, L. V. *et al.* Control of amino-acid transport by antigen receptors
726 coordinates the metabolic reprogramming essential for T cell
727 differentiation. *Nature immunology* **14**, 500-508, doi:10.1038/ni.2556
728 (2013).
- 729 35 Nakaya, M. *et al.* Inflammatory T cell responses rely on amino acid
730 transporter ASCT2 facilitation of glutamine uptake and mTORC1 kinase
731 activation. *Immunity* **40**, 692-705, doi:10.1016/j.immuni.2014.04.007
732 (2014).
- 733 36 Wang, R. *et al.* The transcription factor Myc controls metabolic
734 reprogramming upon T lymphocyte activation. *Immunity* **35**, 871-882,
735 doi:10.1016/j.immuni.2011.09.021 (2011).

- 736 37 Ghoneim, H. E. *et al.* De Novo Epigenetic Programs Inhibit PD-1 Blockade-
737 Mediated T Cell Rejuvenation. *Cell* **170**, 142-157 e119,
738 doi:10.1016/j.cell.2017.06.007 (2017).
- 739 38 Siska, P. J. *et al.* Mitochondrial dysregulation and glycolytic insufficiency
740 functionally impair CD8 T cells infiltrating human renal cell carcinoma. *JCI*
741 *Insight* **2**, doi:10.1172/jci.insight.93411 (2017).
- 742 39 Schurich, A. *et al.* Distinct Metabolic Requirements of Exhausted and
743 Functional Virus-Specific CD8 T Cells in the Same Host. *Cell reports* **16**,
744 1243-1252, doi:10.1016/j.celrep.2016.06.078 (2016).
- 745 40 Fiscaro, P. *et al.* Targeting mitochondrial dysfunction can restore
746 antiviral activity of exhausted HBV-specific CD8 T cells in chronic
747 hepatitis B. *Nature medicine* **23**, 327-336, doi:10.1038/nm.4275 (2017).
- 748 41 Wherry, E. J. & Kurachi, M. Molecular and cellular insights into T cell
749 exhaustion. *Nature reviews. Immunology* **15**, 486-499,
750 doi:10.1038/nri3862 (2015).
- 751 42 Blackburn, S. D., Shin, H., Freeman, G. J. & Wherry, E. J. Selective expansion
752 of a subset of exhausted CD8 T cells by alphaPD-L1 blockade. *Proceedings*
753 *of the National Academy of Sciences of the United States of America* **105**,
754 15016-15021, doi:10.1073/pnas.0801497105 (2008).
- 755 43 Oldstone, M. B. Anatomy of viral persistence. *PLoS pathogens* **5**,
756 e1000523, doi:10.1371/journal.ppat.1000523 (2009).
- 757 44 Chi, H. Regulation and function of mTOR signalling in T cell fate decisions.
758 *Nature reviews. Immunology* **12**, 325-338, doi:10.1038/nri3198 (2012).
- 759 45 Rao, R. R., Li, Q., Odunsi, K. & Shrikant, P. A. The mTOR kinase determines
760 effector versus memory CD8+ T cell fate by regulating the expression of
761 transcription factors T-bet and Eomesodermin. *Immunity* **32**, 67-78,
762 doi:10.1016/j.immuni.2009.10.010 (2010).
- 763 46 Rao, R. R., Li, Q., Gubbels Bupp, M. R. & Shrikant, P. A. Transcription factor
764 Foxo1 represses T-bet-mediated effector functions and promotes memory
765 CD8(+) T cell differentiation. *Immunity* **36**, 374-387,
766 doi:10.1016/j.immuni.2012.01.015 (2012).
- 767 47 Mueller, S. N. & Ahmed, R. High antigen levels are the cause of T cell
768 exhaustion during chronic viral infection. *Proceedings of the National*
769 *Academy of Sciences of the United States of America* **106**, 8623-8628,
770 doi:10.1073/pnas.0809818106 (2009).
- 771 48 Staron, M. M. *et al.* The transcription factor FoxO1 sustains expression of
772 the inhibitory receptor PD-1 and survival of antiviral CD8(+) T cells
773 during chronic infection. *Immunity* **41**, 802-814,
774 doi:10.1016/j.immuni.2014.10.013 (2014).
- 775 49 Blackburn, S. D. *et al.* Coregulation of CD8+ T cell exhaustion by multiple
776 inhibitory receptors during chronic viral infection. *Nature immunology*
777 **10**, 29-37, doi:10.1038/ni.1679 (2009).
- 778 50 Oestreich, K. J., Yoon, H., Ahmed, R. & Boss, J. M. NFATc1 regulates PD-1
779 expression upon T cell activation. *Journal of immunology* **181**, 4832-4839
780 (2008).
- 781 51 Martinez, G. J. *et al.* The transcription factor NFAT promotes exhaustion of
782 activated CD8(+) T cells. *Immunity* **42**, 265-278,
783 doi:10.1016/j.immuni.2015.01.006 (2015).

784 52 Patsoukis, N. *et al.* PD-1 alters T-cell metabolic reprogramming by
785 inhibiting glycolysis and promoting lipolysis and fatty acid oxidation.
786 *Nature communications* **6**, 6692, doi:10.1038/ncomms7692 (2015).

787 53 Odorizzi, P. M., Pauken, K. E., Paley, M. A., Sharpe, A. & Wherry, E. J.
788 Genetic absence of PD-1 promotes accumulation of terminally
789 differentiated exhausted CD8+ T cells. *The Journal of experimental*
790 *medicine* **212**, 1125-1137, doi:10.1084/jem.20142237 (2015).

791 54 Barber, D. L. *et al.* Restoring function in exhausted CD8 T cells during
792 chronic viral infection. *Nature* **439**, 682-687 (2006).

793 55 McKinney, E. F. & Smith, K. G. T cell exhaustion and immune-mediated
794 disease-the potential for therapeutic exhaustion. *Current opinion in*
795 *immunology* **43**, 74-80, doi:10.1016/j.coi.2016.09.005 (2016).

796 56 Frauwirth, K. A. *et al.* The CD28 signaling pathway regulates glucose
797 metabolism. *Immunity* **16**, 769-777 (2002).

798 57 Klein Geltink, R. I. *et al.* Mitochondrial Priming by CD28. *Cell* **171**, 385-397
799 e311, doi:10.1016/j.cell.2017.08.018 (2017).

800 58 Wieman, H. L., Wofford, J. A. & Rathmell, J. C. Cytokine stimulation
801 promotes glucose uptake via phosphatidylinositol-3 kinase/Akt
802 regulation of Glut1 activity and trafficking. *Mol Biol Cell* **18**, 1437-1446,
803 doi:10.1091/mbc.E06-07-0593 (2007).

804 59 Zheng, Y., Delgoffe, G. M., Meyer, C. F., Chan, W. & Powell, J. D. Anergic T
805 cells are metabolically anergic. *Journal of immunology* **183**, 6095-6101,
806 doi:10.4049/jimmunol.0803510 (2009).

807 60 Aubert, R. D. *et al.* Antigen-specific CD4 T-cell help rescues exhausted CD8
808 T cells during chronic viral infection. *Proceedings of the National Academy*
809 *of Sciences of the United States of America* **108**, 21182-21187,
810 doi:10.1073/pnas.1118450109 (2011).

811 61 McKinney, E. F., Lee, J. C., Jayne, D. R. W., Lyons, P. A. & Smith, K. G. C. T-cell
812 exhaustion, co-stimulation and clinical outcome in autoimmunity and
813 infection. *Nature* **523**, 612-616, doi:10.1038/nature14468 (2015).

814 62 Pollizzi, K. N. *et al.* Asymmetric inheritance of mTORC1 kinase activity
815 during division dictates CD8(+) T cell differentiation. *Nature immunology*
816 **17**, 704-711, doi:10.1038/ni.3438 (2016).

817 63 Pollizzi, K. N. *et al.* mTORC1 and mTORC2 selectively regulate CD8(+) T
818 cell differentiation. *The Journal of clinical investigation* **125**, 2090-2108,
819 doi:10.1172/JCI77746 (2015).

820 64 Hedrick, S. M., Hess Michelini, R., Doedens, A. L., Goldrath, A. W. & Stone, E.
821 L. FOXO transcription factors throughout T cell biology. *Nature reviews.*
822 *Immunology* **12**, 649-661, doi:10.1038/nri3278 (2012).

823 65 Ouyang, W. *et al.* Foxo proteins cooperatively control the differentiation of
824 Foxp3+ regulatory T cells. *Nature immunology* **11**, 618-627,
825 doi:10.1038/ni.1884 (2010).

826 66 Ouyang, W. *et al.* Novel Foxo1-dependent transcriptional programs
827 control T(reg) cell function. *Nature* **491**, 554-559,
828 doi:10.1038/nature11581 (2012).

829 67 Ouyang, W., Beckett, O., Flavell, R. A. & Li, M. O. An essential role of the
830 Forkhead-box transcription factor Foxo1 in control of T cell homeostasis
831 and tolerance. *Immunity* **30**, 358-371, doi:10.1016/j.immuni.2009.02.003
832 (2009).

833 68 Kerdiles, Y. M. *et al.* Foxo1 links homing and survival of naive T cells by
834 regulating L-selectin, CCR7 and interleukin 7 receptor. *Nature*
835 *immunology* **10**, 176-184, doi:10.1038/ni.1689 (2009).

836 69 Riou, C. *et al.* Convergence of TCR and cytokine signaling leads to FOXO3a
837 phosphorylation and drives the survival of CD4+ central memory T cells.
838 *The Journal of experimental medicine* **204**, 79-91 (2007).

839 70 You, H. *et al.* FOXO3a-dependent regulation of Puma in response to
840 cytokine/growth factor withdrawal. *The Journal of experimental medicine*
841 **203**, 1657-1663, doi:10.1084/jem.20060353 (2006).

842 71 Brunet, A. *et al.* Akt promotes cell survival by phosphorylating and
843 inhibiting a Forkhead transcription factor. *Cell* **96**, 857-868 (1999).

844 72 Sullivan, J. A., Kim, E. H., Plisch, E. H., Peng, S. L. & Suresh, M. FOXO3
845 regulates CD8 T cell memory by T cell-intrinsic mechanisms. *PLoS*
846 *pathogens* **8**, e1002533, doi:10.1371/journal.ppat.1002533 (2012).

847 73 Sullivan, J. A., Kim, E. H., Plisch, E. H. & Suresh, M. FOXO3 regulates the
848 CD8 T cell response to a chronic viral infection. *Journal of virology* **86**,
849 9025-9034, doi:10.1128/JVI.00942-12 (2012).

850 74 van Grevenynghe, J. *et al.* Transcription factor FOXO3a controls the
851 persistence of memory CD4(+) T cells during HIV infection. *Nature*
852 *medicine* **14**, 266-274, doi:10.1038/nm1728 (2008).

853 75 Zhang, X. *et al.* FOXO1 is an essential regulator of pluripotency in human
854 embryonic stem cells. *Nat Cell Biol* **13**, 1092-1099, doi:10.1038/ncb2293
855 (2011).

856 76 Hess Michelini, R., Doedens, A. L., Goldrath, A. W. & Hedrick, S. M.
857 Differentiation of CD8 memory T cells depends on Foxo1. *The Journal of*
858 *experimental medicine* **210**, 1189-1200, doi:10.1084/jem.20130392
859 (2013).

860 77 Ferber, E. C. *et al.* FOXO3a regulates reactive oxygen metabolism by
861 inhibiting mitochondrial gene expression. *Cell Death Differ* **19**, 968-979,
862 doi:10.1038/cdd.2011.179 (2012).

863 78 Zhang, W. *et al.* FoxO1 regulates multiple metabolic pathways in the liver:
864 effects on gluconeogenic, glycolytic, and lipogenic gene expression. *The*
865 *Journal of biological chemistry* **281**, 10105-10117,
866 doi:10.1074/jbc.M600272200 (2006).

867 79 Greer, E. L. *et al.* The energy sensor AMP-activated protein kinase directly
868 regulates the mammalian FOXO3 transcription factor. *The Journal of*
869 *biological chemistry* **282**, 30107-30119, doi:10.1074/jbc.M705325200
870 (2007).

871 80 Wu, Z. *et al.* Mechanisms controlling mitochondrial biogenesis and
872 respiration through the thermogenic coactivator PGC-1. *Cell* **98**, 115-124,
873 doi:10.1016/S0092-8674(00)80611-X (1999).

874 81 Finck, B. N. & Kelly, D. P. PGC-1 coactivators: inducible regulators of
875 energy metabolism in health and disease. *The Journal of clinical*
876 *investigation* **116**, 615-622, doi:10.1172/JCI27794 (2006).

877 82 Daitoku, H., Yamagata, K., Matsuzaki, H., Hatta, M. & Fukamizu, A.
878 Regulation of PGC-1 promoter activity by protein kinase B and the
879 forkhead transcription factor FKHR. *Diabetes* **52**, 642-649 (2003).

- 880 83 Olmos, Y. *et al.* Mutual dependence of Foxo3a and PGC-1alpha in the
881 induction of oxidative stress genes. *The Journal of biological chemistry*
882 **284**, 14476-14484, doi:10.1074/jbc.M807397200 (2009).
- 883 84 Long, S. A. *et al.* Partial exhaustion of CD8 T cells and clinical response to
884 teplizumab in new-onset type 1 diabetes. *Sci Immunol* **1**,
885 doi:10.1126/sciimmunol.aai7793 (2016).
- 886 85 Freitag, J., Berod, L., Kamradt, T. & Sparwasser, T. Immunometabolism and
887 autoimmunity. *Immunol Cell Biol* **94**, 925-934, doi:10.1038/icb.2016.77
888 (2016).
- 889 86 Zhao, J. *et al.* A missense variant in NCF1 is associated with susceptibility
890 to multiple autoimmune diseases. *Nature genetics* **49**, 433-437,
891 doi:10.1038/ng.3782 (2017).
- 892 87 Gelderman, K. A., Hultqvist, M., Holmberg, J., Olofsson, P. & Holmdahl, R. T
893 cell surface redox levels determine T cell reactivity and arthritis
894 susceptibility. *Proceedings of the National Academy of Sciences of the*
895 *United States of America* **103**, 12831-12836,
896 doi:10.1073/pnas.0604571103 (2006).
- 897 88 Tkachev, V. *et al.* Programmed death-1 controls T cell survival by
898 regulating oxidative metabolism. *Journal of immunology* **194**, 5789-5800,
899 doi:10.4049/jimmunol.1402180 (2015).
- 900 89 Lee, J. C. B., D.; Roberts, R; Gearry, R.; Mansfield, J.C.; Ahmad, T.; Prescott,
901 N.; Satsangi, J.; Wilson, D.; Anderson, C.; UK IBD Genetics Consortium; Lyons,
902 P.A.; Parkes, M.; Smith, K.G.C. Evidence for distinct genetic contributions to
903 prognosis and susceptibility in Crohn's disease. *Nature (submitted)*
904 (2015).
- 905 90 Lee, J. C. *et al.* Human SNP links differential outcomes in inflammatory and
906 infectious disease to a FOXO3-regulated pathway. *Cell* **155**, 57-69,
907 doi:10.1016/j.cell.2013.08.034 (2013).
- 908