

Elsevier Editorial System(tm) for The
Veterinary Journal
Manuscript Draft

Manuscript Number: YTVJL-D-14-00884R4

Title: Lack of association between arterial oxygen tensions in horses during exploratory coeliotomy and post-operative incisional complications: A retrospective study

Article Type: Original Article

Keywords: Equine; Anaesthesia; Coeliotomy; Incisional complications; Hypoxaemia

Corresponding Author: Ms. Katherine Robson,

Corresponding Author's Institution: University of Liverpool

First Author: Katherine Robson

Order of Authors: Katherine Robson; Peter Cripps; David Bardell

Abstract: The aim of this retrospective study was to determine if there was an association between the lowest arterial blood oxygen tensions (PaO₂) measured during anaesthesia and post-operative incisional complications in horses. Clinical records of 278 horses undergoing ventral midline coeliotomy from 1 January 2010 to 31 December 2013 were examined.

The frequency of incisional complications was 32.0% (n = 89). In a multivariable model, intra-operative arterial blood oxygen tensions (PaO₂) was not significantly associated with development of an incisional complication (P = 0.351). Using hypertonic (7.2%) saline (P = 0.028, OR 3.167, 95% CI 1.132-8.861), increasing total plasma protein concentration (TP) (P = 0.002, OR 1.061 per g/L, 95% CI 1.021-1.102), an intestinal resection (P < 0.001, OR 4.056, 95% CI 2.231-9.323), increasing body mass (P = 0.004, OR 1.004 per kg, 95% CI 1.001-1.006) and the use of penicillin alone compared with penicillin and gentamicin pre-operatively (P = 0.009, OR 4.145, 95% CI 1.568-10.958) increased the risk of incisional complications. The study was unable to demonstrate a link between low intra-operative PaO₂ and increased risk of post-operative incisional complications.

1 **Lack of association between arterial oxygen tensions in horses during exploratory**
2 **coeliotomy and post-operative incisional complications: A retrospective study**

3

4

5 Katherine Robson *, Peter Cripps, David Bardell

6

7 *School of Veterinary Science, University of Liverpool, Leahurst Campus, Chester High Road,*
8 *Neston, CH64 7TE, UK*

9

10

11

12

13 * Corresponding author. Tel: +44 151 7956100.

14 *E-mail address:* knrobson@liv.ac.uk (K. Robson).

15 **Abstract**

16 The aim of this retrospective study was to determine if there was an association
17 between the lowest arterial blood oxygen tensions (PaO₂) measured during anaesthesia and
18 post-operative incisional complications in horses. Clinical records of 278 horses undergoing
19 ventral midline coeliotomy from 1 January 2010 to 31 December 2013 were examined.

20

21 The frequency of incisional complications was 32.0% ($n = 89$). In a multivariable
22 model, intra-operative arterial blood oxygen tensions (PaO₂) was not significantly associated
23 with development of an incisional complication ($P = 0.351$). Using hypertonic (7.2%) saline
24 ($P = 0.028$, OR 3.167, 95% CI 1.132-8.861), increasing total plasma protein concentration
25 (TP) ($P = 0.002$, OR 1.061 per g/L, 95% CI 1.021-1.102), an intestinal resection ($P < 0.001$,
26 OR 4.056, 95% CI 2.231-9.323), increasing body mass ($P = 0.004$, OR 1.004 per kg, 95% CI
27 1.001-1.006) and the use of penicillin alone compared with penicillin and gentamicin pre-
28 operatively ($P = 0.009$, OR 4.145, 95% CI 1.568-10.958) increased the risk of incisional
29 complications. The study was unable to demonstrate a link between low intra-operative PaO₂
30 and increased risk of post-operative incisional complications.

31

32 *Keywords:* Equine; Anaesthesia; Coeliotomy; Incisional complications; Hypoxaemia

33 **Introduction**

34 The survival of horses undergoing emergency abdominal surgery has improved in
35 recent decades (Freeman et al., 2000). Post-operative complications at the incision site, such
36 as drainage or oedema, are still common (Phillips and Walmsley, 1993; Mair and Smith,
37 2005; Freeman et al., 2012). In a previous study, 16% of horses undergoing emergency
38 abdominal surgery developed an incisional complication (Proudman et al., 2002); other
39 reported rates of incisional complication range from 7.4% to 42.2% (Freeman et al., 2000;
40 Mair and Smith, 2005; Torfs et al., 2010; Durward-Akurst et al., 2013).

41
42 Risk factors associated with an increased likelihood of incisional complications
43 include increasing heart rate on admission (French et al., 2002), increasing body mass and
44 increasing age (Wilson et al., 1995), performing an enterotomy (Honnas and Cohen, 1997),
45 use of polyglactin 910 to suture the linea alba (Honnas and Cohen, 1997), using staples rather
46 than suture material for skin closure (Torfs et al., 2010), anaesthetic duration > 110 min
47 (Smith et al., 2007) and poor anaesthetic recovery (Freeman et al., 2012). Factors thought to
48 reduce the incidence of incisional complications include using two rather than three layers of
49 sutures to close the abdomen (Colbath et al., 2014) and using a stent (Tnibar et al., 2013) or
50 an abdominal bandage to cover the incision (Smith et al., 2007).

51
52 In a study by Costa-Farre et al. (2014), horses with intra-operative arterial blood
53 oxygen tensions (PaO_2) < 80 mmHg (10.6 kPa) were significantly more likely to develop
54 post-operative incisional complications than those with $\text{PaO}_2 \geq 80$ mmHg (10.6 kPa). A direct
55 correlation has also been found between arterial and subcutaneous partial pressures of O_2 and
56 incisional infection rates in human patients undergoing colorectal surgery (Greif et al., 2000).
57 Higher rates of wound infection have been demonstrated in rabbits exposed to hypoxic

58 conditions (14% inspired oxygen) for 21 days postoperatively, compared with normoxic
59 conditions (21% inspired oxygen) (Hunt et al. 1975). We hypothesised that PaO₂ measured
60 during anaesthesia would be associated with the development of post-operative incisional
61 complications in horses.

62

63 **Materials and methods**

64 *Inclusion criteria*

65 Clinical records of horses admitted to the Philip Leverhulme Equine Hospital for
66 investigation of colic from 1 January 2010 to 31 December 2013 were examined. Cases were
67 included if they had undergone ventral midline coeliotomy for correction of gastrointestinal
68 pathology and survived for at least 7 days post surgery. Any horses that underwent repeat
69 coeliotomy, midline coeliotomy for a cause not related to the gastrointestinal tract, or where
70 anaesthetic records did not include PaO₂ values, were excluded. Anaesthetic, surgical and
71 post-operative management varied between cases and was determined by the anaesthetist and
72 surgeon involved.

73

74 Ethical approval was granted from Veterinary Research Ethics Committee University
75 of Liverpool, UK, on 15 October 2013, approval number VREC155

76

77 *Definitions*

78 An incisional complication was defined as positive bacterial culture from an incision
79 site swab and/or the presence of the following clinical signs: marked oedema at the incision
80 site, purulent discharge from the incision site and hypoechoic areas around the incision site on
81 ultrasonographic examination, with accompanying pyrexia.

82

83 *Data collected*

84 The following data were recorded from each case: age, body mass, breed, sex, heart
85 rate (HR), packed cell volume (PCV) and total plasma protein concentration (TP) at
86 admission, clinical signs of systemic inflammatory response syndrome (SIRS) at admission
87 (defined as HR > 50 beats per min (bpm) and PCV > 0.5 L/L), American Society of
88 Anesthesiologists (ASA) grade, pre-anaesthetic medication, anaesthetic induction and
89 maintenance agents and peri-operative administration of antimicrobial agents. Intra-operative
90 data included use of vasopressors and inotropes, fluid therapy, analgesic infusions,
91 dexamethasone administration, mean arterial blood pressure (MAP), highest arterial partial
92 pressure of carbon dioxide (PaCO₂), lowest pH, lowest PaO₂ and duration of anaesthesia.
93 Recovery score, anaesthetist, surgeon, whether surgery was performed out of hours, type of
94 pathology, whether an intestinal resection and/or enterotomy was performed, suture materials
95 for abdominal closure, presence of an incisional complication before discharge from the
96 hospital and how this was defined were also recorded.

97

98 *Anaesthetic and surgical technique*

99 Anaesthetic protocol varied and was determined by the anaesthetist involved. After
100 induction of anaesthesia, all horses had an orotracheal tube placed, were hoisted onto a
101 padded operating table, positioned in dorsal recumbency and the endotracheal tube connected
102 to a large animal circle breathing system. Inhalational anaesthetic agents were vaporised in
103 100% oxygen and all horses were mechanically ventilated throughout.

104

105 In all cases, the surgical site was clipped and prepared aseptically with chlorhexidine,
106 followed by surgical spirit, and an adhesive drape was placed over the incision site. The
107 pattern of abdominal closure and the suture material used to close the abdomen was

108 determined by the surgeon involved; however, in all cases, suture material rather than staples
109 were used to close the skin. All horses had an adhesive dressing and an abdominal bandage
110 for recovery.

111

112 At the end of surgery, horses were disconnected from the anaesthetic breathing
113 system, hoisted into a padded recovery box and positioned in right lateral recumbency.
114 Oxygen was provided via a demand valve until extubation and subsequently at 15 L/min via a
115 nasopharyngeal tube. This was continued until horse head movement dislodged the tubing.

116

117 Until discharge from the hospital, incision sites were examined at least once daily. In
118 horses showing clinical signs of infection, the decision to swab the incision site for bacterial
119 culture and sensitivity was determined by the attending clinician.

120

121 *Blood gas analysis*

122 During anaesthesia, arterial blood samples were taken through a cannula placed in the
123 mandibular branch of the facial artery, which also permitted invasive arterial blood pressure
124 monitoring. Samples were collected anaerobically into heparinised syringes (PICO50;
125 Radiometer) following withdrawal of approximately 1 mL of blood that was discarded, and
126 analysed immediately using a bench top blood gas analyser (ABL77; Radiometer). The timing
127 and frequency of blood gas analysis was not standardised, and the lowest recorded PaO₂ from
128 each horse was used for data analysis.

129

130 *Statistical analysis*

131 Statistical analyses were performed using Minitab 16 (Minitab) and Stata 13
132 (Statacorp). Data were tested for normality using a Ryan-Joiner test. Parametric data are

133 presented as means \pm standard deviations (SDs); non-parametric data are presented as
134 medians with interquartile ranges (IQR). Univariable analysis was performed with Pearson's
135 χ^2 analysis for categorical variables and binary logistic regression for continuous variables.
136 Variables with $P < 0.2$ were offered to a multivariable logistic regression model, using both
137 forwards and backwards stepwise entry, with a P value of 0.20 for entry and 0.21 for removal,
138 although the final model chosen by both procedures was the same. The possible effect of the
139 interaction of anaesthetic duration and PaO₂ was tested by forcing it into the final model. The
140 statistical significance of entering terms into regression models was assessed by changes in
141 the Likelihood Ratio and the Wald Statistic used to obtain the P value of coefficients in the
142 model, together with their confidence intervals (CIs). A P value < 0.05 was considered to be
143 significant.

144

145 **Results**

146 Of 278 horses that met the inclusion criteria from 1 January 2010 to 31 December
147 2013 (Fig. 1), 89 (32%) developed an incisional complication; swabs were taken from 64 of
148 these and all were positive on bacterial culture.

149

150 *Demographic data*

151 The ages of horses ranged from 2 months to 30 years, with a mean \pm SD of 12.2 ± 5.7
152 years, which was not significantly different between groups. Body mass ranged from 48 to
153 750 kg, with a median of 532 (466-600) kg. The most commonly represented breeds were
154 Thoroughbreds ($n = 35$; 12.6%) and Cobs ($n = 35$; 12.6%), followed by Welsh ponies ($n = 33$;
155 11.9%) and Warmbloods ($n = 31$; 11.2%); the remaining 144 horses consisted of multiple
156 different breeds. There were 154 (55.4%) neutered males, 112 (40.3%) entire females and 12
157 (4.3%) entire males; these proportions were not significantly different between groups. The

158 median pre-operative HR was 48 (40-60) bpm, the median PCV was 0.39 (0.34-0.43) L/L and
159 the median TP was 68 (62-72) g/L; overall 21/278 (7.6%) horses had clinical signs of SIRS
160 on presentation.

161

162 *Anaesthetic and surgical management*

163 Horses were graded ASA 4 or 5 in 89/278 (32%) cases. The most common pre-
164 anaesthetic medication was xylazine and morphine (211/278; 75.9%). In 233/278 (83.8%)
165 cases, anaesthesia was induced with ketamine and diazepam or midazolam. Maintenance of
166 anaesthesia was with isoflurane (150/278; 54.0%), sevoflurane (126/278; 45.3%) or
167 desflurane (2/278; 0.7%); none of these variables were significantly different between groups.

168

169 Peri-operative penicillin was used in 199/278 (71.6%) cases, while 44/278 (15.8%)
170 cases received penicillin and gentamicin. Most horses, (242/278; 87%) were treated with
171 vasopressors and/or positive inotropes due to hypotension; the agent most frequently
172 administered was dobutamine (210/278; 75.6%). Drugs used were not significantly different
173 between groups. All horses received intravenous fluid therapy, including Hartmann's solution
174 (165/278; 59.3%), Hartmann's solution with supplemental potassium chloride (112/278;
175 40.3%) or dextrose in saline (1/278; 0.4%); in addition, 24/278 (8.6%) received hypertonic
176 (7.2%) saline and 64/278 (23.0%) received synthetic colloids. Continuous infusion with
177 lidocaine was used in 184/278 (66.2%) cases and dexamethasone was administered to 34/278
178 (12.2%) cases; there were no significant differences between groups.

179

180 Similarly, there were no significant differences between groups in the lowest MAP,
181 the mean MAP, the lowest pH or the highest PaCO₂ measured during anaesthesia. The
182 median duration of anaesthesia was 106 (90-132) min, the minimum duration was 50 min and

183 the maximum duration was 240 min; recovery score was not significantly different between
184 groups. The majority of horses dislodged the nasal tubing supplying oxygen in the early stage
185 of recovery although exact timings and numbers of horses were not recorded.

186

187 Twelve anaesthetists and 10 surgeons were involved with the surgical procedures,
188 Pathology affected the small intestine in 168/278 (60.4%) cases and the large intestine in
189 110/278 (39.6%) cases. Enterotomy was performed in 117/278 (63.7%) cases and resection
190 was performed in 75/278 (27.0%) cases. Braided lactomer (Polysorb) was used to suture the
191 linea alba in 270/278 (97.1%) cases and polypropylene (Prolene) was used to suture the skin
192 in 263/278 (94.6%) cases.

193

194 *Blood gas analysis*

195 One to five arterial blood gas analyses were performed in each case; in 51/278
196 (18.4%) horses, only one arterial blood gas sample was taken; in all cases, the first sample
197 was taken within 40 min of anaesthetic induction. There was no significant difference in the
198 number (mean \pm SD) of blood gas analyses between cases which developed incisional
199 complications (2.57 ± 1.08) and cases which did not develop incisional complications ($2.42 \pm$
200 1.04 ; $P = 0.305$). Of the horses sampled more than once, 45/233 (19.3%) initially had a PaO_2
201 ≥ 100 mmHg (13.3 kPa), which reduced to < 100 mmHg on subsequent samples.

202

203 *Univariable and multivariable analysis*

204 Categorical and continuous variables associated with outcome at $P < 0.2$ are shown in
205 Tables 1 and 2, respectively. Table 3 lists the variables chosen in the final regression model.
206 The stepwise model did not require PaO_2 and, when forced into the final model, was not

207 significant ($P = 0.351$). The odds ratio (OR) for each unit increase in PaO₂ was 0.999 (95%
208 confidence interval, CI, 0.996-1.002).

209

210 Anaesthetic duration was > 2 h in 97/278 (34.9%) cases and was not associated with
211 an increased risk of incisional complication ($P = 0.427$). If lowest PaO₂ was divided into
212 categories of < 80 mmHg (10.6 kPa) and ≥ 80 mmHg (10.6 kPa) (Costa-Farre et al., 2014),
213 this also was not associated with outcome ($P = 0.379$). However, when the two terms were
214 combined with a new categorical variable with four levels (PaO₂ < 80 mmHg and anaesthetic
215 duration ≤ 2 h; PaO₂ < 80mmHg and anaesthetic duration > 2 h; PaO₂ ≥ 80 mmHg and
216 anaesthetic duration ≤ 2 h; PaO₂ ≥ 80 mmHg and anaesthetic duration > 2 h), this variable
217 was statistically significant ($P = 0.041$), but did not materially alter the direction, size and
218 statistical significance of the other coefficients. The adjusted coefficients for these terms
219 suggested that, regardless of the PaO₂ level, the highest risks occurred for anaesthetic duration
220 ≤ 2 h and these were significantly greater than the risks of anaesthetic duration > 2 h ($P <$
221 0.05).

222

223 **Discussion**

224 This study found 32.0% of horses developed an incisional complication following
225 ventral midline coeliotomy and gastrointestinal surgery, but we did not find a link between
226 low PaO₂ during anaesthesia and increased post-operative incisional complications. Oxygen is
227 essential for normal wound healing; it is needed for collagen production, angiogenesis,
228 fibroblast production and epithelialisation (Gottrup et al., 2004). Reactive oxygen species
229 mediate destruction of bacteria within leucocytes and this effect is correlated with increasing
230 tissue partial pressure of oxygen (Hopf and Rollins, 2007). Tissue hypoxia will reduce the
231 effectiveness of leucocytes, resulting in decreased production of interleukins 2 and 8 (Gottrup

232 et al., 2004), which may contribute to the development of wound infections. Human surgical
233 patients who were hyperoxaemic in the peri-operative period were less likely to develop
234 incision site infections (Qadan et al., 2009). However, we were unable to demonstrate the
235 same effect in horses undergoing gastrointestinal surgery.

236

237 In human studies, an increased fraction of inspired oxygen (FiO_2) is often provided
238 post-operatively. Greif et al. (2000) showed that human patients receiving FiO_2 0.8 (PaO_2 206
239 mmHg, 27.5 kPa) intra-operatively and for 6 h post-operatively had significantly fewer
240 surgical wound infections compared with patients receiving FiO_2 0.31 (PaO_2 114 mmHg, 15.2
241 kPa). In contrast, supplemental oxygen was provided post-operatively in the early recovery
242 period, initially via demand valve at the end of the endotracheal tube and after extubation via
243 nasal insufflation. The majority of horses dislodged the nasal tube supplying oxygen early on
244 during recovery. Provision of supplemental oxygen after horses leave the recovery box
245 presents significant practical problems and was not provided during this study. Therefore one
246 potential reason for the lack of association found in our study is that horses were not provided
247 with a high enough FiO_2 post-operatively to affect development of incisional complications.
248 This may be compounded by the fact that horses are frequently hypoxaemic in the immediate
249 post-operative period (Mason et al., 1987; McMurphy and Cribb, 1989), due to impaired
250 pulmonary function secondary to atelectasis (Nyman et al., 1990), coupled with reduced FiO_2 .
251 However PaO_2 was not measured in the post-operative period in any of the horses in our
252 study.

253

254 Bacterial contamination of the wound may occur in the early post-operative period
255 (Ingle-Fehr et al., 1997), indicating that post-operative arterial and tissue oxygen tensions
256 may be important in preventing incisional infections. However, these data were not available,

257 since it was standard practice to remove the arterial cannula at the end of surgery. Prospective
258 studies where arterial blood gases are analysed during recovery and in the post-operative
259 period could be performed to investigate this further.

260

261 Costa-Farre et al. (2014) studied horses undergoing general anaesthesia for
262 exploratory coeliotomy and found that those with a $\text{PaO}_2 < 80$ mmHg (10.6 kPa), combined
263 with an anaesthetic duration > 2 h, had the highest risk of surgical site infections. To compare
264 our findings with Costa-Farre et al. (2014), we performed an additional analysis, in which the
265 effect of the interaction of $\text{PaO}_2 < 80$ mmHg and anaesthetic duration > 2 h was forced into
266 our multivariable model. Our statistical model did not require the main effects of $\text{PaO}_2 < 80$
267 mmHg or anaesthetic duration > 2 h, but the interaction term was statistically significant at P
268 = 0.041. However, in contrast to the study by Costa-Farre et al. (2014), our findings suggest
269 that a longer anaesthesia time is protective against incisional infections. This is unexpected,
270 since previous veterinary studies have reported increased incisional complication rates for
271 longer durations of anaesthesia (Costa-Farre et al., 2014; Smith et al., 2007) and similar
272 findings have also been reported in the human literature (Curry et al., 2014).

273

274 The stress response to anaesthesia and the properties of anaesthetic drugs can
275 adversely affect the immune response (Anderson et al., 2014). Longer anaesthesia may be
276 associated with hypothermia, which is associated with increased incisional complication rates
277 in human beings (Kurz et al., 1996). However, an anaesthesia time > 2 h alone was not
278 significantly associated with outcome and became significant only when forced into a
279 statistical model that did not require its inclusion in combination with PaO_2 . Therefore, the
280 clinical relevance of this finding is questionable.

281

282 Tissue partial pressure of oxygen depends on PaO₂, but this relationship can be
283 affected by factors reducing perfusion, such as hypovolaemia, hypotension and peripheral
284 vasoconstriction (Chang et al., 1983). In hypovolaemia, tissue oxygen tensions will be lower
285 for a given PaO₂ compared with normovolaemia, as demonstrated experimentally in dogs
286 (Gottrup et al., 1987). Horses undergoing colic surgery are likely to be hypovolaemic;
287 therefore, PaO₂ values obtained during surgery may not reflect tissue oxygen tensions.
288 Hypovolaemia occurs in these horses due to SIRS and sequestration of extracellular fluid into
289 the intestinal lumen (Mair and Edwards, 2003). Despite aggressive fluid therapy during
290 anaesthesia, volume deficits may not be totally resolved before recovery. In our study, risk
291 factors indicating that horses were hypovolaemic were significantly associated with incisional
292 complications, including increased TP and administration of hypertonic (7.2%) saline.

293
294 Hypertonic saline was most often used immediately pre-operatively to improve
295 circulating volume in horses showing clinical signs of hypovolaemia. Higher than normal TP
296 is also thought to be suggestive of hypovolaemia (Mair and Edwards, 2003). It is possible that
297 these horses had low tissue oxygen tensions (regardless of PaO₂), which contributed to the
298 development of incisional complications; however, since tissue oxygen was not measured,
299 this cannot be proven. Other factors indicative of hypovolaemia, such as PCV, HR (Mair and
300 Edwards, 2003) and MAP were not significantly associated with incisional complications in
301 the multivariable model. Intra-operative MAP may have been affected by pre-operative fluid
302 administration, intra-operative vasopressor and inotrope use, and ventilation; therefore values
303 may not accurately indicate volume status. Although PCV and HR alone were not associated
304 with incisional complications, clinical signs of SIRS, which includes cases with HR > 50 bpm
305 and PCV > 0.5 L/L was significantly associated with incisional complications ($P < 0.001$) in

306 univariable analysis. It did not remain significant following multivariable analysis, which may
307 be due to the low numbers of animals with SIRS included ($n = 21$).

308

309 Horses that had intestinal resection were at increased risk of incisional complications.
310 Bacterial contamination of the abdominal incision site following incision into the intestinal
311 lumen may increase the risk of surgical site infection (Honnas and Cohen, 1997). Resection is
312 often required for strangulating intestinal lesions. Ischaemic insult following vascular
313 occlusion leads to compromise of the mucosal barrier and translocation of bacteria and
314 endotoxin into the bloodstream (Moore et al., 1981), which may lead to SIRS and a greater
315 degree of hypovolaemia than with lesions not requiring resection.

316

317 Increasing body mass was found to be associated with an increased risk of developing
318 incisional complications, as reported previously (Wilson et al., 1995). Using penicillin alone
319 compared to penicillin and gentamicin for peri-operative antimicrobial treatment increased the
320 frequency of incisional complications (OR 4.145, 95% CI 1.568-10.958.). The combination of
321 penicillin and gentamicin has an extended spectrum of activity compared with penicillin
322 alone, particularly against Gram negative organisms (Haggett and Wilson, 2008), which may
323 have reduced the frequency of incisional complications. Only 15.8% of horses received this
324 combination, with the majority receiving penicillin alone (71.6%), it therefore may be useful
325 to consider penicillin-gentamicin as perioperative antimicrobial treatment for horses
326 undergoing exploratory coeliotomy and gastrointestinal surgery.

327

328 The retrospective nature of this study results in a number of major limitations.
329 Varying anaesthetic protocols were employed, which may have affected the results, although
330 no significant differences were found in pre-anaesthetic medication, induction or maintenance

331 agents between groups. A wide variety of breeds were included and therefore breed could not
332 be subjected to meaningful statistical analysis, in contrast with Costa-Farre et al. (2014), who
333 studied predominantly Andalusian horses (35/84; 41.7%) and had a standardised anaesthetic
334 protocol.

335

336 The number and timing of arterial blood gas analyses in our study was not
337 standardised. The first sample was taken within 40 min of anaesthetic induction and the
338 number of samples taken ranged from one to five. When multiple analyses were performed,
339 45/233 (19.3%) horses initially had PaO₂ values > 100 mmHg (13.3 kPa), which subsequently
340 decreased to < 100 mmHg (13.3 kPa). Since 51/278 (18.4%) horses were sampled only once,
341 a proportion of these may have developed hypoxaemia that went undetected, potentially
342 affecting the results.

343

344 Some potentially important factors that could contribute to post-operative incisional
345 complications, such as body temperature, were not analysed due to the lack of data.
346 Hypothermia increases surgical site infections in human patients (Kurz et al., 1996) and body
347 temperature decreases in horses under general anaesthesia (Edner et al., 2007). Future
348 prospective studies should include measurement of body temperature.

349

350 The number of layers of sutures used to close the abdomen was not consistently
351 reported in the clinical records. Use of two layer modified subcuticular closure of the
352 abdomen significantly reduces incisional complication rates compared with three layer
353 closure (Colbath et al., 2014). However, a previous study using data from the same hospital as
354 our study found no difference in incisional complication rates between two and three layer
355 closure (Coomer et al., 2007).

356

357 The definition of incisional complications was not limited to those cases that had a
358 positive bacterial culture from an incision site swab. Previous studies have defined incisional
359 infection as incisional drainage persisting > 12 h after surgery (Durward-Akurst, 2013),
360 purulent discharge with heat, pain or swelling around the incision site (Mair and Smith, 2005)
361 and serosanguineous/purulent discharge, with or without bacterial culture (Torfs et al., 2010).
362 Whilst these may not all represent true bacterial infections, wound suppuration and oedema
363 have been shown to increase the risk of further complications, such as incisional herniation
364 and dehiscence (Gibson et al., 1989; French et al., 2002). Therefore, it was considered to
365 include horses with these case definitions even if bacterial culture was not performed.

366

367 **Conclusions**

368 This study was unable to demonstrate a link between low intra-operative PaO₂ and
369 increased risk of developing post-operative incisional complications in horses. Other factors
370 that could indicate poor tissue oxygen tension including use of hypertonic saline, increasing
371 TP and requiring an intestinal resection were associated with an increased risk. Further
372 prospective studies are required to investigate an association between tissue oxygen indices
373 during the early post-operative period and the incidence of post-operative incisional
374 complications.

375

376 **Conflict of interest statement**

377 None of the authors of this paper has a financial or personal relationship with other
378 people or organisations that could inappropriately influence or bias the content of the paper.

379

380 **References**

381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430

Anderson, S.L., Duke-Novakovski, T., Singh B., 2014. The immune response to anaesthesia: Part 2 sedatives, opioids, and injectable anaesthetic agents. *Veterinary Anaesthesia and Analgesia* 41, 553-566.

Chang, N., Goodson, W.H., Gottrup, F., Hunt, T.K., 1983. Direct measurement of wound and tissue oxygen tension in postoperative patients. *Annals of Surgery* 197, 470-478.

Colbath, A.C., Patipa, L., Berghaus, R.D., Parks, A.H., 2014. The influence of suture pattern on the incidence of incisional drainage following exploratory laparotomy. *Equine Veterinary Journal* 46, 156-160.

Coomer, R.P.C., Mair, T.S., Edwards, G.B.E., Proudman, C.J., 2007. Do subcutaneous sutures increase risk of laparotomy wound suppuration? *Equine Veterinary Journal* 39, 396-399.

Costa-Farre, C., Prades, M., Ribera, T., Valero, O., Taura, P., 2014. Does intraoperative low arterial partial pressure of oxygen increase the risk of surgical site infection following emergency exploratory laparotomy in horses? *The Veterinary Journal* 200, 175-180.

Curry, C.S., Smith, K.A., Allyn, J.W., 2014. Evaluation of anesthetic technique on surgical site infections (SSIs) at a single institution. *Journal of Clinical Anesthesia* 26, 601-605.

Durward-Akhurst, S.A., Mair, T.S., Boston, R., Dunkel, B., 2013. A comparison of two antimicrobial regimens on the prevalence of incisional infections after colic surgery. *Veterinary Record* 172, 287

Edner, A., Nyman, G., Essén-Gustavsson, B., 2007. Metabolism before, during and after anaesthesia in colic and healthy horses. *Acta Veterinaria Scandinavica* 49, 34.

Freeman, D.E., Hammock, P., Baker, G.J., Goetz, T., Foreman, J.H., Schaeffer, D.J., Richter, R.A., Inoue, O., Magid, J.H., 2000. Short- and long-term survival and prevalence of postoperative ileus after small intestinal surgery in the horse. *Equine Veterinary Journal Supplement* 32, 42-51.

Freeman, K.D., Southwood, L.L., Lane, J., Lindborg, S., Aceto, H.W., 2012. Post operative infection, pyrexia and perioperative antimicrobial drug use in surgical colic patients. *Equine Veterinary Journal* 44, 476-481

French, N.P., Smith, J., Edwards, G.B., Proudman, C.J., 2002. Equine surgical colic: Risk factors for postoperative complications. *Equine Veterinary Journal* 34, 444-449.

Gibson, K.T., Curtis, C.R., Turner, A.S., McIlwraith, C.W., Aanes, W.A., Stashak, T.S., 1989. Incisional hernias in the horse incidence and predisposing factors. *Veterinary Surgery* 18, 360-366.

Gottrup, F., Firmin, R., Rabkin, J., Halliday, B.J., Hunt, T.K., 1987. Directly measured tissue oxygen tension and arterial oxygen tension assess tissue perfusion. *Critical Care Medicine* 15, 1030-1036.

431
432 Gottrup, F., Gellet, S., Kirkegaard, L., Hansen, E.S., Johannsen, G., 2004. Continuous
433 monitoring of tissue oxygen tension during hyperoxia and hypoxia: Relation of
434 subcutaneous, transcutaneous and conjunctival oxygen tension to hemodynamic
435 variables. *Critical Care Medicine* 16, 1229-1234.
436
437 Greif, R., Akca, O., Horn, E.P., Kurz, A., Sessler, D.I., 2000. Supplemental perioperative
438 oxygen to reduce the incidence of surgical-wound infection. *New England Journal of*
439 *Medicine* 342,161-167
440
441 Haggett, E.F., Wilson, W.D., 2008. Overview of the use of antimicrobials for the treatment of
442 bacterial infections in horses. *Equine Veterinary Education* 20, 433-448.
443
444 Hopf, H.W., Rollins, M.D., 2007. Wounds: An overview of the role of oxygen. *Antioxidants*
445 *and Redox Signalling* 9, 1183-1192.
446
447 Honnas, C.M., Cohen, N.D., 1997. Risk factors for wound infections following celiotomy in
448 horses. *Journal of the American Veterinary Medical Association* 210, 78-81.
449
450 Hunt, T.K., Linsey, M., Grislis, G., Sonne, M., Jawetz, E., 1975. The effect of differing
451 ambient oxygen tensions on wound infection. *Annals of Surgery* 181, 35-39.
452
453 Ingle-Fehr, J.E., Baxter, G.M., Howard, R.D., Trotter, G.W., Stashak, T.S., 1997. Bacterial
454 culturing of ventral median celiotomies for prediction of postoperative incisional
455 complications in horses. *Veterinary Surgery* 26, 7-13.
456
457 Kurz, A., Sessler, D.I., Lenhardt, R., 1996. Perioperative normothermia to reduce the
458 incidence of surgical-wound infection and shorten hospitalization. *The New England*
459 *Journal of Medicine* 334, 1209-1215.
460
461 Mair, T.S., Edwards, G.B., 2003. Strangulating obstructions of the small intestine. *Equine*
462 *Veterinary Education* 15, 192-199.
463
464 Mair, T.S., Smith, L.J., 2005. Survival and complication rates in 300 horses undergoing
465 surgical treatment of colic. Part 2: Short-term complications. *Equine Veterinary*
466 *Journal* 37, 303-309.
467
468 Mason, D.E., Muir, W.W., Wade, A., 1987. Arterial blood gas tensions in the horse during
469 recovery from anaesthesia. *Journal of the American Veterinary Medical Association*
470 190, 989-994.
471
472 McMurphy, R.M., Cribb, P.H., 1989. Alleviation of postanesthetic hypoxemia in the horse
473 *Canadian Veterinary Journal* 30, 37-41.
474
475 Moore, J.N., White, N.A., Berg, J.N., Trim, C.E., Garner, H.E., 1981. Endotoxemia following
476 experimental intestinal strangulation obstruction in ponies. *Canadian Journal of*
477 *Comparative Medicine* 45, 330-332.
478
479 Nyman, G., Funkquist, B., Kvarn, C., Frostell, C., Tokics, L., Strandberg, A., Lundquist, H.,
480 Lundh, B., Brismar, B., Hedenstierna, G., 1990. Atelectasis causes gas exchange

481 impairment in the anaesthetised horse. *Equine Veterinary Journal* 22, 317-324.
482
483 Phillips, T.J., Walmsley, J.P., 1993. Retrospective analysis of the results of 151 exploratory
484 laparotomies in horses with gastrointestinal disease. *Equine Veterinary Journal* 25,
485 427-431.
486
487 Proudman, C.J., Smith, J.E., Edwards, G.B., French, N.P., 2002. Long-term survival of equine
488 surgical colic cases. Part 1: Patterns of mortality and morbidity. *Equine Veterinary*
489 *Journal* 34, 432-437.
490
491 Qadan, M., Akca, O., Mahid, S.S., Hornung, C.A., Polk Jr, H.C., 2009. Perioperative
492 supplemental oxygen therapy and surgical site infection, a meta-analysis of
493 randomized controlled trials. *Archives of Surgery* 144, 359-366.
494
495 Smith, L.J., Mellor, D.J., Marr, C.M., Reid, S.W.J., Mair, T.S., 2007. Incisional complications
496 following exploratory celiotomy: Does an abdominal bandage reduce the risk? *Equine*
497 *Veterinary Journal* 39, 277-283
498
499 Tnibar, A., Grubbe, Lin K., Thuroe, Nielsen K., Christophersen, M.T., Lindegaard, C.,
500 Martinussen, T., Ekstrom, C.T., 2013. Effect of a stent bandage on the likelihood of
501 incisional infection following exploratory celiotomy for colic in horses: A
502 comparative retrospective study. *Equine Veterinary Journal* 45, 564-569.
503
504 Torfs, S., Levet, T., Delesalle, C., Dewulf, J., Vlaminck, I., Pille, F., Lefere, L., Martens, A.,
505 2010. Risk factors for incisional complications after exploratory celiotomy in horses:
506 Do skin staples increase the risk? *Veterinary Surgery* 39, 616-620
507
508 Wilson, D.A., Baker, G.J., Boero, M.J., 1995. Complications of celiotomy incisions in horses.
509 *Veterinary Surgery* 24, 506-514.

510 **Table 1**

511 Categorical variables associated with incisional complications at $P < 0.2$ in 278 horses

512 undergoing general anaesthesia and coeliotomy.

513

	Incisional complication ($n = 89$)	No complication ($n = 189$)	P
Clinical signs of SIRS			
No	74 (83.2)	183 (96.8)	< 0.001
Yes	15 (16.8)	6 (3.2)	
Location of pathology			
Small	64 (71.9)	104 (55.0)	0.007
Large	25 (28.1)	85 (45.0)	
Enterotomy			
No	46 (51.7)	55 (29.1)	< 0.001
Yes	43 (48.3)	134 (70.9)	
Resection and anastomosis			
No	48 (53.9)	155 (82.0)	< 0.001
Yes	41 (46.1)	34 (18.0)	
Surgeon			
Surgeon 1	22 (24.7)	30 (15.8)	0.010
Surgeon 2	8 (9.0)	27 (14.3)	
Surgeon 3	3 (3.4)	28 (14.8)	
Surgeon 4	26 (29.2)	47 (24.9)	
Surgeon 5	2 (2.3)	16 (8.5)	
Surgeon 6	10 (11.8)	11 (5.8)	
Surgeon 7	9 (10.1)	18 (9.5)	
Other surgeons	9 (10.1)	12 (6.4)	
GGE for induction			
No	75 (84.3)	170 (90.0)	0.172
Yes	14 (15.7)	19 (10.0)	
Hypertonic (7.2%) saline			
No	74 (83.2)	180 (95.2)	0.001
Yes	15 (16.8)	9 (4.8)	
Antibiotics			
Procaine penicillin	68 (76.4)	131 (69.3)	0.192
Procaine penicillin + gentamicin	8 (9.0)	36 (19.0)	
Other	5 (5.6)	9 (4.8)	
None	8 (9.0)	13 (6.9)	

514

515 Data are expressed as number (%).

516 SIRS, systemic inflammatory response syndrome; GGE, guaiacol glycerine ether.

517 **Table 2**518 Continuous variables associated with incisional complications at $P < 0.2$ in 278 horses

519 undergoing general anaesthesia and coeliotomy.

520

	Incisional complication ($n = 89$)	No complication ($n = 189$)	P
Body mass (kg)	550 (495-614)	526 (453-581)	0.006
Initial HR (bpm)	52 (40-64)	48 (40-60)	0.196
Initial PCV (L/L)	0.39 (0.33-0.44)	0.39 (0.34-0.42)	0.131
Initial TP (g/L)	70 (64-78)	68 (62-72)	0.001
PaO ₂ lowest (mmHg)	101 (65-190)	109 (70-238)	0.040
PaO ₂ lowest (kPa)	13.3 (8.6-25.0)	14.3 (9.2-31.3)	0.040
Duration of anaesthesia (min)	115 (90-145)	105 (90-130)	0.117

521

522 Data are expressed as median (interquartile range).

523 HR, heart rate; PCV, packed cell volume; TP, total protein; PaO₂, partial pressure of arterial oxygen; bpm, beats

524 per min.

525 **Table 3**

526 Results of the final multivariable logistic regression model of risk factors associated with
 527 incisional complications in horses undergoing general anaesthesia and coeliotomy.

528

	OR	95% CI	<i>P</i>
Enterotomy	0.564	0.300-1.058	0.075
Resection	4.056	2.231-9.323	<0.001
Hypertonic (7.2%) saline	3.167	1.132-8.861	0.028
Initial TP (g/L)	1.061 ^a	1.021-1.102	0.002
Body mass (kg)	1.004 ^a	1.001-1.006	0.004
Antibiotics ^b			
Penicillin	4.145	1.568-10.958	0.009
Other	1.317	0.270-6.437	0.734
None	3.071	0.960-14.341	0.057

529

530 OR, odds ratio; 95% CI, 95% confidence intervals; TP, total protein.

531 ^a Odds ratio per unit increase in value of continuous variable.

532 ^b Referent to penicillin and gentamicin.


533 **Figure legend**

534 Fig. 1. Flow diagram of case enrolment and drop out

535 GA, general anaesthesia; GI, gastrointestinal.

536

Figure 1



Highlights

- Clinical records of horses undergoing general anaesthesia for colic were studied.
- No association was found between intra-operative PaO₂ and incisional complications in a multivariable model.
- Hypertonic saline, total protein, body mass and intestinal resections were associated with incisional complications.
- Use of penicillin was associated with more incisional complications than use of penicillin-gentamicin in combination.

*Revision Note

I have addressed the editor's comments in the main manuscript, highlighting my changes in blue. I was copying the style of the editor's comments whereby they highlighted the areas they wanted me to address in yellow. I hope this is acceptable, if not please let me know and I can revise the manuscript differently.