

1 **Changing the stability conditions in a back squat: the effect on maximum load lifted**
2 **and erector spinae muscle activity**

3

4 *Iain M. Fletcher PhD

5

6 University of Bedfordshire

7 Department of Sport and Exercise Science

8 Polhill Avenue

9 Bedford

10 MK41 9EA

11 UK

12 Tel . +44 (0)1234 793291

13 Email iain.fletcher@beds.ac.uk

14

15 Ashley Bagley BSc

16 University of Bedfordshire

17 Department of Sport and Exercise Science

18 Polhill Avenue

19 Bedford

20 MK41 9EA

21 UK

22 ashley.bagley@beds.ac.uk

23

24 *Corresponding Author

25

26 **Running head: Effects of various stability conditions in back squat**

27

28

29

30

31

32

33

34

35 **Abstract**

36 The aim of this study was to identify how changes in the stability conditions of a back
37 squat affect maximal loads lifted and erector spinae muscle activity. Fourteen male
38 participants performed a Smith Machine squat (SM), the most stable condition, a Barbell
39 back squat (BB) and Tendo-Destabilising Bar squat (TBB), the least stable condition. A
40 one repetition max (1-RM) was established in each squat condition, before
41 electromyography (EMG) activity of the erector spinae was measured at 85% of 1-RM.
42 Results indicated that the SM squat 1-RM load was significantly ($p = 0.006$) greater
43 (10.9%) than BB squat, but no greater than TBB squat. EMG results indicated
44 significantly greater ($p < 0.05$) muscle activation in the TBB condition compared to other
45 conditions. The BB squat produced significantly greater ($p = 0.036$) EMG activity
46 compared to the SM squat. A greater stability challenge applied to the torso seems to
47 increase muscle activation. The maximum loads lifted in the most stable and unstable
48 squats were similar. However, the lift with greater stability challenge required greatest
49 muscle activation. The implications of this study may be important for training
50 programmes; coaches wishing to challenge trunk stability, while their athletes lift
51 maximal loads designed to increase strength.

52

53 **Key Words:** electromyography, *torso instability, squat performance, muscle activity*

54 Word Count 200

55

56

57

58 **Introduction**

59 Resistance training has long been considered the most effective way of increasing
60 muscular strength for the human muscular skeletal system in general (Peterson, Rhea, &
61 Alvar, 2005). It has also been proposed that ‘functional training’, where natural
62 movements are executed in multiple planes, is superior to isolation of individual limb
63 movements (Norwood, Anderson, Gaetz, & Twist, 2007). This is exemplified by the
64 popularity of the back squat as a way of loading the lower extremities in
65 flexion/extension patterns, common to many sporting actions. In addition, the back squat
66 places a demand on the torso musculature and in particular the erector spinae to maintain
67 a neutral spine (Schwanbeck, Chilibeck, & Binsted, 2009). Consequently, the load
68 applied to the torso causes a stability challenge for the athlete to control.

69
70 Increasing the stability challenge for athletes has been popularised by increasing the
71 instability of specific movements in order to promote trunk muscle activation in whole
72 body actions. This has involved performing exercises on unstable support surfaces such
73 as Swiss Balls or Bosu Balls (Anderson & Behm, 2004; Drake et al., 2006; Vera-Garcia,
74 Elvira, Brown, & McGill, 2007) in the belief that this will increase neuromuscular
75 recruitment (Lehman, 2007). Although studies (Anderson, & Behm, 2005; Norwood et
76 al., 2007) have found that the use of unstable surfaces is linked to increased
77 neuromuscular activity, these findings could be misleading. Anderson and Behm (2005)
78 and Norwood et al. (2007) used the same load within their stable and unstable conditions,
79 despite the maximal loads lifted in unstable conditions being reported to be substantially
80 less than the same actions in stable conditions (McBride, Larkin, Doyne, Haines, &

81 Kirby, 2010). Indeed, the maximum force agonistic muscles can produce in unstable
82 exercises has been reported to be reduced to less than 70% of the force produced in
83 comparable stable activities (Drake, Fischer, Brown, & Callaghan, 2006; Drinkwater,
84 Pritchett, & Behm, 2007; Santana, Vera-Garcia, & McGill, 2007). Therefore, if studies
85 on unstable actions are using a higher percentage of one repetition max (1-RM), then it is
86 unsurprising that greater electromyography (EMG) activity is suggested to be linked to
87 exercises performed on unstable surfaces. Moreover, when research has compared the
88 same relative loads, an increase in neuromuscular activity in the stable rather than the
89 unstable condition has been established (Hamlyn, Behm, & Young, 2007; McBride et al.,
90 2010). The validity of exercising on unstable surfaces can also be criticised. Movement
91 in real sports situations often involves athletes having a stable surface (the ground) to
92 apply force to, while overcoming an unstable resistance such as an opponent or an
93 external load (Kohler et al., 2010). In these situations greater trunk stability is required
94 and simply using an unstable platform has limited value within ground-based sports.

95
96 A more valid resistance training modality could be the use of lifting unstable loads. This
97 type of training, where the mass shifts randomly during the lift action, has become
98 popular in many training facilities (Behm & Colado, 2012). This is exemplified by the
99 use of water-filled logs (Langford, McCurdy, Ernest, Doscher, & Walters, 2007) within
100 training regimes. These lifts are purported to simulate more functionally valid
101 conditioning regimes for athletes whose sport requires dealing with unstable loads. Past
102 work would suggest that lifting an unstable load can lead to greater EMG output in the
103 trunk musculature (Lee, & Lee, 2002; Van Dieen, Kingma, & Van der Bug, 2001) and an

104 increase in force (Van Dieen, Dekkers, Joris, Groen, Toussaint, & Meijer, 2001)
105 compared to stable loads of the same mass. However, these lifts have not been examined
106 in any depth within the scientific literature. The loads examined have been light (e.g. 10
107 kg [Van Dieen et al., 2001] or 18 kg [Lee & Lee, 2002]) which do not represent loads
108 typically used by athletes to develop strength. To the authors' knowledge, no study has
109 explored the back squat while lifting an unstable load when a suitable strength training
110 intensity (85% or higher of 1-RM) is applied.

111

112 Therefore, the aim of this study was to identify how changes in the stability conditions
113 applied to a barbell during a back squat affect maximal load lifted and erector spinae
114 muscle activity. It was hypothesised that less stable scenarios would induce greater
115 EMG activity, while decreasing the maximal load lifted.

116

117 **Methods**

118 *Participants*

119 Fourteen healthy males with a minimum free weight squat experience of one year and no
120 history of back pain (age = 21.7 ± 2.6 years, height = 1.79 ± 0.07 m and body mass =
121 83.2 ± 14.1 kg, 1-RM back squat = 123.5 ± 35.5 kg, relative strength = 1.56 ± 0.43 kg
122 lifted/kg of body mass) volunteered for this study. Participants were collegiate games
123 players who trained or played a minimum of three times per week and took part in at least
124 one squat-based training session per week. Participants were tested during their
125 transition phase before pre-season training commenced. Procedures were approved by
126 the University of Bedfordshire's Institute of Sport and Physical Activity Research

127 Committee for Ethics in accordance with the Helsinki Declaration of 1983. Before
128 written consent was obtained, participants completed a health screen and were provided
129 with written and oral information regarding the experimental protocol and the possible
130 risks of participation. Participants were required not to consume alcohol or perform any
131 physical activity in the 24 hours prior to each testing session. Participants did not
132 consume food or any caffeine products for four hours prior to testing. Participants were
133 instructed to continue their regular training throughout the experimental period; this was
134 monitored via a training diary and only maintenance dosages of exercise stimulus were
135 involved. Power calculations (G*Power3, Erdfelder, Faul, & Buchner, 1996) using
136 participant numbers (n=14) and the alpha level achieved during main effect calculations,
137 found a 1-RM statistical power of 0.944 and EMG power of 0.984.

138

139 *Experimental Design*

140 Three different squat techniques were explored in this randomized, counter-balanced
141 repeated measures study. Three different 1-RM lifts were performed, categorised by
142 level of stability. The most stable condition was the Smith Machine (Pullum, Luton, UK)
143 squat, where the bar lifted was stabilized by two parallel tracks allowing movement in
144 only the sagittal plane. The mid-stability condition was the Barbell (Pullum, Luton, UK)
145 squat, where the bar is able to freely move in all three planes. The least stable condition
146 was the Tendo-Destabilizing Bar (Tendo Sports Machine, Basingstoke, UK) squat. The
147 Tendo system uses a normal barbell, with a 30-kg exercise load hung below the bar on
148 two 3.5 kg springs. It should be noted that while the hung load in the Tendo system can
149 be changed, 30-kg is the maximum mass recommended by the manufacturer. A load of

150 30-kg was chosen as it was felt that this would have the largest impact on lift
151 performance and EMG, establishing if the system had any potential benefit to athletes.
152 The system swings in an anterior and posterior direction, while oscillating vertically
153 during the lift, creating a stability challenge to the torso musculature (see Figure 1). The
154 EMG activity of the erector spinae was recorded at 85% of the participant's 1-RM,
155 individually calculated from the separate 1-RM test for each squat conditions examined.

156

157 **Figure 1 about here**

158

159 *Procedures*

160 Participants were required to perform a high bar back squat in each of the test conditions
161 which involved positioning of the feet shoulder width apart, with the barbell across the
162 shoulders resting on the trapezius and slightly above the posterior aspect of the deltoids.
163 The squat consisted of hip and knee flexions until the top of the thighs were parallel to
164 the floor, followed by an immediate extension of the hips and knees. Participants kept
165 their backs in a neutral curve, with their heels on the floor and knees in line with their
166 toes throughout each lift (Gullett, Tillman, Gutierrez, & Chow, 2009). All squat actions
167 were videoed to ensure required technique was maintained, any deviation from this
168 resulted in that particular squat being removed from the data set. Each squat's timing
169 was standardised using a metronome (MIE Medical Research Ltd, Leeds, UK) set at 40
170 beats/min, with participants instructed to lift at the same tempo for the downward and
171 upward phase of the squat to prevent bouncing at the bottom and top of the squat action,
172 which could cause an increase in the oscillation of the Tendo device. Squat depth was

173 established during a familiarisation session, with a gravity dependent goniometer (MIE
174 Medical Research Ltd, Leeds, UK) used to indicate when the top of the thighs were
175 parallel to the floor. This position was recorded and standardised in all the following
176 experimental sessions. Appropriate squat depth was achieved when participants touched
177 their ischial tuberosities on a bar, held by two clamp stands, at the bottom of each squat.
178 Foot position was standardised as shoulder width, with the amount of hip external
179 rotation self selected. This stance position was recorded during the familiarisation
180 session and tape markers placed on the floor to keep the foot position standard. Foot
181 position was constant within the three squat derivatives examined, however it is
182 acknowledged that the Smith Machine squat torso position was more upright than the
183 other two squats lifts, due to the fixed nature of the bar within the Smith Machine.

184

185 *Familiarisation and 1 Repetition Maximum Protocol (1-RM)*

186 Independent 1-RM test sessions were conducted in a randomised counter balanced
187 fashion for each of the three back squat interventions utilised in this study. A ten-minute
188 warm-up on a cycle ergometer (Monark Ergomedic, Monark Exercise, 874E, Vansbro,
189 Sweden) performed at 100 W was completed before foot position and squat depth were
190 established. Participants then performed ten squats using only the bar at the required
191 metronome rate. When this rate could be reproduced, a 1-RM test was performed, to
192 establish the maximum load each participant could lift for each squat type. This involved
193 increasing load incrementally, until failure to perform a squat with good form to a
194 parallel position was established. Three minutes seated rest was enforced between each
195 squat attempt. The heaviest load lifted correctly was used as a measure of 1-RM

196 (Baechle, Earle, & Wathen, 2003). The three test sessions were performed seven days
197 apart, to prevent the impact of fatigue on test scores, and at the same time of day to avoid
198 any diurnal variations. A qualified strength and conditioning coach supervised this and all
199 subsequent exercise sessions. The 1-RM protocol was used for the Smith Machine squat,
200 the barbell squat and the destabilizing bar squat.

201

202 *Experimental Protocol*

203 Participants performed three repetitions of each squat type in a random order, using 85%
204 of the 1-RM achieved in each of the three squat 1-RM tests. A five minute warm-up at
205 100 W was performed on a cycle ergometer, followed by eight squats with just the bar to
206 establish the correct movement pattern and test velocity. Three repetitions were then
207 performed at 50% of 1-RM, followed by three minutes seated rest, before three
208 repetitions at 85% of 1-RM were performed. The same warm-up was performed prior to
209 each squat type. Seventy two hours rest, where no resistance exercise was performed,
210 was enforced between the three test sessions.

211

212 *Electromyographical Analysis*

213 EMG recordings were collected for each squat type and for all three repetitions
214 performed at 85% of 1-RM. Participants were fitted with 40 mm silver/silver chloride
215 electromyographic electrodes (EMG electrodes, Cardiocare Limited, Romford, UK), after
216 the skin was shaved and cleaned with an alcohol swab to minimise electrical impedance
217 (Hamlyn et al., 2007). Electrodes were attached onto the skin on the dominant and non-
218 dominant side of the erector spinae longissimus, positioned on the midpoint of the muscle

219 belly and two fingers width lateral from the lumbar vertebrae L1 (SENIAM,
220 <http://seniam.org/>), with an inter-electrode distance of 2 cm, aligned parallel to the
221 direction of the underlying fibres (Clarys & Cabri, 1993). A reference electrode was
222 positioned on the cervical vertebrae C7 (Seniam, 1997). Electrodes were attached with
223 participants lying prone, with their lumbar vertebral columns slightly flexed (SENIAM,
224 <http://seniam.org/>).

225

226 EMG activity was recorded at a sampling frequency of 2,000 Hz, with the high-pass filter
227 set at 20 Hz and the low-pass filter at 500 Hz and a mains notch filter utilised (Enoka,
228 2002). The EMG signals were recorded using a Powerlab isolated amplifier (Powerlab
229 AD Instruments 4/25T, AD Instruments, Chalgrove, UK). The data were analysed using a
230 computer program (Chart version 5.4.1, AD instruments, Chalgrove, UK). The raw EMG
231 signal was processed by full wave rectification, integrated and averaged (average
232 rectified value) for each squat repetition within each test condition.

233

234 EMG signals were normalized by measuring the average rectified value of a maximal
235 voluntary isometric contraction (MVC) performed after the main squat test (Burden &
236 Bartlett, 1999; Fletcher, 2010). MVCs were performed after the main test battery to limit
237 any possible effects linked to post-activation potentiation (PAP) which could be caused
238 by maximal contractions and could result in either increased performance, by stimulating
239 the nervous system, or decreased performance, by causing a level of fatigue (Chiu,
240 Schilling, Johnson, & Weiss, 2004). In order to prevent the squat repetitions from
241 affecting the MVC value due to fatigue (Burden, Trew, Baltzopoulos, 2003; DeLuca,

242 1997), a five-minute seated rest was enforced prior to MVC measurement. It was
243 assumed that any fatigue from the squats performed would be the same for each squat
244 condition, therefore if the MVC's were decreased by being performed post squat test this
245 would be similar for each test condition. The MVC involved participants performing an
246 isometric squat against an immovable barbell at a knee angle of 135° for three seconds,
247 (Burden et al., 2003; DeLuca, 1997), with foot position standardised to mimic that of all
248 three squat conditions. MVC's were performed three times with two-minute rest between
249 contractions (DeLuca, 1997). This measurement represented the EMG activation of
250 erector spinae at an MVC relative to the squatting action only. Greater EMG output may
251 be possible by isolating spinal extension through dynamometry, but would not represent
252 the potential activation during a squat movement. This activation data was used for
253 comparisons between the EMG values for each squat and for the dominant and non-
254 dominant erector spinae longissimus musculature. This process allowed the calculation
255 of EMG activation as a % of the participant's MVC (Burden et al., 2003). With the
256 exhibited reliability (intra-class correlation coefficient [ICC] = 0.952, 95% confidence
257 interval [CI] = 0.610-0.999), the first MVC was used to normalize EMG values. This
258 satisfactory level of MVC reliability allowed the first MVC performed to be used to
259 normalize EMG values. The dependent variables explored were the loads lifted and the
260 EMG activity during the squat actions, with the independent variables being the three
261 different squat conditions utilised.

262

263

264

265 *Statistical Procedures*

266 All data were considered to be normally distributed, as the Shapiro-Wilks test for
267 normality was found to have an alpha level of $p > 0.05$. Main effects were examined
268 with a two (dominant and non-dominant erector spinae activity) x three (squat conditions)
269 repeated measures ANOVA. A one-way repeated measures ANOVA explored
270 differences in the 1-RM loads lifted for each squat condition. Following the ANOVAs, a
271 pairwise comparison post-hoc test was performed (Bonferroni) to explore differences
272 between individual variables. Effect size was calculated using partial eta-squared (η^2).
273 Statistical analysis was performed using SPSS version 17 for Windows (SPSS Inc,
274 Chicago, IL, USA) with the alpha level set at $p \leq 0.05$. Reliability of the EMG measures
275 was assessed using ICC and 95% CI to compare repeated test measures. Reliability was
276 calculated as ICC = 0.87 (95% CI = 0.724-0.966) for the Smith Machine squat, ICC =
277 0.85 (95% CI = 0.68-0.941) for the barbell squat condition and ICC = 0.80 (95% CI =
278 0.56-0.929) within the destabilizing bar squat.

279

280 **Results**

281 **Table 1 about here**

282

283 When the mass lifted for the 1-RM attempt for each squat condition was explored (Table
284 1), a significant main effect was noted ($F = 6.952, p = 0.01, \eta^2 = 0.537$; large effect size
285 [Olejnik & Algina, 2003]). Pairwise comparisons indicated that the Smith Machine squat
286 load was significantly ($p = 0.006, 95\% \text{ CI} = 4.376 - 25.766$) higher than the barbell squat
287 (10.9%), with a marginal increase (2.5%) compared to the destabilizing bar squat, found

288 to be non-significant ($p = 1.000$). The destabilizing bar squat load was greater than the
289 barbell squat load (8.7%), but was found to be non-significant ($p = 0.100$).

290

291 When the normalized EMG values for the dominant and non-dominant erector spinae
292 were combined, a significant main effect was found when squat types were compared (F
293 $= 5.852$, $p = 0.017$, $\eta^2 = 0.517$; large effect size). Pairwise comparisons indicated that the
294 destabilizing bar squat was linked to significantly greater EMG activity compared to the
295 Smith Machine squat ($p = 0.011$, 95% CI = 7.422 – 58.654) and the barbell squat ($p =$
296 0.022 , 95% CI = 2.130 – 30.111). The barbell squat also produced a significantly greater
297 EMG value compared to the Smith Machine squat ($p = 0.036$, 95% CI = 1.020 – 32.815).

298

299 When the EMG values were explored in greater depth, to investigate differences between
300 the dominant and non-dominant sides (Table 1), a greater mean EMG value was recorded
301 for the non-dominant side in all squat conditions, with the destabilizing bar squat
302 showing the largest difference (14.5%). However, none of these patterns was found to be
303 significant ($p > 0.05$).

304

305 **Discussion and Implications**

306 The results from the present study indicate a significant increase in the load lifted in the
307 Smith Machine squat compared to the barbell squat. This pattern of response was
308 expected and is supported by Cotterman et al. (2005) who also found a significant
309 decrease in 1-RM load in barbell squats compared to Smith Machine squats. The
310 decrease in maximal load in the barbell intervention is likely to be due to the barbell

311 squat offering instability in three planes of motion, thereby forcing the lifter to produce
312 force in all three planes (Schick et al., 2010). The Smith Machine squat, although a
313 similar movement, is considered to be an easier action as the barbell is stabilised in two
314 parallel tracks. This allows greater attention to force production by prime movers, as the
315 mass being lifted is largely being stabilized by the Smith Machine itself, rather than the
316 athlete (Schick et al., 2010).

317

318 The results of this study confirm that the Smith Machine squat offers lifters the
319 opportunity to overcome heavier loads. However, there is a potential problem with
320 regard to the transfer of this force to more dynamic sporting/exercise situations. Schick et
321 al. (2010) consider the barbell squat to be a superior exercise compared to the Smith
322 Machine squat, as the muscles contract in a more natural fashion, ensuring balance in
323 three planes of motion. Barbell actions cause a higher demand on the lifter to stabilize
324 the load and control the movement while overcoming the chosen resistance (Langford et
325 al., 2007). This increases the stress on the lifter to coordinate the activity of more
326 synergist, fixator and antagonistic muscle groups (Behm, & Colado, 2012).

327 Consequently, the transfer of strength gains to more unstable conditions (e.g. sports and
328 exercise) is increased to a greater extent than through Smith Machine actions (Langford
329 et al., 2007).

330

331 If Langford et al. (2007) are correct, in terms of the transfer of a training stimulus to
332 sports actions being greater when the load lifted needs to be stabilised, then the results of
333 the destabilizing bar squat within the present study could be of particular interest to

334 coaches and athletes. The loads lifted in this condition were only marginally less than in
335 the Smith Machine condition, while there was an average increase in the load of 11.7 kg
336 compared to the barbell squat. It should be remembered that past research which has
337 induced instability at the foot/ground interface has found a significant decrease in force
338 production in the unstable condition (Behm, Anderson, & Curnew, 2002, McBride,
339 Cormie, & Dean, 2006, McBride et al., 2010). Shifting the instability focus to the
340 athlete's torso could better replicate more random sports situations, where coordination
341 and muscle synergy are vital, without involving a decrease in the load being lifted.
342 Interestingly, although the increase in destabilizing bar compared to bar bell load was not
343 statistically significant, future research may warrant a more in-depth look at maximal
344 loads lifted in these squat conditions. This is particularly pertinent as the subjects used
345 within this study were all experienced in barbell and Smith Machine squat actions, but
346 had not used the Tendo Bar system in their training. Thus, the possibility that the
347 destabilizing bar squat load could increase with greater familiarisation is plausible,
348 particularly in light of Cotterman et al. (2005) finding that lifters experienced in Smith
349 Machine and barbell squats showed no difference in performance, while inexperienced
350 lifters could lift greater loads in the more stable (Smith Machine) squat condition.

351

352 The destabilizing bar squat was classified as the least stable condition examined. The
353 subjects commented on this, in terms of the difficulty they experienced in controlling the
354 load during lifts (it was noted visually that torso perturbations were substantially greater
355 in the destabilizing bar squat than in the other test conditions). However, this stability
356 challenge did not decrease the 1-RM lifted and there seem to be two possible

357 explanations for this. Firstly, the Tendo system positioned part of the load (37 kg) below
358 the barbell, decreasing the height of the lifter and the loads combined centre of gravity
359 compared to either the Smith Machine or barbell squat conditions, (where the mass lifted
360 is held at the level of the shoulder girdle). This could increase the global stability felt at
361 the foot/ground interface, due to the fact that a lower centre of gravity is linked to greater
362 human stability (Hall, 2003). This could offset the local stability issues experienced by
363 subjects around the torso and shoulder girdle, allowing increases in the load being lifted
364 compared to the barbell squat. This was commented on by subjects who felt that their
365 balance at ground contact was unaffected by the Tendo device, while their torso stability
366 was greatly challenged.

367 A second possible theory is linked to the Tendo device being set on springs. This causes
368 a decrease in bar stability, as the load attached to the bar ‘bounces’ thus making it harder
369 to stabilize the mass lifted as the torso attempts to dampen the oscillations it is
370 experiencing. However, the system’s springs may also stretch in the descent phase of the
371 squat, storing strain energy. The strain energy is subsequently utilised in the upward
372 phase, as the springs return to their resting length, thus helping the lifter overcome a
373 greater load than would normally be lifted. At present this is only a supposition, but
374 could be worthy of further investigation as this type of device becomes a more popular
375 method of training athletes.

376

377 The EMG results from the present study indicate that as the stability of the squat exercise
378 decreases, the erector spinae longissimus muscle activity increases. The destabilizing bar
379 EMG activity was significantly higher than in the other squat conditions, with the barbell

380 EMG significantly higher than the Smith Machine activation. The Smith Machine squat
381 EMG activity was lower than in the other actions studied, due to the Smith Machine
382 providing stability in two planes, thus requiring the postural muscles to work less to
383 maintain a neutral torso. However, the differences between the barbell and destabilizing
384 bar conditions are less clear. The destabilizing bar system positions part of the load
385 below the bar, thus decreasing the centre of gravity of the load lifted and causing less
386 torque to be applied to the torso when compared to the barbell squat, where the entire
387 load is positioned on the shoulders. Theoretically, this would cause a greater need to
388 stabilise the torso in the barbell action and therefore a greater need to recruit torso
389 musculature compared to the destabilizing bar action. However, this was not found in the
390 present study, with EMG activity greatest in the destabilizing bar squat. It could be that
391 the backward and forward movement and the oscillating process produced by the
392 destabilizing bar spring system generate a greater stability challenge to the trunk
393 musculature, by making it harder to balance the load being lifted. This seems to cause
394 greater motor unit recruitment of the back muscles to keep an upright neutral posture and
395 maintain spinal stability through the lift, regardless of the load's centre of gravity and its
396 effect on torque.

397

398 Past literature has shown different patterns of response with regard to trunk muscle
399 activation compared to the present study's findings. Anderson and Behm (2005) found
400 increased erector spinae activity when a barbell squat was performed on two balance
401 discs compared to a Smith Machine or normal barbell squat. However, normal barbell
402 squat produced no greater EMG activity of the lumbo-sacral erector spinae or abdominal

403 stabilisers compared to the Smith Machine action. Schwanbeck et al. (2009) also found
404 no difference in activation of the erector spinae and rectus abdominus when Smith
405 Machine and barbell squats were compared. Interestingly, erector spinae activity was
406 higher than rectus abdominus activity in both squats, highlighting the importance of the
407 erector spinae muscle recruitment in trunk stabilisation while squatting. However, it
408 must be acknowledged that this previous research is fundamentally different to the
409 present study. Anderson and Behm (2005) lifted the same sub-maximal load for the
410 Smith Machine and barbell squat, which is a potential problem, given that the present
411 study's findings showed a significantly greater load lifted in the Smith Machine
412 condition. Schwanbeck et al. (2009) used 8-RM as their experimental load rather than an
413 exercise intensity designed to develop strength ($\geq 85\%$), as the present study utilised.
414 Therefore, the maximal loads used to develop strength have not been explored in terms of
415 trunk activation until now. The present study aimed to produce instability in a more
416 functional fashion, with a stable surface used for all squats and instability being generated
417 by the lifting modality adopted (Kohler et al., 2010). It may be argued that this may have
418 more transfer and greater benefits for athletes' training regimes than either lifting very
419 stable loads, or lifting on unstable platforms.

420

421 Interestingly, non-dominant erector spinae activity was higher in each squat condition
422 when compared to the dominant side, with the destabilizing bar squat producing larger
423 differences between the dominant/non-dominant sides. It is acknowledged that this is a
424 tentative conclusion as statistical significance was not met, (probably due to the large
425 standard deviation found in the EMG data set), but it does warrant further investigation,

426 especially considering evidence exists to show that muscle imbalances are linked to
427 decreases in performance (Young, et al., 2002) and an increased likelihood of injury
428 (Orchard, et al., 1997). In particular, it would be interesting to investigate if muscle
429 imbalances can be alleviated if a stability challenge is added to a squat exercise.

430

431 **Conclusion**

432 The aim of this study was to identify how changes in the stability conditions applied to a
433 barbell during a back squat affect maximal load lifted and erector spinae muscle activity.
434 This was achieved by examining squats with different stability challenges. Smith
435 Machine (most stable), barbell and destabilizing bar (least stable) squats 1-RM loads and
436 EMG activity while lifting 85% of 1-RM were compared. The study found a significant
437 increase in erector spinae EMG activity linked to a decrease in squat stability. When load
438 lifted was recorded, though the Smith Machine squat was significantly greater than the
439 barbell squat, no difference between the destabilizing bar and Smith Machine squat was
440 found.

441

442 **References**

- 443 Anderson, K.G., & Behm, D.G. (2004). Maintenance of EMG activity and loss of force
444 output with instability. *Journal of Strength and Conditioning Research*, 18, 637-
445 640.
- 446 Anderson, B., & Behm, D.G. (2005). Trunk muscle activity increases with unstable
447 squat movements. *Applied Physiology, Nutrition and Metabolism*, 30, 33-45.
- 448 Baechle, T.R., Earle, R.W., & Wathen, D. (2003). Resistance training. In T.R. Baechle

449 and R.W. Earle (Eds.), *Essentials of Strength Training and Conditioning*. 3rd ed.
450 (pp. 395-400). Champaign, IL: Human Kinetics.

451 Behm, D.G., Anderson, K., & Curnew, R.S. (2002). Muscle force and activation under
452 stable and unstable conditions. *Journal of Strength and Conditioning Research*,
453 *16*, 416-422.

454 Behm, D., & Colado, J.C. (2012). The effectiveness of resistance training using unstable
455 surfaces and devices for rehabilitation. *International Journal of Sports Physical
456 Therapy*, *7*, 226-241.

457 Burden, A., & Bartlett, R. (1999). Normalisation of EMG amplitude: an evaluation and
458 comparison of old and new methods. *Medical Engineering Physics*, *21*, 247-257.

459 Burdon, A.M., Trew, M., & Baltzopoulos, V. (2003). Normalisation of gait EMGs: a re-
460 examination. *Journal of Electromyography and Kinesiology*, *13*, 519-532.

461 Chiu, L.Z., Fry, A.C., Schilling, B.K., Johnson, E.J., & Weiss, L.W. (2004).
462 Neuromuscular fatigue and potentiation following two successive high intensity
463 resistance exercise sessions. *European Journal of Applied Physiology*, *92*, 385-
464 392.

465 Clarys, J.P., & Cabri, J. (1993). Electromyography and the study of sports movements: a
466 review. *Journal of Sport Science*, *11*, 379-448.

467 Cotterman, M.L., Darby, L.A., & Skelly, W.A. (2005). Comparison of muscle force
468 production using the smith machine and free weights for bench press and squat
469 exercises. *Journal of Strength and Conditioning Research*, *19*, 169-176.

470 DeLuca, C.J. (1997). The use of surface electromyography in biomechanics. *Journal of
471 Applied Biomechanics*, *13*, 135-163.

472 Drake, J.D.M., Fischer, S.L., Brown, S.H.M., & Callaghan, J.P. (2006). Do exercise balls
473 provide a training advantage for trunk extensor exercises? A biomechanical
474 evaluation. *Journal of Manipulative Physical Therapy*, 29, 354-362.

475 Drinkwater, E.J., Pritchett, E.J., & Behm, D.G. (2007). Effect of instability and resistance
476 on unintentional squat-lifting kinetics. *International Journal of Sports Physiology
477 and Performance*, 2, 400-413.

478 Enoka, R.M. (2002). *Neuromechanics of Human Movement (3rd Edition)*.
479 Champaign, Il., Human Kinetics.

480 Erdfelder, E., Faul, F., & Buchner, A. (1996). Gpower: a general power analysis
481 program. *Behavior Research Methods, Instruments and Computers*, 28, 1-11.

482 Fletcher, I.M. (2010). The effect of different dynamic stretch velocities on jump
483 performance. *European Journal of Applied Physiology*, 109, 491-498.

484 Gullett, J.C., Tillman, M.D., Gutierrez, G.M., & Chow, J.W. (2009). A biomechanical
485 comparison of back and front squats in healthy trained individuals. *Journal of
486 Strength and Conditioning Research*, 23, 284-292.

487 Hall, S.J. (2003). *Basic Biomechanics (4th Edition)*, London UK, McGraw-Hill.

488 Hamlyn, N., Behm, D.G., & Young, W.B. (2007). Trunk muscle activation during
489 dynamic weight-training exercises and isometric instability activities. *Journal of
490 Strength and Conditioning Research*, 21, 1108-1112.

491 Kohler, J.M., Flanagan, S.P., & Whiting, W.C. (2010). Muscle activation Patterns while
492 lifting stable and unstable loads on stable and unstable surfaces. *Journal of
493 Strength and Conditioning Research*, 24, 313-321.

494 Langford, G.A., McCurdy, K.W., Ernest, J.M., Doscher, M.W., & Walters, S.D. (2007).

495 Specificity of machine, barbell, and water-filled log bench press resistance
496 training on measures of strength. *Journal of Strength and Conditioning Research*,
497 21, 1061-1066.

498 Lee, Y., & Lee, T. (2002). Human muscular and postural responses in unstable load
499 lifted. *Spine*, 27, 1881-1886.

500 Lehman, G.L. (2007). An unstable support surface is not a sufficient condition for
501 increases in muscle activity during rehabilitation exercise. *Journal of Canadian*
502 *Chiropractic Association*, 51, 139-143.

503 McBride, J.M., Cormie, P., & Deane, R. (2006). Isometric squat force output and
504 muscle activity in stable and unstable conditions. *Journal of Strength and*
505 *Conditioning Research*, 20, 915-918.

506 McBride, J.M., Larkin, T.R., Doyne, A.M., Haines, T.L., & Kirby, T.J. (2010). Effect of
507 absolute and relative loading on muscle activity during stable and unstable
508 squatting. *International Journal of Physiology and Performance*, 5, 177-183.

509 Norwood, J., Anderson, G.S., Gaetz, M., & Twist, P. (2007). Electromyographic activity
510 of the trunk stabilizers during stable and unstable bench press. *Journal of Strength*
511 *and Conditioning Research*, 21, 497-502.

512 Olejnik, S., & Algina, J. (2003) Generalized eta and omega squared statistics: Measures
513 of effect size for common research designs. *Psychology Methods*, 8, 434-447.

514 Orchard, J., Marsden, J., Lord, S., & Galick, D. (1997). Pre season hamstring muscle
515 weakness associated with hamstring muscle injury in Australian footballers.
516 *American Journal of Sports Medicine*, 25, 81-85.

517 Peterson, M.D., Rhea, M.R., & Alvar, B.A. (2005). Applications of the dose-response for

518 muscular strength development; a review of meta-analytic efficacy and reliability
519 for designing training prescription. *Journal of Strength and Conditioning*
520 *Research, 19*, 950-958.

521 Santana, J.C., Vera-Garcia, F.J., & McGill, S.M. (2007). A kinetic and
522 electromyographic comparison of the standing cable press and bench press.
523 *Journal of Strength and Conditioning Research, 21*, 1271-1277.

524 Schick, E.E., Coburn, J.W., Brown, L.E., Judelson, D.A., Khamoui, A.V., Tran, T.T., &
525 Uribe, B.P. (2010). A Comparison of muscle activation between a smith machine
526 and free weight bench press. *Journal of Strength and Conditioning Research, 24*,
527 779-784.

528 Schwanbeck, S., Chilibeck, P.D., & Binsted, G. (2009). A comparison of free weight
529 squat to smith machine squat using electromyography. *Journal of Strength and*
530 *Conditioning Research, 23*, 2588-2591.

531 Van Dieen, J.H., Dekkers, M., Joris, J., Groen, V., Toussaint, H.M., & Meijer, O.G.
532 (2001). Within subject variability in low back load in a repetitively performed,
533 mildly constrained lifting task. *Spine, 26*, 1799-1804.

534 Van Dieen, J.H., Kingma, I., Van der Bug, J.C.E. (2001). Evidence for a role of
535 antagonistic cocontraction in controlling trunk stiffness during lifting. *Journal of*
536 *Biomechanics, 36*: 1829-1836.

537 Vera-Garcia, F.J., Elvira, J.L., Brown, S.H., & McGill, S.M. (2007). Effects of abdominal
538 stabilization maneuvers on the control of spine motion and stability against
539 sudden trunk perturbations. *Journal of Electromyography and Kinesiology, 17*,
540 556-567.

541 Young, W.B., James, R., & Montgomery, I. (2002). Is muscle power related to running
542 speed with changes of direction? *Journal of Sports Medicine and Physical*
543 *Fitness, 42*, 282-288.

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564 **Tables**

565 Table 1. Summary of Erector Spinae EMG Activity and Maximal Loads Lifted (n=14).

566 Squat	Dominant Side	Non Dominant Side	1-RM Load
567	(% MVC)	(% MVC)	(kg)
568 Smith Machine	91.8 ± 35.9	95.7 ± 39.1	138.5 ± 35.0
569 Barbell	107.8 ± 38.1	113.5 ± 37.1*	123.5 ± 35.5*
570 Destabilizing Bar	119.5 ± 39.5* [§]	134.1 ± 55.4* [§]	135.0 ± 38.0

571 * Significantly different from the Smith Machine squat condition ($p \leq 0.05$). [§]

572 Significantly different from the barbell squat condition ($p \leq 0.05$).

573

574

575

576

577

578

579

580

581

582

583

584


585

586

587 **Figure Captions**

588 Figure 1. Tendo-Destabilising Bar System: posterior view (A) and lateral view (B).

589



590
591

592

593

594

595