

An excess of star-forming galaxies in the fields of high-redshift QSOs

J. A. Stevens,^{1★} Matt J. Jarvis,¹ K. E. K. Coppin,² M. J. Page,³ T. R. Greve,⁴
F. J. Carrera⁵ and R. J. Ivison^{6,7}

¹Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, Herts AL10 9AB

²Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE

³Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT

⁴Max-Planck Institute für Astronomie, Königstuhl 17, Heidelberg 69117, Germany

⁵Instituto de Física de Cantabria (CSIC-UC), Avenida de los Castros, 39005 Santander, Spain

⁶Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ

⁷Scottish Universities Physics Alliance, Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ

Accepted 2010 March 8. Received 2010 March 7; in original form 2009 December 14

ABSTRACT

We present submillimetre (submm) and mid-infrared (MIR) imaging observations of five fields centred on quasi-stellar objects (QSOs) at $1.7 < z < 2.8$. All five QSOs were detected previously at submm wavelengths. At 850 (450) μm , we detect 17 (11) submillimetre galaxies (SMGs) in addition to the QSOs. The total area mapped at 850 μm is ~ 28 arcmin² down to rms noise levels of 1 – 2 mJy beam⁻¹, depending on the field. Integral number counts are computed from the 850 - μm data using the same analytical techniques adopted by ‘blank-field’ submm surveys. We find that the ‘QSO-field’ counts show a clear excess over the blank-field counts at deboosted flux densities of ~ 2 – 4 mJy; at higher flux densities, the counts are consistent with the blank-field counts. Robust MIR counterparts are identified for all four submm detected QSOs and ~ 60 per cent of the SMGs. The MIR colours of the QSOs are similar to those of the local ultraluminous infrared galaxy (ULIRG)/active galactic nuclei (AGN) Mrk 231 if placed at $1 < z < 3$ whilst most of the SMGs have colours very similar to those of the local ULIRG Arp 220 at $1 < z < 3$. MIR diagnostics therefore find no strong evidence that the SMGs host buried AGN although we cannot rule out such a possibility. Taken together our results suggest that the QSOs sit in regions of the early universe which are undergoing an enhanced level of major star formation activity, and should evolve to become similarly dense regions containing massive galaxies at the present epoch. Finally, we find evidence that the level of star formation activity in individual galaxies appears to be lower around the QSOs than it is around more powerful radio-loud AGN at higher redshifts.

Key words: galaxies: evolution – galaxies: formation – galaxies: starburst – galaxies: high redshift – submillimetre: galaxies.

1 INTRODUCTION

The popular hierarchical model of galaxy formation predicts that elliptical galaxies found today in the cores of rich clusters formed at high redshifts and at rare high-density peaks of the dark matter distribution. Within these regions, gas-rich proto-galaxies merge together rapidly, forming stars at a high rate. The same reservoir of gas used to build the stellar mass can also fuel the growth of the supermassive black holes (SMBH) found dormant in the centre of the galaxies at low redshift (e.g. Kauffmann & Haehnelt 2000; Page et al. 2001; Hopkins et al. 2008). Since massive star formation is known to be a dusty phenomenon, the optical light from many young galaxies should be highly obscured. Dust enshrouded star

formation is, however, a luminous phenomenon in the millimetre (mm) through far-infrared (FIR) wavebands where the starlight absorbed by dust grains is re-emitted. Indeed, in the last decade, deep surveys at submillimetre (submm) (e.g. Smail, Ivison & Blain 1997; Hughes et al. 1998; Eales et al. 1999; Scott et al. 2002; Coppin et al. 2006) and mm (e.g. Greve et al. 2004; Bertoldi et al. 2007) wavelengths have identified a high-redshift population of massive, dusty galaxies that are undergoing extreme starbursts (e.g. Smail et al. 2002). Likewise, hotter dust heated in the circumnuclear environment of an active galactic nucleus (AGN) will emit at mid-infrared (MIR) wavelengths. Therefore, submm and MIR observations can be combined to study activity in forming galaxies due to dust obscured starbursts and AGN.

To date, there has been no published survey at (sub)mm or FIR wavelengths of a randomly selected region of sky over an area sufficiently large to ensure that very rare structures in the early

★E-mail: j.a.stevens@herts.ac.uk

universe are contained within it. Although large surveys of this type will be available soon from instruments such as SCUBA-2 on the James Clerk Maxwell Telescope (JCMT) and SPIRE/PACS on-board the *Herschel Space Observatory*, published surveys have so far been limited to an area of $\sim 0.5 \text{ deg}^2$ (Coppin et al. 2006; Weiß et al. 2009; Austermann et al. 2010). One of the largest published submm surveys is the SCUBA Half-Degree Extragalactic Survey (SHADES; Mortier et al. 2005; Coppin et al. 2006) which, due to technical problems, mapped only 0.2 deg^2 discovering ~ 100 submillimetre galaxies (SMGs). Whilst this survey provided one of the largest and most uniformly selected samples of SMGs ever assembled, the area of sky surveyed is extremely small by most standards. For example, Λ cold dark matter simulations predict that a dark matter halo of mass greater than a few times $10^{14} M_{\odot}$ at $z = 2$ will evolve to become a very rich cluster of galaxies by $z = 0$ (e.g. Farrah et al. 2006). However, since the surface density of such massive haloes at $z \geq 2$ is only of the order of 0.01 deg^{-2} (Evrard et al. 2002), a wide-area survey would be required in order to contain one by chance.

One promising method of locating rare structures is to target the fields of high-redshift AGN. Given their huge luminosities, such objects must contain a SMBH even at very high redshifts, and therefore must represent some of the most massive objects in existence at their epochs. They should thus act as signposts to rare high-density regions of the early universe, and can be observed efficiently with existing technology. Several groups have exploited this technique to search for proto-clusters forming around high-redshift radio galaxies (e.g. Ivison et al. 2000; Kurk et al. 2000; Pentericci et al. 2000; Smail et al. 2003; Stevens et al. 2003; De Breuck et al. 2004; Greve et al. 2007; Venemans et al. 2007) and quasi-stellar objects (QSOs; Stevens et al. 2004; Priddey, Ivison & Isaak 2008). Most of these studies have concentrated on extreme objects, those with the highest luminosities and at the highest redshifts, and all have found overdensities of star-forming galaxies (typically Ly α emitters or SMGs) in the fields of the AGN. Here, we extend this work to investigate the environments of typical QSOs at submm and MIR wavelengths. The QSOs are selected over the redshift range $1.7 < z < 2.8$ where the star formation rate (SFR) density and accretion luminosity density of the universe peak. This work is motivated by the desire to study the coupled formation of stellar mass and black holes in galaxies that should evolve to become typical ellipticals in moderately rich clusters by the present day. Since the bulk of the field SMG population are also found at these redshifts (Chapman et al. 2003, 2005), we can also compare the properties of galaxies found in different environments.

This paper presents submm and MIR imaging data for five QSO fields. We extract submm catalogues and integral number counts, and compare these with results from blank-field surveys. We then

characterize the MIR properties of the QSOs and the SMGs discovered in their fields.

A Hubble constant $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and density parameters $\Omega_{\Lambda} = 0.7$ and $\Omega_m = 0.3$ are assumed throughout.

2 SAMPLE SELECTION

Our targets are drawn from the sample of 28 hard X-ray selected QSOs described by Page, Mittaz & Carrera (2001). The objects have X-ray spectra harder than the X-ray background, consistent with moderate absorption [$20.5 < \log(N_{\text{H}}/\text{cm}^{-2}) < 23$] and display broad optical emission lines. In previous works (Page et al. 2001; Stevens et al. 2005), we have observed 20 of these objects using SCUBA in photometry mode. The remaining eight objects were not targeted with SCUBA due to observing constraints, so this sample should be representative of the population. Our sample for subsequent mapping comprises those QSOs with photometry mode detections of $> 5 \text{ mJy beam}^{-1}$. This flux density threshold is based on limitations imposed by current sensitivity limits and by confusion noise, but also means that any source detected at $z > 1$ must be either an ultraluminous or hyperluminous FIR galaxy if the emission is from dust, and if the temperature of this dust is typical of such objects detected at low redshift. The observations of RX J094144.51+385434.8 were published by Stevens et al. (2004) but are included here for completeness. The five QSOs are listed in Table 1 along with some basic parameters. They are selected to (i) span the redshift range $1.7 < z < 2.8$ and (ii) have 0.5–2 keV X-ray luminosities within ~ 0.7 dex of L_{X}^* , the break in the X-ray luminosity function.

3 OBSERVATIONS AND DATA REDUCTION

3.1 SCUBA observations

SCUBA jiggle-map observations (Holland et al. 1999) at 450 and 850 μm were made at the JCMT between 2003 July and 2004 December. These observations are summarized in Table 2. We used a 64-point jiggle pattern to fully sample the image plane at both wavelengths but employed a number of different chop/nod strategies. For RX J163308, RX J094144 and RX J121803, we chopped and nodded east–west with throws of 30, 30 and 45 arcsec, respectively. For these objects the off-positions fall at a fixed position on the sky; a real source thus has negative images of itself to the east and west, separated by twice the chop throw but with half the amplitude. For RX J005734 and RX J124913, we chopped and nodded in azimuth by 30 arcsec, so the off-positions rotate with the field throughout the observation.

Each mapping observation, taking typically one hour to complete, was bracketed by pointing observations on bright point-like

Table 1. The sample of QSOs mapped with SCUBA.

Name	RA (J2000.0)	Dec. (J2000.0)	z	$\log(L_{\text{X}})$ [$\log(\text{erg s}^{-1})$]	$\log(L_{\text{FIR}})$ [$\log(L_{\odot})$]
RX J005734.78–272827.4	00 ^h 57 ^m 34 ^s .94	–27°28′28″.0	2.19	44.85 ^{+0.14} _{–0.14}	13.32 ^{+0.08} _{–0.06}
RX J094144.51+385434.8	09 ^h 41 ^m 44 ^s .61	+38°54′39″.1	1.82	44.83 ^{+0.12} _{–0.14}	13.41 ^{+0.05} _{–0.07}
RX J121803.82+470854.6	12 ^h 18 ^m 04 ^s .54	+47°08′51″.0	1.74	44.69 ^{+0.17} _{–0.15}	13.10 ^{+0.08} _{–0.10}
RX J124913.86–055906.2	12 ^h 49 ^m 13 ^s .85	–05°59′19″.4	2.21	45.10 ^{+0.10} _{–0.13}	13.11 ^{+0.09} _{–0.11}
RX J163308.57+570258.7	16 ^h 33 ^m 08 ^s .59	+57°02′54″.8	2.80	45.24 ^{+0.11} _{–0.10}	13.00 ^{+0.08} _{–0.11}

Note. Coordinates are those of the optical QSO. Absorption corrected X-ray luminosities are given for the 0.5–2 keV band whilst FIR luminosities are calculated from submm photometric observations at 850 μm (see Stevens et al. 2005 for more details on how these luminosities were calculated).

Table 2. Observing log of the SCUBA mapping observations.

Name	Date (yy/mm/dd)	N_{map}	τ_{wvm}
RX J005734.78–272827.4	2003/10/08	20	0.07–0.08
	2003/12/09	40	0.06–0.08
	2003/12/12	20	0.03–0.04
	2003/12/15	10	0.04–0.05
	2003/12/26	20	0.04–0.05
	2004/01/08	20	0.04–0.05
	2004/01/09	10	0.05–0.06
	2004/08/26	20	0.06–0.08
	2004/08/29	20	0.07–0.08
	2004/12/30	20	0.04–0.05
RX J121803.82+470854.6	2004/12/18	40	0.07–0.08
	2004/12/29	53	0.05–0.06
	2004/12/30	100	0.03–0.08
RX J124913.86–055906.2	2003/12/24	20	0.04–0.05
	2003/12/26	40	0.04–0.05
	2004/01/07	60	0.04–0.06
	2004/01/08	60	0.04–0.05
	2004/01/09	60	0.05–0.06
RX J163308.57+570258.7	2004/01/10	40	0.05–0.06
	2003/07/08	80	0.06–0.08
	2003/07/09	80	0.08–0.09
	2003/07/11	80	0.07–0.08
	2003/07/12	60	0.05–0.07
	2004/01/09	20	0.04–0.05
	2004/01/10	60	0.05–0.06
	2004/01/11	20	0.05–0.06

Note. For each field, we give the date of observation, the number of 64-pt jiggle maps where 10 maps is equivalent to ~ 20 min of integration, and the opacity range (τ_{wvm}) as given at 225 GHz from measurements with the JCMT water vapour radiometer. Observations of RX J094144.51+385434.8 were published by Stevens et al. (2004).

objects. Pointing offsets were in most cases < 2 arcsec and in all cases < 3 arcsec. Focus was checked throughout the night but particularly around sunset and sunrise when rapid changes are known to occur.

Atmospheric transmission was determined with skydips and checked against the JCMT water vapour radiometer. Maps of Uranus, Mars and/or CRL618 were used for flux density calibration while maps of bright blazars were used to measure the beam profile. At 850 μm , the calibration was found to be stable enough to apply a single calibration to all observations; our combined flux calibration factor (FCF) was $221 \pm 15 \text{ Jy/V beam}^{-1}$ giving a calibration uncertainty of about 7 per cent. At 450 μm , the FCF varies with the atmospheric conditions because the filter profile contains a relatively strong water line, but also depends quite strongly on the thermal state of the antenna (see Jenness et al. 2002). At 450 μm , we thus applied our best estimate of the FCF to each observation in turn; FCFs varied from 220 to 490 $\text{Jy V}^{-1} \text{ beam}^{-1}$. The calibration uncertainty at 450 μm is ~ 20 per cent.

Data were reduced with standard STARLINK SURF routines up to the rebinning stage (extinction correction, flat-fielding and de-spiking). For each mapping observation the individual bolometer timestreams were inspected and noisy integrations/noisy bolometers were blanked. We then used adapted versions of the SURF tasks SETBOLWT and REBIN to make bolometer noise weighted mean signal and noise maps on a 2 arcsec pixel scale (see Stevens et al. 2004 for more details). These maps were then convolved with a Gaussian of full width half-maximum (FWHM) 4 pixels (850 μm) or 3 pixels (450 μm) giving final resolutions of 14.2 (850 μm) and 8.6 arcsec

(450 μm). The convolved maps with signal-to-noise ratio (S/N) contours are shown in Fig. 1.

3.2 Spitzer observations

We observed each field with the InfraRed Array Camera (IRAC) (at 4.5 and 8.0 μm) and the Multiband Imaging Photometer for SIRTf (MIPS) (at 24 μm) cameras on-board the *Spitzer Space Telescope*. IRAC observations were made between 2005 March and July while the MIPS observations were made between 2005 March and December. For the IRAC observations, we used 5-pt Gaussian dithers with frame times of 100 or 200s depending on the predicted brightness of the target. Frame times for the MIPS observations were between 3 and 10 s. These observations are summarized in Table 3.

The IRAC frames were processed by the Spitzer Science Center using the standard pipeline version S14.0.0. The images were then cropped to only include the region of constant exposure time; this allows catalogues to be constructed from regions with similar noise properties. Source extraction was performed with SEXTRACTOR (Bertin & Arnouts 1996) with a fixed aperture of 3.5 arcsec and a local background fit. Aperture corrections of 1.6 and 2.13 were applied to the 4.5- and 8.0- μm flux densities, respectively, in accord with the values determined from the Spitzer First Look Survey (Lacy et al. 2005).

For MIPS observations, the 24 μm Basic Calibrated Data (BCD) images were fitted with a plane after first masking out the bright objects. This plane was then subtracted from each BCD frame resulting in a flat background image. These images were then combined with the IRAF task IMCOMBINE using the astrometric solution from each individual BCD image. This procedure resulted in a much improved 24- μm image, free of the jail-bar pattern common to the Post Basic Calibrated Data (PBCD) images supplied from the Spitzer pipeline. Again, only the central $3 \times 3 \text{ arcmin}^2$ region of the images was used to detect sources. We used SEXTRACTOR to find sources and the PHOT script in IDL to measure their flux density within a 13 arcsec radius aperture. All flux densities were then subjected to an aperture correction of 1.167 (see Seymour et al. 2007).

4 THE SUBMILLIMETRE CATALOGUES

Our philosophy is to match our analysis as closely as possible to that performed by the SHADES consortium (Coppin et al. 2006), thus allowing a direct comparison of the number counts determined for blank fields and those in the highly biased regions of the Universe surrounding our QSOs. Most of the analysis in this section pertains to the 850- μm data. For the 450- μm data, we extract a catalogue at a fixed S/N threshold but do not compute number counts.

4.1 Source extraction

We adopt the source extraction algorithm described by Scott et al. (2002) which performs a maximum-likelihood fit to the flux densities of all potentially significant peaks identified on the convolved maps. For each source, we took the convolved S/N map and searched for all peaks with $S/N > 1.5$. A peak is found if a 10×10 arcsec grid centred on any pixel fails to find another pixel in the grid with higher S/N. In practice, this means that real sources separated by less than a beamwidth can be missed by the extraction algorithm. This is the case for RX J094144 where we fail to recover two sources which are clearly real because we see them in the higher resolution 450 μm data but are blended together at 850 μm . We then took the unconvolved signal map and convolved it with appropriately scaled beam maps centred at these locations. The amplitudes




Figure 1. SCUBA maps of (a) RX J005734.87–272827.4, (b) RX J094144.51+385434.8, (c) RX J121803.82+470854.6, (d) RX J124913.86–055906.2 and (e) RX J163308.57+570256.7 at 850 (left) and 450 μm (right). The grey-scales show flux density while the contours, which show S/N, are calculated using the noise maps (see text). Contour levels are 2, 3, 4, 5 and 6σ except for RX J124913.86–055906.2 where they are 2, 3, 4, 5, 6, 8, 10 and 12σ . These contours refer to the significance levels on the pre-extracted maps as shown. Note that real sources will have higher significance when extracted with the beam profile because emission in the off-positions is also factored into the total S/N.

of these beam maps were then adjusted until the overall χ^2 of the fit was minimized. For two sources with azimuth chop/nod schemes, the beam was reconstructed by going back to the observing logs and calculating the position angle of the chop throw on the sky at

the mid-point of each observation. We then added these together, weighted by the integration time for each observation.

This method is suitable for extracting sources in fields where confusion is likely to be a problem; it also has the advantage of




Figure 1 – continued

Table 3. Observing log of the *Spitzer* observations.

Name	IRAC 4.5/8.0 μm		MIPS 24 μm	
	Date (yy/mm/dd)	T_{int} (s)	Date (yy/mm/dd)	T_{int} (s)
RX J005734.78–272827.4	2005/07/17	2000	2005/12/02	432
RX J094144.51+385434.8	2005/05/06	3000	2005/04/12	336
RX J121803.82+470854.6	2005/05/06	2000	2005/05/19	210
RX J124913.86–055906.2	2005/07/15	5000	2005/06/27	636
RX J163308.57+570258.7	2005/03/26	1000	2005/03/06	970

Note. For each field, we give the date of observation and the on-source integration time (T_{int}) for each instrument.

using the off-positions to distinguish between real sources and noise spikes, and increases the S/N by considering the flux density in the off-positions. The method was used for all four independent SHADES reductions although note that ‘Reduction A’ of Coppin et al. (2006) searched for peaks on the convolved signal maps rather than the convolved S/N maps. We tried both methods but found that peaks towards the map edges were better recovered from the S/N maps.

Uncertainties in submm positions measured on SCUBA maps have been discussed extensively by Ivison et al. (2007) who show that the 1σ error circle is given by $0.9\theta_A(S/N^2 - 3)^{-1/2}$ where θ_A is the FWHM of the beam and S/N is the raw map S/N. A typical

source with $S/N \sim 3-4$ at $850 \mu\text{m}$ would thus have a 1σ positional accuracy of 3–5 arcsec.

4.2 Flux density deboosting

A well-known phenomenon common to all deep submm surveys is that of flux density boosting. Real sources falling in regions of positive noise will have their intrinsic flux densities boosted, raising the likelihood of inclusion in the catalogue. This effect becomes increasingly significant as the intrinsic flux density of the source approaches the limit of the survey. If not corrected for, flux density

Table 4. The 850- μm catalogue.

Field	ID#	RA (J2000.0)	Dec. (J2000.0)	S_{850} (mJy beam $^{-1}$)	S_{850} (deb.) (mJy beam $^{-1}$)	Map S/N	Note
RX J005734.78–272827.4	1	00 ^h 57 ^m 35 ^s .21	–27°28′27″.9	8.2 ± 1.2	7.8 ^{+1.0} _{–0.9}	7.9	QSO
	2	00 ^h 57 ^m 37 ^s .79	–27°28′57″.9	5.9 ± 1.6	4.3 ^{+1.7} _{–2.1}	3.8	
RX J094144.51+385434.8	1	09 ^h 41 ^m 44 ^s .84	+38°54′36″.9	12.7 ± 1.4	12.4 ^{+1.1} _{–1.0}	11.4	QSO
	2	09 ^h 41 ^m 46 ^s .53	+38°53′54″.7	7.4 ± 1.9	5.7 ^{+2.0} _{–2.1}	4.1	
	3	09 ^h 41 ^m 42 ^s .80	+38°55′34″.8	5.5 ± 1.7	3.1 ^{+2.1} _{–2.2}	3.3	
RX J121803.82+470854.6	1	12 ^h 18 ^m 05 ^s .51	+47°08′55″.1	4.3 ± 1.3	3.1 ^{+1.3} _{–1.6}	3.5	
	2	12 ^h 18 ^m 07 ^s .66	+47°08′45″.1	4.5 ± 1.4	2.8 ^{+1.6} _{–1.7}	3.3	
	3	12 ^h 18 ^m 05 ^s .53	+47°08′36″.9	3.6 ± 1.2	2.1 ^{+1.3} _{–1.3}	3.1	
	4	12 ^h 18 ^m 02 ^s .76	+47°08′24″.9	3.5 ± 1.2	2.1 ^{+1.3} _{–1.2}	3.1	
	5	12 ^h 18 ^m 04 ^s .74	+47°08′59″.0	3.6 ± 1.2	1.9 ^{+1.5} _{–1.2}	3.0	
RX J124913.86–055906.2	1	12 ^h 49 ^m 13 ^s .99	–05°59′19″.3	11.2 ± 1.1	11.0 ^{+0.7} _{–0.7}	13.9	QSO
	2	12 ^h 49 ^m 15 ^s .45	–05°58′37″.3	4.6 ± 1.1	3.8 ^{+1.1} _{–1.1}	4.3	
	3	12 ^h 49 ^m 17 ^s .06	–05°59′33″.3	3.1 ± 0.9	2.3 ^{+0.9} _{–1.0}	3.5	
	4	12 ^h 49 ^m 12 ^s .37	–05°58′33″.5	3.2 ± 1.0	2.3 ^{+1.0} _{–1.0}	3.4	
	5	12 ^h 49 ^m 11 ^s .56	–05°59′59″.3	3.2 ± 1.0	2.1 ^{+1.1} _{–1.1}	3.2	
RX J163308.57+570258.7	1	16 ^h 32 ^m 59 ^s .99	+57°03′18″.7	10.7 ± 1.4	10.4 ^{+1.0} _{–1.1}	9.5	
	2	16 ^h 33 ^m 08 ^s .82	+57°02′58″.7	7.1 ± 0.9	6.9 ^{+0.7} _{–0.7}	9.3	QSO
	3	16 ^h 33 ^m 12 ^s .74	+57°03′26″.7	4.5 ± 1.0	3.8 ^{+1.0} _{–1.0}	4.5	
	4	16 ^h 33 ^m 10 ^s .78	+57°02′10″.7	4.0 ± 1.1	3.0 ^{+1.2} _{–1.3}	3.7	
	5	16 ^h 33 ^m 03 ^s .18	+57°02′58″.7	4.0 ± 1.2	2.8 ^{+1.3} _{–1.4}	3.5	
	6	16 ^h 33 ^m 12 ^s .61	+57°02′32″.0	3.4 ± 1.1	2.3 ^{+1.1} _{–1.1}	3.4	

Note. Coordinates give the extracted position of the source. Also listed are extracted flux density and S/N together with deboosted flux density (see text for details). Quoted uncertainties on the raw flux densities include contributions from the calibration uncertainty and the extracted S/N values while those on the deboosted flux densities are purely statistical.

boosting will therefore populate the survey catalogue with an excess of low flux-density sources.

We use the Bayesian flux density deboosting recipe of Coppin et al. (2005, 2006) which calculates a posterior flux density probability distribution for each source assuming a prior distribution of real flux densities constructed from existing data. The prior is constructed using artificial skies which are sampled with the same observing technique used for the real data. Since we used two different chop/nod schemes, we employ two priors: a two position chop scheme for the three fields observed with east–west chop throws and the SHADES observing pattern for the two fields that were chopped in azimuth. SHADES employed a six position chop/nod scheme, chopping in both right ascension and declination with chop throws of 30, 44 and 68 arcsec. We note that while this method is not a true match to an azimuth chop scheme, where the chop position angle rotates on the sky throughout the observation, it is nevertheless a better approximation than a two chop scheme. In any case, running the deboosting algorithm with either simulated observing technique resulted in the same catalogue give or take 1 or 2 sources close to the selection threshold. Deboosted flux densities are listed in Table 4.

4.3 Catalogue membership

As a first cut, we took all sources found by the extraction algorithm with a S/N > 2.0. This catalogue was then trimmed to include only those sources which have less than 5 per cent of their posterior probability distribution below 0 mJy. This cut is exactly that used by Coppin et al. (2006) for the SHADES catalogue. In Fig. 2, we plot extracted 850- μm S/N against extracted 850- μm




Figure 2. Extracted 850- μm flux density against extracted 850- μm S/N (uncorrected for flux density boosting). All sources extracted with S/N > 2 are plotted; those that fall above the solid line have >95 per cent posterior probability that their flux density is >0 mJy. These sources make it into our catalogue. Dotted lines show noise levels of 1 and 2 mJy beam $^{-1}$.

flux density. Those sources falling above the solid line, and that remain there after deboosting, meet the above criterion and are included in the final catalogue (Table 4). In the five fields, we find 21 sources at 850 μm including four of the five QSOs. At 450 μm ,

Table 5. The 450- μm catalogue.

Field	ID#	RA (J2000.0)	Dec. (J2000.0)	S_{450} (mJy beam $^{-1}$)	Map S/N	Note
RX J005734.78–272827.4	1	00 ^h 57 ^m 35 ^s .08	–27°28′34″.1	32.4 ± 10.2	4.1	QSO
	2	00 ^h 57 ^m 33 ^s .13	–27°28′40″.1	25.9 ± 9.6	3.2	
RX J094144.51+385434.8	1	09 ^h 41 ^m 45 ^s .06	+38°54′36″.7	44.7 ± 10.2	9.1	QSO
	2	09 ^h 41 ^m 46 ^s .73	+38°53′56″.7	41.2 ± 10.0	7.3	850#2
	3	09 ^h 41 ^m 46 ^s .73	+38°54′14″.9	33.5 ± 8.8	5.9	
	4	09 ^h 41 ^m 44 ^s .16	+38°54′22″.7	28.8 ± 7.8	5.5	
RX J121803.82+470854.6	1	12 ^h 18 ^m 05 ^s .51	+47°08′47″.1	35.1 ± 8.9	6.4	850#1
	2	12 ^h 18 ^m 00 ^s .80	+47°08′34″.9	31.9 ± 10.7	3.7	
	3	12 ^h 18 ^m 05 ^s .11	+47°08′35″.1	30.4 ± 11.0	3.3	850#3
	4	12 ^h 18 ^m 10 ^s .02	+47°08′41″.1	24.3 ± 8.8	3.3	
	5	12 ^h 18 ^m 07 ^s .08	+47°08′52″.9	21.9 ± 8.3	3.1	850#2
RX J124913.86–055906.2	1	12 ^h 49 ^m 13 ^s .97	–05°59′21″.3	34.9 ± 8.1	8.7	QSO
	2	12 ^h 49 ^m 11 ^s .03	–05°58′37″.5	21.6 ± 7.7	3.4	
	3	12 ^h 49 ^m 11 ^s .03	–05°59′57″.5	33.7 ± 12.5	3.2	850#5

Note. Coordinates give the extracted position of the source. Also listed are extracted flux density and S/N. Quoted uncertainties on the raw flux densities include contributions from the calibration uncertainty and the extracted S/N values. The last column gives the likely ID if the 450- μm position falls within 10 arcsec of either the QSO or one of the objects extracted at 850 μm .

we simply list those sources extracted with $S/N > 3$ (Table 5) and do not attempt to deboost their flux densities. We assume that an 850- μm source is also detected at 450 μm if their extracted positions are within 10 arcsec. Using this criterion, we recover eight of the sources detected at 850 μm including three of the QSOs. A further six sources detected at 450 μm are not detected at 850 μm . Two of these objects lie in the RX J094144 field and were missed at 850 μm due to blending as discussed in Section 4.1. We note that the QSO RX J121803.82+470854.6 is not formally detected at either 850 or 450 μm even though it was detected by Page et al. (2001) using SCUBA in photometry mode. This QSO is discussed further in Appendix A.

4.4 Catalogue completeness

Extensive simulations were performed to determine the completeness as a function of flux density. For each field, we added false sources to the signal maps by scaling the normalized beam map to have an amplitude of 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16 and 18 mJy. We then attempted to recover each source with the extraction algorithm and repeated this procedure 1000 times per field. We consider a source detected if its recovered flux density is within a factor of 2 of its input flux density, if its recovered position is within 7 arcsec of its input position and if it makes it into our catalogue as described in the previous section. These criteria are those adopted by the SHADES survey. We plot the fractional completeness for each field as a function of input flux density in Fig. 3. The fits to the data have the form $\text{Completeness} = (S^a)/(b + cS^a)$ where S is the 850- μm flux density and a, b, c are free parameters. It is apparent that the completeness is a function of both the average noise level of the maps and the structure that they contain. We note that the fractional completeness converges to ~ 0.85 – 0.9 rather than 1 at high-flux densities. This is due to the failure of the extraction algorithm to recover input sources which fall very close to the field edges. The effect will be worse towards the top and bottom centres of the fields where both off-positions fall outside the area used for extraction. We confirmed that this is the case by performing an 18 mJy simulation over the centre region of one of the maps. This simulation returned a fractional completeness of 1.0. However, since

real sources may also fall on the map edges in exactly the same way that randomly generated false sources do, we do not correct for this edge effect. One other feature of these simulations worthy of remark is that the shape of the completeness curve appears to be sensitive to the adopted chopping scheme. Those fields observed with an azimuthal chop/nod scheme have relatively high completeness at low flux densities, presumably due to an absence of the deep negative off-positions that are produced by chopping at a fixed position on the sky (most fields have a bright QSO at their centre), but they are relatively less complete at high flux densities where the absence of these off-positions becomes more of a hindrance.

5 INTEGRAL NUMBER COUNTS

Number counts at 850 μm were calculated from the deboosted flux densities after first making a correction for the completeness of the catalogue for each field using the curves shown in Fig. 3. The total area of the survey is 28.4 arcmin 2 . We follow Coppin et al. (2006) so that our counts are calculated in the same manner as the SHADES counts. The method uses the deboosted flux density probability distributions as input for a modified bootstrapping simulation which estimates the number counts and their errors. This method allows the flux density of an individual source to be shared appropriately between adjacent bins. We quote counts in bins of width 2 mJy (QSOs included; Table 6) and 1 mJy (QSOs removed; Table 7). These counts represent the mean value in each bin with the quoted error being the frequentist 68 per cent confidence interval in each flux bin based on bootstrapped Monte Carlo simulations, including Poisson counting errors. While including the QSOs gives a biased representation of the overdensity in these fields, this method gives the total number of submm detected galaxies. This approach is also analogous to that adopted by previous workers on QSO overdensities (e.g. Priddey et al. 2008), although we stress that the counts are only appropriate for X-ray absorbed QSOs that are pre-selected to be submm luminous.

Quoted number counts exclude the contribution from sources 3, 4 and 5 in the RX J121803 field. All three have deboosted flux densities close to 2 mJy where the completeness estimate from our simulations is very low (< 1 per cent) and therefore subject to a large




Figure 3. Fractional completeness as a function of 850- μ m flux density for each of the five fields. The curves are fits to the data (see text for details). The left-hand panel shows the completeness for RX J163308 (squares), RX J121803 (diamonds) and RX J094144 (triangles) while the right-hand panel shows the completeness for RX J124913 (asterisks) and RX J005734 (diamonds).

Table 6. Combined integral number counts for the five QSO fields including the QSOs.

Flux density (mJy)	850- μ m integral counts, $N(> S)$ (deg $^{-2}$)	
	This work	SHADES
2.0	11 072 $^{+6019}_{-2791}$	2506 $^{+407}_{-407}$
4.0	2074 $^{+1060}_{-470}$	844 $^{+117}_{-117}$
6.0	953 $^{+507}_{-305}$	362 $^{+61}_{-59}$
8.0	552 $^{+278}_{-258}$	150 $^{+35}_{-35}$
10.0	375 $^{+190}_{-237}$	68 $^{+21}_{-21}$
12.0	120 $^{+85}_{-105}$	33 $^{+16}_{-15}$

Note. Bins are quoted at the lower flux density bound. The bins are of width 2 mJy. SHADES counts are taken from Coppin et al. (2006).

Table 7. Combined integral number counts for the five QSO fields excluding the QSOs.

Flux density (mJy)	850- μ m integral counts, $N(> S)$ (deg $^{-2}$)
	This work
2.0	10 420 $^{+3676}_{-3584}$
3.0	3574 $^{+1129}_{-1211}$
4.0	1423 $^{+524}_{-540}$
5.0	636 $^{+308}_{-300}$
6.0	343 $^{+205}_{-190}$
7.0	227 $^{+156}_{-167}$
8.0	175 $^{+146}_{-151}$
9.0	139 $^{+130}_{-131}$

Note. Bins are quoted at the lower flux density bound. The bins are of width 1 mJy.

uncertainty. The implication is that number counts in the lowest flux density bins will be underestimated. If we assume that the true completeness of this field at 2 mJy is 5 per cent (a reasonable value from inspection of Fig. 3) and that the three sources contribute




Figure 4. Integral number counts plotted in 1 mJy bins. The QSOs have been removed from the counts. The dotted line is a best-fitting Schechter function to the SHADES counts as described in Coppin et al. (2006) while the dashed line is the result of multiplying this fit by 4 and is shown for reference.

entirely to the lowest flux density bin, then the number counts in this bin would increase by 70 per cent. However, no correction is applied to the quoted counts.

The counts (QSOs removed) are plotted in Fig. 4 which also shows number counts determined for the SHADES survey (Coppin et al. 2006). We note that counts determined at 870 μ m from a slightly larger survey conducted with the Large APEX Bolometer Camera (LABOCA) are consistent with those measured for SHADES at flux densities below 3 mJy and are about a factor of 2 below the SHADES counts at higher flux densities (Weiß et al. 2009). The dashed line is not a fit to the data but is simply the best-fitting Schechter function to the SHADES counts multiplied by 4. From this plot, we can see that the observed overdensity of sources around the QSOs arises almost

entirely from galaxies with deboosted flux densities in the range of $\sim 2\text{--}4$ mJy. At higher flux densities, the counts in the QSO fields are matched very well to those determined from SHADES data for the field.

6 MID-INFRARED COUNTERPARTS

Likely MIR counterparts to the submm galaxies are given in Table 8. For each field, we searched within fixed radii of the extracted submm

Table 8. Candidate *Spitzer* IRAC and MIPS counterparts to the submm galaxies.

Field	ID	λ (μm)	Cat#	S_λ (μJy)	RA (J2000.0)	Dec. (J2000.0)	$\Delta\theta$ (arcsec)	P	Note
RX J005734.78–272827.4	850#1	4.5	272	476 ± 15	00 ^h 57 ^m 34 ^s .93	–27°28′27″.6	3.69	0.005	QSO
	850#1	8.0	203	1534 ± 31	00 ^h 57 ^m 34 ^s .94	–27°28′27″.6	3.67	0.003	QSO
	850#1	24.0	10	4107 ± 111	00 ^h 57 ^m 34 ^s .90	–27°28′27″.3	4.12	0.002	QSO
	450#2	4.5	231	26.5 ± 3.6	00 ^h 57 ^m 33 ^s .06	–27°28′39″.7	0.97	0.009	
	450#2	8.0	180	18.4 ± 4.4	00 ^h 57 ^m 33 ^s .05	–27°28′39″.6	1.19	0.011	
RX J094144.51+385434.8	850#1	4.5	242	713 ± 19	09 ^h 41 ^m 44 ^s .69	+38°54′39″.8	3.38	0.003	QSO
	850#1	8.0	180	2147 ± 36	09 ^h 41 ^m 44 ^s .70	+38°54′39″.9	3.41	0.001	QSO
	850#1	24.0	1	5254 ± 182	09 ^h 41 ^m 44 ^s .66	+38°54′39″.5	3.30	0.001	QSO
	850#2	4.5	191	87.9 ± 6.6	09 ^h 41 ^m 46 ^s .34	+38°53′56″.5	2.80	0.013	
	850#2	8.0	155	65.7 ± 6.9	09 ^h 41 ^m 46 ^s .33	+38°53′55″.6	2.45	0.008	
	850#2	24.0	6	1041 ± 166	09 ^h 41 ^m 46 ^s .38	+38°53′57″.0	2.89	0.004	
	850#3	4.5	339	74.7 ± 6.0	09 ^h 41 ^m 42 ^s .66	+38°55′36″.8	2.53	0.012	
	850#3	8.0	241	63.9 ± 6.9	09 ^h 41 ^m 42 ^s .68	+38°55′36″.9	2.50	0.009	
	450#3	4.5	205	19.0 ± 3.1	09 ^h 41 ^m 47 ^s .00	+38°54′11″.2	4.90	0.124	
	450#3	4.5	206	13.1 ± 2.6	09 ^h 41 ^m 46 ^s .39	+38°54′13″.9	4.11	0.131	
	450#3	8.0	159	24.9 ± 4.8	09 ^h 41 ^m 46 ^s .95	+38°54′10″.9	4.69	0.055	
	450#3	8.0	160	22.3 ± 4.7	09 ^h 41 ^m 46 ^s .39	+38°54′13″.9	4.11	0.052	
RX J121803.82+470854.6	850#2	4.5	223	47.8 ± 4.9	12 ^h 18 ^m 07 ^s .57	+47°08′45″.2	0.92	0.008	
	850#2	8.0	175	35.9 ± 5.3	12 ^h 18 ^m 07 ^s .57	+47°08′45″.0	0.90	0.004	
	850#2	24.0	6	476 ± 104	12 ^h 18 ^m 07 ^s .60	+47°08′46″.8	1.76	0.001	
	850#4	4.5	256	4.5 ± 1.5	12 ^h 18 ^m 02 ^s .54	+47°08′28″.3	4.07	0.317	
	850#4	8.0	188	16.7 ± 4.0	12 ^h 18 ^m 02 ^s .67	+47°08′27″.2	2.48	0.038	
	850#5	24.0	5	3200 ± 160	12 ^h 18 ^m 04 ^s .46	+47°08′50″.6	8.87	0.006	Likely false ID
RX J124913.86–055906.2	850#1	4.5	261	3671 ± 42	12 ^h 49 ^m 13 ^s .88	–05°59′19″.0	1.71	< 0.001	QSO
	850#1	8.0	200	12764 ± 88	12 ^h 49 ^m 13 ^s .90	–05°59′18″.9	1.46	< 0.001	QSO
	850#1	24.0	5	26311 ± 125	12 ^h 49 ^m 13 ^s .86	–05°59′18″.4	2.13	0.001	QSO
	850#4	4.5	342	18.1 ± 3.0	12 ^h 49 ^m 12 ^s .56	–05°58′34″.9	3.18	0.093	
	850#4	4.5	343	79.1 ± 6.2	12 ^h 49 ^m 12 ^s .26	–05°58′30″.6	3.34	0.018	
	850#4	8.0	265	55.9 ± 7.5	12 ^h 49 ^m 12 ^s .26	–05°58′31″.2	2.82	0.015	
	850#5	4.5	324	25.5 ± 3.6	12 ^h 49 ^m 11 ^s .39	–05°59′57″.9	2.89	0.061	
	850#5	8.0	236	33.7 ± 6.6	12 ^h 49 ^m 11 ^s .44	–05°59′58″.4	2.00	0.023	
	450#2	4.5	404	2.2 ± 1.1	12 ^h 49 ^m 10 ^s .95	–05°58′34″.7	3.06	0.194	
RX J163308.57+570258.7	850#1	4.5	142	21.3 ± 3.3	16 ^h 32 ^m 59 ^s .74	+57°03′21″.5	3.48	0.076	
	850#1	8.0	134	30.0 ± 5.3	16 ^h 32 ^m 59 ^s .74	+57°03′21″.3	3.29	0.037	
	850#1	24.0	17	275 ± 55	16 ^h 32 ^m 59 ^s .65	+57°03′20″.8	3.47	0.017	
	850#2	4.5	242	83.3 ± 6.4	16 ^h 33 ^m 08 ^s .55	+57°02′54″.6	4.67	0.038	QSO
	850#2	8.0	226	153 ± 10	16 ^h 33 ^m 08 ^s .52	+57°02′54″.5	4.86	0.023	QSO
	850#2	24.0	18	705 ± 40	16 ^h 33 ^m 08 ^s .54	+57°02′53″.0	6.18	0.004	QSO
	850#5	4.5	174	19.1 ± 3.1	16 ^h 33 ^m 03 ^s .17	+57°02′58″.6	0.13	< 0.001	
	850#5	8.0	168	17.0 ± 4.5	16 ^h 33 ^m 03 ^s .18	+57°02′58″.9	0.16	< 0.001	
	850#5	24.0	19	256 ± 32	16 ^h 33 ^m 03 ^s .09	+57°02′57″.6	1.27	0.002	
	850#6	4.5	290	2.9 ± 1.3	16 ^h 33 ^m 12 ^s .45	+57°02′34″.9	3.20	0.172	
	850#7	4.5	225	25.3 ± 3.6	16 ^h 33 ^m 05 ^s .23	+57°04′17″.1	See Appendix A
	850#7	4.5	243	17.1 ± 2.9	16 ^h 33 ^m 06 ^s .23	+57°04′09″.5	See Appendix A
	850#7	4.5	260	29.8 ± 3.9	16 ^h 33 ^m 06 ^s .30	+57°04′03″.2	See Appendix A
	850#7	8.0	232	58.7 ± 6.7	16 ^h 33 ^m 06 ^s .29	+57°04′03″.3	See Appendix A
	850#7	8.0	229	21.5 ± 4.8	16 ^h 33 ^m 06 ^s .24	+57°04′09″.9	See Appendix A
	850#7	8.0	212	20.4 ± 4.7	16 ^h 33 ^m 05 ^s .34	+57°04′17″.6	See Appendix A
	850#7	8.0	211	18.6 ± 4.6	16 ^h 33 ^m 04 ^s .96	+57°04′15″.7	See Appendix A
	850#7	8.0	216	13.6 ± 4.3	16 ^h 33 ^m 05 ^s .52	+57°04′11″.8	See Appendix A
850#7	24.0	8	249 ± 49	16 ^h 33 ^m 05 ^s .05	+57°04′14″.5	See Appendix A	
850#7	24.0	10	270 ± 49	16 ^h 33 ^m 06 ^s .28	+57°04′02″.6	See Appendix A	

Note. Column 2 gives the ID number of the submm galaxy as listed in Table 4 or Table 5; we searched for counterparts to all 450- μm detected objects that were not detected at 850 μm . Column 4 gives the catalogue number for each ID; for reference, these numbers are also shown on Figs A1 and A2. Column 8 gives the angular offset between the SCUBA position and the *Spitzer* position. Column 9 gives the corrected P statistic (see text).




Figure 5. Colour–colour plot for those sources detected in two or three of the observed *Spitzer* bands (24, 8.0 and 4.5 μm). The QSOs are plotted as diamonds while the submm-selected galaxies discovered in their fields are shown as squares. Lines show the redshifted colour tracks of three local ultraluminous infrared galaxies (ULIRGs) with (Mrk 231) and without (Arp 220 and NGC 6240) obvious signatures of powerful AGNs at MIR wavelengths. Redshifts of 1.0, 2.0 and 3.0 are marked on each of these tracks.

positions. These radii of 5 arcsec at 4.5 and 8.0 μm and 10 arcsec at 24 μm were chosen to approximately match the theoretical error circles on the submm positions added in quadrature with the uncertainty on the MIR positions, which are significant at 24 μm (see Ivison et al. 2007 for a full discussion). The corrected P statistic (Downes et al. 1986) is given for each source found within the search radii. This quantity is given by $P = 1 - \exp[-P_*\{1 + \ln(P_c/P_*)\}]$ where $P_c = \pi r_s^2 N_T$, $P_* = \pi r^2 N$, r_s is the search radius, N_T is the surface density of objects at the limiting flux density of the catalogue, N is the surface density of objects brighter than S where S is the flux density of a potential counterpart galaxy located a distance, r , away from the submm position.

A value of $P \leq 0.05$ is typically used as a threshold to select robust radio IDs (e.g. Ivison et al. 2002, 2007) to submm galaxies. We select all MIR sources found within our search radii as potential counterparts and use the P statistic to distinguish between competing IDs when more than one galaxy is found. We note that almost all SMGs listed in Table 8 have MIR counterparts for which the computed P value is < 0.05 in at least one waveband, and can therefore be considered robust based on this criterion.

The four QSOs detected at submm wavelengths have counterparts in all three *Spitzer* bands. In addition, we find counterparts in at least one waveband for ~ 60 per cent of the SMGs. Postage stamp images of the MIR counterparts are shown in Figs A1 and A2 which can be found in Appendix A.

7 MID-INFRARED COLOURS

MIR colours provide a powerful means of distinguishing between galaxies which contain AGN and those that do not (e.g. Egami et al. 2004; Ivison et al. 2004, 2007). Luminous galaxies at $1 < z < 3$ hosting AGN have power-law MIR spectra, whereas those without

AGN are dominated by starlight in the IRAC bands and dust/PAH (polycyclic aromatic hydrocarbon) emission at 24 μm .

In Fig. 5, we plot the ratio of 24- and 8- μm flux density against the ratio of 8- and 4.5- μm flux density for those objects detected in at least two *Spitzer* bands. We superimpose redshift tracks for the local ULIRGs Mrk 231 [AGN-like spectral energy distribution (SED)] and Arp 220 and NGC 6240 (starburst-like SEDs). The QSOs and SMGs clearly inhabit different regions of this diagram. As expected, the QSOs have colours more similar to Mrk 231 than to the starburst galaxies. Three QSOs have remarkably similar MIR colours given that they were selected on their X-ray properties. The other QSO, RX J163308, has significantly different IRAC colours but we note from the postage stamp image (Fig. A1) that it has a close ‘companion’ galaxy which is blended with the QSO in our extracted catalogue and will thus contaminate the colours. While the other three QSOs have 8.0/4.5 μm flux density ratios roughly consistent with the Mrk 231 template at their known redshifts we note that their 24/8.0 μm flux-density ratios appear to be lower than that of Mrk 231 by a factor of ~ 2 which might be related to the relative strength of emission/absorption features. All but one of the SMGs on the other hand have MIR colours consistent with Arp 220 and/or NGC 6240 at $1 < z < 3$. The one exception has colours consistent with Arp 220 at $0.1 < z < 0.5$ but its faintness and non-detection at 24 μm are more consistent with a Mrk 231-like object at $1 < z < 3$. It is thus the only SMG in our sample for which MIR colours indicate significant emission from an AGN. We thus find no strong evidence for buried AGN in the SMGs. Having said this, we note that deep X-ray (e.g. Vignati et al. 1999; Netzer et al. 2005) and IR spectroscopy (e.g. Lutz et al. 2003; Armus et al. 2006) observations of NGC 6240 show that this galaxy does in fact host a highly obscured AGN but that the energetics are dominated by the starburst. This leaves open the possibility that many of the SMGs

contain buried AGN which are not powerful enough to dominate over the starburst in the MIR.

8 DISCUSSION AND CONCLUSIONS

We have shown that the number counts of SMGs with flux densities $\sim 2\text{--}4$ mJy in the ~ 2 arcmin diameter fields surrounding our X-ray absorbed QSOs is higher than that found for a blank field. Most of these galaxies have MIR colours very similar to the local ULIRG Arp 220 if it were placed at $1 < z < 3$. Fitting the mm to FIR spectrum of Arp 220 with a greybody and then integrating to determine the FIR luminosity, we find $L_{\text{FIR}} = (1.3 \pm 0.1) \times 10^{12} L_{\odot}$ which corresponds to a SFR of $224 \pm 17 M_{\odot} \text{ yr}^{-1}$ after applying the usual scaling-law (Kennicutt 1998). Using the same greybody fit and applying the appropriate k -correction leads to predicted 850- μm flux densities of 1.73, 1.44 and 1.42 mJy for Arp 220 at $z = 1, 2$ and 3, respectively. The SMGs in our sample have 850- μm flux densities ranging from about 3–10 mJy implying that their SFRs range between 400–1300 $M_{\odot} \text{ yr}^{-1}$ if Arp 220 is indeed a good analogue. Therefore, if the starburst can be sustained for a few hundred million years or so then a stellar mass equivalent to that of the bulge of a large galaxy will be assembled.

It is natural to speculate that an overdense region of space at high redshift containing galaxies that are capable of building large, homogeneous populations of stars will evolve over cosmic time to become clusters of galaxies containing a population of massive elliptical galaxies. Since it is well established that elliptical galaxies contain dormant SMBHs (e.g. Magorrian et al. 1998), and that these must have assembled at high redshifts where the accretion luminosity density peaks, we expect that our SMGs should contain growing black holes, i.e. AGN. How the major growth phase of the black hole coincides temporally with the luminous starburst that builds up the stellar mass is a current topic of debate. Indeed, models predict a range of possibilities (e.g. Archibald et al. 2002; Di Matteo, Springel & Hernquist 2005), but there must be a link between the formation of the stars in the bulge and the SMBH because of the tight correlation between their masses found for local galaxies (Ferrarese & Merritt 2000). After a number of early null results using moderate depth X-ray data (e.g. Fabian et al. 2000; Waskett et al. 2003), deep X-ray (Alexander et al. 2005a,b, 2008) and MIR spectroscopy (Menéndez-Delmestre et al. 2007; Pope et al. 2008) observations show that SMGs found in the field often do contain AGN but that the masses of their black holes may be much smaller than those found in high-redshift QSOs, and consequently that the bulk of the FIR emission is due to the starburst. Moreover, estimates of the host galaxy masses suggest that the growth of the black hole lags the growth of the spheroid (Borys et al. 2005; Alexander et al. 2008; see also Coppin et al. 2009). While this may be the case for SMGs, analysis of massive galaxies selected on other criteria find evidence to the contrary, suggesting that more than one scenario is possible for the formation of a massive galaxy and its SMBH. For example, studies of powerful radio sources (McLure et al. 2006) and luminous QSOs (Peng et al. 2006) suggest that the black hole mass to bulge luminosity ratio increases towards high redshifts, thus indicating that the black hole reaches its critical mass before the galaxy bulge is fully formed.

The MIR colours of our SMGs also argue against the presence of a powerful AGN in these objects unless they are very highly obscured. They are thus either intrinsically different objects to the QSO or they are caught at a different time in their evolution. Page et al. (in preparation) argue that the X-ray absorption in the QSOs has its origin in an ionized absorber associated with a powerful

wind capable of terminating the star formation. Such feedback is a popular ingredient of models since it leads naturally to the relationship between SMBH mass and spheroid mass discussed above (e.g. Fabian 1999). If the SMGs in the same field evolve to form a cluster of ellipticals, then they would presumably go through a similar evolutionary sequence to the QSO but are caught at an earlier stage of their life cycle at $1 < z < 3$. Estimates of the stellar masses would be useful to put further constraints on their evolutionary status. To this end, we are currently fitting models to optical through IR data to determine photometric redshifts and restframe K -band luminosities; an analysis of the RXJ 094144 field is currently in progress (Carrera et al. in preparation).

One difference worthy of note between the findings of this paper and those for high-redshift (typically $z > 3$) radio galaxies (HzRGs) concerns very bright SMGs in their fields. Two of the seven HzRG fields imaged with SCUBA contain SMGs with $S_{850} > 20$ mJy (Stevens et al. 2003), whereas the brightest object found around the QSOs has $S_{850} \sim 10$ mJy. Although the samples are too small to give a statistically significant result, it appears that the SMGs detected around the lower redshift, less powerful, radio-quiet AGN have lower S_{850} . A direct comparison is not informative because the sources were extracted with different S/N criteria and by different methods but comparing flux densities significantly above the catalogue cut-offs yields eight SMGs with $S_{850} \geq 6$ mJy in the 7 HzRG fields while only two are detected around the five QSOs.

Given the effectively flat relationship between S_{850} and redshift for $1 < z < 10$, this difference in flux density leads to a proportional difference in SFRs (all other things being equal). There is thus reasonably good evidence for a difference in star formation activity in individual galaxies in these fields indicating that the galaxy formation process might be dependent on the magnitude of the density peak and/or its redshift. For example, in the case of HzRGs, it was found that the detection rate increased dramatically at $z > 2.5$ (Archibald et al. 2001; Reuland et al. 2003) which is significantly higher than the average redshift of our QSO sample. Another possibility is that the radio properties of the central AGN are in some way linked to environmental density (e.g. Kauffmann, Heckman & Best 2008). Indeed, recent work by Falder et al. (2010) finds evidence for larger overdensities of galaxies around radio-loud objects (based on work carried out at 3.6 μm). Perhaps radio synchrotron luminosity is enhanced in regions where the intergalactic medium is more dense.

An obvious next step will be to conduct a statistically meaningful survey of fields around AGN over the full range of redshifts in order to track star formation activity as a function of cosmic time. The next generation bolometer camera on the JCMT (SCUBA-2) and SPIRE/PACS on-board the *Herschel Space Observatory* are instruments with good enough sensitivity to make such a campaign practical. It would also allow a bigger area to be mapped around each AGN, better matching the size of proto-cluster regions predicted by numerical simulations.

ACKNOWLEDGMENTS

We thank Ian Smail for extensive comments on the draft manuscript and Mark Thompson for useful discussions. The JCMT is operated by The Joint Astronomy Centre on behalf of the Science and Technology Facilities Council of the United Kingdom, the Netherlands Organisation for Scientific Research and the National Research Council of Canada. JCMT data were taken under project IDs M03AU46, M03BU32 and M04BU14. This work is based (in part) on observations made with the *Spitzer Space Telescope*, which

is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Support for this work was provided by NASA through an award issued by JPL/Caltech. JAS, MJP and FJC acknowledge support from the Royal Society. FJC acknowledges further support from the Spanish Ministerio de Educación y Ciencia under project ESP2006-13608.

REFERENCES

- Alexander D. M., Smail I., Bauer F. E., Chapman S. C., Blain A. W., Brandt W. N., Ivison R. J., 2005a, *Nat*, 434, 738
- Alexander D. M., Bauer F. E., Chapman S. C., Smail I., Blain A. W., Brandt W. N., Ivison R. J., 2005b, *ApJ*, 632, 736
- Alexander D. M. et al., 2008, *AJ*, 135, 1968
- Archibald E. N., Dunlop J. S., Hughes D. H., Rawlings S., Eales S. A., Ivison R. J., 2001, *MNRAS*, 323, 417
- Archibald E. N., Dunlop J. S., Jimenez R., Friaça A. C. S., McLure R. J., Hughes D. H., 2002, *MNRAS*, 336, 353
- Armus L. et al., 2006, *ApJ*, 640, 204
- Austermann J. E. et al., 2010, *MNRAS*, 401, 160
- Bertin E., Arnouts S., 1996, *A&AS*, 117, 393
- Bertoldi F. et al., 2007, *ApJS*, 172, 132
- Borys C., Smail I., Chapman S. C., Blain A. W., Alexander D. M., Ivison R. J., 2005, *ApJ*, 635, 853
- Chapman S. C., Blain A. W., Ivison R. J., Smail I. R., 2003, *Nat*, 422, 695
- Chapman S. C., Blain A. W., Smail I., Ivison R. J., 2005, *ApJ*, 622, 772
- Coppin K., Halpern M., Scott D., Borys C., Chapman S., 2005, *MNRAS*, 357, 1022
- Coppin K. et al., 2006, *MNRAS*, 372, 1621
- Coppin K. et al., 2009, *MNRAS*, 389, 45
- De Breuck C. et al., 2004, *A&A*, 424, 1
- Di Matteo T., Springel V., Hernquist L., 2005, *Nat*, 433, 604
- Downes A. J. B., Peacock J. A., Savage A., Carrie D. R., 1986, *MNRAS*, 218, 31
- Eales S. A., Lilly S., Gear W., Dunne L., Bond J. R., Hammer F., Le Fèvre O., Crampton D., 1999, *ApJ*, 515, 518
- Egami E. et al., 2004, *ApJS*, 154, 130
- Evrard A. E. et al., 2002, *ApJ*, 573, 7
- Fabian A. C., 1999, *MNRAS*, 308, L39
- Fabian A. C. et al., 2000, *MNRAS*, 315, L8
- Falder J. T. et al., 2010, *MNRAS*, in press (arXiv:1001.5419)
- Farrah D. et al., 2006, *ApJ*, 643, L139
- Ferrarese L., Merritt D., 2000, *ApJ*, 539, L9
- Greve T. R., Ivison R. J., Bertoldi F., Stevens J. A., Dunlop J. S., Lutz D., Carilli C. L., 2004, *MNRAS*, 354, 779
- Greve T. R., Stern D., Ivison R. J., De Breuck C., Kovács A., Bertoldi F., 2007, *MNRAS*, 382, 48
- Holland W. S. et al., 1999, *MNRAS*, 303, 659
- Hopkins P. F., Hernquist L., Cox T. J., Kereš D., 2008, *ApJS*, 175, 356
- Hughes D. H. et al., 1998, *Nat*, 394, 241
- Ivison R. J., Dunlop J. S., Smail I., Dey A., Lui M. C., Graham J. R., 2000, *ApJ*, 542, 27
- Ivison R. J. et al., 2002, *MNRAS*, 337, 1
- Ivison R. J. et al., 2004, *ApJS*, 154, 124
- Ivison R. J. et al., 2007, *MNRAS*, 380, 199
- Jenness T., Stevens J. A., Archibald E. N., Economou F., Jessop N. E., Robson E. I., 2002, *MNRAS*, 336, 14
- Kauffmann G., Haehnelt M., 2000, *MNRAS*, 311, 576
- Kauffmann G., Heckman T. M., Best P. N., 2008, *MNRAS*, 384, 953
- Kennicutt R. C., 1998, *ApJ*, 498, 541
- Kurk J. D. et al., 2000, *A&A*, 358, L1
- Lacy M. et al., 2005, *ApJS*, 161, 41
- Lutz D., Sturm E., Genzel R., Spoon H. W. W., Moorwood A. F. M., Netzer H., Sternberg A., 2003, *A&A*, 409, 867
- Magorrian J. et al., 1998, *AJ*, 115, 2285
- Menéndez-Delmestre K. et al., 2007, *ApJ*, 655, L65
- McLure R. J., Jarvis M. J., Targett T. A., Dunlop J. S., Best P. N., 2006, *MNRAS*, 368, 1395
- Mortier A. M. J. et al., 2005, *MNRAS*, 363, 563
- Netzer H., Lemze D., Kaspi S., George I. M., Turner T. J., Lutz D., Boller T., Chelouche D., 2005, *ApJ*, 629, 739
- Page M. J., Mittaz J. P. D., Carrera F. J., 2001a, *MNRAS*, 325, 575
- Page M. J., Stevens J. A., Mittaz J. P. D., Carrera F. J., 2001b, *Sci*, 294, 2516
- Peng C. Y., Impey C. D., Rix H.-W., Kochanek C. S., Keeton C. R., Falco E. E., Lehar J., McLeod B. A., 2006, *ApJ*, 649, 616
- Pentericci L. et al., 2000, *A&A*, 361, L25
- Pope A. et al., 2008, *ApJ*, 675, 1171
- Priddey R. S., Ivison R. J., Isaak K. G., 2008, *MNRAS*, 383, 289
- Reuland M., Röttgering R., van Breugel W., De Breuck C., 2003, *MNRAS*, 353, 377
- Scott S. E. et al., 2002, *MNRAS*, 331, 817
- Seymour N. et al., 2007, *ApJS*, 171, 353
- Smail I., Ivison R. J., Blain A. W., 1997, *ApJ*, 490, L5
- Smail I., Ivison R. J., Blain A. W., Kneib J.-P., 2002, *MNRAS*, 331, 495
- Smail I., Scharf C. A., Ivison R. J., Stevens J. A., Bower R. G., Dunlop J. S., 2003, *ApJ*, 599, 86
- Stevens J. A. et al., 2003, *Nat*, 425, 264
- Stevens J. A., Page M. J., Ivison R. J., Smail I., Carrera F. J., 2004, *ApJ*, 604, L17
- Stevens J. A., Page M. J., Ivison R. J., Carrera F. J., Mittaz J. P. D., Smail I., McHardy I. M., 2005, *MNRAS*, 360, 610
- Venemans B. P. et al., 2007, *A&A*, 461, 823
- Vignati P. et al., 1999, *A&A*, 349, L57
- Waskett T. J. et al., 2003, *MNRAS*, 341, 1217
- Weiß A. et al., 2009, *ApJ*, 707, 1201

APPENDIX A: NOTES AND POSTAGE STAMP IMAGES

In this section, we present and discuss postage stamp images of the QSOs and SMGs (Figs A1 and A2). For the RX J094144 field, we compare the MIR counterparts to the SMGs with those identified previously at optical and near-infrared wavelengths. We also discuss cases where likely MIR counterparts were not catalogued because they fell just outside our formal search radii, and one case where an obvious SMG was not found by our extraction algorithm; these sources were not included in the analysis described in the main part of this paper.

A1 The RX J094144.51+385434.8 field

The SCUBA data for this field were published previously by Stevens et al. (2004). In the present paper, we have treated the data slightly differently in the final reduction stages (rebinning and smoothing). We also apply a formal source extraction algorithm in this work resulting in only two sources at 850 μm and four at 450 μm , whereas Stevens et al. (2004) simply identified peaks on the maps above a 3σ S/N threshold, identifying six sources. As already noted in Section 4.1, the source extraction algorithm fails to detect sources that are blended, which has undoubtedly affected the 850- μm extraction. At 450 μm , we can see that sources 450#1 and 450#3 appear more elongated than the other two objects hinting that they may be resolved. It was these objects that Stevens et al. (2004) identified as blends.

Inspection of the postage stamps shows that source 850#2/450#2 is coincident with a bright MIR source detected in all three *Spitzer* channels. This object was one of a pair of sources detected in the *R* and *K* bands by Stevens et al. (2004). The new *Spitzer* data show that the correct identification is the object with the redder *R* – *K* colour since this object becomes progressively more prominent with





Figure A1. Postage stamps of 25×25 arcsec cut out at the extracted right ascension and declination of the $850\text{-}\mu\text{m}$ sources from Table 4. The panels show (left) MIPS $24\text{-}\mu\text{m}$ contours (black) overlaid on the IRAC $4.5\text{-}\mu\text{m}$ data (grey-scale and white contours) and (right) SCUBA $850\text{-}\mu\text{m}$ contours (black) overlaid on the IRAC $8.0\text{-}\mu\text{m}$ data (grey-scale and white contours). Contour levels at $850\text{ }\mu\text{m}$ start at $\sim 2\sigma$ and increase in $\sim 1\sigma$ steps while the *Spitzer* contours are chosen to highlight faint sources that might not have been formally extracted. The numbering corresponds to the catalogue numbers given in Table 8.

Figure A1 – *continued*

increasing wavelength compared to its neighbour and is detected at 24 μm .

Source 450#3 falls in a crowded field. We find two possible identifications in the IRAC channels. Of these sources, 206/160 lies closer to the 450- μm source centroid and is also coincident with faint (formally undetected) emission at 24 μm . Source 206/160 was not one of the possible identifications discussed by Stevens et al. (2004) since it is very faint/undetected in the K band. We conclude that the likely identification for this SMG is the very red object 206/160.

Source 450#4 was matched with an extremely red object ($R - K = 5.8$) by Stevens et al. (2004). This source is not for-

mally recovered in this work because it lies outside our 5 arcsec IRAC search radius but is visible in the postage stamp to the west of the 450- μm centroid as a bright source at 4.5 and 8.0 μm . Given its red colours, we consider this object as a likely identification. Measured flux densities are 57.0 ± 5.3 and 48.5 ± 6.1 μJy at 4.5 and 8.0 μm , respectively, giving a colour similar to Arp 220 at the redshift of the QSO. The 4.5- μm centroid is 6.6 arcsec from the extracted 450- μm position at $09^{\text{h}}41^{\text{m}}43^{\text{s}}.61$, $+38^{\circ}54'24''.5$ while the 8.0- μm centroid is 6.4 arcsec from the extracted 450- μm position at $09^{\text{h}}41^{\text{m}}43^{\text{s}}.64$, $+38^{\circ}54'24''.6$. The calculated P -statistic is 0.089 and 0.064 at 4.5 and 8.0 μm , respectively.




Figure A2. Postage stamps of 25×25 arcsec cut out at the extracted right ascension and declination of the 450- μm sources from Table 5. The panels show (left) MIPS 24- μm contours (black) overlaid on the IRAC 4.5- μm data (grey-scale and white contours) and (right) SCUBA 450- μm contours (black) overlaid on the IRAC 8.0- μm data (grey-scale and white contours). Contour levels at 450- μm start at $\sim 2\sigma$ and increase in $\sim 1\sigma$ steps while the *Spitzer* contours are chosen to highlight faint sources that might have not been formally extracted. The numbering corresponds to the catalogue numbers given in Table 8.

A2 The RX J121803.82+470854.6 field

In Section 4.3, we noted that the QSO in this field was not found by our extraction algorithm but was detected by Page et al. (2001). The latter observations used SCUBA in ‘photometry’ mode where the central bolometer of each array is centred on the target but no image is made. Inspection of Fig. 1 shows that while there is submm emission at the location of the QSO (located below source 5 on the 850- μm image), the surrounding structure is complex. It is thus possible that the source extraction algorithm failed to locate a significant peak because of blending. It is also likely that the photometry observation of this object reported by Page et al. (2001) is contaminated by emission from the SMGs close to the QSO.

In Table 8, we noted that object 850#5 has a likely false identification at 24 μm . This can also be attributed to the crowded field close to the QSO since it is the QSO itself that has fallen within the search radius at 24 μm while it was not picked up in the smaller search radii adopted for the IRAC channels.

A3 The RX J124913.86–055906.2 field

Source 850#2 has no listed identification in Table 8 but the postage stamp images show bright 4.5- and 8.0- μm sources close to the 850- μm centroid. The closest 4.5- μm source is at $12^{\text{h}}49^{\text{m}}15^{\text{s}}.19$, $-05^{\circ}58'40''.9$ which at 5.3 arcsec from the SMG is only just outside our formal search radius. This object has a flux density of $17.7 \pm 3.0 \mu\text{Jy}$ and is coincident with an 8.0- μm source although the latter is not detected by *SEXTRACTOR*. Inspection of the 8.0- μm image reveals that the diffraction spike from the bright QSO

at the field centre falls on top of this object probably explaining the non-detection. Although the P -statistic for the 4.5- μm source is high at 0.259, we consider this object as a likely identification because of its detection in both IRAC bands.

Sources 850#4 and 850#4 have an obvious identification in the IRAC bands as well as faint (formally undetected) emission at 24 μm .

A4 The RX J163308.57+570258.7 field

Source 850#4 has no listed identification in Table 8 but the postage stamp images show a bright 4.5- and 8.0- μm source close to the 850- μm centroid along with faint (formally undetected) emission at 24 μm . The 4.5- μm source is located at $16^{\text{h}}33^{\text{m}}10^{\text{s}}.15$, $+57^{\circ}02'10''.2$ only 5.2 arcsec from the SMG, has a flux density of 12.9 ± 2.6 and a P -statistic of 0.266. The 8.0- μm source is located at $16^{\text{h}}33^{\text{m}}10^{\text{s}}.08$, $+57^{\circ}02'09''.3$ only 5.9 arcsec from the SMG, has a flux density of 17.6 ± 4.5 and a P -statistic of 0.198. This object has IRAC colours similar to Arp 220 at the redshift of the QSO. The proximity of this object to the SMG together with its detection in both IRAC bands make it a likely identification.

Inspection of the 850- μm map for this field (Fig. 1) shows submm emission north of the QSO at approximately $16^{\text{h}}33^{\text{m}}05^{\text{s}}.75$, $+57^{\circ}04'11''.7$. There is a negative region to the east of this position which is consistent with a negative image of the source located at one of the off-positions. The western off-position would fall off the map. We believe that this is a real SMG (probably two sources) which was missed by the extraction algorithm

because of its proximity to the edge of the map. Inspection of the postage stamp images reinforces this belief since we find two 24- μm sources coincident with the SMG position and at the same approximate position angle as the 850- μm emission. The IRAC data reveal three and five sources at 4.5 and 8.0 μm , respectively. The southern 24-/850- μm source has two possible IRAC identifications. Of these, source 260/232 is the brightest and reddest object; its 8.0-/4.5- μm flux density ratio is 1.96 ± 0.22 . If source 10 is the 24- μm counterpart, then the ratio of flux densities at 24 and 8.0 μm is 4.60 ± 0.99 which would place this object on the Mrk 231 curve shown in Fig. 5 indicating that it hosts a buried AGN, possibly at a lower redshift than RX J163308. The northern 24-/850- μm source has one coun-

terpart at 4.5 μm and 3 at 8.0 μm although the 4.5- μm emission is extended in the same manner as that at 8.0 μm implying that this SMG could have multiple MIR counterparts. Inspection of the flux densities presented in Table 8 shows that the IRAC colours are similar to those of Arp 220 at the redshift of the QSO, so this source is unlikely to host an AGN. Since this system is one of the most interesting in our sample, we include it in Table 8 as source 850#7 although it is not included in the 850 μm number counts analysis because it was not found by the extraction algorithm.

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.