

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

USDA National Wildlife Research Center - Staff  
Publications

U.S. Department of Agriculture: Animal and Plant  
Health Inspection Service

---

2017

# Wide but Variable Distribution of a Hypervirulent *Campylobacter jejuni* Clone in Beef and Dairy Cattle in the United States

Yizhi Tang

Iowa State University

Richard J. Meinersmann

USDA-ARS, rick.meinersmann@ars.usda.gov

Orhan Sahin

Iowa State University, osahin@iastate.edu

Zuowei Wu

Iowa State University, wuzw@iastate.edu

Lei Dai

Iowa State University, ldai@iastate.edu

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.unl.edu/icwdm\\_usdanwrc](https://digitalcommons.unl.edu/icwdm_usdanwrc)



Part of the [Life Sciences Commons](#)

---

Tang, Yizhi; Meinersmann, Richard J.; Sahin, Orhan; Wu, Zuowei; Dai, Lei; Carlson, James C.; Lawrence, Jodie Plumblee; Genzlinger, Linda; LeJeune, Jeffrey T.; and Zhang, Qijing, "Wide but Variable Distribution of a Hypervirulent *Campylobacter jejuni* Clone in Beef and Dairy Cattle in the United States" (2017). *USDA National Wildlife Research Center - Staff Publications*. 2059.

[https://digitalcommons.unl.edu/icwdm\\_usdanwrc/2059](https://digitalcommons.unl.edu/icwdm_usdanwrc/2059)

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Yizhi Tang, Richard J. Meinersmann, Orhan Sahin, Zuowei Wu, Lei Dai, James C. Carlson, Jodie Plumblee Lawrence, Linda Genzlinger, Jeffrey T. LeJeune, and Qijing Zhang



---

U.S. Department of Agriculture  
Animal and Plant Health Inspection Service  
Wildlife Services

U.S. Government Publication

---

Tang, Y., R.J. Meinersmann, O. Sahin, Z. Wu, L. Dai, J. Carlson, J. Plumblee Lawrence, L. Genzlinger, J.T. LeJeune, and Q. Zhang. 2017. Wide but variable distribution of an hypervirulent *Campylobacter jejuni* clone in beef and dairy cattle in the United States. *Applied and Environmental Microbiology* 83(24): e01425-17. doi: 10.1128/AEM.01425-17



# Wide but Variable Distribution of a Hypervirulent *Campylobacter jejuni* Clone in Beef and Dairy Cattle in the United States

Yizhi Tang,<sup>a</sup> Richard J. Meinersmann,<sup>b</sup> Orhan Sahin,<sup>a,c</sup> Zuowei Wu,<sup>a</sup> Lei Dai,<sup>a</sup> James Carlson,<sup>d</sup> Jodie Plumblee Lawrence,<sup>d</sup> Linda Genzlinger,<sup>d</sup> Jeffrey T. LeJeune,<sup>e</sup> Qijing Zhang<sup>a</sup>

Departments of Veterinary Microbiology and Preventive Medicine<sup>a</sup> and Veterinary Diagnostic and Production Animal Medicine,<sup>c</sup> Iowa State University, Ames, Iowa, USA; Russell Research Center, USDA Agricultural Research Service, BEAR-RU, Athens, Georgia, USA<sup>b</sup>; National Wildlife Research Center, USDA APHIS, Fort Collins, Colorado, USA<sup>d</sup>; Food Animal Health Research Program, The Ohio State University, Wooster, Ohio, USA<sup>e</sup>

**ABSTRACT** *Campylobacter jejuni* clone SA is the major cause of sheep abortion and contributes significantly to foodborne illnesses in the United States. Clone SA is hypervirulent because of its distinct ability to produce systemic infection and its predominant role in clinical sheep abortion. Despite the importance of clone SA, little is known about its distribution and epidemiological features in cattle. Here we describe a prospective study on *C. jejuni* clone SA prevalence in 35 feedlots in 5 different states in the United States and a retrospective analysis of clone SA in *C. jejuni* isolates collected by National Animal Health Monitoring System (NAHMS) dairy studies in 2002, 2007, and 2014. In feedlot cattle feces, the overall prevalence of *Campylobacter* organisms was 72.2%, 82.1% of which were *C. jejuni*. Clone SA accounted for 5.8% of the total *C. jejuni* isolates, but its prevalence varied by feedlot and state. Interestingly, starlings on the feedlots harbored *C. jejuni* in feces, including clone SA, suggesting that these birds may play a role in the transmission of *Campylobacter*. In dairy cattle, the overall prevalence of clone SA was 7.2%, but a significant decrease in the prevalence was observed from 2002 to 2014. Whole-genome sequence analysis of the dairy clone SA isolates revealed that it was genetically stable over the years and most of the isolates carried the tetracycline resistance gene *tet(O)* in the chromosome. These findings indicate that clone SA is widely distributed in both beef and dairy cattle and provide new insights into the molecular epidemiology of clone SA in ruminants.

**IMPORTANCE** *C. jejuni* clone SA is a major cause of small-ruminant abortion and an emerging threat to food safety because of its association with foodborne outbreaks. Cattle appear to serve as a major reservoir for this pathogenic organism, but there is a major gap in our knowledge about the epidemiology of clone SA in beef and dairy cattle. By taking advantage of surveillance studies conducted on a national scale, we found a wide but variable distribution of clone SA in feedlot cattle and dairy cows in the United States. Additionally, the work revealed important genomic features of clone SA isolates from cattle. These findings provide critically needed information for the development of preharvest interventions to control the transmission of this zoonotic pathogen. Control of *C. jejuni* clone SA will benefit both animal health and public health, as it is a zoonotic pathogen causing disease in both ruminants and humans.

**KEYWORDS** *Campylobacter jejuni*, cattle, clone SA, prevalence

Received 28 June 2017 Accepted 23 September 2017

Accepted manuscript posted online 29 September 2017

**Citation** Tang Y, Meinersmann RJ, Sahin O, Wu Z, Dai L, Carlson J, Plumblee Lawrence J, Genzlinger L, LeJeune JT, Zhang Q. 2017. Wide but variable distribution of a hypervirulent *Campylobacter jejuni* clone in beef and dairy cattle in the United States. *Appl Environ Microbiol* 83:e01425-17. <https://doi.org/10.1128/AEM.01425-17>.

**Editor** Edward G. Dudley, The Pennsylvania State University

**Copyright** © 2017 Tang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Qijing Zhang, zhang123@iastate.edu.

*Campylobacter jejuni* is a major zoonotic bacterial pathogen and primarily causes foodborne enteritis in humans (1, 2). The organism is widely distributed across a broad range of animal species, including livestock, poultry, and wildlife, and is transmitted to humans mainly via consumption of contaminated food, water, and milk (2). As reported by the Centers for Disease Control and Prevention's (CDC) FoodNet surveillance program in 2016, *Campylobacter* ranked second (12.97 per 100,000 population) among the causes of laboratory-confirmed bacterial foodborne illnesses in the United States (3). Poultry, especially market age broiler chickens, are frequently colonized by *C. jejuni*, resulting in carcass contamination in processing plants (4, 5). Consequently, poultry meat is considered a major source of infection for human campylobacteriosis.

In addition to poultry, cattle also serve as an important reservoir for *Campylobacter*. Bovine *Campylobacter* contributes significantly to both outbreak and sporadic cases of campylobacteriosis in humans (6, 7). *Campylobacter* can be transmitted from cattle to humans via multiple routes, including direct contact (e.g., petting zoo and occupational exposure), consumption of unpasteurized milk (and associated dairy products), and environmental contamination (water, produce, etc.) (8–10). Molecular typing of *C. jejuni* isolates using multilocus sequence typing (MLST) attributed approximately 40% of sporadic human cases to cattle sources in the United Kingdom (11). The contribution of bovine *Campylobacter* to outbreaks of human campylobacteriosis is even more prominent because *Campylobacter* from cattle feces frequently contaminates raw milk (9, 12–14). Ruminant *Campylobacter* may also contaminate water supplies via agricultural runoff, leading to large waterborne outbreaks (8). Of note, red meat is infrequently contaminated by *Campylobacter* (15) and does not appear to play a major role in the transmission of *Campylobacter* to humans. Additionally, ruminants are an integral part of *Campylobacter* ecology and may serve as a source of *Campylobacter* transmission to the environment and other farm animals, such as poultry. Thus, poultry and cattle are the two most important animal reservoirs for this zoonotic pathogen.

*Campylobacter* is highly prevalent in both beef and dairy cattle in the United States and worldwide (10, 16–20). In cattle, *Campylobacter* is mainly carried in the intestinal tract and less frequently can be isolated from the rumen, gallbladder, and bile (11, 21). The predominant *Campylobacter* species isolated from cattle is *C. jejuni*, followed by *C. coli* (10, 22–25). Isolation rates vary with country, herd size and type, age of animals, season, and confinement level (10, 25). In the United States, several nationwide surveillance studies of cattle (National Animal Health Monitoring System [NAHMS] Dairy 1996, 2002, and 2007 and Feedlot'99) indicated that fecal carriage rates ranged from 15 to 50% and the majority of the tested operations (herds/farms/feedlots) were positive for *Campylobacter* (17, 24, 26). Several other studies conducted in different states in the United States also revealed a similar range of prevalence (between 20 and 60% at the fecal sample level) of *Campylobacter* in feedlot cattle and dairy cattle (18, 22, 23, 25, 27).

Although *Campylobacter* mainly colonizes in the gastrointestinal (GI) tract in animals, it may translocate across the intestinal epithelial barrier, leading to systemic infection, such as bacteremia and abortion in small ruminants and occasionally in humans (28). Indeed, *Campylobacter* infection is one of the most prevalent causes of ovine abortion in the United States and worldwide, with an overall abortion rate of 5% to 50% (average, 23.2%) in affected flocks (29). Historically, *Campylobacter fetus* subsp. *fetus* was the major cause of *Campylobacter*-associated ovine abortion. However, studies conducted during late 1980s and early 1990s in the United States revealed a progressive increase in isolation of *C. jejuni* from aborted sheep placentas (30, 31). Recently, our studies demonstrated that a single hypervirulent tetracycline-resistant *C. jejuni* clone (named clone SA) has emerged as the predominant cause of *Campylobacter*-associated ovine abortions and is responsible for >90% of the clinical abortion cases in the United States (29, 32, 33). The hypervirulence of clone SA is related to its ability to translocate across the intestinal epithelium, producing systemic infection and clinical abortion (33). Additionally, clone SA was also associated with bovine and goat abortion cases in the United States (29, 34). Importantly, *C. jejuni* clone SA has

**TABLE 1** Prevalence of *Campylobacter jejuni* and *C. jejuni* clone SA in feces of feedlot cattle and starlings in the United States

State	% prevalence of <i>Campylobacter</i> in cattle (no. of isolates/total no. of samples)	No. (%) of cattle isolates		% prevalence of <i>Campylobacter</i> in birds (no. of isolates/total no. of samples)	No. (%) of bird isolates	
		<i>C. jejuni</i> <sup>a</sup>	Clone SA <sup>b</sup>		<i>C. jejuni</i> <sup>a</sup>	Clone SA <sup>b</sup>
Iowa	69.2 (554/800)	487 (87.9)	16 (3.3)	NA <sup>c</sup>	NA	NA
Texas	71.9 (414/576)	367 (88.6)	42 (11.4)	25.8 (23/89)	23 (100)	1 (4.3)
Missouri	70.0 (210/300)	191 (91.0)	15 (7.9)	90.9 (10/11)	10 (100)	0
Colorado	78.2 (593/758)	438 (73.9)	14 (3.2)	55.0 (11/20)	10 (91.0)	0
Kansas	70.3 (527/750)	403 (76.5)	23 (5.7)	23.3 (7/30)	7 (100)	0
Total	72.2 (2,298/3,184)	1,886 (82.1)	110 (5.8)	34.0 (51/150)	50 (98.0)	1 (2.0)

<sup>a</sup>The percentage is the proportion of *C. jejuni* isolates among the *Campylobacter* isolates.

<sup>b</sup>The percentage is the proportion of clone SA isolates among the *C. jejuni* isolates.

<sup>c</sup>NA, starling samples were not available.

been implicated in a number of cases of foodborne illnesses, both outbreaks and sporadic cases, in the United States (34). These findings clearly indicated that *C. jejuni* clone SA is an important pathogen for both animal health and food safety in the United States and suggest that cattle may serve as a major reservoir for its zoonotic transmission.

Despite the obvious significance of *C. jejuni* clone SA to ruminant health and food safety, little information is available about its distribution in beef and dairy cattle, which represents an important knowledge gap in our understanding of the overall epidemiology and this particular zoonotic risk. To close this knowledge gap and facilitate the control of *C. jejuni* clone SA, we conducted a before-after controlled impact (BACI) study, with repeated sampling of 35 feedlots located in various geographical regions on two different occasions. Additionally, we analyzed the *Campylobacter* isolates in the collections of the NAHMS 2002, 2007, and 2014 dairy studies (17, 35). The purposes of this work were to (i) investigate the overall prevalence of *Campylobacter* in feedlot cattle and evaluate the effect of starling control intervention on the occurrence and spread of *Campylobacter* in feedlot operations and (ii) determine the occurrence and distribution of *C. jejuni* clone SA in feedlot and dairy cattle.

## RESULTS

**Overall prevalence of *Campylobacter* in feedlot cattle.** In total, 2,298 (72.1%) out of 3,184 fecal samples from feedlots were positive for *Campylobacter*. The overall prevalence rates of *Campylobacter* were 69.2% (554/800), 71.9% (414/576), 70.0% (210/300), 78.2% (593/758), and 70.3% (527/750) in Iowa, Texas, Missouri, Colorado, and Kansas, respectively. The *Campylobacter* prevalence rates among the states were not statistically different ( $P > 0.05$ ). Of the 2,298 *Campylobacter* isolates, 1,886 (82.1%) were determined to be *C. jejuni* by PCR. In each of the states, 487 (87.9%), 367 (88.6%), 191 (91.0%), 438 (73.9%), and 403 (76.5%) were identified as *C. jejuni*, respectively (Table 1), indicating that *C. jejuni* was the predominant *Campylobacter* species isolated from cattle feces.

**Effect of starling intervention on *Campylobacter* prevalence in feedlot cattle.** Delineation of the prevalence data pre- and post-starling control intervention is shown in Table 2. The intervention program did not appear to significantly affect the overall prevalences of *Campylobacter* ( $P = 0.10$ ) and *C. jejuni* ( $P = 0.29$ ) in the feedlot cattle. Before intervention, the overall prevalence of *Campylobacter* (69.1% [1,044/1,510 samples tested]) and the relative prevalence of *C. jejuni* (79.3% [828/1,044]) were comparable to those observed postintervention (74.9% and 84.4%, respectively). Also, analysis of data by each state separately indicated no significant differences pre- and postintervention in the prevalence rates for overall *Campylobacter* and *C. jejuni* (Table 2).

**Prevalence of *C. jejuni* clone SA in feedlot cattle.** Initial screening of the *C. jejuni* isolates for putative clone SA using PCR revealed that 8.7% (164/1,886) of the isolates were positive by PCR. As this PCR is not 100% specific for clone SA, pulsed-field gel electrophoresis (PFGE) was performed to confirm their identity as clone SA. Of the 164

**TABLE 2** Prevalence of *Campylobacter* isolated from fecal samples of feedlot cattle before and after starling intervention

State	Preintervention				Postintervention			
	No. of samples tested	No. <i>Campylobacter</i> positive (%)	No. of <i>C. jejuni</i> isolates (%) <sup>a</sup>	No. of clone SA isolates (%) <sup>b</sup>	No. of samples tested	No. <i>Campylobacter</i> positive (%)	No. of <i>C. jejuni</i> isolates (%) <sup>a</sup>	No. of clone SA isolates (%) <sup>b</sup>
Iowa	400	280 (70.0)	245 (87.5)	7 (2.9)	400	273 (68.3)	242 (88.6)	9 (3.7)
Texas	250	170 (68.0)	144 (84.7)	12 (8.3)	326	244 (74.8)	223 (91.4)	30 (13.5)
Missouri	150	85 (56.7)	72 (84.7)	5 (6.9)	150	125 (83.3)	119 (95.2)	10 (8.4)
Colorado	360	274 (76.1)	202 (80.3)	6 (3.0)	398	319 (80.2)	236 (74.0)	8 (3.4)
Kansas	350	235 (67.1)	165 (63.8)	6 (3.6)	400	292 (73.0)	238 (81.5)	17 (7.1)
Total	1,510	1,044 (69.1)	828 (79.3)	36 (4.3)	1,674	1,253 (74.9)	1,058 (84.4)	74 (7.0)


<sup>a</sup>The percentage is the proportion of *C. jejuni* isolates among the *Campylobacter* isolates.

<sup>b</sup>The percentage is the proportion of clone SA isolates among the *C. jejuni* isolates.

isolates initially identified by the PCR as putative clone SA, 110 (67.1%) had patterns that matched to the known subtypes of clone SA—I and II (Fig. 1)—which is in accordance with our previously published results (29). Additionally, MLST was performed on a subset of these isolates in both PFGE subtypes, which identified all of them as sequence type (genotype) 8 (ST-8), which confirmed the PFGE typing result. MLST was also performed on seven isolates with non-clone SA PFGE patterns, two of which with one band difference from clone SA were also identified as ST-8 genotype, while the remaining five isolates were identified with sequence types different from ST-8, including ST-2876, ST-93, ST-239, and ST-14.

Based on the genotyping results, a relative prevalence of 5.8% (110 out of 1,886 *C. jejuni* isolates) and an absolute prevalence of 3.5% (110 out of 3,184 total samples tested) were calculated for clone SA occurrence in the feedlot cattle surveyed in this study. The absolute prevalence rates of clone SA varied by state: 1.8% (14/758) in Colorado, 2.0% (16/800) in Iowa, 3.1% (23/750) in Kansas, 5.0% (15/300) in Missouri, and 7.3% (42/576) in Texas (Table 1). The chi-square test revealed that at least one state is significantly different from the rest ( $P < 0.0001$ ). The relative prevalence rates of clone SA also varied by state: Iowa, 2.9% (16/554); Texas, 10.1% (42/414); Missouri, 7.1% (15/210); Colorado, 2.4% (14/593); and Kansas, 4.4% (23/527) ( $P < 0.0001$ ). However, starling intervention did not affect the prevalence of clone SA on the surveyed farms (Table 2). Clone SA strains were isolated from at least half of the feedlots surveyed in each state, in the range of 1 to 28 isolates per feedlot (result not shown). Although PFGE was performed only on putative clone SA isolates identified by PCR, the non-clone SA *C. jejuni* isolates showed diverse PFGE patterns (Fig. 1), suggesting the overall genetic diversity of *C. jejuni* isolates from feedlot cattle. These findings indicate that *C. jejuni* clone SA is widely distributed and constitutes a substantial portion (~6%) of the total *C. jejuni* population in feedlot cattle.

**Presence of *Campylobacter*, including clone SA, in starlings.** European starlings are commonly found on farms, serving as a potential transmission vehicle for *Campylobacter* (36). To investigate whether they may be a source of farm cattle infection of *C. jejuni* clone SA, fecal samples from European starlings present on 7 feedlots were tested for *Campylobacter* occurrence. Of note, the same feedlots were also sampled for cattle feces at or about the time of the starling survey. Of the 150 total starling fecal samples tested, 51 (34%) were positive for *Campylobacter*; 50 (98%) isolates were identified as *C. jejuni* by PCR, and the remaining isolate was of a species other than *C. jejuni* or *C. coli*. Initial screening using PCR identified 1 of the 50 *C. jejuni* isolates to be a putative clone SA (Table 1). This isolate and an additional 14 randomly chosen *C. jejuni* isolates were analyzed by PFGE, which confirmed that the putative clone SA isolate identified by PCR had a PFGE pattern indistinguishable from IA3902 of clone SA (Fig. 1). MLST analysis further identified this starling isolate as ST-8, indicating that it was a clone SA isolate. All together, these results indicate that starlings carry diverse *C. jejuni* strains and can serve as a vector for transmission of *Campylobacter*, including clone SA, within and between farms.



**FIG 1** Dendrogram showing the PFGE patterns (KpnI) of *C. jejuni* isolates from feces of feedlot cattle and starlings. The clone SA strains are represented by two closely associated PFGE patterns (I and II), as was the case for sheep clone SA isolates (29). IA3902 is a known isolate of clone SA and was used as a reference. The isolates' names are listed on the right of the dendrogram. Starling isolates are marked with stars. TX, Texas; CO, Colorado; MO, Missouri; IA, Iowa; KS, Kansas; FC, feedlot cattle; ST, starling. The numbers in the names of the isolates are arbitrary numbers assigned to feedlots and samples. Please note that the starling isolates represent the total *C. jejuni* population isolated from the birds, while the cattle isolates included in the PFGE analysis were preselected for putative clones SA by the PCR.



**TABLE 3** Occurrence and characteristics of *C. jejuni* clone SA isolates in dairy cows


Dairy study <sup>a</sup>	No. of <i>C. jejuni</i> isolates tested	No. (%) clone SA by:		No. of isolates in which pVir was present	No. of isolates with <i>tet</i> (O) located on:	
		PCR	WGS		Chromosome	pTet
2002	205	23 (11.2)	21 (10.2)	2	13	3
2007	627	66 (10.5)	55 (8.8)	1	40	9
2014	576	39 (6.8)	26 (4.5)	2	15	1
Total	1,408	128 (9.1)	102 (7.2)	5	68	13

<sup>a</sup>NAHMS national surveillance studies.

**Prevalence of clone SA in dairy cattle.** A previous study reported that raw milk was the main source of foodborne illness outbreaks caused by *C. jejuni* clone SA (34), suggesting the presence of clone SA in dairy cattle. Thus, we performed an analysis of the retrospective collections of *Campylobacter* isolates derived from dairy cattle by NAHMS. In 2002, 2007, and 2014, the NAHMS conducted national surveillance studies on *Campylobacter* prevalence in dairy cattle (17, 35). In total, 205, 627, and 576 *C. jejuni* isolates collected in 2002, 2007, and 2014, respectively, were available for clone SA screening. Of these *C. jejuni* collections, 11.2% (23/205), 10.5% (66/627), and 6.8% (39/576) of isolates were initially identified as putative clone SA by PCR ( $n = 128$  total), respectively. All but three (one from the Dairy 2007 study and two from the Dairy 2014 study) of the putative clone SA isolates were subjected to whole-genome sequencing (WGS) analysis. Overall, 16 STs were identified among the genome-sequenced isolates (Table S1). Of the 125 isolates sequenced, 102 (81.6%) were confirmed as clone SA, which gave a relative prevalence of 7.2% (102/1,408) for clone SA among the *C. jejuni* isolates from U.S. dairy cattle. These clone SA isolates included 21 (10.2%) from the Dairy 2002 study, 55 (8.8%) from the Dairy 2007 study, and 26 (3.2%) from the Dairy 2014 study (Table 3). The differences between the earlier time points and the 2014 data were statistically significant ( $P < 0.05$ ). Among those non-clone SA isolates that were PCR positive and whole-genome sequenced, 15 STs were identified (Table S1), five of which were novel sequence types (i.e., they had not been reported previously). Of the 15 STs, 11 were represented by one isolate each, 2 STs were represented by two isolates, 1 ST was represented by five isolates, and 1 ST was represented by three isolates (Table S1).

As carrying tetracycline resistance gene *tet*(O) chromosomally is one of the key features of clone SA isolates from sheep (18), its presence was investigated in the dairy clone SA isolates. Results showed that 81 (79.4%) of the dairy clone SA isolates contained the *tet*(O) gene either in the chromosome ( $n = 68$ ) or on plasmid pTet ( $n = 13$ ), including 13 (61.9%) isolates from the Dairy 2002 study, 51 (92.7%) isolates from the Dairy 2007 study, and 17 (65.4%) isolates from the Dairy 2014 study. In contrast, of the 23 non-clone SA isolates with the whole genome sequenced, 9 isolates harbored a *tet*(O) gene in the pTet plasmid but none of them had *tet*(O) in the chromosome. The pVir plasmid was also found in some of the NAHMS Dairy 2002 ( $n = 2$ ), 2007 ( $n = 1$ ), and 2014 ( $n = 2$ ) study isolates (Table 3).

Previously we have determined the whole-genome sequences of clone SA isolates derived from sheep abortion (33). To investigate the genomic relationship between the clinically abortifacient isolates from sheep and the clone SA isolates from dairy cattle feces, maximum likelihood phylogenetic trees were constructed based on the pan-genome (Fig. 2a) and core genome (Fig. 2b) of the clone SA isolates. The trees were constructed with 170 clone SA isolates, including 72 isolates from sheep abortions collected previously (33) and 98 dairy isolates sequenced in this study (the genomic sequences of 3 dairy isolates were excluded due to poor quality). The 72 ovine isolates represented historical and contemporary isolates of clone SA in the United States over the last 2 decades, while the 98 bovine isolates were selected from the NAHMS studies (2002 to 2014). In both trees, clone SA isolates from sheep and cattle were intermixed and formed clusters irrespective of their host species, indicating that clone SA isolates



**FIG 2** Maximum likelihood phylogenetic tree constructed with the pan-genome (a) and core genome (b) differences among 170 *C. jejuni* clone SA isolates from sheep and cattle. The clone SA strains are intermixed between sheep and cattle and among the isolation years (2002, 2007, and 2014). The isolates are color-coded based on their source hosts and isolation years: red for sheep, blue for NAHMS Dairy 2002 cattle, green for NAHMS Dairy 2007 cattle, and black for NAHMS Dairy 2014 cattle.

were not host specific. There were some discrete clusters in the trees (Fig. 2). The isolates from the same feedlot and collected in the same year tended to be clustered together, but each of the clusters contained isolates from different feedlots. In addition, we did not observe any specific evolution patterns from the genomic data over the 12-year time span, suggesting that the genome of clone SA was fairly stable.

## DISCUSSION

Results from this study revealed a high prevalence (72.2%) of *Campylobacter* spp. in feedlot cattle and the distribution of *C. jejuni* clone SA in both feedlot cattle and dairy cattle in the United States. The identification of *C. jejuni* as the predominant *Campylobacter* species in cattle is consistent with previous findings reported by others (17, 26). Considering that genetically diverse *C. jejuni* strains are present in cattle (32), the prevalence of clone SA (5.8% in feedlot cattle and 7.2% in dairy cattle) is substantial, suggesting that clone SA is well adapted in cattle, similar to the situation in sheep (34).

Additionally, we found that European starlings on cattle farms carry *C. jejuni*, including clone SA, and may serve as a vehicle for the transmission of *Campylobacter* on farms. Furthermore, WGS analysis of the clone SA isolates collected from dairy cattle during the period from 2002 to 2014 revealed the high genomic stability of the isolates. These findings provide new information on the epidemiology of *C. jejuni* clone SA in both beef and dairy cattle. Our study documents the distribution of *C. jejuni* clone SA in beef and dairy cattle, and this work has closed a major knowledge gap regarding the ecology of this zoonotic pathogen in animal reservoirs.

In this study, initial preliminary identification of clone SA was done with a rapid PCR method that targets CJSA\_1356, which is one of the variable genes in the capsule locus and is quite specific for clone SA isolates. Previous work has shown the utility of this PCR method for initial screening for clone SA isolates (37). However, this method is not 100% specific for clone SA, which requires further confirmation of the putative clone SA isolates by other methods. For the prospective study on feedlots, we used PFGE and MLST to confirm the identity of clone SA. In the absence of whole-genome sequences, PFGE and MLST are considered the “gold standards” for establishing clonality in *Campylobacter* isolates (34, 38), and their utility in identifying clone SA was further proven by WGS analysis (33). For the retrospective analysis of the dairy isolates from NAHMS studies, WGS was used to confirm the identity of the clone SA isolates initially identified by PCR. The use of multiple approaches ensured the accuracy of detecting clone SA from a large number of samples.

An interesting finding is that the prevalence rate of clone SA varied significantly in feedlots of different states, highest (7.3%) in Texas and lowest (2.4%) in Colorado. Even within a single state, the prevalence varied from farm to farm. For example, the highest prevalence of clone SA was detected with the no. 4 feedlot in Texas, for which 22 of 47 isolates tested were identified as clone SA, including the clone SA isolate from a starling. The exact reasons for the variable prevalences in different feedlots and states are unknown, but it is possible that the variations are related to differences in management practices that influence transmission and persistence of clone SA in cattle feedlots.

The NAHMS examined *Campylobacter* prevalence in dairy cows by analyzing individual fecal samples in three separate studies: Dairy 2002 (17), Dairy 2007 (35), and Dairy 2014. By taking advantage of the NAHMS's collections of *Campylobacter* isolates, we were able to determine the prevalence of clone SA in dairy cattle on a national scale. The availability of isolates from studies conducted in three different years (2002, 2007, and 2014) allowed us to examine the temporal changes in clone SA prevalence over the years. Interestingly, the prevalences of clone SA in 2002 and 2007 were comparable: 10.2% and 8.8%, respectively. However, in 2014, the prevalence decreased to 4.5%, which is significantly different from those in the previous two studies. What is responsible for the decrease of clone SA in dairy cattle is interesting and remains to be determined in future studies.

It was found in this study that 34% of starling fecal samples were *Campylobacter* positive, with *C. jejuni* identified as the predominant *Campylobacter* species. This prevalence rate is within the range of 11.1% to 50.4% previously reported in the United States and outside the United States (36, 39–41). PCR screening and molecular typing identified one clone SA isolate in the starling samples. Additionally, PFGE analysis of selected starling isolates revealed genetically diverse strains (Fig. 1), consistent with previous findings for starlings (41, 42). Despite the genetic diversity, two isolates (including a clone SA isolate) showed PFGE patterns indistinguishable from those of the cattle isolates (Fig. 1), suggesting that starlings may play a role in spreading *Campylobacter* on cattle farms. It should be pointed out that PFGE analysis of the cattle isolates was biased toward putative clone SA isolates and did not represent the entire genetic profiles of the cattle isolates. Thus, the matching between the cattle and starling isolates might be even higher if more cattle isolates (non-clone SA) were analyzed by PFGE. Regardless, results from this study demonstrated frequent isolation of *Campylobacter* from European starlings on cattle farms and suggest possible two-

way transmission of *Campylobacter* between the two animal species. Interestingly, starling intervention on farms did not affect the overall prevalence of *Campylobacter* (Table 2), suggesting that starling control alone does not appear to be an effective intervention strategy to reduce cattle fecal shedding of *Campylobacter* or clone SA in feedlot cattle. This may be due to the fact that multiple interacting factors contribute to the transmission of *Campylobacter* on cattle farms and control of a single factor has limited impact on its prevalence. Regardless, starlings can be a source for clone SA and can move these isolates between otherwise separate feedlots visited by foraging starlings.

The advances in next-generation sequencing technologies have made it possible to perform high-resolution molecular typing of bacterial isolates. We conducted WGS analysis of the putative clone SA isolates from NAHMS dairy studies, not only for identification of clone SA but also for understanding evolution of clone SA over the 12-year period (2002 to 2014). The WGS analysis confirmed that 102 of the 128 putative clone SA isolates identified by PCR were true clone SA isolates. The genomic data were further used for maximum likelihood phylogenetic tree construction, which revealed that the clone SA isolates derived from 2002 to 2014 are genetically stable and a clear pattern of evolution was not detected as indicated by lack of clustering of the isolates by isolation years (Fig. 2). Inclusion of sheep clone SA isolates (33) in the phylogenetic analysis also revealed that the sheep and cattle isolates are mixed in clustering (Fig. 2), suggesting that the genomic sequences of clone SA isolates are not uniquely associated with host species and the possibility of interspecies (cattle and sheep) transmission of clone SA. These genomic features and the identified wide distribution of clone SA in both beef and dairy cattle suggest that bovine clone SA may serve as an important reservoir for the source of infection in sheep, and clinical abortion induced by *C. jejuni* clone SA continues to be a significant burden for sheep producers (34).

Tetracycline resistance is an important feature of *C. jejuni* clone SA isolated from sheep, and acquisition of this resistance trait is likely due to antibiotic selection pressure, as tetracyclines are frequently used for control of sheep abortion on farms in the United States (34, 43). The *tet(O)* gene is the only tetracycline resistance determinant identified in *Campylobacter* so far. Although *tet(O)* is typically carried by plasmids, it is predominantly located in chromosome in clone SA (34). In this study, we found that 79.4% (Table 3) of the dairy clone SA isolates carried the *tet(O)* gene, and in most of the isolates (68/81) it was located on chromosome. However, the *tet(O)* gene in the non-clone SA isolates was all carried by a plasmid. These results are consistent with our previous findings for the sheep *Campylobacter* isolates (43) and further indicate the advantage of *C. jejuni* clone SA in dealing with the selection pressure from tetracycline antibiotics. The pVir plasmid was also identified in a small number (5/102) of the clone SA isolates in this study. This plasmid is not required for abortion induction by clone SA (44) and is also infrequently present in sheep clone SA isolates (33). Thus, pVir is not unique to clone SA and its *in vivo* function is still unknown.

In summary, this study revealed detailed molecular and epidemiological features of *C. jejuni* clone SA in beef and dairy cattle, as well as in European starlings present on cattle farms. These findings underscore the importance of cattle and wild birds in the overall ecology of *C. jejuni* clone SA in animal reservoirs and provide critically needed information for development of intervention strategies. For example, the high prevalence of *C. jejuni* clone SA in cattle explains why many of the clone SA-associated foodborne disease outbreaks were attributed to consumption of raw milk (34) and highlights the need to reduce fecal contamination of milk and pasteurize milk before consumption to prevent the transmission of clone SA to humans. Additionally, the variable distribution of clone SA on cattle farms suggests that production practices and/or environmental factors may influence its prevalence and may be managed to control clone SA in cattle. Furthermore, our findings also suggest that control of *Campylobacter*-induced abortion in small ruminants should consider intervention strategies that mitigate the transmission of clone SA from the cattle reservoir. These findings provide directions for designing future studies to evaluate intervention strategies.

Considering the significance of *C. jejuni* clone SA in ruminant health and food safety, reducing its prevalence on cattle farms will benefit both animal health and public health.

## MATERIALS AND METHODS

**Sample collection and bacterial isolation.** In the prospective BACI study, a total of 3,184 cattle fecal samples were collected from 35 different feedlot herds located in Iowa, Texas, Colorado, Missouri, and Kansas on two different occasions during the period from December 2012 to March 2013. Collection of cattle fecal samples followed methods described previously (45). A sample was collected from a fecal pat only after a cow was observed defecating. Freshly voided fecal pats were scraped with sterile cotton-tipped swabs, and the swabs were immediately placed in 10-ml glass tubes containing *Campylobacter* thioglycolate broth (CAMPY-THIO). All cattle fecal samples were shipped priority overnight to the testing laboratory. All samples received the next day of collection were accepted and processed to culture *Campylobacter* as described in a previous study (46). Of note, the fecal samples were collected during a European starling intervention program taking place on the farms (39). The intervention program was designed to examine the role of invasive European starlings in the spread of antibiotic-resistant bacteria in feedlots. During the intervention, Wildlife Services biologists baited starlings using a 2% solution of DRC-1339 (3-chloro-*p*-toluidine hydrochloride) on treated corn chop. Technical DRC-1339 powder was mixed with water to create a 2% solution. Treated corn chop was soaked in the 2% solution and screen dried. The bait was applied at a concentration of 1:10 treated to untreated corn chop. All DRC-1339 applications were implemented in accordance with label requirements “Compound DRC-1339 Concentrate—Feedlots” (EPA registration 56228-10). In order to determine the effect of this control program on *Campylobacter* prevalence, approximately one-half of the samples was obtained before the intervention, while the other half was obtained after the intervention.

In addition to cattle fecal specimens, we collected 150 starlings from 7 feedlots (from which postintervention cattle samples were also tested concurrently) within cattle pens and pen lanes during February and March of 2013. All starlings were collected with shotguns, and no birds were collected outside the feedlots. Starling collections followed the methods conforming to agency policy as stated in U.S. Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), Wildlife Service Directive 2.505 and were approved by the National Wildlife Research Center’s (NWRC) Internal Animal Care and Use Committee (NWRC protocol QA-1919). All specimens were individually bagged in sterile Whirl-Paks and stored in coolers until shipping. European starlings were shipped to the NWRC in Fort Collins, CO. All samples received the next day of collection were accepted and processed. All European starling dissections occurred at the NWRC and were conducted using published methods (47). Starling lower gastrointestinal (GI) tracts (duodenum to the cloaca) were removed and placed in sterile Whirl-Paks. To reduce the risk of cross-contamination, we cleaned the starling carcasses, scissors, scalpels, and lab stations with 70% ethanol before the removal of each starling GI tract. Lab mats and gloves were replaced after processing of each starling. The starling GI samples were macerated for 120 s at 200 rpm using a Stomacher 80 paddle blender (Seward Laboratory Systems, Bohemia, NY). Fecal material from the macerated starling GI tracts was squeezed by hand to one corner of the bag and an aliquot was extracted using sterile cotton swabs, making sure to completely saturate the tip of the swab.

In the laboratory, 1 ml of the transport medium containing a fecal swab was added into a tube containing 9 ml of *Campylobacter* enrichment broth, which was then incubated at 42°C for 48 h under microaerobic conditions (5% O<sub>2</sub>, 10% CO<sub>2</sub>, and 85% N<sub>2</sub>). The enrichment medium was Mueller-Hinton (MH) broth supplemented with *Campylobacter*-specific selective agents (SR084E and SR117E; Oxoid). From the enrichment culture, an inoculum of 100 μl was streaked onto an MH agar plate containing the same supplements, which was further incubated for 48 h at 42°C under microaerobic conditions. A single *Campylobacter*-like colony from each sample was subpassaged onto a plain MH agar plate and the pure culture was stored in glycerol stocks at –80°C until further use.

To determine the distribution of *C. jejuni* clone SA in dairy cattle feces, retrospective collections of *Campylobacter* isolates from the NAHMS 2002, 2007, and 2014 dairy studies (17, 35, 48) were screened for clone SA. Respectively, 205, 627, and 576 *C. jejuni* isolates from the three studies were screened for putative clone SA using a specific PCR (see below). Further confirmation of the putative clone SA isolates was performed via whole-genome sequence (WGS) analysis.

**DNA extraction and PCR identification.** DNA was extracted from *Campylobacter* colonies using single-cell lysis buffer (49) and was used as the template for PCRs. In order to detect and/or differentiate *C. jejuni*, *C. coli*, and *C. jejuni* clone SA, three sets of previously published primers were used. The first primer pair (CCCJ-F, 5'-AAT CTA ATG GCT TAA CCA TTA-3', and CCCJ-R, 5'-GTA ACT AGT TTA GTA TTC CGG-3'), targeting 16S rRNA, was designed to coidentify *C. jejuni* and *C. coli* (50). The second primer pair (mapA-F, 5'-GAG TGC TTG TGC AAC TAA AC-3', and mapA-R, 5'-ATA GCA TCT TGA GTT GCT CC-3') was specific for *C. jejuni* (51). The third PCR primer pair (CJSA\_1356F, 5'-TCC CAT TTG GAT GTT GTT GA-3', and CJSA\_1356R, 5'-CAG AAC CTG GCC ACA AAC TT-3') was used for identification of putative *C. jejuni* clone SA as described previously (37). *C. jejuni* IA3902, a clinical isolate of clone SA, was used as a positive control for the PCR, whereas reactions with no DNA template were used as negative controls. Each PCR amplification was carried out in a 25-μl volume containing 16 μl of distilled water, 2.0 μl of template DNA, 10 pmol of each primer, and 5 μl of GoTaq (Promega) green master mix by following the cycling conditions described previously (37, 50, 51).

**PFG.** Pulsed-field gel electrophoresis (PFGE) analysis of *C. jejuni* isolates was performed using KpnI by following the PulseNet protocol (Centers for Disease Control and Prevention [CDC]), with minor

modifications (29). Briefly, fresh cultures of *Campylobacter* were embedded in 1% SeaKem Gold agarose (Fisher Scientific, Fair Lawn, NJ) and lysed with proteinase K for 1 h at 55°C in a water bath shaker. The gel plugs were digested with KpnI for 4 h at 37°C. Digested plugs were embedded into 1% agarose and separated by electrophoresis in 0.5× Tris-borate-EDTA (TBE) buffer (Promega) at 14°C for 18 h using a Chef Mapper electrophoresis system (Bio-Rad, Hercules, CA). Gels were stained with ethidium bromide for 30 min and then photographed by using Chemilmager 5500 (Alpha Innotech, CA). The PFGE patterns were analyzed by the GelCompare II v.6.5 software program (Applied Maths, Kortrijk, Belgium) using the Dice similarity coefficient and unweighted-pair group method with arithmetic averages (UPGMA) with 0.5% optimization and 1.5% position tolerance. *C. jejuni* IA3902 was used as a control for identification of *C. jejuni* clone SA isolates. A lambda DNA ladder (Bio-Rad) was used as the molecular size marker.

**MLST.** To confirm the PFGE results, multilocus sequence typing (MLST), originally developed by Dingle et al. (52), was performed on 11 representative *C. jejuni* isolates (10 from cattle and 1 from a starling) from the prospective study on feedlots. Of the 11 isolates chosen, 4 (3 from cattle and 1 from a starling) had PFGE profiles indistinguishable from that of the positive control (*C. jejuni* IA3902), 4 had minor differences in PFGE patterns, and 3 showed totally different PFGE profiles. The seven housekeeping genes from these 11 *C. jejuni* isolates were amplified and sequenced using the primer sets described at the *C. jejuni* MLST website (<http://pubmlst.org/campylobacter/>), which was developed by Keith Jolley and Man-Suen Chan at the University of Oxford (53). Allele numbers were assigned to the isolates by performing BLAST searches for the assembled sequences using the single-locus query function, whereas sequence types were assigned using the allelic profile query function in the MLST database. Sequences that were identical to existing alleles in the MLST database were assigned the corresponding allele numbers. Novel allele profiles ( $n = 5$ ) were assigned new sequence types (STs) within the MLST database.

**WGS analysis.** The putative clone SA isolates identified by PCR screening from the retrospective NAHMS dairy studies were subjected to WGS. Total DNA was extracted from each isolate using the Wizard Genomic DNA purification kit (Promega) and then used for WGS. The library was constructed using the NEXT Ultra DNA library prep kit (New England Biolabs), and 250-bp paired-end reads were obtained using an Illumina HiSeq2500 (Bionova 42 Biotech Co.). A draft assembly of the sequences of each genome was generated using the *de novo* short-read assembler Velvet (54) and VelvetOptimiser (<http://www.vicbioinformatics.com/software/velvetoptimiser.shtml>). Draft genome sequences were aligned and the core genome phylogenetic tree was constructed using the single nucleotide polymorphisms (SNPs) by Parsnp in the Harvest package (55), while the pan-genome phylogenetic tree was constructed using binary accessory nucleotide data by Panseq (56). The phylogenetic tree was visualized with FigTree (<http://tree.bio.ed.ac.uk/software/figtree>). MLST of these isolates was performed using the sequences of the 7 housekeeping genes according to the scheme from PubMLST (<https://pubmlst.org/campylobacter/>).

**Statistical analysis.** A chi-square test was used to compare the prevalences of *Campylobacter* before and after starling intervention as well as the prevalences of clone SA in different states for the feedlot cattle and in different years for the dairy cattle. The prevalence of *Campylobacter* was taken as the response, and the intervention, the state, and the year were taken as the factors. The independence between the factors and the response was tested by SPSS (version 17.0; SPSS Inc., Chicago, IL). *P* values less than 0.05 were considered significant.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AEM.01425-17>.

**SUPPLEMENTAL FILE 1**, PDF file, 0.6 MB.

## ACKNOWLEDGMENTS

We thank Nada Pavlovic for technical assistance.

This project was supported by Agriculture and Food Research Initiative Competitive Grant no. 2012-67005-19614 and grant no. 2013-67015-20368 from the USDA National Institute of Food and Agriculture.

We have no conflicts of interest to declare.

## REFERENCES

1. Fitzgerald C. 2015. *Campylobacter*. Clin Lab Med 35:289–298. <https://doi.org/10.1016/j.cll.2015.03.001>.
2. Chowdhury MN. 1984. *Campylobacter jejuni* enteritis; a review. Trop Geogr Med 36:215–222.
3. CDC. 2016. Infection with pathogens transmitted commonly through food and the effect of increasing use of culture-independent diagnostic tests on surveillance—Foodborne Diseases Active Surveillance Network, 10 U.S. sites, 2012–2015. MMWR Morb Mortal Wkly Rep 65:368–371.
4. Elvers KT, Morris VK, Newell DG, Allen VM. 2011. Molecular tracking, through processing, of *Campylobacter* strains colonizing broiler flocks. Appl Environ Microbiol 77:5722–5729. <https://doi.org/10.1128/AEM.02419-10>.
5. Berrang ME, Bailey JS, Altekruze SF, Patel B, Shaw WK, Jr, Meinersmann RJ, Fedorka-Cray PJ. 2007. Prevalence and numbers of *Campylobacter* on broiler carcasses collected at rehanging and postchill in 20 U.S. processing plants. J Food Prot 70:1556–1560. <https://doi.org/10.4315/0362-028X-70.7.1556>.
6. Hannon SJ, Taboada EN, Russell ML, Allan B, Waldner C, Wilson HL, Potter A, Babiuk L, Townsend HG. 2009. Genomics-based molecular epidemiology of *Campylobacter jejuni* isolates from feedlot cattle and from

- people in Alberta, Canada. *J Clin Microbiol* 47:410–420. <https://doi.org/10.1128/JCM.01432-08>.
7. Mullner P, Shadbolt T, Collins-Emerson JM, Midwinter AC, Spencer SE, Marshall J, Carter PE, Campbell DM, Wilson DJ, Hathaway S, Pirie R, French NP. 2010. Molecular and spatial epidemiology of human campylobacteriosis: source association and genotype-related risk factors. *Epidemiol Infect* 138:1372–1383. <https://doi.org/10.1017/S0950268809991579>.
  8. Clark CG, Price L, Ahmed R, Woodward DL, Melito PL, Rodgers FG, Jamieson F, Ciebin B, Li A, Ellis A. 2003. Characterization of waterborne outbreak-associated *Campylobacter jejuni*, Walkerton, Ontario. *Emerg Infect Dis* 9:1232–1241. <https://doi.org/10.3201/eid0910.020584>.
  9. Centers for Disease Control and Prevention. 2009. *Campylobacter jejuni* infection associated with unpasteurized milk and cheese—Kansas, 2007. *MMWR Morb Mortal Wkly Rep* 57:1377–1379.
  10. Stanley K, Jones K. 2003. Cattle and sheep farms as reservoirs of *Campylobacter*. *J Appl Microbiol* 94(Suppl):1045–1135.
  11. Wilson DJ, Gabriel E, Leatherbarrow AJ, Cheesbrough J, Gee S, Bolton E, Fox A, Fearnhead P, Hart CA, Diggle PJ. 2008. Tracing the source of campylobacteriosis. *PLoS Genet* 4:e1000203. <https://doi.org/10.1371/journal.pgen.1000203>.
  12. Heuvelink AE, van Heerwaarden C, Zwartkruis-Nahuis A, Tilburg JJ, Bos MH, Heilmann FG, Hoffhuis A, Hoekstra T, de Boer E. 2009. Two outbreaks of campylobacteriosis associated with the consumption of raw cows' milk. *Int J Food Microbiol* 134:70–74. <https://doi.org/10.1016/j.ijfoodmicro.2008.12.026>.
  13. Lejeune JT, Rajala-Schultz PJ. 2009. Food safety: unpasteurized milk: a continued public health threat. *Clin Infect Dis* 48:93–100. <https://doi.org/10.1086/595007>.
  14. Newkirk R, Hedberg C, Bender J. 2011. Establishing a milkborne disease outbreak profile: potential food defense implications. *Foodborne Pathog Dis* 8:433–437. <https://doi.org/10.1089/fpd.2010.0731>.
  15. Jacobs-Reitsma WLU, Wagenaar J. 2008. *Campylobacter* in the food supply, p 627–644. In Nachamkin I, Szymanski CM, Blaser MJ (ed), *Campylobacter*, 3rd ed. ASM Press, Washington, DC.
  16. Horrocks SM, Anderson RC, Nisbet DJ, Ricke SC. 2009. Incidence and ecology of *Campylobacter jejuni* and coli in animals. *Anaerobe* 15: 18–25. <https://doi.org/10.1016/j.anaerobe.2008.09.001>.
  17. Englen MD, Hill AE, Dargatz DA, Ladely SR, Fedorka-Cray PJ. 2007. Prevalence and antimicrobial resistance of *Campylobacter* in US dairy cattle. *J Appl Microbiol* 102:1570–1577. <https://doi.org/10.1111/j.1365-2672.2006.03189.x>.
  18. Sato K, Bartlett PC, Kaneene JB, Downes FP. 2004. Comparison of prevalence and antimicrobial susceptibilities of *Campylobacter* spp. isolates from organic and conventional dairy herds in Wisconsin. *Appl Environ Microbiol* 70:1442–1447. <https://doi.org/10.1128/AEM.70.3.1442-1447.2004>.
  19. Oporto B, Esteban JI, Aduriz G, Juste RA, Hurtado A. 2007. Prevalence and strain diversity of thermophilic campylobacters in cattle, sheep and swine farms. *J Appl Microbiol* 103:977–984. <https://doi.org/10.1111/j.1365-2672.2007.03328.x>.
  20. Châtre P, Haenni M, Meunier D, Botrel MA, Calavas D, Madec JY. 2010. Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from cattle between 2002 and 2006 in France. *J Food Prot* 73:825–831. <https://doi.org/10.4315/0362-028X-73.5.825>.
  21. Krueger NA, Anderson RC, Krueger WK, Horne WJ, Wesley IV, Callaway TR, Edrington TS, Carstens GE, Harvey RB, Nisbet DJ. 2008. Prevalence and concentration of *Campylobacter* in rumen contents and feces in pasture and feedlot-fed cattle. *Foodborne Pathog Dis* 5:571–577. <https://doi.org/10.1089/fpd.2007.0059>.
  22. Halbert LW, Kaneene JB, Ruegg PL, Warnick LD, Wells SJ, Mansfield LS, Fossler CP, Campbell AM, Geiger-Zwald AM. 2006. Evaluation of antimicrobial susceptibility patterns in *Campylobacter* spp isolated from dairy cattle and farms managed organically and conventionally in the midwestern and northeastern United States. *J Am Vet Med Assoc* 228: 1074–1081. <https://doi.org/10.2460/javma.228.7.1074>.
  23. Bae W, Kaya KN, Hancock DD, Call DR, Park YH, Besser TE. 2005. Prevalence and antimicrobial resistance of thermophilic *Campylobacter* spp. from cattle farms in Washington State. *Appl Environ Microbiol* 71: 169–174. <https://doi.org/10.1128/AEM.71.1.169-174.2005>.
  24. Wesley IV, Wells SJ, Harmon KM, Green A, Schroeder-Tucker L, Glover M, Siddique I. 2000. Fecal shedding of *Campylobacter* and *Arcobacter* spp. in dairy cattle. *Appl Environ Microbiol* 66:1994–2000. <https://doi.org/10.1128/AEM.66.5.1994-2000.2000>.
  25. Sanad YM, Kassem II, Abley M, Gebreyes W, LeJeune JT, Rajashekara G. 2011. Genotypic and phenotypic properties of cattle-associated *Campylobacter* and their implications to public health in the USA. *PLoS One* 6:e25778. <https://doi.org/10.1371/journal.pone.0025778>.
  26. Englen MD, Fedorka-Cray PJ, Ladely SR, Dargatz DA. 2005. Antimicrobial resistance patterns of *Campylobacter* from feedlot cattle. *J Appl Microbiol* 99:285–291. <https://doi.org/10.1111/j.1365-2672.2005.02609.x>.
  27. Besser TE, Lejeune JT, Rice DH, Berg J, Stilborn RP, Kaya K, Bae W, Hancock DD. 2005. Increasing prevalence of *Campylobacter jejuni* in feedlot cattle through the feeding period. *Appl Environ Microbiol* 71: 5752–5758. <https://doi.org/10.1128/AEM.71.10.5752-5758.2005>.
  28. Smith JL. 2002. *Campylobacter jejuni* infection during pregnancy: long-term consequences of associated bacteremia, Guillain-Barre syndrome, and reactive arthritis. *J Food Prot* 65:696–708. <https://doi.org/10.4315/0362-028X-65.4.696>.
  29. Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, Wang L, Yaeger MJ, Hoffman LJ, Zhang Q. 2008. Emergence of a tetracycline-resistant *Campylobacter jejuni* clone associated with outbreaks of ovine abortion in the United States. *J Clin Microbiol* 46: 1663–1671. <https://doi.org/10.1128/JCM.00031-08>.
  30. Delong WJ, Jaworski MD, Ward AC. 1996. Antigenic and restriction enzyme analysis of *Campylobacter* spp associated with abortion in sheep. *Am J Vet Res* 57:163–167.
  31. Kirkbride CA. 1993. Diagnoses in 1,784 ovine abortions and stillbirths. *J Vet Diagn Invest* 5:398–402. <https://doi.org/10.1177/104063879300500316>.
  32. Wu Z, Sippy R, Sahin O, Plummer P, Vidal A, Newell D, Zhang Q. 2014. Genetic diversity and antimicrobial susceptibility of *Campylobacter jejuni* isolates associated with sheep abortion in the United States and Great Britain. *J Clin Microbiol* 52:1853–1861. <https://doi.org/10.1128/JCM.00355-14>.
  33. Wu Z, Periaswamy B, Sahin O, Yaeger M, Plummer P, Zhai W, Shen Z, Dai L, Chen SL, Zhang Q. 2016. Point mutations in the major outer membrane protein drive hypervirulence of a rapidly expanding clone of *Campylobacter jejuni*. *Proc Natl Acad Sci U S A* 113:10690–10695. <https://doi.org/10.1073/pnas.1605869113>.
  34. Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, Plummer PJ, Han J, Yaeger MJ, Zhang Q. 2012. Molecular evidence for zoonotic transmission of an emergent, highly pathogenic *Campylobacter jejuni* clone in the United States. *J Clin Microbiol* 50:680–687. <https://doi.org/10.1128/JCM.06167-11>.
  35. USDA. 2011. *Salmonella, Listeria, and Campylobacter on U.S. dairy operation, 1996–2007*. USDA, APHIS, VS, CEAH, Fort Collins, CO.
  36. Colles FM, McCarthy ND, Howe JC, Devereux CL, Gosler AG, Maiden MC. 2009. Dynamics of *Campylobacter* colonization of a natural host, *Sturnus vulgaris* (European starling). *Environ Microbiol* 11:258–267. <https://doi.org/10.1111/j.1462-2920.2008.01773.x>.
  37. Luo Y, Sahin O, Dai L, Sippy R, Wu Z, Zhang Q. 2012. Development of a loop-mediated isothermal amplification assay for rapid, sensitive and specific detection of a *Campylobacter jejuni* clone. *J Vet Med Sci* 74: 591–596. <https://doi.org/10.1292/jvms.11-0462>.
  38. Lucarelli C, Dionisi AM, Trezzi L, Farina C, Passera M, Karki T, D'Ancona F, Luzzi I. 2016. Molecular and epidemiological analysis of a *Campylobacter jejuni* outbreak in Northern Italy in November 2013. *Foodborne Pathog Dis* 13:490–494. <https://doi.org/10.1089/fpd.2015.2104>.
  39. Sanad YM, Closs G, Jr, Kumar A, LeJeune JT, Rajashekara G. 2013. Molecular epidemiology and public health relevance of *Campylobacter* isolated from dairy cattle and European starlings in Ohio, USA. *Foodborne Pathog Dis* 10:229–236.
  40. Waldenström J, Broman T, Carlsson I, Hasselquist D, Achterberg RP, Wagenaar JA, Olsen B. 2002. Prevalence of *Campylobacter jejuni*, *Campylobacter lari*, and *Campylobacter coli* in different ecological guilds and taxa of migrating birds. *Appl Environ Microbiol* 68:5911–5917. <https://doi.org/10.1128/AEM.68.12.5911-5917.2002>.
  41. Sippy R, Sandoval-Green CM, Sahin O, Plummer P, Fairbanks WS, Zhang Q, Blanchong JA. 2012. Occurrence and molecular analysis of *Campylobacter* in wildlife on livestock farms. *Vet Microbiol* 157:369–375. <https://doi.org/10.1016/j.vetmic.2011.12.026>.
  42. French NP, Midwinter A, Holland B, Collins-Emerson J, Pattison R, Colles F, Carter P. 2009. Molecular epidemiology of *Campylobacter jejuni* isolates from wild-bird fecal material in children's playgrounds. *Appl Environ Microbiol* 75:779–783. <https://doi.org/10.1128/AEM.01979-08>.
  43. Wu Z, Sahin O, Shen Z, Liu P, Miller WG, Zhang Q. 2013. Multi-omics approaches to deciphering a hypervirulent strain of *Campylobacter*

- jejuni*. *Genome Biol Evol* 5:2217–2230. <https://doi.org/10.1093/gbe/evt172>.
44. Shen Z, Patil RD, Sahin O, Wu Z, Pu XY, Dai L, Plummer PJ, Yaeger MJ, Zhang Q. 2016. Identification and functional analysis of two toxin-antitoxin systems in *Campylobacter jejuni*. *Mol Microbiol* 101:909–923. <https://doi.org/10.1111/mmi.13431>.
  45. Carlson JC, Engeman RM, Hyatt DR, Gilliland RL, DeLiberto TJ, Clark L, Bodenchuk MJ, Linz GM. 2011. Efficacy of European starling control to reduce *Salmonella enterica* contamination in a concentrated animal feeding operation in the Texas panhandle. *BMC Vet Res* 7:9. <https://doi.org/10.1186/1746-6148-7-9>.
  46. Tang Y, Sahin O, Pavlovic N, LeJeune J, Carlson J, Wu Z, Dai L, Zhang Q. 2017. Rising fluoroquinolone resistance in *Campylobacter* isolated from feedlot cattle in the United States. *Sci Rep* 7:494. <https://doi.org/10.1038/s41598-017-00584-z>.
  47. Carlson JC, Hyatt DR, Ellis JW, Pipkin DR, Mangan AM, Russell M, Bolte DS, Engeman RM, DeLiberto TJ, Linz GM. 2015. Mechanisms of antimicrobial resistant *Salmonella enterica* transmission associated with starling-livestock interactions. *Vet Microbiol* 179:60–68. <https://doi.org/10.1016/j.vetmic.2015.04.009>.
  48. USDA. 2016. Dairy cattle management practices in the United States, 2014. USDA, APHIS, VS, CEAH, NAHMS, Fort Collins, CO.
  49. Olah PA, Doetkott C, Fakhr MK, Logue CM. 2006. Prevalence of the *Campylobacter* multi-drug efflux pump (CmeABC) in *Campylobacter* spp. isolated from freshly processed turkeys. *Food Microbiol* 23:453–460.
  50. Linton D, Lawson AJ, Owen RJ, Stanley J. 1997. PCR detection, identification to species level, and fingerprinting of *Campylobacter jejuni* and *Campylobacter coli* direct from diarrheic samples. *J Clin Microbiol* 35:2568–2572.
  51. Linton D, Owen RJ, Stanley J. 1996. Rapid identification by PCR of the genus *Campylobacter* and of five *Campylobacter* species enteropathogenic for man and animals. *Res Microbiol* 147:707–718. [https://doi.org/10.1016/S0923-2508\(97\)85118-2](https://doi.org/10.1016/S0923-2508(97)85118-2).
  52. Dingle KE, Colles FM, Wareing DR, Ure R, Fox AJ, Bolton FE, Bootsma HJ, Willems RJ, Urwin R, Maiden MC. 2001. Multilocus sequence typing system for *Campylobacter jejuni*. *J Clin Microbiol* 39:14–23. <https://doi.org/10.1128/JCM.39.1.14-23.2001>.
  53. Jolley KA, Chan MS, Maiden MC. 2004. mlstdbNet—distributed multilocus sequence typing (MLST) databases. *BMC Bioinformatics* 5:86. <https://doi.org/10.1186/1471-2105-5-86>.
  54. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
  55. Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol* 15:524. <https://doi.org/10.1186/s13059-014-0524-x>.
  56. Laing C, Buchanan C, Taboada EN, Zhang Y, Kropinski A, Villegas A, Thomas JE, Gannon VP. 2010. Pan-genome sequence analysis using Panseq: an online tool for the rapid analysis of core and accessory genomic regions. *BMC Bioinformatics* 11:461. <https://doi.org/10.1186/1471-2105-11-461>.



Table. S1. Distribution of sequence types and clonal complexes of selected *C. jejuni* isolates from dairy cattle.

CC	ST	No. of isolates	aspA	glnA	gltA	glyA	pgm	tkt	uncA	Percent (%)
21	8	102	2	1	1	3	2	1	6	88
21	21	1	2	1	1	3	2	1	5	0.8
42	42	1	1	2	3	4	5	9	3	0.8
508	132	1	1	6	22	24	12	28	1	0.8
21	262	5	2	1	1	3	2	1	3	4.0
682	682	1	26	2	9	51	8	46	21	0.8
403	933	1	10	1	59	19	10	5	7	0.8
828	1068	1	33	39	30	78	104	43	17	0.8
61	1244	1	1	1	2	2	225	3	17	0.8
21	2876	3	2	2	1	3	2	1	6	2.4
179	3598	2	1	6	137	176	423	32	3	1.6
685	<b>8535</b>	1	1	1	2	4	2	3	6	0.8
21	<b>8536</b>	2	40	1	1	3	2	1	6	1.6
21	<b>8537</b>	1	2	24	1	3	676	1	6	0.8
NA	<b>8538</b>	1	4	6	137	176	40	478	3	0.8
353	<b>8539</b>	1	7	17	234	2	10	3	6	0.8

NA: not assigned to any clonal complex; CC: clonal complex; ST: sequence type; bold numbers: new sequence types identified in this study.