

University of Nebraska - Lincoln  
**DigitalCommons@University of Nebraska - Lincoln**

---

USGS Northern Prairie Wildlife Research Center      Wildlife Damage Management, Internet Center for

---

2011

# Wolves, *Canis lupus*, Carry and Cache the Collars of Radio-collared White-tailed Deer, *Odocoileus virginianus*, they Killed


Michael E. Nelson

U.S. Geological Survey, [Michael\\_nelson@usgs.gov](mailto:Michael_nelson@usgs.gov)

L. David Mech

USGS Northern Prairie Wildlife Research Center, [david\\_mech@usgs.gov](mailto:david_mech@usgs.gov)

Follow this and additional works at: <https://digitalcommons.unl.edu/usgsnpwrc>

 Part of the [Animal Sciences Commons](#), [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Environmental Policy Commons](#), [Recreation, Parks and Tourism Administration Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

---

Nelson, Michael E. and Mech, L. David, "Wolves, *Canis lupus*, Carry and Cache the Collars of Radio-collared White-tailed Deer, *Odocoileus virginianus*, they Killed" (2011). *USGS Northern Prairie Wildlife Research Center*. 357.  
<https://digitalcommons.unl.edu/usgsnpwrc/357>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Northern Prairie Wildlife Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Wolves, *Canis lupus*, Carry and Cache the Collars of Radio-collared White-tailed Deer, *Odocoileus virginianus*, they Killed

MICHAEL E. NELSON<sup>1,2</sup> and L. DAVID MECH<sup>1,3</sup>

<sup>1</sup> U.S. Geological Survey, Northern Prairie Wildlife Research Center, 8711-37<sup>th</sup> Street, SE, Jamestown, North Dakota 58401 USA

<sup>2</sup> Mailing address: 7332 Tracy Road, Duluth, Minnesota N 55803 USA; e-mail: Michael\_nelson@usgs.gov

<sup>3</sup> Mailing address: The Raptor Center, 1920 Fitch Avenue, University of Minnesota, St. Paul, Minnesota 55108 USA

Nelson, Michael E., and L. David Mech. 2011. Wolves, *Canis lupus*, carry and cache the collars of radio-collared White-tailed Deer, *Odocoileus virginianus*, they killed. *Canadian Field-Naturalist* 125(1): 67–68.

Wolves (*Canis lupus*) in northeastern Minnesota cached six radio-collars (four in winter, two in spring-summer) of 202 radio-collared White-tailed Deer (*Odocoileus virginianus*) they killed or consumed from 1975 to 2010. A Wolf bedded on top of one collar cached in snow. We found one collar each at a Wolf den and Wolf rendezvous site, 2.5 km and 0.5 km respectively, from each deer's previous locations.

Key Words: Wolf, *Canis lupus*, White-tailed Deer, *Odocoileus virginianus*, caching, radio-collars, den, homesite, Minnesota.

Wolves (*Canis lupus*) cache food in summer when they travel alone or in small groups (Murie 1944, Cowan 1947, Magoun 1976; Mech and Adams 1999). Caching stores excess food, protects it from scavengers and maggots, and provides a buffer during periods of prey scarcity (Mech 1970). However, caching by free-ranging Wolves has not been reported during winter (Peterson and Cuicci 2003). This is not surprising because much of the Wolf literature is based on forest-dwelling Wolves that have primarily been observed from small aircraft during winter, conditions that have precluded detailed observations of caching. Wolves do return to old kills in winter and conceivably may uncover caches made earlier (Mech 1966).

In a long-term radio-telemetry study of White-tailed Deer (*Odocoileus virginianus*) preyed upon by Wolves in northeastern Minnesota (48°N, 91°W; Nelson and Mech 1986), retrieving the collars of radio-collared deer killed by Wolves revealed that Wolves cached radio-collars. Aerially locating deer 1-3 days/week and retrieving radio-collars of killed deer within 24 hours of discovery presented an unbiased method to document caching in winter, as well as during other seasons.

Of 202 radio-collared deer killed and/or consumed by Wolves during 1975-2010, Wolves cached the collars of six (3%) deer, one each in January, April, August and November, and two in March. Wolves buried four collars in 25-76 cm of snow and two in soil covered by duff and leaf litter. The collar was still attached to the neck and head of research animals for two collars buried in snow, and one of those was beneath a Wolf bed. We found one collar 200 m from the kill site and another in a Wolf summer homesite 500 m from the deer's location just before being killed, although the kill site was not found. Of the remaining 196 radio-collared deer, Wolves in April and May (0-15 cm snow) carried and dropped two collars 300 and 400 m from kill sites. We found a third collar dropped at a Wolf den the first week in April, 2.5 km from the deer's previous

26 January–March locations, and like the previous homesite observation, the kill site was unknown.

Our observations document that Wolves make caches during winter and confirm that Wolves will bed on top of their caches, behavior reported for the Red Fox (*Vulpes vulpes*, Mech 1967) but only suspected for Wolves (Mech 1970).

Presumably caching a radio-collar must be far less frequent than caching food because a collar has no nutritional value. As a nonfood item, it is unclear why Wolves would cache a radio-collar, although, deer hair, blood, scent, and bone-like rigidity of the collar could conceivably play a role in eliciting caching behavior. However, well-fed captive Wolves carry away and cache nonfood objects belonging to their caretakers (Schmidt, International Wolf Center, personal communication). Carrying a radio-collar for considerable distances appears similar to this behavior, which suggests that aspects of caching behavior may be more complex than simply a response to nutritional pressures. Awareness of and attraction to physical objects may be a deep-seated trait in the genome. Hiestand (1989) found that Wolves spontaneously oriented to physical objects in fewer experimental trials than dogs (*Canis familiaris*), a finding corroborated by observations of captive Wolves (Fentress 1967, 1992; MacDonald 1980, Zimen 1981). The selective breeding of dogs for retrieving objects for humans is evidence that possession of objects plays a role in canid social behavior. Because Wolves are the progenitors of dogs, the propensity for this trait must have its origins in the Wolf genome.

### Acknowledgements

This research was supported by the U.S. Geological Survey, United States Department of Agriculture North Central Research Station, and the Special Projects Foundation. We also acknowledge volunteer wildlife technicians who assisted in live capture of deer and who helped retrieve collars. We also thank the pilots

of the U.S. Forest Service for skillful and safe flying. We followed the American Society of Mammalogists Guidelines (Animal Care and Use Committee 1998) for animal handling and care in conducting research.

### Literature Cited

- Animal Care and Use Committee.** 1998. Guidelines for the capture, handling, and care of mammals as approved by the American Society of Mammalogists. *Journal of Mammalogy* 79: 1416-1431.
- Cowan, I. M.** 1947. The timber wolf in the Rocky Mountain national parks of Canada. *Canadian Journal of Research D* 25: 139-74.
- Fentress, J. C.** 1967. Observations on the behavioral development of a hand-reared male timber wolf. *American Zoologist* 7: 339-51.
- Fentress, J. C.** 1992. The covalent animal: On bonds and their boundaries in behavioral research. Pages 44-71 in *The inevitable bond: Examining scientist-animal interactions*. Edited by H. Davis and D. Balfour. Cambridge University Press, Cambridge.
- Hiestand, N. L.** 1989. A comparison of problem-solving and spatial orientation in the wolf (*Canis lupus*) and dog (*Canis familiaris*). Ph.D. dissertation, University of Connecticut, Storrs.
- Magoun, A.** 1976. Summer scavenging activity in northeastern Alaska. Master's thesis, University of Alaska, Fairbanks. 168 pages.
- MacDonald, K. B.** 1980. Activity patterns in a captive wolf pack. *Carnivore* 3:62-64.
- Mech, L. D.** 1966. *The Wolves of Isle Royale*. National Parks Fauna Series Number 7. U.S. Government Printing Office. 210 pages. [Reprinted 2002. University of the Pacific, Honolulu Hawaii].
- Mech, L. D.** 1967. Telemetry as a technique in the study of predation. *Journal of Wildlife Management* 31: 492-496.
- Mech, L. D.** 1970. *The wolf: The ecology and behavior of an endangered species*. Natural History Press, Garden City, New York.
- Mech, L. D., and L. G. Adams.** 1999. Killing of a muskox, *Ovibos mochatus*, by two wolves, *Canis lupus*, and subsequent caching. *Canadian Field-Naturalist* 113: 673-75.
- Murie, A.** 1944. *The wolves of Mount McKinley*. U.S. National Park Service Fauna Series, no. 5. U.S. Government Printing Office, Washington, D.C. 238 pages.
- Nelson, M. E., and L. D. Mech.** 1986. Mortality of white-tailed deer in northeastern Minnesota. *Journal of Wildlife Management* 50: 691-98.
- Peterson, R. O., and P. Ciucci.** 2003. The wolf as a carnivore. Pages 104-130 in *Wolves, behavior, ecology, and conservation*. Edited by Mech, L. D. and L. Boitani. University of Chicago Press, Chicago, Illinois. 448 pages.
- Zimen, E.** 1981. *The wolf: A species in danger*. Delacorte Press, New York.

Received 27 October 2010

Accepted 16 November 2010