

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

2014

West Nile Virus Isolated from a Virginia Opossum (*Didelphis virginiana*) in Northwestern Missouri, USA, 2012

Angela Bosco-Lauth

Centers for Disease Control and Prevention-Fort Collins, mopargal@rams.colostate.edu

Jessica R. Harmon

Centers for Disease Control and Prevention-Atlanta

R. Ryan Jash

Centers for Disease Control and Prevention-Atlanta

Sonja Weiss

Centers for Disease Control and Prevention-Atlanta

Stanley Langevin

Sandia National Laboratories

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc

 Part of the [Life Sciences Commons](#)

Bosco-Lauth, Angela; Harmon, Jessica R.; Jash, R. Ryan; Weiss, Sonja; Langevin, Stanley; Savage, Harry M.; Godsey, Marvin S. Jr.; Burkhalter, Kristen; Root, J. Jeffrey; Gidlewski, Thomas; Nicholson, William L.; Brault, Aaron C.; and Komar, Nicholas, "West Nile Virus Isolated from a Virginia Opossum (*Didelphis virginiana*) in Northwestern Missouri, USA, 2012" (2014). *USDA National Wildlife Research Center - Staff Publications*. 2084.

https://digitalcommons.unl.edu/icwdm_usdanwrc/2084

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Angela Bosco-Lauth, Jessica R. Harmon, R. Ryan Jash, Sonja Weiss, Stanley Langevin, Harry M. Savage, Marvin S. Godsey Jr., Kristen Burkhalter, J. Jeffrey Root, Thomas Gidlewski, William L. Nicholson, Aaron C. Brault, and Nicholas Komar

West Nile Virus Isolated from a Virginia Opossum (*Didelphis virginiana*) in Northwestern Missouri, USA, 2012

Angela Bosco-Lauth,^{1,5} Jessica R. Harmon,² R. Ryan Lash,² Sonja Weiss,² Stanley Langevin,³ Harry M. Savage,¹ Marvin S. Godsey Jr.,¹ Kristen Burkhalter,¹ J. Jeffrey Root,⁴ Thomas Gidlewski,⁴ William L. Nicholson,² Aaron C. Brault,¹ and Nicholas Komar¹ ¹Division of Vector-Borne Diseases, Arboviral Diseases Branch, Centers for Disease Control and Prevention, 3156 Rampart Road, Foothills Campus, Fort Collins, Colorado 80521, USA; ²Division of Vector-Borne Diseases, Rickettsial Zoonoses Branch, Centers for Disease Control and Prevention, 1600 Clifton Road, N.E., Atlanta, Georgia 30333, USA; ³Sandia National Laboratories, 7011 East Ave., Livermore, California 94550, USA; ⁴US Department of Agriculture, National Wildlife Research Center, 4101 LaPorte Avenue, Fort Collins, Colorado 80521, USA; ⁵Corresponding author (email: mopargal@rams.colostate.edu)

ABSTRACT: We describe the isolation of West Nile virus (WNV; *Flaviviridae*, *Flavivirus*) from blood of a Virginia opossum (*Didelphis virginiana*) collected in northwestern Missouri, USA in August 2012. Sequencing determined that the virus was related to lineage 1a WNV02 strains. We discuss the role of wildlife in WNV disease epidemiology.

West Nile virus (WNV; *Flaviviridae*, *Flavivirus*) was introduced into the US in 1999 and has since become the leading cause of domestically acquired arboviral disease in humans (Reimann et al. 2008). The virus circulates in enzootic cycles between *Culex* mosquito vectors and certain competent avian hosts, and causes epidemics/epizootics when it infects large numbers of disease-susceptible hosts, such as humans and horses. Passerine birds are the major WNV amplification hosts and humans serve as incidental hosts, incapable of serving as an infection source for mosquitoes. However, some mammals may play a role in WNV maintenance or amplification. Few WNV isolates have been obtained from sampling wild mammals (Root 2013). Here we describe the isolation of WNV from a Virginia opossum (*Didelphis virginiana*; order Marsupialia) collected in northwestern Missouri.

Free-ranging mammals and birds were sampled in northwestern Missouri, USA, as part of a larger arbovirus surveillance study, which will be described elsewhere. In August 2012, blood samples were collected from 16 Virginia opossums trapped using Tomahawk live traps (81×25×30 cm; Tomahawk Live Traps,

LLC, Hazelhurst, Wisconsin, USA). Opossums were anesthetized with isoflurane in customized anesthesia chambers similar to those described by Bentler et al. (2012) or by intramuscular injection of ketamine/xylazine (60 mg/kg of 5:1 mixture), bled from a peripheral vein, individually marked with ear tags, and subsequently released at their locations of capture. Blood samples were centrifuged for separation of serum, frozen on dry ice, and transported to the Centers for Disease Control, Division of Vector-Borne Diseases, in Fort Collins, Colorado, USA. Samples were thawed and serum tested for virus isolation by plaque assay on Vero cells (Beatty et al. 1995). One of 16 serum samples (sample MO12-W059 from an adult opossum in Nodaway County, Missouri, USA, 10 August 2012) produced viral plaques (2.5 log₁₀ pfu/mL); several plaques were harvested and RNA was extracted for RT-PCR analysis using a One-Step RT-PCR kit according to manufacturer's instructions (Qiagen, Valencia, California, USA). Full-length genome sequencing using Illumina (San Diego, California, USA) next-generation sequencing technology and genetic analysis demonstrated that the isolate was a member of lineage 1a viruses, related to other WNV02 strains as determined by screening for diagnostic residues and comparing to DNA sequences in GenBank. The closest match was an isolate from mosquitoes in Texas in August 2012 with 99.6% shared nucleotide identity and 99.7% amino acid identity (Duggal et al. 2013).

All other serum samples were screened for prior WNV exposure using a plaque reduction neutralization test and none were positive for WNV-neutralizing antibodies (Beaty et al. 1995).

Missouri experienced extreme drought conditions in August 2012 (US Drought Monitor 2012), and very few mosquitoes were present during the sampling period. Following collection by CDC light traps (BioQuip Products, Rancho Dominguez, California, USA) baited with dry ice, mosquitoes were identified, sorted, and pooled by species. The RNA was extracted from pools using a BioRobot Universal (Qiagen) and quantitative reverse transcriptase-PCR was used for WNV RNA detection. A total of 679 mosquitoes were collected in the study region during early August 2012, of which the dominant species ($n=570$) was *Culex erraticus*. All pools were negative for WNV RNA (Savage et al. 2013).

Sylvatic transmission of WNV is poorly understood, but serologic evidence suggests that multiple species of vertebrates are exposed and therefore are potentially involved in viral maintenance or amplification. The roles of mammals other than humans and horses are less clear. Passerine birds of the Missouri rural forests, such as Northern Cardinal (*Cardinalis cardinalis*), Blue Jay (*Cyanocitta cristata*), and American Robin (*Turdus migratorius*), are probable amplification hosts (Bowen and Nemeth 2007). However, many other species of birds, mammals, and even ectotherms are exposed to WNV by a variety of vectors and could play roles in the epizootiology of WNV, or be at risk of developing disease. Large wild carnivores and mesocarnivores, including striped skunk (*Mephitis mephitis*), raccoon (*Procyon lotor*), and black bear (*Ursus americanus*), have been found to be antibody positive for WNV (Bentler et al. 2007; Root 2013). Experimental infection of raccoons has demonstrated moderately high viremias and viral fecal shedding up to 10 days postinfection (Root et al. 2010).

Similarly, some species in the order Rodentia, including fox squirrel (*Sciurus niger*), eastern gray squirrel (*Sciurus carolinensis*), and eastern chipmunk (*Tamias striatus*), as well as eastern cottontail (*Sylvilagus floridanus*; order Lagomorpha), can develop viremias $>10^5$ pfu/mL serum. This level indicates that some mosquitoes could become infected through viremic blood meal acquisition from these mammals (Tiawsirisup et al. 2005; Root et al. 2006; Platt et al. 2008).

In spite of a very low mosquito presence in the region at the time of collection, WNV was isolated for the first time from a Virginia opossum, although neutralizing antibody has been detected in this species previously (Dietrich et al. 2005; Bentler et al. 2007). According to the Arbonet reporting system of the Centers for Disease Control and Prevention, 2012 ranked as the second highest year for human WNV cases in the US, with 2003 being the most severe outbreak year, so it is not surprising that sylvatic WNV activity was evident as well. The predominant mosquito present at the time of sampling, *C. erraticus*, is not considered a primary WNV vector, although it appears to be important for WNV transmission in Alabama swamps and is known to feed preferentially on birds (Hassan et al. 2003; Cupp et al. 2007). Of the other mosquito species present, albeit in low numbers, *Culex tarsalis*, *Culex salinarius*, and *Culex restuans* are all potential WNV vectors. Detection of WNV in a Virginia opossum is a reminder that the diversity of hosts infected with WNV is high and that investigations into novel hosts will be needed to fully assess the potential for alternative transmission cycles that could serve to maintain viral circulation during climactic anomalies such as the drought observed in Missouri in 2012.

Sandia National Laboratories is a multi-program laboratory managed and operated by Sandia Corporation, a wholly owned subsidiary of Lockheed Martin

Corporation, for the US Department of Energy's National Nuclear Security Administration under contract DE-AC04-94AL85000.

LITERATURE CITED

- Beaty BJ, Calisher CH, Shope RE. 1995. Diagnostic procedures for viral, rickettsial, and chlamydial infections. In: *Arboviruses*, Lennette EH, Lennette DA, Lennette ET, editors. American Public Health Association, Washington, DC, pp. 189–212.
- Bentler KT, Hall JS, Root JJ, Klenk K, Schmit B, Blackwell BF, Ramey PC, Clark L. 2007. Serologic evidence of West Nile virus exposure in North American mesopredators. *Am J Trop Med Hyg* 76:173–179.
- Bentler KT, Gossett DN, Root JJ. 2012. A novel isoflurane anesthesia induction system for raccoons. *Wildl Soc Bull* 36:807–812.
- Bowen RA, Nemeth NM. 2007. Experimental infections with West Nile virus. *Curr Opin Infect Dis* 20:293–297.
- Cupp EW, Hassan HK, Yue X, Oldland WK, Lilley BM, Unnasch TR. 2007. West Nile virus infection in mosquitoes in the mid-south USA, 2002–2005. *J Med Entomol* 44:117–125.
- Dietrich G, Monteneri JA, Panella NA, Langevin S, Lasater SE, Klenk K, Kile JC, Komar N. 2005. Serologic evidence of West Nile virus infection in free-ranging mammals, Slidell, Louisiana, 2002. *Vector Borne Zoonotic Dis* 5:288–292.
- Duggal NK, D'Anton M, Xiang J, Seiferth R, Day J, Nasci R, Brault AC. 2013. Sequence analyses of 2012 West Nile virus isolates from Texas fail to associate viral genetic factors with outbreak magnitude. *Am J Trop Med Hyg* 89:205–210.
- Hassan HK, Cupp EW, Hill GE, Katholi CR, Klingler K, Unnasch TR. 2003. Avian host preference by vectors of eastern equine encephalomyelitis virus. *Am J Trop Med Hyg* 69:641–647.
- Platt KB, Tucker BJ, Halbur PG, Blitvich BJ, Fabiosa FG, Mullin K, Parikh GR, Kitikoon P, Bartholomay LC, Rowley WA. 2008. Fox squirrels (*Sciurus niger*) develop West Nile virus viremias sufficient for infecting select mosquito species. *Vector Borne Zoonotic Dis* 8:225–233.
- Reimann CA, Hayes EB, DiGuseppi C, Hoffman R, Lehman JA, Lindsey NP, Campbell GL, Fischer M. 2008. Epidemiology of neuroinvasive arboviral disease in the United States, 1999–2007. *Am J Trop Med Hyg* 79:974–979.
- Root JJ. 2013. West Nile virus associations in wild mammals: A synthesis. *Arch Virol* 158:735–752.
- Root JJ, Oesterle PT, Nemeth NM, Klenk K, Gould DH, McLean RG, Clark L, Hall JS. 2006. Experimental infection of fox squirrels (*Sciurus niger*) with West Nile virus. *Am J Trop Med Hyg* 75:697–701.
- Root JJ, Bentler KT, Nemeth NM, Gidlewski T, Spraker TR, Franklin AB. 2010. Experimental infection of raccoons (*Procyon lotor*) with West Nile virus. *Am J Trop Med Hyg* 83:803–807.
- Savage HM, Godsey MS Jr, Lambert A, Panella NA, Burkhalter KL, Harmon JR, Lash RR, Ashley DC, Nicholson WL. 2013. First detection of Heartland virus (*Bunyaviridae: Phlebovirus*) from field collected arthropods. *Am J Trop Med Hyg* 89:445–452.
- Tiawsirisup S, Platt KB, Tucker BJ, Rowley WA. 2005. Eastern cottontail rabbits (*Sylvilagus floridanus*) develop West Nile virus viremias sufficient for infecting select mosquito species. *Vector Borne Zoonotic Dis* 5:342–350.
- US Drought Monitor. 2012. *The National Drought Mitigation Center at the University of Nebraska-Lincoln*, <http://droughtmonitor.unl.edu/MapsAndData/MapArchive.aspx>. Accessed November, 2013.

Submitted for publication 26 September 2013.

Accepted 25 March 2014.