

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2000

Computer Model for Simulating Population Dynamics of the Predator *Lyctocoris campestris* (Heteroptera: Anthocoridae) in Stored Shelled Corn

James E. Throne

USDA-ARS, Manhattan, KS, james.throne@ars.usda.gov

Megha N. Parajulee

Texas Agricultural Experiment Station

Thomas W. Phillips

Oklahoma State University

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Throne, James E.; Parajulee, Megha N.; and Phillips, Thomas W., "Computer Model for Simulating Population Dynamics of the Predator *Lyctocoris campestris* (Heteroptera: Anthocoridae) in Stored Shelled Corn" (2000). *Publications from USDA-ARS / UNL Faculty*. 2013.
<https://digitalcommons.unl.edu/usdaarsfacpub/2013>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Computer Model for Simulating Population Dynamics of the Predator *Lyctocoris campestris* (Heteroptera: Anthracoridae) in Stored Shelled Corn

JAMES E. THRONE, MEGHA N. PARAJULEE,¹ AND THOMAS W. PHILLIPS²

Grain Marketing and Production Research Center, USDA-ARS, 1515 College Avenue, Manhattan, KS 66502

Environ. Entomol. 29(6): 1236-1243 (2000)

ABSTRACT A computer model for simulating population dynamics of *Lyctocoris campestris* (F.), a predator of stored-product insects, was developed using data from the literature and newly collected data for adult longevity and fecundity. Immature development time and survival, adult longevity, fecundity, and cannibalism were simulated. The model was validated at ambient environmental conditions during three seasons in Wisconsin. The model accurately simulated development of the predator feeding on larvae of the Indianmeal moth, *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae), in the validation studies. The model will be useful in optimizing mass-rearing and augmentative releases of this predator for biological control of stored-product insect pests.

KEY WORDS *Lyctocoris campestris*, stored products, predation, simulation model, biological control, fecundity

Lyctocoris campestris (F.) is a predator of insects in stored grain and other stored commodities. Little was known about the biology of *L. campestris* until recent studies determined its potential as a biological control agent in stored-product ecosystems. Procedures for mass rearing *L. campestris* and developmental parameters at 30°C and 60–70% RH were reported by Parajulee and Phillips (1992). Development from egg to adult takes 30 d at 30°C, and females can lay >250 eggs during a 40-d oviposition period. Both laboratory and field studies indicated that high moisture aided population increase. Adults readily fed on beetle and moth larvae that were as large as 20 mm long. Prey are immobilized within seconds of being bitten, which suggested that a venom was used to overcome prey. The bug's omnivorous feeding habits and ability to feed on large prey indicated potential for use in biological control programs.

A test of 30 species of Coleoptera, Hymenoptera, and Lepidoptera as potential prey for *L. campestris* indicated that there were few immature insects unsuitable as prey (Parajulee and Phillips 1993). However, only a few adults, all stored-product beetles, were attacked and killed by *L. campestris*. Prey species most affected nymphal survivorship and fecundity in life history studies. Both male and female *L. campestris* were able to kill up to 10 lepidopteran or coleopteran

prey in 24 h in laboratory functional response studies (Parajulee et al. 1994). Rate of predation doubled when grain was not present in the search arenas, as might occur in a warehouse storage situation. Rate of predation was lower in oats than in corn or wheat (Meagher and Locke 1997). *L. campestris* is cannibalistic, and cannibalism increases with density (Parajulee and Phillips 1995). Adult longevity is reduced by ≈50–70% when water or prey are not provided (Parajulee and Phillips 1995). Duration of development and survivorship of immature stages of *L. campestris* are affected by temperature, but not relative humidity (Parajulee et al. 1995).

A computer model for simulating population dynamics of *L. campestris* would aid in determining the potential of this predator for controlling stored-product insect pests and in optimizing control programs using this predator. We report on the effects of temperature and relative humidity on adult longevity and fecundity, the development of a computer model for simulating *L. campestris* population dynamics, on studies conducted under ambient outdoor conditions to collect data to validate the model, and on the validity of the model.

Materials and Methods

Insects. *L. campestris* were from a laboratory colony initiated with insects collected from a grain storage in Dane County, WI, in 1991. Additional *L. campestris* collected from grain storages in Dane County were added to the colony each year. The colony was maintained using techniques developed by Parajulee and Phillips (1992). Indianmeal moth, *Plodia interpunc-*

Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA for its use.

¹ Texas Agricultural Experiment Station, 11708 Highway 70 South, P.O. Box 1658, Vernon, TX 76385.

² Oklahoma State University, Department of Entomology and Plant Pathology, 127 Noble Research Center, Stillwater, OK 74078-0464.




Fig. 1. Effects of temperature on life history of adult *Lyctocoris campestris*.

tella (Hübner), larvae from a laboratory colony (rearing procedures in Silhacek and Miller 1972) were used as prey. Voucher specimens of *L. campestris* were placed in the Oregon State University and University of Wisconsin-Madison insect collections.

Adult Longevity and Fecundity. Longevity and fecundity data were obtained at 12 temperature-relative humidity combinations—four temperatures (17, 21, 25, and 29°C) and three relative humidities (43, 58, and 75%)—at a photoperiod of 16:8 (L:D) h, using methods as in Parajulee and Phillips (1992). Adults used in the fecundity study at different temperature-relative humidity regimes were obtained from immatures reared at the respective regimes. Newly laid eggs from the laboratory colony were incubated at each temperature-relative humidity combination. Upon eclosion, newly emerged first instars were individually reared on fifth-instar *P. interpunctella* simultaneously in all 12 environmental conditions. Upon adult emergence, adults were confined as single mating pairs ($n = 10$ –20 pairs per treatment) in individual petri dishes (100 by 15 mm) with one or two fifth-instar *P. interpunctella* as prey. *P. interpunctella* larvae used as prey

for nymphs and adults were frozen before use to kill them, and they were replaced daily. A stack of three filter papers (9-cm-diameter Whatman No. 1, Kent, UK) moistened with distilled water was provided in each petri dish for oviposition. Eggs were harvested and counted from each pair every day until the last female died. Longevity and fecundity data were analyzed using analysis of variance (PROC GLM, SAS Institute 1990). Data for male longevity and duration of female preoviposition and oviposition period were transformed ($\lambda = 0.27, 0.03,$ and $0.13,$ respectively) before analysis using the Box-Cox transformation (Box and Cox 1964) to homogenize variances. Fecundity data were transformed before analysis using a logarithmic transformation.

Model Development. TableCurve 2D software (Jandel Scientific 1996) was used to fit equations describing the effects of temperature on development time and survivorship of immature stages using data from Parajulee et al. (1995) and on adult longevity and fecundity. We did not use the equations developed by Parajulee et al. (1995) because differences between sexes were small and there was no obvious pattern to




Fig. 2. Effects of temperature on mean \pm SD duration of development of immature *Lyctocoris campestris*.

mental conditions and survivorship caused by cannibalism.

Validation. Validation studies were conducted in a screened shed on the University of Wisconsin-Madison campus. Validation tests were set up on 26 May, 7

July, and 18 August 1994, and each test lasted 16 wk. The design for each test was the same. The bottom of each of 16 cans (\approx 8 liters, 24 cm diameter by 19.5 cm high) was covered to a depth of 1 cm with shelled corn (*Zea mays* L.), and five pairs of newly emerged




Fig. 3. Relationship between density and proportion survivorship as a result of cannibalism in *Lyctocoris campestris*.

L. campestris (0–24 h old) adults were placed in each can. Twenty frozen fifth-instar *P. interpunctella* were placed in each can as prey every other day for the first 4 wk, and then 40 larvae were placed in each can every other day for the remainder of the study. An oviposition substrate consisting of a stack of three filter papers (9-cm-diameter Whatman No. 1) moistened with distilled water was also placed in each can every other day, and all filter papers remained in the cans throughout the study. Four cans were destructively sampled every 4 wk, and all live and dead nymphs and adults in the cans were counted. Temperature and




Fig. 4. Environmental conditions recorded hourly during validation tests in a screened shed in Madison, WI. Lines below the graph indicate the duration of each of the three validation studies.




Fig. 5. Observed (circles, mean and 95% confidence intervals) and simulated (solid line) population development of *L. campestris* in the first validation test conducted from 26 May to 15 September in a screened shed in Madison, WI.

relative humidity were monitored in an uninfested can throughout the study (Datapod DP220, Omnidata International, Logan, UT).

Simulations were started with five pairs of newly emerged adults (0–24 h old). Recorded hourly temperatures from the validation study were input and used to determine population development. Our goal was for predictions to be within the 95% CL of population levels in the validation studies.

Results

Adult Longevity and Fecundity. Male longevity varied with temperature ($F = 33.8$; $df = 3, 107$; $P < 0.01$) (Fig. 1), but not relative humidity ($F = 1.8$; $df = 2, 107$; $P = 0.18$; interaction $F = 1.1$; $df = 6, 107$; $P = 0.39$). Duration of female preoviposition period varied with temperature ($F = 29.7$; $df = 3, 111$; $P < 0.01$) (Fig. 1) and relative humidity ($F = 3.9$; $df = 2, 111$; $P = 0.02$; interaction $F = 1.6$; $df = 6, 111$; $P = 0.17$). However,




Fig. 6. Observed (circles, mean and 95% confidence intervals) and simulated (solid line) population development of *L. campestris* in the second validation test conducted from 7 July to 27 October in a screened shed in Madison, WI.




Fig. 7. Observed (circles, mean and 95% confidence intervals) and simulated (solid line) population development of *L. campestris* in the third validation test conducted from 18 August to 8 December in a screened shed in Madison, WI.

there was no obvious pattern to the relationship between duration of preoviposition period and relative humidity, therefore we did not include relative humidity in the model. Duration of female oviposition period varied with temperature ($F = 37.3$; $df = 3, 111$; $P < 0.01$) (Fig. 1), but not relative humidity ($F = 1.5$; $df = 2, 111$; $P = 0.23$; interaction $F = 1.4$; $df = 6, 111$; $P = 0.23$). Similarly, fecundity varied with temperature ($F = 4.5$; $df = 3, 111$; $P < 0.01$) (Fig. 1), but not relative humidity ($F = 1.1$; $df = 2, 111$; $P = 0.35$; interaction $F = 1.3$; $df = 6, 111$; $P = 0.25$).

Model Development. Equations for describing the effects of temperature on life history of *L. campestris* and the effects of density on mortality caused by cannibalism are presented in Table 1 (Figs. 1–3). Although lack-of-fit was significant for some equations, we used these equations because other equations that resulted in a nonsignificant lack-of-fit did not have a form that one would expect based on biological principles.

Validations. In the first validation test, which ran through the summer (26 May to 15 September), temperatures in the cans were between 9.5 and 33°C and relative humidities were between 23 and 92% (Fig. 4). Simulated number of nymphs was always within the 95% confidence intervals for observed numbers of nymphs (Fig. 5). However, simulated numbers of nymphs and adults appeared to lag slightly behind observed population levels, and simulated number of adults at the end of the study was slightly higher than the upper confidence limit for the observed data.

Population levels of both adults and nymphs were always within the 95% confidence intervals for observed numbers of nymphs and adults in the second and third validation tests (Figs. 6 and 7). The second validation test was conducted from midsummer into autumn (7 July to 27 October). Temperatures were between 2.5 and 30°C, and relative humidities were between 54 and 95% (Fig. 4). The third validation test was conducted from late summer into winter (18 Au-

gust to 8 December). Temperatures were between -14 and 29°C, and relative humidities were between 34 and 100% (Fig. 4). No adults were produced in the third validation test because of low winter temperatures. The model accurately simulated *L. campestris* population development under a series of very different environmental conditions.

Discussion

Although simulated population levels were almost always within 95% confidence intervals for observed population levels, population development may be slightly slow in the model. Simulated number of adults appeared to lag behind observed number of adults, although simulated numbers were usually within observed 95% CL. Temperatures to which the insects were actually exposed may have been slightly different than in the can in which we monitored temperature, or fluctuating temperatures in the validation studies may have affected insect development slightly differently than did constant temperatures in the laboratory studies used to collect data to develop the model (Liu et al. 1995). It is also possible that *L. campestris* that were cannibalizing other *L. campestris* in the validation studies may have developed more rapidly than *L. campestris* feeding on *Plodia*. Duration of the nymphal stage varied by 6 d, depending on prey species, in a previous study (Parajulee and Phillips 1993). As with most insects, effects of extreme temperatures on development of *L. campestris* have not been determined. We made assumptions in the model about development at extreme temperatures, and inaccuracies in those assumptions could be partly responsible for the possible time lag in the simulations. Also, rate of cannibalism may vary with temperature, particularly at extreme temperatures, but did not vary with temperature in the model.

Observed numbers of adults dropped more rapidly at the end of the first validation study than in the simulation. Again, this difference may have been a result of slightly slower development in the simulation or to inaccuracy of simulation of cannibalism at extreme temperatures (temperatures were high at the end of the first validation study). However, this may also indicate that cannibalism may not be accurately simulated in the model at high population densities. The population densities at the end of the first validation study were much higher than densities in the studies in which the cannibalism data were collected (Parajulee and Phillips 1995). These population densities would not be expected in a field situation where *L. campestris* would be able to disperse as population levels increased.

The lack of a significant effect of relative humidity in the longevity and fecundity studies was probably a result of providing moist filter papers for oviposition. In an earlier study, eggs laid in *P. interpunctella* larvae in the laboratory did not hatch if the larvae desiccated, and most *L. campestris* trapped in a field study were in portions of the grain mass with high moisture (Parajulee and Phillips 1992). Dependence on moisture for

oviposition and egg hatch may be a limiting factor for use of this predator for biological control of insect pests of grain and stored commodities, but it may be possible to overcome this limitation by providing a contained source of moisture or by repeated augmentative releases of nymphs or adults.

This model is the first developed for a predator of stored-product insects. Models for simulating population development of natural enemies of stored-product insect pests are generally lacking (Throne 1995), although there has been recent work in this area (Flinn and Hagstrum 1995). Such models are useful for understanding population ecology and optimizing pest management strategies.

Lyctocoris campestris has potential for controlling insect pests in both grain storages and in warehouses and other facilities where processed commodities are stored. *L. campestris* is a general predator (Parajulee and Phillips 1993) and quite fecund (Parajulee and Phillips 1992). *L. campestris* is also able to develop at relatively low temperatures (17°C or lower, Parajulee et al. 1995), and therefore it may be suitable for use with aeration (blowing cool air through grain to lower the temperature and help control insect pests, Arthur et al. 1998) or grain chilling with refrigeration (Maier et al. 1996). Use of this simulation model should enhance efforts to optimize mass rearing and release of this species for augmentative biological control.

Acknowledgments

We thank Richard T. Arbogast (USDA-ARS, Gainesville, FL), Robert L. Meagher, Jr. (USDA-ARS, Gainesville, FL), and Lincoln Smith (USDA-ARS, Sidney, MT) for reviewing the manuscript.

References Cited

- Abkin, M. H., and C. Wolf. 1976. Distributed delay routines: DEL, DELS, DELF, DELLF, DELVF, DELLVF. Department of Agricultural Economics Computer Library for Agricultural Systems Simulation Document 8. Michigan State University, East Lansing.
- Arthur, F. H., J. E. Throne, D. E. Maier, and M. D. Montross. 1998. Feasibility of aeration for management of maize weevil populations in corn stored in the southern United States: model simulations based on recorded weather data. *Am. Entomol.* 44: 118-123.
- Box, G.E.P., and D. R. Cox. 1964. An analysis of transformations. *Proc. R. Stat. Soc.* 26: 211-252.
- Draper, N. R., and H. Smith. 1981. *Applied regression analysis*, 2nd ed. Wiley, New York.
- Flinn, P. W., and D. W. Hagstrum. 1995. Simulation model of *Cephalonomia waterstoni* (Hymenoptera: Bethyilidae) parasitizing the rusty grain beetle (Coleoptera: Cucujidae). *Environ. Entomol.* 24: 1608-1615.
- Jandel Scientific. 1996. TableCurve 2D automated curve fitting and equation discovery, version 4. Jandel Scientific, San Rafael, CA.
- Liu, S. S., G. M. Zhang, and J. Zhu. 1995. Influence of temperature variations on rate of development in insects: analysis of case studies from entomological literature. *Ann. Entomol. Soc. Am.* 88: 107-119.
- Maier, D. E., W. H. Adams, J. E. Throne, and L. J. Mason. 1996. Temperature management of the maize weevil,

- Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) in three locations in the United States. *J. Stored Prod. Res.* 32: 255–273.
- Manetsch, T. J. 1976. Time-varying distributed delays and their use in aggregative models of large systems. *IEEE Trans. Syst. Man Cybern.* SMC-6: 547–553.
- Meagher, R. L., Jr., and L. A. Locke. 1997. Predation of Indianmeal moth larvae by *Lyctocoris campestris* (F.) (Hemiptera: Anthocoridae) in different stored commodities. *J. Entomol. Sci.* 32: 271–280.
- Parajulee, M. N., and T. W. Phillips. 1992. Laboratory rearing and field observations of *Lyctocoris campestris* (Hemiptera: Anthocoridae), a predator of stored-product insects. *Ann. Entomol. Soc. Am.* 85: 736–743.
- Parajulee, M. N., and T. W. Phillips. 1993. Effects of prey species on development and reproduction of the predator *Lyctocoris campestris* (Hemiptera: Anthocoridae). *Environ. Entomol.* 22: 1035–1042.
- Parajulee, M. N., and T. W. Phillips. 1995. Survivorship and cannibalism in *Lyctocoris campestris* (Hemiptera: Anthocoridae): effects of density, prey availability, and temperature. *J. Entomol. Sci.* 30: 1–8.
- Parajulee, M. N., T. W. Phillips, and D. B. Hogg. 1994. Funct. response of *Lyctocoris campestris* (F.) adults: effects of predator sex, prey species, and experimental habitat. *Biol. Control* 4: 80–87.
- Parajulee, M. N., T. W. Phillips, J. E. Throne, and E. V. Nordheim. 1995. Life history of immature *Lyctocoris campestris* (Hemiptera: Anthocoridae): effects of constant temperatures and relative humidities. *Environ. Entomol.* 24: 889–897.
- SAS Institute. 1990. SAS/STAT user's guide, version 6, 4th ed. SAS Institute, Cary, NC.
- Silhacek, D. L., and G. L. Miller. 1972. Growth and development of the Indian meal moth, *Plodia interpunctella* (Lepidoptera: Phycitidae), under laboratory mass-rearing conditions. *Ann. Entomol. Soc. Am.* 65: 1084–1087.
- Throne, J. E. 1989. Effects of noncatastrophic control technologies that alter life history parameters on insect population growth: a simulation study. *Environ. Entomol.* 18: 1050–1055.
- Throne, J. E. 1995. Computer modeling of the population dynamics of stored-product pests, pages 169–195. In D. S. Jayas, N.D.G. White, and W. E. Muir [eds.], *Stored-grain ecosystems*. Marcel Dekker, New York.

Received for publication 19 November 1999; accepted 15 August 2000.
