

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USGS Staff -- Published Research

US Geological Survey

8-2-2017

Cytonuclear discordance in the Florida Everglades invasive Burmese python (*Python bivittatus*) population reveals possible hybridization with the Indian python (*P. molurus*)

Margaret E. Hunter

Wetland and Aquatic Research Center, mhunter@usgs.gov

Nathan A. Johnson

Wetland and Aquatic Research Center

Brian J. Smith

Wetland and Aquatic Research Center

Michelle C. Davis

Wetland and Aquatic Research Center

John S.S. Butterfield

Wetland and Aquatic Research Center

Follow this and additional works at: <http://digitalcommons.unl.edu/usgsstaffpub>



Part of the [Geology Commons](#), [Oceanography and Atmospheric Sciences and Meteorology Commons](#), [Other Earth Sciences Commons](#), and the [Other Environmental Sciences Commons](#)

Hunter, Margaret E.; Johnson, Nathan A.; Smith, Brian J.; Davis, Michelle C.; Butterfield, John S.S.; Snow, Ray W.; and Hart, Kristen M., "Cytonuclear discordance in the Florida Everglades invasive Burmese python (*Python bivittatus*) population reveals possible hybridization with the Indian python (*P. molurus*)" (2017). *USGS Staff -- Published Research*. 1043.
<http://digitalcommons.unl.edu/usgsstaffpub/1043>

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Margaret E. Hunter, Nathan A. Johnson, Brian J. Smith, Michelle C. Davis, John S.S. Butterfield, Ray W. Snow, and Kristen M. Hart

ORIGINAL RESEARCH

Cytonuclear discordance in the Florida Everglades invasive Burmese python (*Python bivittatus*) population reveals possible hybridization with the Indian python (*P. molurus*)

Margaret E. Hunter¹  | Nathan A. Johnson¹  | Brian J. Smith² |
Michelle C. Davis¹ | John S. S. Butterfield¹ | Ray W. Snow³ | Kristen M. Hart⁴ 

¹U.S. Geological Survey, Wetland and Aquatic Research Center, Gainesville, Florida

²Wetland and Aquatic Research Center, Cherokee Nation Technologies, Davie, Florida

³U.S. National Park Service, Everglades National Park, Homestead, Florida

⁴U.S. Geological Survey, Wetland and Aquatic Research Center, Davie, Florida

Correspondence

Margaret E. Hunter, U.S. Geological Survey, Wetland and Aquatic Research Center, FL Gainesville, Florida.
Email: mhunter@usgs.gov

Funding information

U.S. Geological Survey

Abstract

The invasive Burmese python (*Python bivittatus*) has been reproducing in the Florida Everglades since the 1980s. These giant constrictor snakes have caused a precipitous decline in small mammal populations in southern Florida following escapes or releases from the commercial pet trade. To better understand the invasion pathway and genetic composition of the population, two mitochondrial (mtDNA) loci across 1,398 base pairs were sequenced on 426 snakes and 22 microsatellites were assessed on 389 snakes. Concatenated mtDNA sequences produced six haplotypes with an average nucleotide and haplotype diversity of $\pi = 0.002$ and $h = 0.097$, respectively. Samples collected in Florida from morphologically identified *P. bivittatus* snakes were similar to published cytochrome oxidase 1 and cytochrome *b* sequences from both *P. bivittatus* and *Python molurus* and were highly divergent (genetic distances of 5.4% and 4.3%, respectively). The average number of microsatellite alleles and expected heterozygosity were $N_A = 5.50$ and $H_E = 0.60$, respectively. Nuclear Bayesian assignment tests supported two genetically distinct groups and an admixed group, not geographically differentiated. The effective population size ($N_E = 315.1$) was lower than expected for a population this large, but reflected the low genetic diversity overall. The patterns of genetic diversity between mtDNA and microsatellites were disparate, indicating nuclear introgression of separate mtDNA lineages corresponding to cytonuclear discordance. The introgression likely occurred prior to the invasion, but genetic information on the native range and commercial trade is needed for verification. Our finding that the Florida python population is comprised of distinct lineages suggests greater standing variation for adaptation and the potential for broader areas of suitable habitat in the invaded range.

KEYWORDS

hybridization, invasive species, mitochondrial marker, nuclear microsatellite marker, phylogenetic population structure

1 | INTRODUCTION

Understanding the processes driving invasion dynamics of non-native species represents an important challenge for biologists and resource managers. Advancements in molecular tools and techniques have allowed for the delimitation of taxonomic units and genetic diversity, and identification of nonnative animals and plants in the absence of reliable morphological data (Bock et al., 2015; Darling, 2015; Serrao, Steinke, & Hanner, 2014). In many cases, only molecular information can elucidate the phylogeographic origin, transportation routes into nonnative ranges, and release history of nonnative species. Further, genetic tools can help identify source-sink population dynamics and movement pathways across invasion ranges for control and eradication efforts. Collectively, genetic characterization can inform management decisions and help to guide targeted removal efforts (Collins, Vazquez, & Sanders, 2002; Ficetola, Miaud, Pompanon, & Taberlet, 2008; Kolbe et al., 2007; McPhee & Turner, 2009; Stepien & Tumeo, 2006; Vidal, García-Berthou, Tedesco, & García-Marin, 2010).

Accurate and efficient identification and classification at the species level are necessary for invasive species management. For example, accurate species identification can indicate the required habitat types, diet (including prey species), intrinsic ecological constraints, and climatic suitability (Chown et al., 2015; Gotelli & Stanton-Geddes, 2015; Pfeiffer, Johnson, Randklev, Howells, & Williams, 2016; Rissler & Apodaca, 2007). Population expansion capabilities or limitations can be assessed through knowledge of the species life history, population growth rates, and susceptibility to diseases. Further, once the invasive species has been correctly identified, putative range expansions can be predicted using ecological niche models based on both the native and invasive species ranges (Ikeda et al., 2017; Mainali et al., 2015).

Understanding the potential for hybridization of invasive species is critical because diversity can be increased through crossing of divergent groups prior to release or during sustained releases over time of genetically divergent individuals. Hybridization events can lead to increased diversity, fitness, and fecundity in the invasive population (Kolbe et al., 2004, 2007; Vidal et al., 2010). Further, hybrid vigor and environmental selection can result in improved adaptation to the novel environment and increased areas of climatic suitability (Hahn & Rieseberg, 2017; Roman & Darling, 2007). Deleterious mutations can also accumulate through outbreeding depression via negative dominance effects (Oakley, Ågren, & Schemske, 2015).

In this study, we investigated putative origins, potential for hybridization with congeners, and population structure within the invasive Burmese python (*Python bivittatus*) population in the Greater Everglades Ecosystem (GEE) in Florida, USA. This giant constrictor snake has been reproducing in southern Florida since approximately the mid-1980s (Willson, Dorcas, & Snow, 2011). The cryptic nature of these snakes has limited detection and control efforts (Hunter et al., 2015; Reed et al., 2011), and the population has now expanded from Everglades National Park (ENP) into the eastern and western coasts of southern Florida and the Florida

Keys (Dove, Snow, Rochford, & Mazzotti, 2011; Pittman et al., 2014; Snow, Brien, Cherkiss, Wilkins, & Mazzotti, 2007). Pythons are impacting the ecosystem through heavy predation on mesomammals, including imperiled species, resulting in extensive declines of formerly common species (Dorcas et al., 2012; McCleery et al., 2015; Reichert et al., 2017; Sovie, McCleery, Fletcher, & Hart, 2016).

Python bivittatus taxonomy and nomenclature have been uncertain in part due to the sympatric distribution with *P. molurus* in the native range and lack of a designated neotype (Jacobs, Auliya, & Böhme, 2009; Schleip & O'Shea, 2010). The species was first recognized by Kuhl (1820), but was then reclassified as a subspecies, *P. molurus bivittatus*, 100 years later. *Python molurus molurus* was differentiated as the other subspecies in the complex using subocular scales (McDiarmid, Campbell, & Touré, 1999). Most recently, *P. bivittatus* was again recognized as a distinct species with populations of *P. molurus* identified sympatrically (shared range) and possibly even syntopically (shared localities; Jacobs et al., 2009; Reynolds, Niemiller, & Revell, 2014; Schleip & O'Shea, 2010). The integrity of the two species and interbreeding avoidance in wild populations is thought to be maintained through resource partitioning of prey and microhabitat usage (O'Shea, 2007). Viable crosses, however, have been produced in captivity (Townson, 1980). Hybridization of the two species in the invasive range could affect climatic suitability and adaptation potential (as discussed previously) and also subsequent genetic analyses such as environmental DNA detection (Ryan et al., 2018; Wilcox et al., 2013). Here, we follow the most recent classification by Schleip and O'Shea (2010) and consider the Burmese python (*P. bivittatus*) and Indian python (*P. molurus*) as distinct species. To date, the GEE population has been morphologically identified as *Python bivittatus* throughout the invasive range.

A previous report of the invasive GEE population found one haplotype in cytochrome *b* (*Cyt b*) and two in the control region and used 10 cross-species microsatellites developed by Jordan, Goodman, and Donnellan (2002) to conclude that the ENP population was not genetically structured (Collins, Freeman, & Snow, 2008). The sequence data and several locus-specific and average genetic diversity values were not provided and therefore cannot be used for further comparison. Invasive Florida population-specific microsatellite markers for *P. bivittatus* were subsequently isolated ($N = 18$) and combined with six cross-species markers to identify 61% average expected heterozygosity (H_E) and 2–6 alleles per locus (N_A ; 3.7 average N_A ; Hunter & Hart, 2013). In comparison, higher levels of genetic diversity ($N_A = 10.88$) were identified for *P. bivittatus* in the native range using eight microsatellites (Duan et al., 2017).

Our goal was to more thoroughly characterize the *P. bivittatus* populations in Florida to inform research and management strategies. We compared two mitochondrial DNA (mtDNA) genes with population-specific nuclear microsatellite markers to investigate diversity, relatedness, effective population size, population structure, and introduction dynamics of *P. bivittatus* captured in Florida (Hunter & Hart, 2013). We further assessed phylogeographic structure and haplotype relationships and compared them with published

sequences in an effort to assess the genetic origin and species composition of introduced pythons in Florida.

2 | METHODS

2.1 | Sample collection and DNA extraction

The molecular analyses were conducted using tail tissue obtained from *P. bivittatus* samples collected January 2001 to September 2012. Samples originated in southern Florida from Everglades National Park, Collier County (including Big Cypress National Preserve), southeastern Miami-Dade County, and the Florida Keys (Figure 1). Burmese pythons were identified by the presence of a subocular scale just below the eye, which differentiates them from *P. molurus*, which possess supralabial scales that extend from the lip to the bottom of the eye (O'Shea, 2007). All tissues were stored at -20°C . DNA was extracted using QIAGEN DNeasy kits (Valencia, CA) or plate isolation protocols (Whitlock, Hipperson, Mannarelli, & Burke, 2008). DNA was quantified by nanophotometer (Implen, Munchen, Germany) and diluted to $10\text{ ng}/\mu\text{l}$.

2.2 | Microsatellite analysis

2.2.1 | Microsatellite DNA analysis

To address fine-scale genetic diversity and population structure in the invasive population, 18 population-specific microsatellites were developed through next-generation sequencing and incorporated with six cross-species loci (Jordan et al., 2002) into eight multiplexes to reduce laboratory effort (Hunter & Hart, 2013). Of these markers, two loci (*MS16* and *MS22*) did not produce consistent scores and were excluded here. To optimize previously published annealing multiplex (MP) temperatures, *Pmb-U21* was reassigned to MP1 and *MS09* was reassigned to MP9. Annealing temperatures and PCR parameters followed Hunter and Hart (2013), except for an annealing temperature of 57°C in MP4. All PCR products were analyzed on an ABI 3130xl (Applied Biosystems, Foster City, CA). Fragment data were scored using GENEMARKER v. 1.97 (Soft Genetics, State College, PA). The majority of individual genotypes ($N = 389$) included all 22 loci, with a small percentage of samples missing \leq seven loci.

2.2.2 | Microsatellite statistical analysis

MICRO-CHECKER (Van Oosterhout, Hutchinson, Wills, & Shipley, 2004) was used to identify loci with evidence of null alleles. GENECAP (Wilberg & Dreher, 2004) calculated the probability of identity ($P_{(ID)}$), which is the probability that two individuals drawn at random from a population will have the same genotype at the assessed loci (Paetkau & Strobeck, 1994) and sibling probability of identity ($P_{(ID)sib}$), a related, more conservative statistic for calculating $P_{(ID)}$ among siblings (Evetts & Weir, 1998). The program additionally searched for duplicate genotypes.

The program STRUCTURE 2.3.4 (Pritchard, Stephens, & Donnelly, 2000) was used to identify the genetic relationships and population

structure of the southern Florida population. STRUCTURE, a model-based clustering algorithm, infers population structure by probabilistically assigning individuals, without a priori geographic or ancestral knowledge, to a specific number (K) of clusters (presumably populations). In determining the number of clusters, the algorithm attempts to minimize deviations from Hardy-Weinberg equilibrium (HWE).

Simulations were conducted using the correlated allele frequency model and admixture model, which assumes that individuals could have some proportion of membership (q) from each of K clusters. Multiple Markov chains can delineate differences within populations; therefore, 20 parallel chains were analyzed for $K = 1-11$, with a run length of 200,000 Markov chain Monte Carlo repetitions, following a burn-in period of 50,000 iterations. The most probable number of groups, K , was assessed using the mean log likelihood ($\ln P(D)$) and by calculating ΔK , an ad hoc quantity related to the change in posterior probabilities between runs of different K values (Evanno, Regnaut, & Goudet, 2005), in STRUCTURE HARVESTER (Figure 2; Cristescu, Sherwin, Handasyde, Cahill, & Cooper, 2010). Individual assignment success was recorded as the highest likelihood of assignment (q), and the percentage of individuals in a cluster with $q \geq 0.90$ was calculated. GENECLASS was used to detect first-generation migrants born in a population other than the one in which they were sampled without *a priori* population categorization (Piry et al., 2004). We used the Paetkau, Calvert, Stirling, and Strobeck's (1995) simulation algorithm and L_h to assess the likelihood of finding individuals in the population in which they were sampled, which is most appropriate when all potential source populations have not been sampled. Migrant detection was assessed using the critical value (0.01; Paetkau, Slade, Burden, & Estoup, 2004).

The following statistical tests were conducted for the population as a whole and to assess the accuracy of the STRUCTURE-identified groups. The genetic diversity was estimated by the H_E and observed heterozygosity (H_O), information index (I), N_A , average effective number of alleles (E_A), and private alleles (P_A) using GENALEX 6.501 (Table 1, Supporting Information Table S1; Peakall & Smouse, 2006). Departures from the expected genotypic frequencies in HWE were tested using the Markov chain method, and linkage disequilibrium expectations were tested using the randomization method of Raymond and Rousset (1995) for all pairs of loci within collections to test for the presence of admixture in GENEPOP 4.0 (dememorization, 1,000; batches, 100; iterations per batch, 1,000; Raymond & Rousset, 1995). Sequential Bonferroni adjustments (Rice, 1989) were used to determine significance for these tests.

To assess genetic differentiation of the clusters identified by STRUCTURE, GENALEX 6.501 was used to calculate F_{ST} and R_{ST} via analysis of molecular variance (AMOVA) within and among clusters and individuals with 9,999 permutations. The statistical significance of the correlation between genetic and geographic distance matrices, or isolation by distance, was assessed with a Mantel randomization test performed with GENALEX 6.5 with 999 permutations comparing pairwise genetic distance (in meters; Paetkau & Strobeck, 1994). The genetic groupings were assessed by the LDNE software (Waples & Do, 2008) to estimate effective population sizes (N_E) using the linkage




FIGURE 1 Map indicating python sample locations in southern Florida, USA. The nuclear Bayesian clustering assignments are shown in color. The samples yielding only mitochondrial DNA (mtDNA) sequences are in gray. Overlapping sample points have been offset to increase resolution

disequilibrium (LD) method at the three lowest allele frequency levels (0.01, 0.02, and 0.05), with 95% confidence intervals (CI) following the bias-corrected method of Waples (2006). The single point estimate method removes the downward bias associated with the true N_E being greater than the sample size used to estimate it (Waples, 2006).

We used BOTTLENECK 1.2.02 to evaluate heterozygote excess of populations under the sign test, one-tailed Wilcoxon's signed-rank test for mutation-drift equilibrium, and the allele frequency

distribution test (Table 2; Piry, Luikart, & Cornuet, 1999). The Garza-Williamson index and modified index were calculated in Arlequin 3.5 (Excoffier & Lischer, 2010). The Garza-Williamson index is the mean ratio of the number of alleles at a given locus to the range in allele size (M ; Garza & Williamson, 2001). It is assumed that during a bottleneck event, the number of alleles decreases faster than the allelic range. A bottleneck is indicated with a critical value of $M < 0.68$, and no reduction of effective population size is indicated at $M > 0.80$.




FIGURE 2 (a) Results of Bayesian clustering analysis ($K = 2$) using 22 microsatellite loci with 389 python genotypes in STRUCTURE 2.4.3 (cluster 1, gray; cluster 2, black). (b) The proportion of membership for $K = 2$ was supported by the mean log likelihood ($\ln P(D)$); denoted by bars and ΔK (diamonds) versus K

TABLE 1 Summary statistics of the 22 polymorphic microsatellite loci for the *P. bivittatus* invasive population grouped by STRUCTURE clusters. Number of individuals (N), average number of alleles (N_A), effective number of alleles (E_A), information index (I), observed heterozygosity (H_O), expected heterozygosity (H_E), and private alleles (P_A). Individual locus information is provided in Supporting Information Table S1

Locus	N	N_A	E_A	I	H_O	H_E	P_A
Cluster 1	263.59	3.18	2.53	0.96	0.58	0.59	1
Cluster 2	36.91	4.95	3.05	1.24	0.68	0.66	19
Admixed	49.23	4.55	2.63	1.06	0.58	0.61	10
Overall	349.73	5.50	2.63	1.05	0.59	0.60	20

TABLE 2 Microsatellite bottleneck analyses and effective population sizes (N_e) for the three invasive python STRUCTURE clusters. Two phase model (TPM) and stepwise mutation model (SMM)

Group	TPM		SMM		Mode-shift	G-W modified index	N_e
	Sign test	Wilcoxon's test	Sign test	Wilcoxon's test			
Cluster1	0.00002	0.00000	0.00021	0.00000	Shifted mode	0.568	236.1
Cluster2	0.24128	0.14511	0.41772	0.48731	L-shaped	0.766	44.3
Admixed	0.05562	0.01506	0.41838	0.23139	L-shaped	0.834	32.4
Total	0.16001	0.62488	0.03070	0.97692	L-shaped	0.723	315.1

GENALEX 6.501 was used to calculate F_{IS} , which is close to zero when the population is undergoing random mating. Mean relatedness values (r_{xy}) were computed for all pairwise relationships via ML-RELATE (Kalinowski, Wagner, & Taper, 2006). Relatedness and individual inbreeding coefficients (F_x) were estimated for all pairwise

relationships via COANCESTRY (Wang, 2011). From mean relatedness values, latent coancestry (Θ_{xy}) was calculated following Lynch and Ritland (1999), where $r_{xy} = 2\Theta_{xy}$. To better understand the capacity to increase genetic diversity in the population, relatedness was also estimated specifically for eight collected hatchlings from a

TABLE 3 Summary statistics for the invasive Florida python concatenated cytochrome *b* and cytochrome oxidase 1 sequences (1,397 bps; $N = 399$) for the *P. bivittatus* (Pb) and *P. molurus* (Pm) invasive haplotypes (H01 to H06). Within the sample groups, N represents the number of sequences; S , number of polymorphic sites; H , number of haplotypes; h , haplotype diversity; π , nucleotide diversity; k , average number of nucleotide differences; TD, Tajima's D , *significant ($p \leq 0.05$)

Sample groups	N	S	H	h	π	k	TD	Pb-FL-H01	Pb-FL-H02	Pb-FL-H03	Pb-FL-H04	Pm-FL-H05	Pb-FL-H06
Cluster 1	228	1	2	0.009	0.000	0.009	-0.938	227	0	0	1	0	0
Cluster 2	27	64	4	0.53	0.014	19.67	0.638	18	2	0	0	5	2
Admixed	38	61	3	0.104	0.002	3.260	-2.810*	36	0	1	0	1	0
Clusters 1, 2, admixed	293	65	6	0.080	0.002	2.535	-2.247*	281	2	1	1	6	2
Total sequenced	399	65	6	0.097	0.002	3.405	-1.918*	379	5	1	1	11	2

Note. Sequences are grouped by STRUCTURE clusters obtained through nuclear microsatellite genotypes ($N = 293$) and sequences ($N = 399$). Note that 106 samples did not contain data for both marker types.

single nest (P0003, P0006, P0008, P0009, P0016, P0020, P0021, P0024) to test for multiple paternity. Where not specified previously, the default parameters were used in the previous analyses.

2.3 | Mitochondrial analysis

2.3.1 | Mitochondrial DNA analysis

Mitochondrial DNA variation was assayed at two protein-coding loci: Cyt *b* (Rawlings, 2001) and cytochrome *c* oxidase I (CO1; Folmer, Black, Hoeh, Lutz, & Vrijenhoek, 1994). The PCR conditions were as follows: 10 ng DNA, 1× PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 0.001% gelatin; Sigma-Aldrich, Inc., St. Louis, MO), 0.8 mM dNTP, 3 mM MgCl₂, 0.24 μM of each primer, 0.04 units of Sigma Jump Start Taq DNA polymerase. PCR cycling profile: 5 min at 94°C; then 35 cycles of 1 min at 94°C, 1 min at 55°C, and 1 min at 72°C; then 10 min at 72°C. Amplified products were purified using ExoSap-IT (Affymetrix, Santa Clara, CA) for PCR cleanup. DNA sequencing was accomplished with the BigDye terminator protocol (Applied Biosystems, Foster City, CA).

2.3.2 | Mitochondrial statistical analysis

Sequences were trimmed to those published in GenBank, checked for quality scores, and aligned in GENEIOUS 5.4.6 (Drummond et al., 2011). Representatives from each haplotype and any ambiguous sequences were sequenced in both directions to ensure the accuracy of nucleotide designations. We calculated summary statistics for the mtDNA by assessing nucleotide diversity (π), haplotype diversity (h), sequence diversity (k), and the standard neutrality test, Tajima's D , using DnaSP v5.0 (Table 3, Supporting Information Tables S2 and S3; Librado & Rozas, 2009). To assess the mtDNA and nuclear data collectively, the individuals containing both concatenated mtDNA sequences and microsatellite genotypes were assessed together ($N = 293$), while the concatenated mtDNA matrix only included those individuals sequenced at both loci ($n = 399$; Table 3). Using the microsatellite-defined STRUCTURE populations described below (cluster 1, cluster 2, or admixed), we calculated pairwise Φ_{ST} (10,000 permutations; p value <0.05 significant; Table 4) and exact tests of population differentiation (100,000 Markov chain steps; 10,000 dememorization steps; p value <0.05 significant) using Arlequin 3.5 (Excoffier & Lischer, 2010).

Invasive samples were compared to GenBank and BOLD published sequences with similar length and quality (see Table 5 for sequence name abbreviations, references, and submission details). Complete mitochondrial DNA genomes were recently published for *P. molurus* (Dubey, Meganathan, & Haque, 2012) and *P. bivittatus* (Liu, Zhang, & Cao, 2013), accompanied by direct submissions of mtDNA sequences in GenBank. Slowinski and Lawson (2002) previously addressed phylogenies of 42 snake species using Cyt *b* and *C-mos* genes; however, the *P. molurus* sequence did not include voucher or origin of sample information. The CO1 sequences published in BOLD contained two *P. molurus* samples with voucher specimens. We selected the longer sequence originating from a sample in a forested area in Maharashtra state in western India to avoid trimming our

TABLE 4 Concatenated sequences grouped by STRUCTURE clusters. Pairwise Φ_{ST} values below the diagonal and exact tests of population differentiation above the diagonal

	Cluster 1	Cluster 2	Admixed
Cluster 1	–	0.000*	0.054
Cluster 2	0.514*	–	0.002*
Admixed	0.074*	0.115*	–

Note. An asterisk (*) denotes significance at $p < 0.05$.

alignment (Sequence ID: ISDB081-13.COI-5P). The second sequence (ISDB016-11.COI-5P) was from a snake housed in a zoo in the same state and differed by four base pairs (bps) from ISDB081-13. *Python regius* (Dong & Kumazawa, 2005) was included as the basal member of the python genus (Reynolds et al., 2014).

Pairwise genetic distances were calculated using Tajima-Nei between the invasive population samples and *P. molurus* sequences published in GenBank and the Barcode of Life Data (BOLD) system (www.barcodinglife.org; Ratnasingham & Hebert, 2007; Tajima & Nei, 1984) using MEGA7 (Kumar, Stecher, & Tamura, 2016; Supporting Information Tables S4 and S5). Polymorphic sites and the corresponding diagnostic sites were determined by the alignment of published haplotypes and haplotypes identified in this study (Supporting Information Tables S6 and S7). Genetic differentiation was tested with an analysis of molecular variance (AMOVA) in Arlequin 3.5, using models of DNA sequence evolution selected by the Akaike information criterion (AIC) and Bayesian information criterion (BIC) in MEGA7 (Darriba, Taboada, Doallo, & Posada, 2012; Excoffier & Lischer, 2010; Guindon & Gascuel, 2003). The T92 model (Tamura, 1992) was selected for CO1 sequences. The TN93 model (Tamura & Nei, 1993) was selected for Cyt *b* and concatenated (CO1 [N = 598] and Cyt *b* [N = 799]) analyses.

TABLE 5 References and GenBank accession number or BOLD sequence ID (www.boldsystems.org) for the published haplotypes used in the study

Name	Acc No/Seq ID	Reference	Direct submission	Country
Pb-Ctb-A	KF010492	Liu et al. (2013)	Yes	China
Pb-Ctb-B	KF293729	Liu et al. (2013)	Yes	China
Pm-Ctb-A	AY099983	Slowinski and Lawson (2002)		
Pm-Ctb-B	GQ225654	Dubey et al. (2009)	Yes	India
Pb-CO1-A	KF010492	Liu et al. (2013)	Yes	China
Pb-CO1-B	KF293729	Liu et al. (2013)	Yes	China
Pb-CO1-C	JX401103	You et al. (2013)		China
Pm-CO1-B	AB920233	Supikamolteni and Srikulnath (2014)	Yes	Thailand
Pm-CO1-A	ISDB081-13	www.boldsystems.org	Yes	India
<i>Python regius</i>	AB177878	Dong and Kumazawa (2005)		

Note. The prefix indicates the most similar species (*P. bivittatus*, Pb; *P. molurus*, Pm), and the gene is identified as either cytochrome *b* (Ctb) or cytochrome oxidase 1 (CO1). Direct submission sequences deposited in the databases are not associated with a publication. Country of origin is indicated for the sample or authors.

We created haplotype networks using PopART (Leigh & Bryant, 2015) to assess the geographic distribution of mtDNA diversity and compare relationships between our samples and those previously published (Figure 3). Default minimum spanning network settings were used to generate a haplotype network with pie charts representative of the proportion of samples grouped by STRUCTURE clusters. The number of base pair discrepancies between haplotypes is provided in parenthesis.

3 | RESULTS

To summarize the results, 11 mtDNA haplotypes (GenBank Accession Number: MH357840-50) were identified with high haplotype diversity in the invasive python samples corresponding to both *P. bivittatus* and *P. molurus*. Nuclear microsatellite markers detected lower diversity and N_E as compared to native range samples, likely related to founding and bottleneck effects. Bayesian clustering analyses identified two distinct nuclear groups and an admixed group with no correlation with geographic distribution. The *P. molurus* haplotypes were more predominantly classified in cluster 2.

3.1 | Microsatellite DNA analysis

Only the MS13 locus in cluster 1 indicated the evidence of null alleles due to homozygote excess (>0.05), but there was no evidence of stuttering, large allele dropout, or linkage disequilibrium. The loci produced an unbiased $P_{(ID)}$ estimate of 5.63 E-15 and a $P_{(ID)}$ sib estimate of 2.99 E-07, indicating that unique individuals can be confidently identified across the region. The Bayesian STRUCTURE Ln P(D) estimates indicated similar values and generally plateaued at $K = 4$ clusters, while the STRUCTURE HARVESTER analysis according to



(a) *Cytochrome-b*(b) *Cytochrome oxidase subunit 1*

FIGURE 3 *Python bivittatus* (Pb) and *P. molurus* (Pm) (a) cytochrome *b* and (b) cytochrome oxidase 1 haplotype networks. Bayesian cluster assignment for invasive haplotypes (H01–H06) is denoted by gray shading. Published sequences' (white circles) references are given in Table 5 and denoted by the species prefix. The area of each pie chart represents the number of haplotypes. Base pair discrepancies are given by the hash marks

Evanno et al. (2005) strongly supported $K = 2$ clusters (Figure 2). The $K = 4$ and next highest ΔK ($K = 6$) were also investigated; however, the majority of the genotypes were “roughly symmetrically” assigned across the four or six populations, respectively, indicating that these values of K are not identifying real population structure (Supporting Information Figure S1; Pritchard et al., 2000). Therefore, as recommended, $K = 2$ was selected and groups were assigned as follows: cluster 1 ($N = 292$), cluster 2 ($N = 42$), and a third, admixed group ($N = 55$), containing $q \leq 90\%$ assignment to the two clusters (Figure 2; Evanno et al., 2005; Pritchard et al., 2000).

Across the 389 samples, Hardy–Weinberg disequilibrium was found for *Pmb-N14* and *Pmb-Z26* ($p \leq 0.002$). After the sequential Bonferroni adjustments, linkage disequilibrium was found for 39 of 231 (16.9%) comparisons, likely due to population substructure tested below. Separate analyses of the three groups identified in STRUCTURE resulted

in HWE for all loci and linkage equilibrium for cluster 1 and admixed. However, linkage disequilibrium was found in cluster 2 for two pairs of loci (*Pmb-S19* and *Pmb-R18*, *Pmb-N14* & *Pmb-K11*). This deviation may be due to inbreeding or cryptic subpopulation structure (i.e., Wahlund effect). GENECLASS detected 16 samples with a probability < 0.01 . These samples were all members of cluster 2 defined by STRUCTURE.

F_{ST} values among the three STRUCTURE-defined clusters were low, but significant ($p \leq 0.017$): cluster 1 versus 2 (0.029), cluster 1 versus admixed (0.004), and cluster 2 versus admixed (0.012). R_{ST} values were not significant, likely owing to the minimal time for mutations to occur. Low levels of nuclear diversity were found for all samples assessed together and grouped by clusters (Table 1). The AMOVA identified moderate variation both among the three STRUCTURE clusters and within individuals (19.12% and 1.49%, respectively). In cluster 2, 19 private alleles were found in 33 samples, while the admixed group

contained 10 private alleles in 14 samples distributed across the q -values. Cluster 1 had a single private allele in five samples. Effective population sizes using the linkage disequilibrium method were similar for the three allele frequencies tested; therefore, the 0.01 frequencies are reported (Table 2). The Mantel test indicated no significant correlations between genetic and geographic distances ($p = 0.27$).

Assessing the 389 samples together, the stepwise mutational model (SMM) of the sign test was significant ($p = 0.03$). However, a normal "L"-shaped allele distribution curve was obtained, indicating a larger proportion of alleles in the low-frequency allele classes (Table 2). All BOTTLENECK tests for cluster 1 were significant, and a bottleneck was also indicated with a Garza–Williamson modified index value below the critical value ($M = 0.568$). Cluster 2 indicated a reduction of effective population size ($M = 0.766$), but was not below the critical value threshold. The admixed group was significant under the Wilcoxon's TPM ($p = 0.015$). Nonsignificant bottleneck test values may have been due to smaller sample sizes.

The inbreeding coefficient, F_{IS} , was 0.194 ($p = 0.000$) over the three groups, which indicates inbreeding and/or a founding effect on the population. Overall, the average number of alleles was 5.50 and H_E was 0.60 (Table 1). Relatedness levels ($r_{xy} = 0.091$) were between first ($r_{xy} = 0.125$) and second ($r_{xy} = 0.0625$) cousins on average, and inbreeding coefficients were also indicative of a cousin relationship. Across the STRUCTURE cluster and all estimators, cluster 1 had the highest level of relatedness, cluster 2 had moderate levels, and admixed had the lowest level. Simulations using population allele frequencies estimated that 20% of the population was related, while observed values estimated that 24% of the samples were related. Analysis of the small number of collected hatchlings indicated that they were likely related at either the half-sibling or full-sibling levels.

3.2 | Mitochondrial DNA analysis

Cytochrome *b* produced six novel haplotypes in 419 sequences across 799 bps with relatively high genetic distance (range 0.13%–4.30%) and number of polymorphic sites ($S = 1$ –36 bps; Supporting Information Tables S4 and S6). Cytochrome oxidase 1 produced five haplotypes in 413 sequences across 585 bps with high genetic distances (0.30%–5.40%) and numbers of polymorphic sites ($S = 1$ –31 bps; Supporting Information Tables S5 and S7). The invasive haplotypes split into two strongly divided groups at COI. The H01 sequences most closely associated with the published *P. bivittatus* mtDNA genomes (>99.14%; Liu et al., 2013) and H05 matching a published *P. molurus* sequence (Supikamolnseni & Srikulnath, 2014, direct NCBI submission). The current published *P. bivittatus* and *P. molurus* sequences were $\geq 94.8\%$ similar (Supporting Information Table S5).

The concatenated sequences produced six novel haplotypes in 399 snakes across 1,397 bps. The majority of samples were found to be a single haplotype (Pb-FL-H01; $N = 379$), with the five other haplotypes represented in lower proportions (Table 3). The Pm-FL-H05 haplotype was found in 11 samples associated with the *P. molurus* mitotype. No phylogeographic pattern was found in accordance with collection sites in southern Florida (Figure 1).

3.3 | Mitochondrial DNA haplotypes partitioned by STRUCTURE clusters

The STRUCTURE cluster 1 contained only the Pb-FL-H01 haplotype in all but one sample, while cluster 2 and the admixed groups contained a mixture of haplotypes (Table 3). The majority of the haplotypes were each assigned to a single cluster (Table 3). Interestingly, although not selected as the correct grouping, many of the samples with *P. molurus* haplotypes were assigned to a single cluster in the $K = 4$ plot (green; Supporting Information Figure S1). Diagnostic sites differentiating the two *Python* species were identified for the two loci: Cyt *b* ($N = 27$) and CO1 ($N = 24$; Supporting Information Tables S6 and S7). The highest differences between concatenated sequences, as measured by Φ_{ST} and exact test values, were between the cluster 1 and 2 STRUCTURE groups ($p < 0.05$; Table 4). The mtDNA AMOVA values within and among the three STRUCTURE groups resulted in relatively high variation (69.96% and 30.04%, respectively). The AMOVA identified variation levels of 48.60% and 51.40% within and among clusters 1 and 2, respectively.

3.4 | Comparison with published sequences

The dominant Cyt *b* haplotype that we found in the invasive range matched all but one nucleotide to one of the published *P. bivittatus* mitochondrial genome sequences (Pb-Ctb-A; Supporting Information Table S4; Liu et al., 2013). The next highest frequency haplotype differed by $\geq 4.13\%$ from Liu et al. (2013), but was only $\leq 0.76\%$ different from the published *P. molurus* sequences (Dubey et al., 2012; Slowinski & Lawson, 2002). A similar pattern was found for the CO1 haplotypes with a dominant haplotype most closely resembling *P. bivittatus*. The second most dominant haplotype matched *P. molurus* or differed by 1.38% (Supporting Information Table S5). No subocular differentiation was found in the available photographs of the snakes containing *P. molurus* haplotypes. There was some disparity between the published sequences and associated species labels, which may relate to the lack of consensus in the nomenclature.

4 | DISCUSSION

The invasive Burmese python population in Florida appears to be derived from multiple genetic sources with strongly divergent mitotypes corresponding to species-level differentiation. The Cyt *b* genetic distance (4.3%) was larger than the distance found in the most recent taxonomic assessment that separated *P. bivittatus* and *P. molurus* into species (2.9%; Reynolds et al., 2014). The CO1 genetic distance (5.4%) was also greater than the distances for the two species published in BOLD (4.1%). In the literature, CO1 nucleotide diversity values lower than 4.1% have been used as a minimum threshold to distinguish intraspecific variation from interspecific divergence (Gomes, Pessali, Sales, Pompeu, & Carvalho, 2015; Ratnasingham & Hebert, 2013). In contrast to the strong

mitochondrial differentiation signals, minimal divergence was detected between the three Bayesian clusters at nuclear microsatellites ($F_{ST} \leq 0.029$).

In vertebrates, cytonuclear discordance is indicated by conflicting signals between mtDNA and nuclear genetic diversity. The nonrecombining mitochondrial genome sequence remains as a distinct cytotypic in an admixed clade until, over time, only either the dominant (parent) or introgressed cytotypic is retained (see Seehausen, 2004). On the contrary, admixed nuclear genotypes will continue to recombine with the dominant alleles until introgressed. During the intermediate phase, nuclear genome hybrids and both cytotypes can be detected. Cytonuclear discordance can be caused by hybridization, incomplete lineage sorting, direct balancing selection, indirect selection, or pseudogenes (Grobler, Jones, Johnson, Neves, & Hallerman, 2011; Thielsch, Knell, Mohammadyari, Petrusek, & Schwenk, 2017). In our data, incomplete lineage sorting is unlikely due to the high sequence divergence among the samples and rapid progression of lineage sorting in mitochondrial loci (Funk & Omland, 2003). Similarly, no evidence for indirect selection or pseudogenes causing cytonuclear discordance was observed here, given the haplotype sequences and divergence levels. It is possible that we detected direct balancing selection of rare ancestral mitochondrial lineages that favor specific environmental conditions; however, the role that mtDNA plays in natural selection is not fully understood (Funk & Omland, 2003). Most likely, the cytonuclear divergence we detected is the result of hybridization between snakes contributing mitochondrial genome sequences from both species.

Past hybridization of *P. molurus* and *P. bivittatus* may have led to the identified cytonuclear discordance in the invasive population. The nuclear F_{ST} values ≤ 0.029 suggest significant introgression of the nuclear genomes and a population-level, as opposed to a species-level, divergence. However, residual *P. molurus* nuclear genomic material may be contributing to the cluster 2 and admixed genotypes. For instance, in cluster 2, 16 first-generation migrants were detected, and 19 private alleles were found in 79% of the individuals (with 10 individuals identified in both analyses). In natural systems, the sigmoid shape of the STRUCTURE plot (Figure 2) can be interpreted as hybridization followed by selection against the hybrid alleles or segregating variation. However, as this population was likely released from the pet trade, the admixed group may indicate separate introductions of two relatively similar gene pools that are now interbreeding over a few generations.

The taxonomic uncertainty regarding species boundaries in the genus *Python* complicates our understanding of the precise mechanism responsible for the cytonuclear discordance. Introgression of the diverse lineages could occur through interbreeding (a) in the native range through sympatric associations or secondary contact (Seehausen, 2004), (b) during secondary contact in captivity, or (c) after release into the invasive range, possibly in part due to unidirectional hybridization or unbalanced sex ratios in hybrid generations (Firmat, Alibert, Losseau, Baroiller, & Schliwen, 2013). The support for scenario 3 is limited, given the low diversification in the nuclear genome suggestive of admixture over numerous generations. More extensive native range phylogeographic sampling of mtDNA and nuclear DNA loci is necessary to confirm whether the cytonuclear

discordance observed in Florida is present in native populations or occurred after capture. Limited introgression of *P. molurus* followed by backcrossing to *P. bivittatus* may have occurred in the large, widely distributed commercial trade populations. Alternatively, intraspecific genetic divergence in the *P. bivittatus* native range may have contributed to the three nuclear groups found in China ($F_{ST} = 0.11$ overall), although assessment of mitochondrial DNA is needed to determine whether divergent mitotypes are also present (Duan et al., 2017).

Field observations in the native range indicate that the two species utilize distinct habitats with some overlapping ranges. *Python bivittatus* prefers riverine forests and flooded grasslands, while *P. molurus* occupies dry, sandy, and woodland areas (Schleip & O'Shea, 2010). Hybridization of the two species could allow for improved acclimatization and adaptability to abiotic stressors or climate change and result in broader or more rapid distributions of the invasive population (Hoffmann & Sgrò, 2011; Mazzotti et al., 2011; Rodda, Jarnevich, & Reed, 2009). Currently, pythons occupy both wetlands in Everglades National Park and drier, sandy pinelands with interspersed wetlands in western Collier County. However, evidence of a panmictic population was found with no temporal or phylogeographic pattern across the sampled range. This is not surprising, given that invasive pythons are known to disperse long distances (Hart et al., 2015).

A bottleneck and/or founding event was indicated 0.2–4 N_E generations ago with a $\geq 90\%$ reduction in population size (Williamson-Natesan, 2005). Given that Burmese pythons require two to five years to reach sexual maturity (Willson, Snow, Reed, & Dorcas, 2014), the population would have undergone approximately four to 10 generations after being founded in the mid-1980s (Willson et al., 2011). The detection of a bottleneck of less than four generations ago may indicate a secondary bottleneck due to novel environmental conditions such as cold-induced mortality (Mazzotti et al., 2011). Alternatively, a lag in the generation times or in the population growth rates may have occurred shortly after the population's founding, possibly due to low propagule pressure (Fujisaki et al., 2010). Reproduction was not documented until the first wild hatchlings were found in the mid-1990s (Meshaka, Loftus, & Steiner, 2000), although detection of pythons has remained low until recently (Hunter et al., 2015; Reed et al., 2011). Parthenogenesis has also been identified in the species, which may allow for population expansion even at low densities and would contribute to reduced genetic diversity (Groot, Bruins, & Breeuwer, 2003).

In comparison with our findings, on average, native range Burmese pythons had nearly twice the number of alleles and higher average heterozygosities, with the exception of the Yunnan population, which had similar allelic values ($N_A = 5$; Duan et al., 2017). In the invasive population, effective population sizes were relatively low, supporting the hypothesis that the population was established by a small number of founders and/or closely related individuals (Willson et al., 2011). Monitoring of the effective population size could help to identify changes in the census size as genetic mutations occur and accumulate, especially in order to assess effective control efforts (Hauser, Adcock, Smith, Bernal Ramàrez, & Carvalho, 2002; Hui & Burt, 2015). More accurate effective population size estimates with

lower variance can be calculated with genetic data collected over multiple generations (Hui & Burt, 2015).

While the genetic diversity in the invasive Burmese python population is lower than that found in the native range, it is likely to increase in the large, rapidly growing invasive population, especially if additional animals are released. Multiple paternity was identified in the invasive population which could also contribute to accelerated increases in diversity. Of note, the genetic confirmation of multiple breeding events by different sires lends support to the Judas control technique in which radio-tagged snakes are used to reveal the location of conspecifics during breeding (Smith et al., 2016). Over time, as the population expands, some genotypes may become isolated or fixed, adapting to certain habitats and creating more population structure.

Recently, eDNA has become an important tool to estimate occurrence and detection probabilities and track the invasion front of the Burmese python populations (Hunter et al., 2015; Piaggio et al., 2013). Highly divergent mtDNA sequences could lead to mispriming of eDNA primers or probes, resulting in false negatives, along with potentially lower detection and occurrence estimates (Wilcox, Carim, McKelvey, Young, & Schwartz, 2015; Wilcox et al., 2013). Deep sampling is necessary to detect intraspecific variation found at low frequencies in the population.

The limited number of well-documented, high-quality published sequences hinders our ability to investigate *P. bivittatus* and *P. molurus* species boundaries. Morphological voucher specimens and broader phylogeographic sampling throughout the native range, including sympatric areas, could improve taxonomic uncertainty. Further, genomic-level assessment and transcriptomic studies could address fine-scale population structure and the Burmese python's adaptation to the novel environment (Castoe et al., 2011; Rodda et al., 2009; Wall et al., 2011). Findings from these endeavors could facilitate management in a variety of ways, including the development of effective monitoring tools (e.g., eDNA assays) and more accurate range expansion predictions.

ACKNOWLEDGMENTS

We would like to thank Gaia Meigs-Friend and Theresa Floyd (USGS) for laboratory support, Michael Cherkiss (USGS) for field assistance, Tylan Dean (National Park Service, Everglades National Park), Paul Andreadis (Denison University) and Ian Bartoszek (Conservancy of SW Florida) for sample coordination, Jason Ferrante and Breanna Caton (USGS) for assistance with data analysis, and Michelle Giles. The protocol was approved by the US Geological Survey, Wetland and Aquatic Research Center Institutional Animal Care and Use Committee (IACUC; Permit Number: USGS/SESC 2013-04). Additionally, all samples were collected under the National Park Service (NPS; Everglades) Permit Number: EVER-2007-SCI-001 and EVER-2009-SCI-001. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US government.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTIONS

MEH designed the research, performed the research, analyzed the data, and wrote the paper. KMH and RWS designed the research, collected samples, and wrote the paper. BJS collected samples and wrote the paper. MCD performed the research, analyzed the data, and wrote the paper. NAJ and JSSB analyzed the data and wrote the paper.

DATA ACCESSIBILITY

The data presented in this manuscript are available from <https://doi.org/10.5066/f7hh6hkj>.

ORCID

Margaret E. Hunter  <http://orcid.org/0000-0002-4760-9302>

Nathan A. Johnson  <http://orcid.org/0000-0001-5167-1988>

Kristen M. Hart  <http://orcid.org/0000-0002-5257-7974>

REFERENCES

- Bock, D. G., Caseys, C., Cousens, R. D., Hahn, M. A., Heredia, S. M., Hübner, S., ... Rieseberg, L. H. (2015). What we still don't know about invasion genetics. *Molecular Ecology*, 24, 2277-2297. <https://doi.org/10.1111/mec.13032>
- Castoe, T. A., Fox, S. E., Jason de Koning, A., Poole, A. W., Daza, J. M., Smith, E. N., ... Pollock, D. D. (2011). A multi-organ transcriptome resource for the Burmese Python (*Python molurus bivittatus*). *BMC Research Notes*, 4, 310. <https://doi.org/10.1186/1756-0500-4-310>
- Chown, S. L., Hodgins, K. A., Griffin, P. C., Oakeshott, J. G., Byrne, M., & Hoffmann, A. A. (2015). Biological invasions, climate change and genomics. *Evolutionary Applications*, 8, 23-46. <https://doi.org/10.1111/eva.12234>
- Collins, T., Freeman, B., & Snow, S. (2008). Final Report: Genetic characterization of populations of the nonindigenous Burmese python in Everglades National Park. In *Final report for the South Florida Water Management District* (p. 30). Miami, FL: Department of Biological Sciences, Florida International University.
- Collins, M. D., Vazquez, D. P., & Sanders, N. J. (2002). Species-area curves, homogenization and the loss of global diversity. *Evolutionary Ecology Research*, 4, 457-464.
- Cristescu, R., Sherwin, W., Handasyde, K., Cahill, V., & Cooper, D. (2010). Detecting bottlenecks using BOTTLENECK 1.2.02 in wild populations: The importance of the microsatellite structure. *Conservation Genetics*, 11, 1043-1049. <https://doi.org/10.1007/s10592-009-9949-2>
- Darling, J. A. (2015). Genetic studies of aquatic biological invasions: Closing the gap between research and management. *Biological Invasions*, 17, 951-971. <https://doi.org/10.1007/s10530-014-0726-x>
- Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods*, 9, 772. <https://doi.org/10.1038/nmeth.2109>
- Dong, S., & Kumazawa, Y. (2005). Complete mitochondrial DNA sequences of six snakes: Phylogenetic relationships and molecular evolution of genomic features. *Journal of Molecular Evolution*, 61, 12-22. <https://doi.org/10.1007/s00239-004-0190-9>
- Dorcas, M. E., Willson, J. D., Reed, R. N., Snow, R. W., Rochford, M. R., Miller, M. A., ... Hart, K. M. (2012). Severe mammal declines coincide with proliferation of invasive Burmese pythons in Everglades National Park. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 1035-1040. <https://doi.org/10.1073/pnas.1116000109>

- States of America*, 109, 2418–2422. <https://doi.org/10.1073/pnas.1115226109>
- Dove, C. J., Snow, R. W., Rochford, M. R., & Mazzotti, F. J. (2011). Birds consumed by the invasive Burmese python (*Python molurus bivittatus*) in Everglades National Park, Florida, USA. *Wilson Journal of Ornithology*, 123, 126–131. <https://doi.org/10.1676/10-092.1>
- Drummond, A. J., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Heled, J., ... Wilson, A. (2011). *Genious v5.4*. Retrieved from <http://www.genious.com>
- Duan, Y., Wang, Y., Bai, S., Tian, X., Rong, K., & Ma, J. (2017). Genetic diversity and population genetic structure of *Python bivittatus* in China. *Journal of Forestry Research*, 28, 621–628. <https://doi.org/10.1007/s11676-016-0308-0>
- Dubey, B., Meganathan, P. R., & Haque, I. (2009). *Molecular phylogeny of Natricinae with special reference to genus Xenochrophis*. NCBI Reference Sequence: GQ225654. Kolkata, India: National DNA Analysis Centre, Central Forensic Science Laboratory.
- Dubey, B., Meganathan, P. R., & Haque, I. (2012). Complete mitochondrial genome sequence from an endangered Indian snake, *Python molurus molurus* (Serpentes, Pythonidae). *Molecular Biology Reports*, 39, 7403–7412. <https://doi.org/10.1007/s11033-012-1572-5>
- Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology*, 14, 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Evtett, I. W., & Weir, B. S. (1998). *Interpreting DNA evidence: Statistical genetics for forensic scientists*. Sunderland, MA: Sinauer Associates Inc.
- Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 10, 564–567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using environmental DNA from water samples. *Biology Letters*, 4, 423–425. <https://doi.org/10.1098/rsbl.2008.0118>
- Firmat, C., Alibert, P., Losseau, M., Baroiller, J.-F., & Schlieven, U. K. (2013). Successive invasion-mediated interspecific hybridizations and population structure in the endangered cichlid *oreochromis mossambicus*. *PLoS ONE*, 8, e63880. <https://doi.org/10.1371/journal.pone.0063880>
- Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294–299.
- Fujisaki, I., Hart, K. M., Mazzotti, F. J., Rice, K. G., Snow, S., & Rochford, M. (2010). Risk assessment of potential invasiveness of exotic reptiles imported to South Florida. *Biological Invasions*, 12, 2585. <https://doi.org/10.1007/s10530-009-9667-1>
- Funk, D. J., & Omland, K. E. (2003). Species-level paraphyly and polyphyly: Frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics*, 34, 397–423. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132421>
- Garza, J. C., & Williamson, E. G. (2001). Detection of reduction in population size using data from microsatellite loci. *Molecular Ecology*, 10, 305–318. <https://doi.org/10.1046/j.1365-294x.2001.01190.x>
- Gomes, L. C., Pessali, T. C., Sales, N. G., Pompeu, P. S., & Carvalho, D. C. (2015). Integrative taxonomy detects cryptic and overlooked fish species in a neotropical river basin. *Genetica*, 143, 581–588. <https://doi.org/10.1007/s10709-015-9856-z>
- Gotelli, N. J., & Stanton-Geddes, J. (2015). Climate change, genetic markers and species distribution modelling. *Journal of Biogeography*, 42, 1577–1585. <https://doi.org/10.1111/jbi.12562>
- Grobler, J. P., Jones, J. W., Johnson, N. A., Neves, R. J., & Hallerman, E. M. (2011). Homogeneity at nuclear microsatellite loci masks mitochondrial haplotype diversity in the endangered Fanshell Pearlymussel (*Cyprogenia stegaria*). *Journal of Heredity*, 102, 196–206. <https://doi.org/10.1093/jhered/esq120>
- Groot, T. V., Bruins, E., & Breeuwer, J. A. (2003). Molecular genetic evidence for parthenogenesis in the Burmese python, *Python molurus bivittatus*. *Heredity*, 90, 130–135. <https://doi.org/10.1038/sj.hdy.6800210>
- Guindon, S., & Gascuel, O. (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52, 696–704. <https://doi.org/10.1080/10635150390235520>
- Hahn, M. A., & Rieseberg, L. H. (2017). Genetic admixture and heterosis may enhance the invasiveness of common ragweed. *Evolutionary Applications*, 10, 241–250. <https://doi.org/10.1111/eva.12445>
- Hart, K. M., Cherkiss, M. S., Smith, B. J., Mazzotti, F. J., Fujisaki, I., Snow, R. W., & Dorcas, M. E. (2015). Home range, habitat use, and movement patterns of non-native Burmese pythons in Everglades National Park, Florida, USA. *Animal Biotelemetry*, 3, 8. <https://doi.org/10.1186/s40317-015-0022-2>
- Hauser, L., Adcock, G. J., Smith, P. J., Bernal Ramarez, J. H., & Carvalho, G. R. (2002). Loss of microsatellite diversity and low effective population size in an overexploited population of New Zealand snapper (*Pagrus auratus*). *Proceedings of the National Academy of Sciences of the United States of America*, 99, 11742–11747. <https://doi.org/10.1073/pnas.172242899>
- Hoffmann, A. A., & Sgro, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, 470, 479–485. <https://doi.org/10.1038/nature09670>
- Hui, T.-Y. J., & Burt, A. (2015). Estimating effective population size from temporally spaced samples with a novel, efficient Maximum-Likelihood Algorithm. *Genetics*, 200, 285–293. <https://doi.org/10.1534/genetics.115.174904>
- Hunter, M. E., & Hart, K. M. (2013). Rapid microsatellite marker development using next generation pyrosequencing to inform invasive Burmese python (*Python molurus bivittatus*) management. *International Journal of Molecular Sciences*, 14, 4793–4804. <https://doi.org/10.3390/ijms14034793>
- Hunter, M. E., Oyler-McCance, S. J., Dorazio, R. M., Fike, J. A., Smith, B. J., Hunter, C. T., ... Hart, K. M. (2015). Environmental DNA (eDNA) sampling improves occurrence and detection estimates of invasive Burmese pythons. *PLoS ONE*, 10, e0121655. <https://doi.org/10.1371/journal.pone.0121655>
- Ikeda, D. H., Max, T. L., Allan, G. J., Lau, M. K., Shuster, S. M., & Whitham, T. G. (2017). Genetically informed ecological niche models improve climate change predictions. *Global Change Biology*, 23, 164–176. <https://doi.org/10.1111/gcb.13470>
- Jacobs, H. J., Auliya, M., & Böhme, W. (2009). Zur Taxonomie des Dunklen Tigerpythons, *Python molurus bivittatus* Kuhl, 1820, speziell der Population von Sulawesi. *Sauria*, 31, 5–16.
- Jordan, P. W., Goodman, A. E., & Donnellan, S. (2002). Microsatellite primers for Australian and New Guinean pythons isolated with an efficient marker development method for related species. *Molecular Ecology Notes*, 2, 78–82. <https://doi.org/10.1046/j.1471-8286.2002.00135.x>
- Kalinowski, S. T., Wagner, A. P., & Taper, M. L. (2006). ML-Relate: A computer program for maximum likelihood estimation of relatedness and relationship. *Molecular Ecology Notes*, 6, 576–579. <https://doi.org/10.1111/j.1471-8286.2006.01256.x>
- Kolbe, J. J., Glor, R. E., Rodriguez Schettino, L., Lara, A. C., Larson, A., & Losos, J. B. (2004). Genetic variation increases during biological invasion by a Cuban lizard. *Nature*, 431, 177–181. <https://doi.org/10.1038/nature02807>
- Kolbe, J. J., Glor, R. E., Schettino, L. R., Lara, A. C., Larson, A., & Losos, J. B. (2007). Multiple sources, admixture, and genetic variation in introduced Anolis lizard populations. *Conservation Biology*, 21, 1612–1625. <https://doi.org/10.1111/j.1523-1739.2007.00826.x>
- Kuhl, H. (1820). *Beiträge zur Zoologie und vergleichenden Anatomie*. Frankfurt, Germany: Hermannsche Buchhandlung.

- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33, 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Leigh, J. W., & Bryant, D. (2015). Popart: Full-feature software for haplotype network construction. *Methods in Ecology and Evolution*, 6, 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- Librado, P., & Rozas, J. (2009). DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Liu, Y., Zhang, L., & Cao, T. (2013). *Python bivittatus mitochondrion, complete genome*, NCBI Reference Sequence: NC_021479.1, Department of Animal Sciences, Hainan University, No. 58 Renmin Road, Haikou, Hainan 570228, China.
- Lynch, M., & Ritland, K. (1999). Estimation of pairwise relatedness with molecular markers. *Genetics*, 152, 1753–1766.
- Mainali, K. P., Warren, D. L., Dhileepan, K., McConnachie, A., Strathie, L., Hassan, G., ... Parmesan, C. (2015). Projecting future expansion of invasive species: Comparing and improving methodologies for species distribution modeling. *Global Change Biology*, 21, 4464–4480. <https://doi.org/10.1111/gcb.13038>
- Mazzotti, F. J., Cherkiss, M. S., Hart, K. M., Snow, R. W., Rochford, M. R., Dorcas, M. E., & Reed, R. N. (2011). Cold-induced mortality of invasive Burmese pythons in south Florida. *Biological Invasions*, 13, 143–151. <https://doi.org/10.1007/s10530-010-9797-5>
- McCleery, R. A., Sovie, A., Reed, R. N., Cunningham, M. W., Hunter, M. E., & Hart, K. M. (2015). Marsh rabbit mortalities tie pythons to the precipitous decline of mammals in the Everglades. *Proceedings of the Royal Society of London B: Biological Sciences*, 282, 20150120. <https://doi.org/10.1098/rspb.2015.0120>
- McDiarmid, R., Campbell, J. A., & Touré, T. A. (1999). *Snake species of the world: A taxonomic and geographic reference*, Vol. 1. Washington, DC: The Herpetologists' League.
- McPhee, M. V., & Turner, T. F. (2009). Genealogical diversity suggests multiple introductions of white suckers (*Catostomus commersonii*) into the Rio Grande, New Mexico. *Southwestern Naturalist*, 54, 485–492. <https://doi.org/10.1894/MRD-02.1>
- Meshaka, W. E., Loftus, W. F., & Steiner, T. (2000). An annotated inventory of the herpetofauna of Everglades National Park, Florida. *Florida Scientist*, 63, 84–103.
- Oakley, C. G., Ågren, J., & Schemske, D. W. (2015). Heterosis and outbreeding depression in crosses between natural populations of *Arabidopsis thaliana*. *Heredity*, 115, 73. <https://doi.org/10.1038/hdy.2015.18>
- O'Shea, M. (2007). *Boas and pythons of the World*. Cape Town, South Africa: New Holland Publishers.
- Paetkau, D., Calvert, W., Stirling, I., & Strobeck, C. (1995). Microsatellite analysis of population structure in Canadian polar bears. *Molecular Ecology*, 4, 347–354. <https://doi.org/10.1111/j.1365-294X.1995.tb00227.x>
- Paetkau, D., Slade, R., Burden, M., & Estoup, A. (2004). Genetic assignment methods for the direct, real-time estimation of migration rate: A simulation-based exploration of accuracy and power. *Molecular Ecology*, 13, 55–65. <https://doi.org/10.1046/j.1365-294X.2004.02008.x>
- Paetkau, D., & Strobeck, C. (1994). Microsatellite analysis of genetic variation in black bear populations. *Molecular Ecology*, 3, 489–495. <https://doi.org/10.1111/j.1365-294X.1994.tb00127.x>
- Peakall, R., & Smouse, P. E. (2006). GENALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, 6, 288–295. <https://doi.org/10.1111/j.1471-8286.2005.01155.x>
- Pfeiffer, J. M. III, Johnson, N. A., Randklev, C. R., Howells, R. G., & Williams, J. D. (2016). Generic reclassification and species boundaries in the rediscovered freshwater mussel 'Quadrula' mitchelli (Simpson in Dall, 1896). *Conservation Genetics*, 17, 279–292. <https://doi.org/10.1007/s10592-015-0780-7>
- Piaggio, A. J., Engeman, R. M., Hopken, M. W., Humphrey, J. S., Keacher, K. L., Bruce, W. E., & Avery, M. L. (2013). Detecting an elusive invasive species: A diagnostic PCR to detect Burmese python in Florida waters and an assessment of persistence of environmental DNA. *Molecular Ecology Resources*, 14, 374–380. <https://doi.org/10.1111/1755-0998.12180>
- Piry, S., Alapetite, A., Cornuet, J. M., Paetkau, D., Baudouin, L., & Estoup, A. (2004). GENECLASS2: A software for genetic assignment and first-generation migrant detection. *Journal of Heredity*, 95, 536–539. <https://doi.org/10.1093/jhered/esh074>
- Piry, S., Luikart, G., & Cornuet, J. M. (1999). BOTTLENECK: a computer program for detecting recent reductions in the effective population size using allele frequency data. *Journal of Heredity*, 90, 502–503. <https://doi.org/10.1093/jhered/90.4.502>
- Pittman, S. E., Hart, K. M., Cherkiss, M. S., Snow, R. W., Fujisaki, I., Smith, B. J., ... Dorcas, M. E. (2014). Homing of invasive Burmese pythons in South Florida: Evidence for map and compass senses in snakes. *Biology Letters*, 10, 20140040. <https://doi.org/10.1098/rsbl.2014.0040>
- Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155, 945–959.
- Ratnasingham, S., & Hebert, P. D. N. (2007). BOLD: The barcode of life data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, 7, 355–364.
- Ratnasingham, S., & Hebert, P. D. N. (2013). A DNA-based registry for all animal species: The Barcode Index Number (BIN) system. *PLoS ONE*, 8, e66213. <https://doi.org/10.1371/journal.pone.0066213>
- Rawlings, L. (2001). *Molecular systematics of Australian and New Guinea pythons*. Adelaide, SA: The University of Adelaide.
- Raymond, M., & Rousset, F. (1995). GENEPOP (version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity*, 86, 248–249. <https://doi.org/10.1093/oxfordjournals.jhered.a111573>
- Reed, R. N., Hart, K. M., Rodda, G. H., Mazzotti, F. J., Snow, R. W., Cherkiss, M., ... Goetz, S. (2011). A field test of attractant traps for invasive Burmese pythons (*Python molurus bivittatus*) in southern Florida. *Wildlife Research*, 38, 114–121. <https://doi.org/10.1071/WR10202>
- Reichert, B. E., Sovie, A. R., Udell, B. J., Hart, K. M., Borkhataria, R. R., Bonneau, M., ... McCleery, R. (2017). Urbanization may limit impacts of an invasive predator on native mammal diversity. *Diversity and Distributions*, 23, 355–367. <https://doi.org/10.1111/ddi.12531>
- Reynolds, G., Niemiller, M. L., & Revell, L. J. (2014). Toward a Tree-of-Life for the boas and pythons: Multilocus species-level phylogeny with unprecedented taxon sampling. *Molecular Phylogenetics and Evolution*, 71, 201–213. <https://doi.org/10.1016/j.ympev.2013.11.011>
- Rice, W. R. (1989). Analyzing tables of statistical tests. *Evolution*, 43, 223–225. <https://doi.org/10.1111/j.1558-5646.1989.tb04220.x>
- Rissler, L. J., & Apodaca, J. J. (2007). Adding more ecology into species delimitation: Ecological Niche models and phylogeography help define cryptic species in the Black Salamander (*Aneides flavipunctatus*). *Systematic Biology*, 56, 924–942. <https://doi.org/10.1080/10635150701703063>
- Rodda, G. H., Jarnevich, C. S., & Reed, R. N. (2009). What parts of the US mainland are climatically suitable for invasive alien pythons spreading from Everglades National Park? *Biological Invasions*, 11, 241–252. <https://doi.org/10.1007/s10530-008-9228-z>
- Roman, J., & Darling, J. A. (2007). Paradox lost: Genetic diversity and the success of aquatic invasions. *Trends in Ecology & Evolution*, 22, 454–464. <https://doi.org/10.1016/j.tree.2007.07.002>
- Ryan, S. F., Deines, J. M., Scriber, J. M., Pfrender, M. E., Jones, S. E., Emrich, S. J., & Hellmann, J. J. (2018). Climate-mediated hybrid zone movement revealed with genomics, museum collection, and simulation modeling. *Proceedings of the National Academy of Sciences of the United States of America*, 115, E2284–E2291. <https://doi.org/10.1073/pnas.1714950115>
- Schleip, W., & O'Shea, M. (2010). Annotated checklist of the recent and extinct pythons (Serpentes, Pythonidae), with notes on nomenclature, taxonomy, and distribution. *ZooKeys*, 66, 29–80. <https://doi.org/10.3897/zookeys.66.683>

- Seehausen, O. (2004). Hybridization and adaptive radiation. *Trends in Ecology & Evolution*, 19, 198–207. <https://doi.org/10.1016/j.tree.2004.01.003>
- Serrao, N. R., Steinke, D., & Hanner, R. H. (2014). Calibrating snakehead diversity with DNA barcodes: Expanding taxonomic coverage to enable identification of potential and established invasive species. *PLoS ONE*, 9, e99546. <https://doi.org/10.1371/journal.pone.0099546>
- Slowinski, J. B., & Lawson, R. (2002). Snake phylogeny: Evidence from nuclear and mitochondrial genes. *Molecular Phylogenetics and Evolution*, 24, 194–202. [https://doi.org/10.1016/S1055-7903\(02\)00239-7](https://doi.org/10.1016/S1055-7903(02)00239-7)
- Smith, B. J., Cherkiss, M. S., Hart, K. M., Rochford, M. R., Selby, T. H., Snow, R. W., & Mazzotti, F. J. (2016). Betrayal: Radio-tagged Burmese pythons reveal locations of conspecifics in Everglades National Park. *Biological Invasions*, 18, 3239–3250. <https://doi.org/10.1007/s10530-016-1211-5>
- Snow, R. W., Brien, M. L., Cherkiss, M. S., Wilkins, L., & Mazzotti, F. J. (2007). Dietary habits of the Burmese python, *Python molurus bivittatus*, in Everglades National Park, Florida. *Herpetological Bulletin*, 101, 5–7.
- Sovie, A., McCleery, R., Fletcher, R., & Hart, K. (2016). Invasive pythons, not anthropogenic stressors, explain the distribution of a keystone species. *Biological Invasions*, 18, 3309–3318. <https://doi.org/10.1007/s10530-016-1221-3>
- Stepien, C. A., & Tumeo, M. A. (2006). Invasion genetics of Ponto-Caspian gobies in the Great Lakes: A 'cryptic' species, absence of founder effects, and comparative risk analysis. *Biological Invasions*, 8, 61–78. <https://doi.org/10.1007/s10530-005-0237-x>
- Supikamolnani, A., & Srikulnath, K. (2014). *Molecular barcoding of some venomous snakes, and species-specific multiplex PCR assay to identify seven antivenom snakes in Thailand*. NCBI Reference Sequence: AB920233. Bangkok, Thailand: Department of Genetics, Kasetsart University.
- Tajima, F., & Nei, M. (1984). Estimation of evolutionary distance between nucleotide sequences. *Molecular Biology and Evolution*, 1, 269–285.
- Tamura, K. (1992). Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C-content biases. *Molecular Biology and Evolution*, 9, 678–687.
- Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10, 512–526.
- Thielsch, A., Knell, A., Mohammadyari, A., Petrusek, A., & Schwenk, K. (2017). Divergent clades or cryptic species? Mito-nuclear discordance in a *Daphnia* species complex. *BMC Evolutionary Biology*, 17, 227. <https://doi.org/10.1186/s12862-017-1070-4>
- Townson, S. (1980). Observations of the reproduction of the Indian Python in captivity, with special reference to the interbreeding of the two subspecies *Python molurus molurus* and *Python molurus bivittatus*. In S. Townson, N. J. Millichamp, D. G. D. Lucas, & A. J. Millwood (Eds.), *The care and breeding of captive reptiles* (pp. 69–80). London, UK: Brit. Herp. Soc.
- Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M., & Shipley, P. (2004). MICRO-CHECKER: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4, 535–538. <https://doi.org/10.1111/j.1471-8286.2004.00684.x>
- Vidal, O., García-Berthou, E., Tedesco, P. A., & García-Marín, J.-L. (2010). Origin and genetic diversity of mosquitofish (*Gambusia holbrooki*) introduced to Europe. *Biological Invasions*, 12, 841–851. <https://doi.org/10.1007/s10530-009-9505-5>
- Wall, C. E., Cozza, S., Riquelme, C. A., McCombie, W. R., Heimiller, J. K., Marr, T. G., & Leinwand, L. A. (2011). Whole transcriptome analysis of the fasting and fed Burmese python heart: Insights into extreme physiological cardiac adaptation. *Physiological Genomics*, 43, 69–76. <https://doi.org/10.1152/physiolgenomics.00162.2010>
- Wang, J. (2011). Coancestry: A program for simulating, estimating and analysing relatedness and inbreeding coefficients. *Molecular Ecology Resources*, 11, 141–145. <https://doi.org/10.1111/j.1755-0998.2010.02885.x>
- Waples, R. S. (2006). A bias correction for estimates of effective population size based on linkage disequilibrium at unlinked gene loci. *Conservation Genetics*, 7, 167–184. <https://doi.org/10.1007/s10592-005-9100-y>
- Waples, R. S., & Do, C. (2008). LDNE: A program for estimating effective population size from data on linkage disequilibrium. *Molecular Ecology Resources*, 8, 753–756. <https://doi.org/10.1111/j.1755-0998.2007.02061.x>
- Whitlock, R., Hipperson, H., Mannarelli, M., & Burke, T. (2008). A high-throughput protocol for extracting high-purity genomic DNA from plants and animals. *Molecular Ecology Resources*, 8, 736–741. <https://doi.org/10.1111/j.1755-0998.2007.02074.x>
- Wilberg, M. J., & Dreher, B. P. (2004). GENECAP: A program for analysis of multilocus genotype data for non-invasive sampling and capture-recapture population estimation. *Molecular Ecology Notes*, 4, 783–785. <https://doi.org/10.1111/j.1471-8286.2004.00797.x>
- Wilcox, T. M., Carim, K. J., McKelvey, K. S., Young, M. K., & Schwartz, M. K. (2015). The dual challenges of generality and specificity when developing environmental DNA markers for species and subspecies of *Oncorhynchus*. *PLoS ONE*, 10, e0142008. <https://doi.org/10.1371/journal.pone.0142008>
- Wilcox, T. M., McKelvey, K. S., Young, M. K., Jane, S. F., Lowe, W. H., Whiteley, A. R., & Schwartz, M. K. (2013). Robust detection of rare species using environmental DNA: The importance of primer specificity. *PLoS ONE*, 8, e59520. <https://doi.org/10.1371/journal.pone.0059520>
- Williamson-Natesan, E. G. (2005). Comparison of methods for detecting bottlenecks from microsatellite loci. *Conservation Genetics*, 6, 551–562.
- Willson, J. D., Dorcas, M. E., & Snow, R. W. (2011). Identifying plausible scenarios for the establishment of invasive Burmese pythons (*Python molurus*) in Southern Florida. *Biological Invasions*, 13, 1493–1504. <https://doi.org/10.1007/s10530-010-9908-3>
- Willson, J. D., Snow, R. W., Reed, R. N., & Dorcas, M. E. (2014). *Python molurus bivittatus* (Burmese python). Minimum size at maturity. *Herpetological Review*, 45, 343–344.
- You, C.-W., Lin, Y.-P., Lai, Y.-H., Chen, Y.-L., Tang, Y., Chou, S.-P., ... Lin, S.-M. (2013). Return of the pythons: First formal records, with a special note on recovery of the Burmese python in the demilitarized Kinmen islands. *Zoological Studies*, 52, 1–11.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Hunter ME, Johnson NA, Smith BJ, et al. Cytonuclear discordance in the Florida Everglades invasive Burmese python (*Python bivittatus*) population reveals possible hybridization with the Indian python (*P. molurus*). *Ecol Evol*. 2018;00:1–14. <https://doi.org/10.1002/ece3.4423>