

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Sociology Department, Faculty Publications

Sociology, Department of

---

2017

# Bi-cultural dynamics for risk and protective factors for cardiometabolic health in an Alaska Native (Yup'ik) population

Jacques Philip

*University of Alaska, Fairbanks, [jphilip@alaska.edu](mailto:jphilip@alaska.edu)*

Toves K. Ryman

*Bill and Melinda Gates Foundation*

Scarlett E. Hopkins

*University of Alaska, Fairbanks*

Diane M. O'Brien

*University of Alaska, Fairbanks*

Andrea Bersamin

*University of Alaska, Fairbanks*

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/sociologyfacpub>



Part of the [Family, Life Course, and Society Commons](#), and the [Social Psychology and Interaction Commons](#)

---

Philip, Jacques; Ryman, Toves K.; Hopkins, Scarlett E.; O'Brien, Diane M.; Bersamin, Andrea; Pomeroy, Jeremy; Thummel, Kenneth E.; Austin, Melissa A.; Boyer, Bert B.; and Dombrowski, Kirk, "Bi-cultural dynamics for risk and protective factors for cardiometabolic health in an Alaska Native (Yup'ik) population" (2017). *Sociology Department, Faculty Publications*. 522.

<http://digitalcommons.unl.edu/sociologyfacpub/522>

This Article is brought to you for free and open access by the Sociology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Sociology Department, Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Jacques Philip, Toves K. Ryman, Scarlett E. Hopkins, Diane M. O'Brien, Andrea Bersamin, Jeremy Pomeroy, Kenneth E. Thummel, Melissa A. Austin, Bert B. Boyer, and Kirk Dombrowski


RESEARCH ARTICLE

# Bi-cultural dynamics for risk and protective factors for cardiometabolic health in an Alaska Native (Yup'ik) population

Jacques Philip<sup>1\*</sup>, Tove K. Ryman<sup>2</sup>, Scarlett E. Hopkins<sup>1</sup>, Diane M. O'Brien<sup>1</sup>, Andrea Bersamin<sup>1</sup>, Jeremy Pomeroy<sup>3</sup>, Kenneth E. Thummel<sup>4</sup>, Melissa A. Austin<sup>5</sup>, Bert B. Boyer<sup>1</sup>, Kirk Dombrowski<sup>6</sup>

**1** Center for Alaska Native Health Research, University of Alaska Fairbanks, Fairbanks, Alaska, United States of America, **2** Bill and Melinda Gates Foundation, Seattle, Washington, United States of America, **3** Marshfield Clinic Research Institute, Marshfield, Wisconsin, United States of America, **4** Department of Pharmaceutics, University of Washington, Seattle, Washington, United States of America, **5** Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, United States of America, **6** Department of Sociology, University of Nebraska, Lincoln, Nebraska, United States of America

\* [jphilip@alaska.edu](mailto:jphilip@alaska.edu)



**OPEN ACCESS**

**Citation:** Philip J, Ryman TK, Hopkins SE, O'Brien DM, Bersamin A, Pomeroy J, et al. (2017) Bi-cultural dynamics for risk and protective factors for cardiometabolic health in an Alaska Native (Yup'ik) population. PLoS ONE 12(11): e0183451. <https://doi.org/10.1371/journal.pone.0183451>

**Editor:** Junwen Wang, Mayo Clinic Arizona, UNITED STATES

**Received:** March 22, 2017

**Accepted:** August 6, 2017

**Published:** November 1, 2017

**Copyright:** © 2017 Philip et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The data presented in this manuscript was obtained through a NIH-funded research collaboration with the Yup'ik people and their healthcare providers in the YK-Delta of Alaska. It was generated through the actions of the research team and in partnership with the Yup'ik people. The tribal leadership has approved publication and dissemination of the data in the format provided in this manuscript. The primary dataset from our study is owned by the people of the YK-delta, in recognition of tribal sovereign authority stipulated by the government

## Abstract

Alaska Native people experience disparities in mortality from heart disease and stroke. This work attempts to better understand the relationships between socioeconomic, behavioral, and cardiometabolic risk factors among Yup'ik people of southwestern Alaska, with a focus on the role of the socioeconomic, and cultural components. Using a cross-sectional sample of 486 Yup'ik adults, we fitted a Partial Least Squares Path Model (PLS-PM) to assess the associations between components, including demographic factors [age and gender], socioeconomic factors [education, economic status, Yup'ik culture, and Western culture], behavioral factors [diet, cigarette smoking and smokeless tobacco use, and physical activity], and cardiometabolic risk factors [adiposity, triglyceride-HDL and LDL lipids, glycemia, and blood pressure]. We found relatively mild associations of education and economic status with cardiometabolic risk factors, in contrast with studies in other populations. The socioeconomic factor and participation in Yup'ik culture had potentially protective associations with adiposity, triglyceride-HDL lipids, and blood pressure, whereas participation in Western culture had a protective association with blood pressure. We also found a moderating effect of participation in Western culture on the relationships between Yup'ik culture participation and both blood pressure and LDL lipids, indicating a potentially beneficial additional effect of bi-culturalism. Our results suggest that reinforcing protective effects of both Yup'ik and Western cultures could be useful for interventions aimed at reducing cardiometabolic health disparities.

of the United States. This data access arrangement was approved by the NIH, who funded this research. Requests for access to the primary data can be made to the Yup'ik tribal leadership. While we cannot speak for them, we believe that all requests would be given due consideration. Such requests can be directed to: Bert Boyer, Ph.D., Professor of Molecular Biology, Director, Center for Alaska Native Health Research, Associate Director of Human Health, Institute of Arctic Biology, University of Alaska Fairbanks, 311 Irving I Building, Fairbanks, AK 99775, 907-474-7733 (phone), 907-474-5700 (fax) \*Call to let me know. [bboyer@alaska.edu](mailto:bboyer@alaska.edu) (forwarded to gmail), <http://canhr.uaf.edu/>.

**Funding:** The study was funded by the following grants from the National Institute of Health: P20RR016430, P30GM103325, R01DK074842, U01GM092676, U54GM115371, U54GM104944.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Over the last three decades, attention to social determinants of health among indigenous communities have focused on individual and community social stressors that stem from the marginal place of indigenous communities in the current global political economy [1–5]. Initially this research focused on the discovery and examination of “risk factors” that influenced individual health outcomes [6–8]. More recently, researchers have also emphasized the importance of indigenous culture in influencing health outcomes [3,9–13]. The potential for local culture to act as protection against social harms and serve as a source of community resiliency, for example by promoting healthy traditional diets, represents what some have identified as significant change of research and intervention ideology [14].

Across the Arctic, culture-focused health interventions contend with a legacy of past, deleterious state-inspired programs such as community relocation and sedentarization [15–18]. These programs, originally intended to promote assimilation and the incorporation of Inuit/Inupiat/Yup'ik peoples in larger national political economies, have resulted in a well-known list of social ills, such as broken family ties or loss of language [19–21]. Emerging literature on ‘cultural-based protective factors’ in the region has focused on the potential for cultural participation to mitigate negative outcomes associated with this history [22], including suicide [23,24], substance abuse [19,25,26], Type 2 diabetes [27], depression [4,28] and other mental health outcomes [29,30].

Importantly, ethnographic (and other) research has shown that cultural protective factors are embedded in complex social and historical practices, in which protective and harmful effects may be intermingled [31–35]. In this view, culture is not a simple independent variable in the sense required for linear regression, nor is it monolithic and consistent in its effects on all individuals across a community or a region [36,37]. To fully enact culture-based interventions, a more complex understanding of the inter-relationships between culture and health behaviors among indigenous peoples is needed.

This study provides an analysis of the impact of cultural participation and other socioeconomic variables in relation to cardiometabolic risk factors within an Alaska Native population. Alaska Native people are more at risk of dying from heart disease or stroke [but not from diabetes] than non-Native people [38], justifying research aimed at addressing this disparity. Prior research with Alaska Native Yup'ik people established that cardiometabolic risk was associated positively with increased consumption of processed foods, and negatively with increased consumption of traditional foods [39]. Yet beyond the material aspect of nutrients brought by food, subsistence practices have a much more profound meaning for Yup'ik people, for example through norms around the respect for animals, rituals, and spirituality that may contribute to lasting socioeconomic relationships that carry indirect health benefits [40]. For example, the Yup'ik concept of Yuuyaraq, “The way of being a human being”, defines how humans, animal, and nature interact in the Yup'ik way of life, which not only includes activities such as hunting, fishing, gathering, dances, and ceremonies, but also cultural values transmitted orally by Elders, such as sharing, humility, or conflict avoidance, and rules strongly tied to the spiritual world, like the correct way of speaking about living things, or how to handle hunted animals [41]. Traditional lifestyle was also found to be associated with higher levels of physical activity, lower perceived stress, and lower body mass index (BMI) [42], all potential counteractions to cardiometabolic risk.

In addition, a recent study showed that *iqmik* use (a traditional form of smokeless tobacco used by Yup'ik people) was negatively associated with cardiometabolic risk, and this association was attenuated after adjustment for BMI, suggesting a mediating effect of adiposity [43]. Here too cultural factors may be involved in this relationship; Wolsko et al [44] describe how

*iqmik* is used by Yup'ik people as a way to cope with stress, which is often perceived as being induced by acculturation to a Western way of life [45]. In contrast, Boden-Albala et al [46] found that bicultural identity decreased the association between age and hypertension, and between education and hypertension, indicating that Yup'ik culture and Western culture can potentially interact in complex and potentially health-protective ways.

Operationalizing the concept of culture for statistical analysis offers significant challenges. In studies aimed at mobilizing cultural factors for health interventions, culture has been at times been defined as participation in activities linked to spirituality and ancient traditions—activities such as hunting, food gathering, the sharing of locally obtained foods, and traditional craft production based in these same activities— [29,47–49]. Others have used self-governance and local control associated to define cultural continuity [50–52]. Still others have looked at patterns of social exchange associated with kinship and indigenous/traditional knowledge as stand-ins for ongoing collective culture [37,53,54]. Here we follow a large number of researchers working in indigenous communities in the north and operationalize culture in both the community and individual sense through ongoing use of the indigenous language of the area and a self-assessment of “following a Yup'ik” lifestyle [28,29,55–57]. Research has shown that indigenous language use is often intrinsic to participation in traditional cultural activities [58,59], in effect capturing a range of health-related behaviors related to cardiometabolic risk. In addition, the term “traditional/Yup'ik lifestyle” is commonly used in Alaska as a signifier of participation in traditional, subsistence activities [36,60]. By looking culture as it has been defined and mobilized for health research by these researchers, we build on findings that demonstrate the potential efficacy of cultural participation for individual physical and mental health [58].

The analysis described below investigates the relationship among more than 20 socioeconomic, cultural, behavioral and cardiometabolic measures. To date we are not aware of studies that have attempted to evaluate such a wide range of potentially protective and deleterious factors in a single analytical framework. In this study, we seek to build on prior factor-specific findings by employing multivariate analytical techniques aimed at evaluating the complex interweaving of cultural, socioeconomic, behavioral and physical factors associated with both cardiometabolic risk and protection, while allowing culture (both Yup'ik and Western) to have interdependent and heterogeneous effects. The long-term goal of this research is to further define the interactions between culture and health so that they may be implemented in interventions aimed at improving the cardiometabolic health of Alaska Native people.

## Methods

The Center for Alaska Native Health Research (CANHR) was created in 2001 at the University of Alaska Fairbanks, with the primary aim to identify behavioral, genetic, and nutritional risk and protective factors contributing to obesity and Type 2 diabetes (T2D) in Yup'ik Alaska Native people residing in Southwest Alaska. Detailed descriptions of study design and recruitment methods have been previously published [61,62]. The majority ethnic identity in the region is Yup'ik, with a minority identifying as Cup'ik, with both groups falling broadly under Yup'ik culture.

## Participants

The analytic sample for this article includes participants recruited from 9 rural communities in the Yukon-Kuskokwim Delta with populations between 250 and 700, plus a hub community of ~6,000 residents. Between September 2009 and May 2013, CANHR investigators made recurrent visits to each community. All community members 14 years or older, self-identified

as Alaska Native descent or married to an Alaska Native individual, and non-pregnant were invited to participate. All eligible participants were enrolled at each visit and thus the data consist of multiple cross-sectional, convenience samples. For this analysis, participants include adults ( $\geq 18$  years of age), who self-identified as either Yup'ik or Cup'ik, who had completed measures for cardiometabolic risk factors, dietary and physical activity assessment, and had been fasting. For participants with multiple visits, the latest one was used. From this sample of 620, an additional 134 participants were excluded because they were on diabetes, hypertension, or dyslipidemia control medication, as a way to reduce the bias related to the fact that their cardio-metabolic risk factors are artificially lowered, bringing the final sample to 486 individuals.

All procedures involving study participants were approved by the University of Alaska Fairbanks Institutional Review Board and the Yukon-Kuskokwim Health Corporation Human Studies Committee. Written informed consent was obtained in English or Yup'ik from participants prior to data collection.

### Measures

At each visit, participants answered questionnaires in English or Yup'ik. These included demographic characteristics, socioeconomic and cultural questions, medical history, medication usage, fasting status, tobacco use behavior, and diet from food frequency questionnaire (FFQ). Weight, height and other anthropometric measurements were collected during a physical exam by trained investigators. Physical activity was measured by an Actiheart system, monitoring movement and heart rate.

**Socio-economic and cultural measures.** The assessment of education, economic status, and culture participation is described in Table 1. The two lifestyle questions were developed by CANHR researchers following focus groups and interviews of Yup'ik participants' conceptualizations of health and wellness [45,63] and based on the orthogonal model of cultural

**Table 1. Variables, questions and options for socioeconomic and cultural measures.**

Variables	Questions	Options/Scale
<b>Education</b>		
Number of school years completed <sup>a</sup>	<i>What is the last grade of school (K-12) you completed?</i>	0–12
	<i>How many years of college did you complete?</i>	Number of years
<b>Economic status</b>		
Runs out of money for food	<i>How often does your money for household utilities (ie water, fuel oil, electricity, etc) run out before the end of the month?</i>	1:Never, 2:Seldom, 3: Sometimes, 4:Most times, 5: Always
Runs out of money for utilities	<i>How often does your money for food run out before the end of the month?</i>	1:Never, 2:Seldom, 3: Sometimes, 4:Most times, 5: Always
<b>Western Culture</b>		
Main language spoken at home is English <sup>b</sup>	<i>Which language do you speak most often at home</i>	Cup'ik, Yup'ik, English, Other
Follows Western lifestyle	<i>How much do you follow the White (or Kass'aq) way of life</i>	1:Not at all, 2:Some, 3:A lot
<b>Yup'ik Culture</b>		
Speaks Yup'ik or Cup'ik <sup>cd</sup>	<i>Which languages do you speak?</i>	Cup'ik, Yup'ik, English, Other
Main language spoken at home is Yup'ik or Cup'ik <sup>d</sup>	<i>Which language do you speak most often at home</i>	Cup'ik, Yup'ik, English, Other
Follows Yup'ik lifestyle	<i>How much do you follow the traditional Yup'ik way of life</i>	1:Not at all, 2:Some, 3:A lot

<sup>a</sup>The score is the sum of scores of the two question.

<sup>b</sup>The score is 2 if the answer is English, 1 otherwise.

<sup>c</sup>Multiple answers allowed.

<sup>d</sup>The score is 2 if the answer includes Yup'ik or Cup'ik, 1 otherwise.

<https://doi.org/10.1371/journal.pone.0183451.t001>

identification, which posits that identification patterns with different cultures are independent of each other, rather than being along a linear continuum [64].

**Lifestyle measures.** Dietary intake was assessed with both self-reported FFQ and objectively measured stable isotope biomarkers. A culturally specific Yup'ik FFQ was developed and validated by CANHR [65,66]. The FFQ data were used to identify three dietary patterns of traditional or subsistence foods (e.g., walrus or seal), processed foods, and fruits and vegetables. The dietary patterns were used to calculate dietary pattern scores representing the relative frequency of consumption for each. Details of the dietary patterns have been published previously [65,66].

Stable isotope ratios were used as dietary biomarkers for fish and marine mammal intake and processed food intake. In this population, the nitrogen isotope ratio ( $^{15}\text{N}/^{14}\text{N}$ , expressed as the  $\delta^{15}\text{N}$  value) is strongly associated with fish and marine mammal intake [67–69]. The carbon isotope ratio ( $^{13}\text{C}/^{12}\text{C}$  or the  $\delta^{13}\text{C}$  value) is elevated in corn and sugar-based foods, and is associated with processed food intake when assessed via FFQ [66].

Physical activity was measured by asking participants to wear a combined heart rate/movement device (Actiheart, CamNtech Ltd, Papworth UK), for four consecutive days. We used the average daily total activity counts as a measure of all physical activity regardless of intensity, the moderate to vigorous activity time classified as time where heart rate was at least 1.75 times resting heart rate, and sedentary time which was classified as minutes with less than 10 accelerometry counts per minute and a valid heart rate less than 1.10 times resting heart rate (rev. coded). All three measures were adjusted for actual wear time of the device, which varied between 10 and 24 hours per day.

For both cigarette smoking and smokeless tobacco use, a categorical measure including the levels of current, past, and never user was defined, with current use representing any use at the time of interview, regardless of the amount. We also defined a continuous variable as the number of cigarettes or chews per day currently used. Smokeless tobacco use among Yup'ik people in this study includes mainly *Iqmik*, a homemade mixture of cured tobacco leaves and ash from *Phellinus igniarius*, a fungus commonly found on birch and alders, or ash from willow or alder [70]. Smokeless tobacco also included to a much smaller extent commercial chew and snuff [44].

**Cardiometabolic risk factors.** Blood was drawn after 12 hours of fasting and processed locally into plasma, serum, buffy coat and red blood cells (RBC). Blood samples were analyzed locally for fasting plasma glucose (FPG) and HbA1C and plasma was sent for analysis to UC Davis University for lipids. Resting systolic and diastolic blood pressure were measured on participants, using the OMRON HEM907 automated blood pressure cuff. The mean of the last 2 out of three measures was used for analysis. Waist circumference (WC) was measured twice using a Gulick II 150 cm anthropometric tape attached with a tension-meter immediately below the lowest lateral portion of the rib case. The average of two measures was used for analysis. BMI was calculated as weight (kg)/height (m<sup>2</sup>). More details on these protocols have been published previously [39].

## Analysis

We adopted a two-step approach for our analysis, with (1) a Principal Components Analysis (PCA) of the cardiometabolic risk factors to reduce those nine variables into fewer components, and (2) a Partial Least Square Path Model (PLS-PM) to explore how demographic, socioeconomic, and behavioral factors predict those risk factors components. PLS-PM is a non-parametric alternative method to structural equation modeling (SEM), commonly used in econometrics [71], which allows for the construction of a path model between a set of latent

variables (inner model), each of which being represented by a block of correlated measured or manifest variables (outer model). Compared to SEM, PLS-PM does not require normally distributed variables, allows easier convergence for small samples relative to the number of variables and cases, and is more adapted for exploring a model than strict testing of strongly established hypothesis [72–74]. The disadvantage over SEM is that PLS-PM produces less precise estimates [75].


All analyses were conducted with the R software, v. 3.2 [76], using the Psych package for PCA, and the plsmpm package for PLS-PM analysis. Missing values were replaced by the median for that variable across participants.

**Principal Component Analysis.** We based our PCA of cardiometabolic risk factors on the idea proposed by several authors that continuous scores of components of the metabolic syndrome (MetS) have better predictive power than its dichotomous standard definition [77–79]. Using commonly accepted criteria defining the MetS, i.e., abdominal obesity, an index of insulin resistance/glucose intolerance, dyslipidemia, and hypertension [80,81], We conducted the PCA on the following variables: Waist Circumference, BMI, LDL-C, - HDL-C (reverse coded), Triglycerides, Systolic BP, Diastolic BP, Glucose, and HbA1C, requiring five components with varimax rotation. Even though our PCA is based on definitions of the MetS, the resulting components do not represent a measure of the MetS, but simply five continuous dimensions of cardiometabolic risk.

**Partial Least Square Path analysis.** A PLS-PM model is defined by an outer and an inner model, as represented by Fig 1. To avoid repeating a very similar figure, Fig 1 is used to both show the topology of the model here, and the significant paths later in the results. In the outer or measurement model each latent variable or component is defined as a weighted linear combination of its manifest variables. For example, the latent variable Yup'ik culture participation is defined by its manifest variables Yup'ik at home, Speaks Yup'ik, and Yup'ik lifestyle. The inner or structural model defines the relationships between latent variables. Those relationships are evaluated as a set of linear regressions, one for each endogenous latent variable (a latent variable that has at least one incoming path). In each regression, that endogenous variable is predicted by all variables that have a path to it. For example, adiposity is predicted by behavioral, socioeconomic, and demographic components. The path coefficients are those regression coefficients and are labeled as effects, in PLS-PM terminology, but should be understood as associations, since our analysis uses PLS-PM in an exploratory fashion. These regressions are fitted iteratively as the weights of the outer model are optimized; convergence is obtained when the residual overall variance is minimized. Confidence intervals can be estimated by non-parametric methods [73,82]. Here we used 1000 bootstrapping repetitions, and confidence level was set at 0.05. We did not use multiple comparison adjustment because we are not formally testing hypothesis, but we are using PLS-PM in an exploratory fashion.

After the components of the cardiometabolic risk factors were identified, we included them in the inner model, i.e., demographic components [Gender and Age], socioeconomic components [Education, Economic Status, Western Culture, and Yup'ik Culture], and behavioral components [traditional diet, processed diet, and fruits and vegetables diet, physical activity, cigarette smoking, and smokeless tobacco use]. We then established directional relationships between components that are theorized to predict others. Thus we defined a path between each demographic and behavioral components and cardiometabolic risk factors components. In addition, we defined a path from adiposity to each of the other cardiometabolic risk factors components given the central role of adiposity the International Diabetes Federation (IDF) definition of the metabolic syndrome [81]. Downstream from the cardiometabolic risk factors, we defined paths from each of the socioeconomic components to the behavioral components





**Fig 1. PLS-PM relationships with total effect coefficients between all latent variables other than gender and age.** Significant effects only ( $\alpha = 0.05$ ). Ellipses are latent variables; rectangles are measured variables composing each latent variable. Plain black lines and red dotted lines represent positive and negative associations respectively.

<https://doi.org/10.1371/journal.pone.0183451.g001>

and from the demographic components to the behavioral components as well as the socio-economic components.

We also defined a model which includes an interaction between Western and Yup'ik cultures, based on previous research suggesting that bi-culturalism may have a protective effect on hypertension [46]. Among the different methods to compute an interaction effect in PLS-PM, we elected to use the two stage approach where the first stage consists of fitting the model without interaction as previously described, and the second of fitting an interaction model, which adds an interaction latent variable made of the score of Western and Yup'ik cultures obtained from the first model [83]. The interaction effect can then be assessed by coefficients of the paths between the interaction and the other latent [outcome] variables, and

visualized by using the model to predict each outcome variable for each level of Western and Yup'ik cultures.

## Results

Participant characteristics are presented in [Table 2](#). In summary, 51% of participants were female and the average age  $\pm$  sd was  $35.8 \pm 15.4$  years. The average education level was slightly below high school graduation at  $11.3 \pm 1.9$  years. Our two economic status variables are relatively balanced around the median. 95.1% of participants speak Yup'ik or Cup'ik and the majority (69%) of participants speak mostly Yup'ik or Cup'ik at home. Participants speaking mostly English are almost the reciprocal except for one who reported speaking neither. Almost all participants report following the Yup'ik lifestyle "some" or "a lot" (50% and 49%, respectively), and most follow the western lifestyle "some" or "a lot" (77% and 20%, respectively). Percentages of never, past and current tobacco users are respectively 34, 32, and 34 for cigarettes, and 16, 10, and 74 for smokeless tobacco. The overall average number of cigarettes smoked per day among current smokers is  $4.7 \pm 4.3$ . The overall average number of chews or snuffs per day among current users is  $4.9 \pm 3.9$ . Dietary intake, activity and risk factor variables had similar distributions than was published previously for the same population [43].

## Principal Component Analysis

[Table 3](#) shows the variance partition and variable loadings of the five components of the PCA of cardiometabolic risk factors. Adiposity explains 23% of the overall variance, blood pressure 19%, LDL lipids 15%, glycemia 14%, and TG-HDL lipids 14%, for a total of 84% of the total variance. The variables with highest loadings were retained for each component, to be used subsequently in the PLS-PM analysis as follows: waist circumference and BMI for adiposity, systolic and diastolic measures for blood pressure, just LDL-C for LDL lipids, glucose and HbA1C for glycemia, and—HDL-C and triglycerides for TG-HDL lipids.

## PLS-PM analysis

**Main effects.** [Table 4](#) displays outer model fitting characteristics which were marginal to good for most variables, after we removed other variables (not showed), because they altered unidimensionality or overall goodness of fit. For example loadings which should ideally be close to or above 0.7, were 0.8, 0.7 and 0.64 for the Yup'ik culture variables (*Yup'ik at Home*, *Yup'ik lifestyle* and *Can speak Yup'ik*, respectively). The variables *Mod./vig. Activity*, *Triglycerides*, and *Glucose* had the lowest loadings. Sensitivity analysis without those variables did not show significant differences in the results or overall goodness of fit. Additionally, crossloadings, which evaluate if each manifest variable always loads higher for its theorized latent variable than for the others, were calculated and were adequate, but are not presented in the table. Overall the fitting of the outer model is acceptable.

[Table 5](#) presents the inner model's coefficients and 95% CI or  $p < .05$  for significant total effects. A coefficient corresponds to a normalized regression slope and is considered statistically significant if its CI does not contain 0. A construct's total effect is the sum of its direct and indirect effects, the latter being obtained by multiplying the paths coefficients of indirect pathways.

For clarity, the total effects in the PLS-PM analysis inner and structural model are split into [Figs 1 and 2](#), and non-significant relationships have been omitted. [Fig 1](#) shows the relationships between the socioeconomic, behavioral and cardiometabolic risk components; the behavioral to cardiometabolic risk components; and adiposity to other cardiometabolic risk components. Among these variables, the strongest associations were Adiposity  $\rightarrow$  TG-HDL

**Table 2. Descriptive statistics of variables.**

	n (%)	Mean ± sd	Skew
<b>Gender</b>			
Male	238 (49.0)		
Female	248 (51.0)		
<b>Age (yrs)</b>			
Continuous	486	35.8 ± 15.4	0.7
<b>Number of school years completed (HS + college) (yrs)</b>			
Continuous	486	11.3 ± 1.9	-1.2
<b>Runs out of money for food<sup>b</sup></b>			
Never	41 (8.4)		
Seldom	37 (7.6)		
Sometimes	159 (32.7)		
Most times	107 (22.0)		
Always	80 (16.5)		
Missing	62 (12.8)		
<b>Runs out of money for utilities<sup>b</sup></b>			
Never	90 (18.5)		
Seldom	61 (12.6)		
Sometimes	132 (27.2)		
Most times	59 (12.1)		
Always	77 (15.8)		
Missing	67 (13.8)		
<b>Main language spoken at home is English</b>			
No	338 (69.5)		
Yes	148 (30.5)		
<b>Follows Western lifestyle</b>			
Not at all	15 (3.1)		
Some	375 (77.2)		
A lot	95 (19.5)		
Missing	1 (0.2)		
<b>Main language spoken at home is Yup'ik or Cup'ik</b>			
No	149 (30.7)		
Yes	337 (69.3)		
<b>Follows Yup'ik lifestyle</b>			
Not at all	5 (1.0)		
Some	242 (49.8)		
A lot	239 (49.2)		
<b>Speaks Yup'ik or Cup'ik</b>			
No	24 (4.9)		
Yes	462 (95.1)		
<b>δ15N isotope level (‰)</b>			
Continuous	486	8.8 ± 1.3	0.8
<b>Subsistence food dietary pattern score<sup>a</sup></b>			
Continuous	486	0.2 ± 2.3	-0.3
<b>δ13C isotope level (‰)</b>			
Continuous	486	-19.7 ± 0.7	-0.4
<b>Processed foods dietary pattern score<sup>a</sup></b>			
Continuous	486	0.8 ± 2.2	-0.4

(Continued)

Table 2. (Continued)

	n (%)	Mean ± sd	Skew
<b>Fruits and vegetables dietary pattern score<sup>a</sup></b>			
Continuous	486	-0.0 ± 2.2	-0.3
<b>Activity counts per day (count)</b>			
Continuous	486	60767.1 ± 37769.1	1.9
<b>Moderate to vigorous activity time per day (minutes)</b>			
Continuous	486	201.9 ± 176.9	1.4
<b>Rest time per day (minutes)<sup>b</sup></b>			
Continuous	486	940.3 ± 163.7	-1.5
<b>Never, past or current smoke tobacco user</b>			
Never Smoker	166 (34.2)		
Past Smoker	153 (31.5)		
Current Smoker	167 (34.4)		
<b>Cigarettes smoked per day (count)</b>			
Continuous	486	1.6 ± 3.4	3.3
Among smokers		4.7 ± 4.3	
<b>Never, past or current smokeless user</b>			
Never Smokeless	79 (16.3)		
Past Smokeless	47 (9.7)		
Current Smokeless	360 (74.1)		
<b>Smokeless tobacco uses per day (count)</b>			
Continuous	486	3.6 ± 3.5	1.3
Among users		4.9 ± 3.9	
<b>Waist circumference (cm)</b>			
Continuous	486	87.6 ± 14.1	1
<b>Body Mass Index (kg/m<sup>2</sup>)</b>			
Continuous	486	26.7 ± 5.7	1.1
<b>Plasmatic HDL cholesterol (mg/Dl) <sup>b</sup></b>			
Continuous	486	62.9 ± 18.1	0.9
<b>Plasmatic triglycerides (mg/Dl)</b>			
Continuous	486	82.4 ± 44.3	3.2
<b>Fasting plasmatic glucose (mg/Dl)</b>			
Continuous	486	90.8 ± 9.8	0.3
<b>Hemoglobin A1C (%)</b>			
Continuous	486	5.5 ± 0.3	0.5
<b>Systolic blood pressure (mm Hg)</b>			
Continuous	486	116.3 ± 13.4	0.7
<b>Diastolic blood pressure (mm Hg)</b>			
Continuous	486	68.0 ± 9.5	0.3
<b>Plasmatic LDL cholesterol (mg/Dl)</b>			
Continuous	486	124.7 ± 34.8	0.7

<sup>a</sup>Pattern scores are calculated from food frequencies [65]

<sup>b</sup>Variable is reverse coded

<https://doi.org/10.1371/journal.pone.0183451.t002>

Lipids (0.39), Adiposity -> Blood Pressure (0.23), Yup'ik culture -> Blood Pressure (-0.21), Yup'ik Culture -> Adiposity (-0.19), Western Culture -> Processed food, and Yup'ik Culture -> Subsistence Food (0.19). Fig 2 shows the relationships of age and gender to all other components. Here we see that age had the strongest associations (coefficient size) with other latent

**Table 3. Components of the metabolic syndrome.**

	Adiposity	Blood Pressure	LDL Lipids	Glycemia	TG-HDL Lipids
<b>Variance partition</b>					
SS loadings	2.03	1.67	1.34	1.27	1.24
Proportion Var	0.23	0.19	0.15	0.14	0.14
Cumulative Var	0.23	0.41	0.56	0.70	0.84
<b>Loadings</b>					
Waist Circ.	<b>0.93</b>	0.14	0.06	0.15	0.18
BMI	<b>0.96</b>	0.09	0.04	0.10	0.13
LDL-C	0.19	0.05	<b>0.83</b>	0.08	0.27
- HDL-C	0.21	-0.06	-0.66	0.04	<b>0.49</b>
Triglycerides	0.20	0.10	0.07	0.10	<b>0.87</b>
Systolic BP	-0.03	<b>0.93</b>	0.06	0.04	-0.02
Diastolic BP	0.28	<b>0.87</b>	0.04	0.02	0.14
Glucose	0.11	0.01	-0.12	<b>0.86</b>	0.25
HbA1C	0.17	0.07	0.44	<b>0.69</b>	-0.19

Five components explain 84% of the variability of 9 variables in a PCA analysis.

<https://doi.org/10.1371/journal.pone.0183451.t003>

variables, e.g., with Processed Food (-0.51), Subsistence Food (0.49), LDL lipids (0.47), and Glycemia (0.46) respectively. Gender also had a strong reverse association with Smoke Tobacco (-0.35).

The results of the model show that (1) education had positive associations with processed food, adiposity, and LDL-lipids, and an inverse association with tobacco smoking; (2) economic status had an inverse association with smokeless tobacco use; (3) Yup'ik culture had positive associations with subsistence food and smokeless tobacco, and inverse associations with TG-HDL Lipids, adiposity, and blood pressure; and (4) Western culture had a positive association with processed food, and negative associations with subsistence food and blood pressure.

We also found significant paths afferent to some of the cardiometabolic risk components: (1) subsistence food had a positive association with LDL lipids; (2) smokeless tobacco and physical activity had inverse associations with adiposity; and (3) adiposity had positive associations with all other cardiometabolic risk components.

The overall goodness-of-fit criterion ( $0 < \text{GoF} < 1$ ; (Tenenhaus et al., 2005) for this model is calculated as the geometric mean of the average communality and the average R2 value, and represents the average predictive power of the model.  $\text{GoF} = .35$  barely falls short of the cut-off value of .36 proposed by Wetzels et al. [84] for a large effect size, but also indicates a substantial proportion of variability not captured by the model.

The complete direct path and total effect coefficients (including non-significant relationships), their standard errors, and 95% confidence intervals or  $p < .05$  level are available in the Table in S1 Table.

**Interaction effects.** Table 6 presents coefficient, SE, and 95% CI of the direct paths and total effects for the interaction of Yup'ik and Western cultures. Fig 3 visualizes the interaction for the two variables where the interaction had a significant total effect. Each plot represents the predicted level of the dependent variable, LDL Lipids or Blood Pressure, as a function of the Yup'ik cultural participation level for the minimum and maximum Western cultural participation levels. We see that for the lowest Western culture level, there is a positive association of Yup'ik culture with LDL Lipids (the slope for the black continuous line increases showing

**Table 4. Weights, loadings, communalities and unidimensionality for outer Partial Least Squares Path Model (n = 486).**

Variables	weight	loading	communality	C.alpha	DG.rho	eig.1st	eig.2nd
<b>Gender</b>				1.000	1.000	1.000	0.000
Gender	1.000	1.000	1.000				
<b>Age</b>					1.000	1.000	1.000
Age	1.000	1.000	1.000				
<b>Education</b>				1.000	1.000	1.000	0.000
Years of HS + college	1.000	1.000	1.000				
<b>Economic status</b>				0.732	0.882	1.577	0.423
Food money shortage (rev)	0.592	0.900	0.810				
Utility money shortage (rev)	0.534	0.875	0.766				
<b>Western culture</b>				0.481	0.794	1.317	0.683
English at home	0.822	0.938	0.879				
Western lifestyle	0.366	0.627	0.393				
<b>Yup'ik culture</b>				0.517	0.757	1.532	0.832
Yup'ik at Home	0.536	0.795	0.632				
Yup'ik lifestyle	0.482	0.693	0.481				
Can speak Yup'ik	0.374	0.642	0.412				
<b>Subsistence food</b>				0.483	0.795	1.318	0.682
δ15N	0.847	0.951	0.904				
SF dietary pattern	0.327	0.596	0.355				
<b>Processed food</b>				0.418	0.775	1.264	0.736
δ13C	0.461	0.666	0.443				
PF dietary pattern score	0.774	0.896	0.802				
<b>Fruits and vegetables</b>				1.000	1.000	1.000	0.000
FV dietary pattern score	1.000	1.000	1.000				
<b>Physical activity</b>				0.544	0.763	1.838	1.008
Counts/day	0.526	0.867	0.752				
Mod./vig. activity	0.497	0.513	0.263				
Rest time (rev)	0.368	0.784	0.615				
<b>Smoke tobacco</b>				0.735	0.883	1.581	0.419
Never/past/current user	0.685	0.936	0.876				
Counts/day	0.432	0.830	0.689				
<b>Smokeless tobacco</b>				0.740	0.885	1.587	0.413
Never/past/current user	0.465	0.849	0.720				
Counts/day	0.653	0.927	0.858				
<b>Adiposity</b>				0.959	0.980	1.921	0.079
Waist circ.	0.517	0.980	0.961				
BMI	0.504	0.979	0.959				
<b>TG-HDL Lipids</b>				0.432	0.779	1.275	0.725
HDL-C (rev)	0.937	0.985	0.971				
Triglycerides	0.177	0.434	0.189				
<b>Glycemia</b>				0.469	0.790	1.307	0.693
Glucose	0.109	0.404	0.163				
HbA1C	0.961	0.995	0.989				
<b>Blood Pressure</b>				0.802	0.910	1.669	0.331
Systolic BP	0.613	0.934	0.873				
Diastolic BP	0.480	0.890	0.793				
<b>LDL Lipids</b>				1.000	1.000	1.000	0.000

(Continued)

Table 4. (Continued)

Variables	weight	loading	communality	C.alpha	DG.rho	eig.1st	eig.2nd
LDL-C	1.000	1.000	1.000				

Note. Weight = relative contribution of each manifest variable to the aligned latent variable; outer model fitting characteristics (recommended values) are: Loading = association of each manifest variable and the aligned latent variable ( $\geq 0.7$ ); Communality = squared loadings representing proportion of variance in each manifest variable captured by the aligned latent variable ( $\geq 0.5$ );  $\alpha$  = Cronbach's *alpha* ( $>0.7$ );  $\rho$  = Dillon-Goldstein's *rho* ( $>0.7$ ); 1Eig = 1<sup>st</sup> eigenvalue ( $>1$ ); 2Eig = 2<sup>nd</sup> eigenvalue (recommended  $< 1$ ). Variables labels in **bold** are latent variables; variables labels in plain text are manifest variables.

<https://doi.org/10.1371/journal.pone.0183451.t004>

that greater Yup'ik cultural participation predicts greater relative LDL Lipids levels), whereas for the highest level of Western cultural participation, the association is inverted (the slope for the red dashed line is negative, showing that greater Yup'ik cultural participation predicts lower relative LDL Lipid levels). Below we discuss this effect as indicating a qualified but significant protective effect of bi-culturalism. The main effect shows a slightly negative association between the two. We observe a similar result for blood pressure. Here the positive slope for lowest level of Western culture is not as steep as the one of LDL Lipids, indicating a weaker association between Yup'ik cultural participation and blood pressure than with LDL Lipids for lowest levels of Western culture.

### Discussion


Most of these results point to complex associations between culture, behavior, and cardiometabolic health. The associations of education (often thought as a source of assimilation [85]) with

Table 5. Total significant effects coefficients, for inner Partial Least Squares Path Model, with Bootstrap Standard Error and 95% confidence intervals (n = 486).

Path	Coeff.	CI		Path	Coeff.	CI	
Gender -> Physical activity	<b>-0.202*</b>	-0.285	-0.121	Education -> Smoke tobacco	<b>-0.069*</b>	-0.138	0.000
Gender -> Smoke tobacco	<b>-0.350*</b>	-0.422	-0.274	Education -> Adiposity	<b>0.100*</b>	0.023	0.184
Gender -> Smokeless tobacco	<b>0.137*</b>	0.048	0.227	Education -> LDL Lipids	<b>0.124*</b>	0.025	0.215
Gender -> Adiposity	<b>0.134*</b>	0.052	0.220	Economic status -> Smokeless tobacco	<b>-0.136*</b>	-0.225	-0.045
Gender -> TG-HDL Lipids	<b>-0.245*</b>	-0.318	-0.174	Western culture -> Subsistence food	<b>-0.175*</b>	-0.260	-0.077
Gender -> Blood Pressure	<b>-0.272*</b>	-0.357	-0.182	Western culture -> Processed food	<b>0.185*</b>	0.066	0.289
Age -> Economic status	<b>-0.169*</b>	-0.264	-0.074	Western culture -> Blood Pressure	<b>-0.120*</b>	-0.241	-0.001
Age -> Western culture	<b>-0.216*</b>	-0.294	-0.135	Yup'ik culture -> Subsistence food	<b>0.185*</b>	0.081	0.294
Age -> Yup'ik culture	<b>0.318*</b>	0.239	0.401	Yup'ik culture -> Smokeless tobacco	<b>0.170*</b>	0.030	0.322
Age -> Subsistence food	<b>0.492*</b>	0.422	0.564	Yup'ik culture -> Adiposity	<b>-0.188*</b>	-0.335	-0.026
Age -> Processed food	<b>-0.513*</b>	-0.574	-0.434	Yup'ik culture -> TG-HDL Lipids	<b>-0.145*</b>	-0.256	-0.036
Age -> Fruits and vegetables	<b>-0.198*</b>	-0.285	-0.112	Yup'ik culture -> Blood Pressure	<b>-0.205*</b>	-0.332	-0.075
Age -> Physical activity	<b>-0.320*</b>	-0.397	-0.234	Subsistence food -> LDL Lipids	<b>0.159*</b>	0.056	0.259
Age -> Smoke tobacco	<b>-0.185*</b>	-0.267	-0.110	Physical activity -> Adiposity	<b>-0.133*</b>	-0.211	-0.050
Age -> Smokeless tobacco	<b>0.116*</b>	0.015	0.199	Smokeless tobacco -> Adiposity	<b>-0.103*</b>	-0.187	-0.018
Age -> Adiposity	<b>0.202*</b>	0.121	0.288	Adiposity -> TG-HDL Lipids	<b>0.393*</b>	0.323	0.465
Age -> TG-HDL Lipids	<b>-0.413*</b>	-0.475	-0.333	Adiposity -> Glycemia	<b>0.169*</b>	0.075	0.275
Age -> Glycemia	<b>0.469*</b>	0.375	0.539	Adiposity -> Blood Pressure	<b>0.232*</b>	0.130	0.330
Age -> Blood Pressure	<b>0.311*</b>	0.230	0.391	Adiposity -> LDL Lipids	<b>0.159*</b>	0.066	0.268
Age -> LDL Lipids	<b>0.469*</b>	0.402	0.539				

\* Significant effects ( $\alpha = 0.05$ )

<https://doi.org/10.1371/journal.pone.0183451.t005>



**Fig 2. PLS-PM relationships with total effect coefficients of gender and age with all other latent variables.** Significant effects only ( $\alpha = 0.05$ ). Note that Fig 1 and Fig 2 represent the same model and are separated only for ease of viewing. Plain black lines and red dotted line represent positive and negative associations respectively.

<https://doi.org/10.1371/journal.pone.0183451.g002>

processed food consumption, adiposity and LDL lipids suggests a potentially deleterious effect of western cultural participation, for example by exposing students to processed food lunches. However the absence of a significant path from processed food to LDL lipids suggests that either our processed food latent variable does not capture the specific processed foods that increase LDL-C, or that education is associated with LDL lipids through another unmeasured mechanism. It is also possible that more educated participants are more likely to have office jobs and have no time to obtain subsistence foods as we adjusted for income. The reverse



**Table 6. Direct paths and total effects coefficients of interaction between Western and Yup'ik culture, for inner Partial Least Squares Path Model, with Bootstrap Standard Error and 95% confidence intervals (n = 486).**


Path	Direct Path				Total Effect			
	Coefficient	Std.Error	CI low	CI High	Coefficient	Std.Error	CI low	CI High
Culture interaction -> Subsistence food	-0.038	0.047	-0.126	0.054	-0.038	0.047	-0.126	0.054
Culture interaction -> Processed food	0.047	0.053	-0.058	0.151	0.047	0.053	-0.058	0.151
Culture interaction -> Fruits and vegetables	-0.013	0.065	-0.132	0.119	-0.013	0.065	-0.132	0.119
Culture interaction -> Physical activity	-0.053	0.071	-0.194	0.084	-0.053	0.071	-0.194	0.084
Culture interaction -> Smoke tobacco	0.060	0.062	-0.067	0.176	0.060	0.062	-0.067	0.176
Culture interaction -> Smokeless tobacco	0.020	0.065	-0.097	0.156	0.020	0.065	-0.097	0.156
Culture interaction -> Adiposity	-0.118	0.068	-0.255	0.018	-0.114	0.068	-0.249	0.016
Culture interaction -> TG-HDL Lipids	-0.054	0.048	-0.151	0.044	-0.092	0.052	-0.194	0.012
Culture interaction -> Glycemia	0.006	0.056	-0.097	0.113	-0.019	0.058	-0.128	0.091
Culture interaction -> Blood Pressure	-0.085	0.055	-0.188	0.022	<b>-0.112*</b>	0.055	-0.222	-0.003
Culture interaction -> LDL Lipids	<b>-0.183*</b>	0.060	-0.296	-0.061	<b>-0.211*</b>	0.060	-0.327	-0.089

\* Significant effects ( $\alpha = 0.05$ )

<https://doi.org/10.1371/journal.pone.0183451.t006>

association of education with smoked tobacco however may represent possible protection from cardiometabolic risk, even though associations between smoked tobacco and cardiometabolic risk are not captured by our model. This could be explained by its relatively low prevalence compared to smokeless tobacco (34.4% of current smokers, vs. 74.1% of smokeless users), because the cardiovascular risk is thought to be lower for smokeless than smoked tobacco [86].

Economic status on the other hand is inversely associated with smokeless tobacco use. The association of smokeless tobacco use with Yup'ik culture and inversely with economic status is suggestive of its role as a coping mechanism in Yup'ik culture [44]. The inverse association of



**Fig 3. Model predictions of LDL and blood pressure for Western and Yup'ik cultures interaction by level of Yup'ik culture.** Black full line represents the effect of Yup'ik culture for the lowest level of Western culture; purple dashed line represents the effect of Yup'ik culture for the highest level of Western culture; Red dotted line represents the main effect.

<https://doi.org/10.1371/journal.pone.0183451.g003>

smokeless tobacco with adiposity, but not with any of the other cardiometabolic risk components, could be interpreted as protective of cardiometabolic risk; however, *iqmik*, the main contributor to smokeless tobacco use for our participants, also has deleterious effects on health because of its high content in nicotine and multiple carcinogens [86,87]. The results show no other significant associations for economic status, including no direct significant relationship with processed foods. This is somewhat surprising as others have found associations with receiving food stamps or other government assistance and the consumption of processed foods [88,89].

Overall we did not find that increased education and economic status provided potential protective effects on cardiometabolic risk, in contrast with the health disparities reported in studies of non-Native populations [90,91], but consistent with findings in populations in early epidemiologic transition [92,93]. This may be due in part to the fact that there is less disparity in education and economic status in our population compared to other Arctic regions [37,94], which may impact access to traditional food sources in Alaska [95]. The Gini coefficient of income inequality in Alaska is the lowest of all states, at 0.41 vs. 0.47 nationwide; similarly, while proportions of people with 8<sup>th</sup> grade education are similar in our study communities and the U.S. (7% and 5% respectively), the proportion of people with college education is much smaller in our study communities (13% vs. 56%) [96]. It is also possible that the communal aspect of the Yup'ik way of life reflected by the importance of community connections [63], where people in need may receive more support than in the general population. Such processes could moderate the deleterious effect of low income and education seen in non-Native studies or studies from other Arctic regions.

The associations of our Yup'ik Culture latent variable to smokeless tobacco and subsistence food consumption are not surprising since those behaviors can be considered part of Yup'ik culture. In addition, we found beneficial associations of Yup'ik culture participation and the TG-HDL lipids latent variable (HDL-C is reverse coded), adiposity, and blood pressure. Yup'ik culture participation has the largest beneficial association with adiposity of all model inputs, and adiposity serves as the key mediating variable for all of the health outcomes in the study. These results are compatible with the previous findings that Yup'ik people have lower prevalence of diabetes than the US average, despite a comparable rate of obesity [61,97,98]. We do not find an association between TG-HDL lipids and any of the dietary variables, possibly because we are evaluating a component combining HDL-C (reversed coded) and triglycerides, unlike other studies that looked at them individually. However, the path Yup'ik culture  $\rightarrow$  TG-HDL Lipids shows a substantial difference between direct and total effects (-0.06, ns direct, and 0.15 total, s), indicating the mediating role of adiposity. The inverse association of Yup'ik culture and TG-HDL lipids that we found was previously undescribed as such, although a strong inverse association between traditional food intake and TG, was described using a biomarker of traditional food intake (the  $\delta^{15}\text{N}$  value; [69] and FFQ dietary patterns [39]. The only potentially deleterious association of Yup'ik culture that we found is indirect, through its association with subsistence food consumption, itself associated with LDL lipids. The positive association of marine based traditional food with LDL-C had been previously described in our reference population, e.g., in [69], and is consistent with reported associations of LDL-C with  $\omega$ -3 fatty acid supplementation [99], and of an experiment on the consumption of  $\omega$ -3 rich fish associated with LDL-C [100]. We did not find the positive association of Yup'ik enculturation with physical activity reported by [42], possibly because we used different measures of physical activity and enculturation.

The association of Western culture with the consumption of processed foods and inverse association with the consumption of subsistence foods were expected as they had been reported previously [65]; in addition, we found an inverse association of Western cultural

participation with blood pressure which may be in part explained by an increased level of health literacy [101] and could have implications for interventions by including a component aimed at increasing health literacy. We did not find an association of the consumption of subsistence foods with blood pressure similar to what was reported by Beaulieu-Jones et al. [102].

Importantly, we found indications of interaction effects between the cultural participation variables, indicating a potential effect of biculturalism on cardiometabolic health. The effects and interactions of differential cultural participation are unevenly distributed; for example, the interaction of Yup'ik and Western cultural participation shows that increasing participation in both potentially has important health benefits including lower LDL Lipids and lower blood pressure. Therefore, the beneficial health effects of following a Yup'ik way of life are most pronounced for those who identify with high levels of both Yup'ik and Western cultural practices. Interestingly, the fact that the association of Yup'ik culture with LDL lipids is in opposite direction for high and low levels of Western culture, represented by a significant interaction, explains why we do not find a significant main effect of Yup'ik culture on LDL lipids. The potentially beneficial role of biculturalism on blood pressure is similar to what was described before [46], but it is a new finding for LDL Lipids.

Finally, the results raise the question of whether and to what extent the associations to cardiometabolic risk factors are underwritten by both life course and gender roles. While female gender is associated with greater subsistence food use *and* adiposity, women show lower levels of both LDL lipids and blood pressure, despite adiposity's overall positive association with both of these factors. A protective effect of estrogens on LDL lipids and blood pressure is possible [103]. Of note, gender on its own did not figure significantly in cultural participation for either Yup'ik or Western culture. Age, on the other hand, was associated with nearly all of the model components, including with Yup'ik culture and inversely with Western culture. The assumed implications of this pattern for food consumption hold. Increasing age (and thus greater adherence to a Yup'ik culture and lesser participation in Western culture) is also significantly associated with large increases in the consumption of subsistence foods, and a decreased consumption of processed foods. These same factors contributed to increases in adiposity, blood pressure and LDL lipids.

Clearly the role of social determinants of health is complex and understanding those complexities could be key to improving cardiometabolic health. Mixed economies in indigenous communities have featured elements of both traditional subsistence food harvests and consumption of commercially available mass-produced foods. While thought to be a transitional state that would bridge indigenous communities' transition to a fully "modern" economy, combined food economies are now seen as a complex and ongoing set of food strategies that are unlikely to leave their "mixed" state at any time soon [37,53,95,104]. This points to the importance of preserving the subsistence activities embedded in Yup'ik culture participation from a community sustainability standpoint. Our results emphasize some of the potential health benefits of bicultural participation that may accompany this situation.

Other studies have also found varied impacts of acculturation on cardiometabolic health. For example, a study on the Inuit population in Greenland found that westernization was associated with the metabolic syndrome in men, attributed to a decrease of physical activity due to diminishing hunting and fishing, and an inverse association for women attributed to increase education [105]. A study in Hawaii found that people identifying highly with traditional, but not western culture, were more likely to have T2D [106]. A meta-analysis found that acculturation among Latinos in the United States had both a negative influence for certain health outcomes like substance abuse, dietary practices, and birth outcomes, but a positive one on health care use and access [107]. Another study found that Hispanics with lower levels of

acculturation were more likely to have poorly controlled LDL cholesterol. The commonality of these results with ours suggests that health literacy is an important factor in the beneficial effect of Western culture. The divergence in the effects of adhering to the minority culture may be due in part to the fact that most studies, including ours, do not take into account the interaction between individuals and the social structure that they are embedded in, reflecting the fact that a relational approach to the agency-structure divide has not gained traction in health research [108]. Some novel frameworks, however, may help, such as substantialist approaches (modeling things rather than relations), a focus on cultural capital [21,109], or capital interplay [110,111].

Strengths of our study include the ability to combine socio-cultural, behavioral, and cardiometabolic risk factors in a single model; the use of an orthogonal model of culture, with both Yup'ik and Western latent variables; and the use of a novel methodology. Limitations include its observational and cross-sectional nature of the study, limiting causal interpretations, and the lack of a measure of stress (perceived or through biomarkers)—although our measure of economic status (running out of food for money or utilities) likely captured some element of stress. Limitations also include relatively limited measures of Yup'ik culture and economic status, a fitting of our outer model which is not perfect, difficulty of generalization even to other Alaska Native cultures, the inability to account for clustering at the community level in PLS-PM, a convenience sample, and the fact that we look at cardiometabolic risk factors rather than disease outcomes. Another important limitation is the choice of Yup'ik language use as a proxy for traditional cultural participation. While this strategy finds support in prior studies [112,113], it remains an indirect measure. In addition, this choice does not address questions of cultural consonance, or the extent to which individuals actually subscribe to identified cultural beliefs [114]. Cultural consonance has been shown to be an important consideration in indigenous health research [115], including work on cardiometabolic-related health [116]. The limitation that is most likely to have changed the interpretation of our results is the lack of measure of stress, since previous research points to the importance of stress both among our reference population [45], and as a determinant of cardiometabolic health in the general population [117]. A sensitivity analysis using multiple imputation rather than median values to replace the missing values, showed very little change in the results.

Overall, our analysis suggests that Yup'ik culture participation is an important determinant of cardiometabolic health in rural communities of southwestern Alaska. Bi-culturalism, demonstrated by the beneficial moderating effect of Western culture participation on the relationship between Yup'ik culture participation and blood pressure as well as LDL-cholesterol, was also significant. Although our analysis did not reproduce the potentially beneficial association between traditional food consumption and cardiometabolic risk factors found in previous studies, it uncovered significant potentially beneficial effects of Yup'ik culture participation on blood pressure, adiposity, and TG-HDL Lipids. Given the current understanding that mixed food economies are a permanent, rather than temporary, condition for many northern indigenous communities, such findings are important for long-term as well as short-term health planning. Our results suggest that reinforcing the protective effects of both Yup'ik and Western cultures could be useful in interventions, but new research involving Yup'ik collaborators, and harnessing knowledge gained from Yup'ik authors, e.g., [40,118], is needed to better understand the mechanisms underlying the effects of Yup'ik culture on cardiometabolic health. Such research would likely begin with questions of how stress can be mitigated by traditional practices and beliefs, and how health literacy may influence cardiometabolic health outcomes in bicultural settings.

## Supporting information

**S1 Table. Complete direct and total path coefficients for inner Partial Least Squares Path Model, with Bootstrap Standard Error and 95% confidence intervals (n = 486).**  
(PDF)

## Acknowledgments

We would like to thank the staff of the Center for Alaska Native Health Research, our tribal partners and community research assistants from the Yukon-Kuskokwim Delta, and most of all, our participants from Yup'ik communities.

## Author Contributions

**Conceptualization:** Jacques Philip, Scarlett E. Hopkins, Kenneth E. Thummel, Bert B. Boyer, Kirk Dombrowski.

**Data curation:** Jacques Philip, Scarlett E. Hopkins.

**Formal analysis:** Jacques Philip, Kirk Dombrowski.

**Funding acquisition:** Scarlett E. Hopkins, Kenneth E. Thummel, Bert B. Boyer.

**Investigation:** Scarlett E. Hopkins, Bert B. Boyer.

**Methodology:** Jacques Philip, Tove K. Ryman, Scarlett E. Hopkins, Diane M. O'Brien, Andrea Bersamin, Jeremy Pomeroy, Melissa A. Austin, Bert B. Boyer, Kirk Dombrowski.

**Project administration:** Jacques Philip, Scarlett E. Hopkins, Bert B. Boyer, Kirk Dombrowski.

**Resources:** Scarlett E. Hopkins, Diane M. O'Brien, Bert B. Boyer.

**Software:** Jacques Philip.

**Supervision:** Scarlett E. Hopkins, Kenneth E. Thummel, Bert B. Boyer, Kirk Dombrowski.

**Validation:** Tove K. Ryman, Scarlett E. Hopkins, Kenneth E. Thummel, Melissa A. Austin, Bert B. Boyer, Kirk Dombrowski.

**Visualization:** Jacques Philip.

**Writing – original draft:** Jacques Philip, Kirk Dombrowski.

**Writing – review & editing:** Jacques Philip, Tove K. Ryman, Scarlett E. Hopkins, Diane M. O'Brien, Andrea Bersamin, Jeremy Pomeroy, Kenneth E. Thummel, Melissa A. Austin, Bert B. Boyer, Kirk Dombrowski.

## References

1. Kirmayer LJ, Boothroyd LJ, Hodgins S. Attempted suicide among Inuit youth: psychosocial correlates and implications for prevention. *Can J Psychiatry*. 1998 Oct; 43[8]:816–22. <https://doi.org/10.1177/070674379804300806> PMID: 9806088
2. Sutton P. *The Politics of Suffering: Indigenous Australia and the End of the Liberal Consensus*. New editio. Carlton, Vic: Melbourne University Publishing; 2011. 304 p.
3. Valeggia CR, Snodgrass JJ. Health of Indigenous Peoples. *Annu Rev Anthropol*. 2015; 44:117–135.
4. Roh S, Burnette CE, Lee KH, Lee Y-S, Easton SD, Lawler MJ. Risk and protective factors for depressive symptoms among American Indian older adults: Adverse childhood experiences and social support. *Aging Ment Health*. 2015;
5. Fraser SL, Geoffroy D, Chachamovich E, Kirmayer LJ. Changing Rates of Suicide Ideation and Attempts Among Inuit Youth: A Gender-Based Analysis of Risk and Protective Factors. *Suicide life-threatening Behav*. 2015; 45[2]:141–156.

6. Silvikén A, Kvernmo S. Suicide attempts among indigenous Sami adolescents and majority peers in Arctic Norway: prevalence and associated risk factors. *J Adolesc.* 2007 Aug; 30[4]:613–26. <https://doi.org/10.1016/j.adolescence.2006.06.004> PMID: 16876860
7. Hautala DS, Sittner KJ, Whitbeck LB. Prospective Childhood Risk Factors for Gang Involvement Among North American Indigenous Adolescents. *Youth Violence Juv Justice.* 2015 May; 15:41204015585173.
8. Whitbeck LB, Crawford DM. Gestational risks and psychiatric disorders among indigenous adolescents. *Community Ment Health J.* 2009; 45[1]:62–72. <https://doi.org/10.1007/s10597-008-9172-5> PMID: 18998209
9. Blue Bird Jernigan V, Peercy M, Branam D, Saunkeah B, Wharton D, Winkleby M, et al. Beyond Health Equity: Achieving Wellness Within American Indian and Alaska Native Communities. *Am J Public Health.* 2015; 105[S3]:S376–S379.
10. Douglas V. Introduction to Aboriginal Health and Health Care in Canada: Bridging Health and Healing. Springer Publishing Company; 2013. 257 p.
11. Gone JP, Trimble JE. American Indian and Alaska Native Mental Health: Diverse Perspectives on Enduring Disparities. *Annu Rev Clin Psychol.* 2012; 8[1]:131–60.
12. Wexler L. Inupiat youth suicide and culture loss: Changing community conversations for prevention. *Soc Sci Med.* 2006 Dec; 63[11]:2938–48. <https://doi.org/10.1016/j.socscimed.2006.07.022> PMID: 16952416
13. Kaufman CE, Beals J, Croy C, Jiang L, Novins DK. Multilevel context of depression in two American Indian tribes. *J Consult Clin Psychol.* 2013; 81[6]:1040. <https://doi.org/10.1037/a0034342> PMID: 24016293
14. Joe J, Young R. Introduction: Diabetes programs and the need for cultural capital. *Am Indian Cult Res J.* UCLA American Indian Studies Center; 2014; 38[1]:11–28.
15. Mitrou F, Cooke M, Lawrence D, Povah D, Mobilia E, Guimond E, et al. Gaps in Indigenous disadvantage not closing: a census cohort study of social determinants of health in Australia, Canada, and New Zealand from 1981–2006. *BMC Public Health.* 2014; 14:201. <https://doi.org/10.1186/1471-2458-14-201> PMID: 24568143
16. Arias E, Xu J, Jim MA. Period Life Tables for the Non-Hispanic American Indian and Alaska Native Population, 2007–2009. *Am J Public Health.* 2014 Apr; 104[S3]:S312–9.
17. Tester FJ, Kulchyski PK. Tammarniit (mistakes): Inuit relocation in the Eastern Arctic, 1939–63. UBC Press; 1994. 443 p.
18. Brice-Bennett C. Dispossessed: The Eviction of Inuit from Hebron Labrador (A report to North Program, Royal Commission on Aboriginal Peoples). Labrador Institute, Memorial University, Newfoundland Canada; 1994.
19. Gone JP, Calf Looking PE. American Indian Culture as substance abuse treatment: Pursuing evidence for a local intervention. *J Psychoactive Drugs.* 2011; 43[4]:291–6. <https://doi.org/10.1080/02791072.2011.628915> PMID: 22400459
20. Moore K, Jiang L, Manson SM, Beals J, Henderson W, Pratte K, et al. Case management to reduce cardiovascular disease risk in American Indians and Alaska Natives with diabetes: results from the Special Diabetes Program for Indians Healthy Heart Demonstration Project. *Am J Public Health.* 2014; 104[11]:e158–e164. <https://doi.org/10.2105/AJPH.2014.302108> PMID: 25211728
21. Joe J, Young R. Promoting cultural capital in a medical camp for American Indian youth with diabetes. *Am Indian Cult Res J.* 2014; 38[1]:123–144.
22. MacDonald JP, Ford JD, Willox AC, Ross NA. A review of protective factors and causal mechanisms that enhance the mental health of Indigenous Circumpolar youth. *Int J Circumpolar Health.* 2013; 72.
23. Allen J, Mohatt G V., Fok CCT, Henry D, Burkett R. A protective factors model for alcohol abuse and suicide prevention among Alaska Native youth. *Am J Community Psychol.* 2014 Sep; 54[1–2]:125–39. <https://doi.org/10.1007/s10464-014-9661-3> PMID: 24952249
24. Wexler L, Gubrium A, Griffin M, DiFulvio G. Promoting positive youth development and highlighting reasons for living in Northwest Alaska through digital storytelling. *Health Promot Pract.* Sage Publications; 2013 Jul 1; 14[4]:617–23.
25. Currie CL, Wild TC, Schopflocher DP, Laing L, Veugelers P. Illicit and prescription drug problems among urban Aboriginal adults in Canada: the role of traditional culture in protection and resilience. *Soc Sci Med.* 2013; 88:1–9. <https://doi.org/10.1016/j.socscimed.2013.03.032> PMID: 23702204
26. Philip J, Ford T, Henry D, Rasmus S, Allen J. Relationship of social network to protective factors in suicide and alcohol use disorder intervention for rural Yup'ik Alaska Native youth. *Psychosoc Interv.* Elsevier; 2016 Jan 1; 25[1]:45–54.

27. Reeve R, Church J, Haas M, Bradford W, Viney R. Factors that drive the gap in diabetes rates between Aboriginal and non-Aboriginal people in non-remote NSW. *Aust N Z J Public Health*. 2014; 38[5]:459–465. <https://doi.org/10.1111/1753-6405.12211> PMID: 25168646
28. Gone JP. “We never was happy living like a whiteman”: Mental health disparities and the postcolonial predicament in American Indian communities. *Am J Community Psychol*. 2007; 40[3–4]:290–300. <https://doi.org/10.1007/s10464-007-9136-x> PMID: 17906926
29. Kirmayer L, Simpson C, Cargo M. Healing traditions: culture, community and mental health promotion with Canadian Aboriginal peoples. *Australas Psychiatry*. 2003; 11:S15–S23.
30. MacDonald JP, Willox AC, Ford JD, Shiwak I, Wood M, Team I. Protective factors for mental health and well-being in a changing climate: Perspectives from Inuit youth in Nunatsiavut, Labrador. *Soc Sci Med*. 2015; 141:133–141. <https://doi.org/10.1016/j.socscimed.2015.07.017> PMID: 26275362
31. Cheadle JE, Whitbeck LB. Alcohol Use Trajectories and Problem Drinking Over the Course of Adolescence A Study of North American Indigenous Youth and Their Caretakers. *J Health Soc Behav*. 2011 Jun; 52[2]:228–45. <https://doi.org/10.1177/0022146510393973> PMID: 21558489
32. Spicer P. Culture and the restoration of self among former American Indian drinkers. *Soc Sci Med*. 2001 Jul; 53[2]:227–40. PMID: 11414389
33. Sider GM. *Skin for Skin: Death and Life for Inuit and Innu*. Duke University Press; 2014.
34. O’Neil TD, Mitchell CM. Alcohol use among American Indian adolescents: The role of culture in pathological drinking. *Soc Sci Med*. 1996 Feb; 42[4]:565–78. PMID: 8643981
35. Mitchell FM. Reframing diabetes in American Indian communities: A social determinants of health perspective. *Health Soc Work*. 2012; 37[2]:71–79. PMID: 23029974
36. Dombrowski K. *Culture Politics: The Story of Native Land Claims in Alaska*. Syron Design Academic Publishing; 2014. 254 p.
37. Dombrowski K, Channell E, Khan B, Moses J, Misshula E. Out on the Land: Income, Subsistence Activities, and Food Sharing Networks in Nain, Labrador. *J Anthropol*. 2013; 2013:1–11.
38. Espey DK, Jim MA, Cobb N, Bartholomew M, Becker T, Haverkamp D, et al. Leading causes of death and all-cause mortality in American Indians and Alaska Natives. *Am J Public Health*. 2014; 104 [SUPPL. 3]:303–11.
39. Ryman TK, Boyer BB, Hopkins S, Philip J, Beresford SAA, Thompson B, et al. Associations between diet and cardiometabolic risk among Yup’ik Alaska Native people using food frequency questionnaire dietary patterns. *Nutr Metab Cardiovasc Dis*. Elsevier B.V; 2015 Dec; 25[12]:1140–5.
40. Ayunerak P, Alstrom D, Moses C, Charlie J, Rasmus S. Yup’ik culture and context in southwest Alaska: Community member perspectives of tradition, social change, and prevention. *Am J Community Psychol*. 2014 Apr 26;
41. Napoleon H. Yuuyaraq: The Way of the Human Being. Madsen EC, editor. *Network, Alaska Native Knowledge*. University of Alaska, Fairbanks, College of Rural Alaska, Center for Cross-Cultural Studies; 1991.
42. Bersamin A, Wolsko C, Luick BR, Boyer BB, Lardon C, Hopkins SE, et al. Enculturation, perceived stress, and physical activity: implications for metabolic risk among the Yup’ik—The Center for Alaska Native Health Research Study. *Ethn Health*. Taylor & Francis; 2014; 19[3]:255–69.
43. Ryman TK, Boyer BB, Hopkins SE, Philip J, Thompson B, Beresford SAA, et al. Association between iq’mik smokeless tobacco use and cardiometabolic risk profile among Yup’ik Alaska Native people. *Ethn Health*. 2017 Jan 24; In Press:1–15.
44. Wolsko C, Mohatt G V, Lardon C, Burket R. Smoking, chewing, and cultural identity: prevalence and correlates of tobacco use among the Yup’ik-The Center for Alaska Native Health Research (CANHR) study. *Cultur Divers Ethnic Minor Psychol*. 2009; 15[2]:165–72. <https://doi.org/10.1037/a0015323> PMID: 19364203
45. Wolsko C, Lardon C, Mohatt G V, Orr E. Stress, coping, and well-being among the Yup’ik of the Yukon-Kuskokwim Delta: the role of enculturation and acculturation. *Int J Circumpolar Health*. 2007 Feb; 66[1]:51–61. PMID: 17451134
46. Boden-Albala B, Roberts ET, Hopkins S, Allen J, Boyer BB. Predictors of risk and protection for hypertension in Yup’ik people from Southwest Alaska. *Ethn Dis*. 2012; 23[4]:484–91.
47. Garrouette EM, Goldberg J, Beals J, Herrell R, Manson SM. Spirituality and attempted suicide among American Indians. *Soc Sci Med*. 2003 Apr; 56[7]:1571–9. PMID: 12614706
48. Parlee B, Berkes F, Gwich’in T. Health of the Land, Health of the People: A Case Study on Gwich’in Berry Harvesting in Northern Canada. *Ecohealth*. 2005; 2[2]:127–37.

49. Stone R a T, Whitbeck LB, Chen X, Johnson K, Olson DM. Traditional practices, traditional spirituality, and alcohol cessation among American Indians. *J Stud Alcohol*. 2006 Mar; 67[2]:236–44. PMID: [16562405](https://pubmed.ncbi.nlm.nih.gov/16562405/)
50. Chandler M, Lalonde CE. Cultural continuity as a hedge against suicide in Canada's First Nations. *Transcult Psychiatry*. 1998;1–20.
51. Hallett D, Chandler MJ, Lalonde CE. Aboriginal language knowledge and youth suicide. *Cogn Dev*. 2007; 22[3]:392–9.
52. Lazrus H. Risk perception and climate adaptation in Tuvalu: a combined cultural theory and traditional knowledge approach. *Hum Organ*. 2015; 74[1]:52–61.
53. Dombrowski K, Khan B, Channell E, Moses J, McLean K, Misshula E. Kinship, Family, and Exchange in a Labrador Inuit Community. *Arctic Anthropol*. 2013; 50[1]:89–104.
54. Moses J, Khan B, Gauthier GR, Ponizovsky V, Dombrowski K. Confounding Culture: Drinking, Country Food Sharing, and Traditional Knowledge Networks in a Labrador Inuit Community. *Hum Organ*. 2017; 76[2]:171–183.
55. Chandler MJ, Lalonde CE, Sokol BW, Hallett D, Marcia JE. Personal persistence, identity development, and suicide: A study of native and non-native North American adolescents. *Monogr Soc Res child Dev*. JSTOR; 2003;i—138.
56. Duran E, Duran B. *Native American Postcolonial Psychology*. SUNY Press; 1995. 252 p.
57. Allen J, Mohatt G V, Beehler S, Rowe HL. People Awakening: Collaborative research to develop cultural strategies for prevention in community intervention. *Am J Community Psychol*. 2014 Jun 6;
58. King M, Smith A, Gracey M. Indigenous health part 2: the underlying causes of the health gap. *Lancet*. 2009; 374[9683]:76–85. [https://doi.org/10.1016/S0140-6736\(09\)60827-8](https://doi.org/10.1016/S0140-6736(09)60827-8) PMID: [19577696](https://pubmed.ncbi.nlm.nih.gov/19577696/)
59. Lehti V, Niemelä S, Hoven C, Mandell D, Sourander A. Mental health, substance use and suicidal behaviour among young indigenous people in the Arctic: a systematic review. *Soc Sci Med*. Elsevier Ltd; 2009; 69[8]:1194–203.
60. Berger TR. *Village journey: the report of the Alaska Native Review Commission*. Hill and Wang; 1985.
61. Mohatt G V., Plaetke R, Klejka J, Luick B, Lardon C, Bersamin A. The Center for Alaska Native Health Research Study : a community -based participatory research study of obesity and chronic disease -related protective and risk factors. *Int J Circumpolar Health*. 2007; 66[1]:8–18. PMID: [17451130](https://pubmed.ncbi.nlm.nih.gov/17451130/)
62. Boyer BB, Mohatt G, Lardon C, Plaetke R, Luick B, Hutchison S, et al. Building a community-based participatory research center to investigate obesity and diabetes in Alaska Natives. *Int J Circumpolar Health*. 2005; 64[3].
63. Wolsko C, Lardon C, Hopkins S, Ruppert E. Conceptions of wellness among the Yup'ik of the Yukon-Kuskokwim Delta: the vitality of social and natural connection. *Ethn Health*. 2006 Nov; 11[4]:345–63. <https://doi.org/10.1080/13557850600824005> PMID: [17060033](https://pubmed.ncbi.nlm.nih.gov/17060033/)
64. Oetting ER, Beauvais F. Orthogonal cultural identification theory: The cultural identification of minority adolescents. *Int J Addict*. Taylor & Francis; 1991; 25[sup5]:655–85.
65. Ryman TK, Boyer BB, Hopkins S, Philip J, O'Brien D, Thummel K, et al. Characterizing the reproducibility and reliability of dietary patterns among Yup'ik Alaska Native people. *Br J Nutr*. 2015; 113 [4]:634–43. <https://doi.org/10.1017/S0007114514003596> PMID: [25656871](https://pubmed.ncbi.nlm.nih.gov/25656871/)
66. Ryman TK, Austin MA, Hopkins S, Philip J, O'Brien D, Thummel K, et al. Using exploratory factor analysis of FFQ data to identify dietary patterns among Yup'ik people. *Public Health Nutr*. 2014 Mar; 17 [3]:510–8. <https://doi.org/10.1017/S1368980012005411> PMID: [23290469](https://pubmed.ncbi.nlm.nih.gov/23290469/)
67. O'Brien D, Kristal AR, Jeannet MA, Wilkinson MJ, Bersamin A, Luick B. Red blood cell delta15N: a novel biomarker of dietary eicosapentaenoic acid and docosahexaenoic acid intake. *Am J Clin Nutr*. 2009 Mar; 89[3]:913–9. <https://doi.org/10.3945/ajcn.2008.27054> PMID: [19176727](https://pubmed.ncbi.nlm.nih.gov/19176727/)
68. Nash SH, Bersamin A, Kristal AR, Hopkins S, Church RS, Pasker RL, et al. Stable nitrogen and carbon isotope ratios indicate traditional and market food intake in an indigenous circumpolar population. *J Nutr*. 2012 Jan; 142[1]:84–90. <https://doi.org/10.3945/jn.111.147595> PMID: [22157543](https://pubmed.ncbi.nlm.nih.gov/22157543/)
69. O'Brien DM, Kristal AR, Nash SH, Hopkins SE, Luick BR, Stanhope KL, et al. A Stable Isotope Biomarker of Marine Food Intake Captures Associations between n-3 Fatty Acid Intake and Chronic Disease Risk in a Yup'ik Study Population, and Detects New Associations with Blood Pressure and. *J Nutr*. 2014; 144[5]:706–13. <https://doi.org/10.3945/jn.113.189381> PMID: [24598880](https://pubmed.ncbi.nlm.nih.gov/24598880/)
70. Renner CC, Enoch C, Patten CA, Ebbert JO, Hurt RD, Moyer TP, et al. Iqmik: a form of smokeless tobacco used among Alaska natives. *Am J Health Behav*. PNG Publications; 2005; 29[6]:588–94.
71. Morgan MS. *The history of econometric ideas*. Cambridge University Press; 1992.
72. Tenenhaus M. *Structural Equation Modelling for small samples*. Groupe HEC; 2008.



73. Tenenhaus M, Vinzi VE, Chatelin Y-M, Lauro C. PLS path modeling. *Comput Stat Data Anal.* 2005 Jan; 48[1]:159–205.
74. Henseler J, Ringle CM, Sarstedt M. Using partial least squares path modeling in advertising research: basic concepts and recent issues. In: *Handbook of research on international advertising.* 2012. p. 252.
75. McIntosh CN, Edwards JR, Antonakis J. Reflections on Partial Least Squares Path Modeling. *Organ Res Methods.* 2014; 17[2]:210–51.
76. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria; 2013. <http://www.r-project.org/>
77. Hanson RL, Imperatore G, Bennett PH, Knowler WC. Components of the “metabolic syndrome” and incidence of type 2 diabetes. *Diabetes.* 2002; 51:3120–7. PMID: [12351457](https://pubmed.ncbi.nlm.nih.gov/12351457/)
78. Wijndaele K, Beunen G, Duvigneaud N, Matton L, Duquet W, Thomis M, et al. A continuous Metabolic Syndrome Risk Score utility for epidemiological analyses. *Diabetes Care.* Am Diabetes Assoc; 2006; 29[10]:2329.
79. Edwards KL, Burchfiel CM, Sharp DS, Curb JD, Rodriguez BL, Fujimoto WY, et al. Factors of the insulin resistance syndrome in nondiabetic and diabetic elderly Japanese-American men. *Am J Epidemiol.* 1998; 147[5]:441–7. PMID: [9525530](https://pubmed.ncbi.nlm.nih.gov/9525530/)
80. Grundy SM, Brewer HB, Cleeman JI, Smith SC, Lenfant C. Definition of Metabolic Syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation.* 2004; 109[3]:433–8. <https://doi.org/10.1161/01.CIR.0000111245.75752.C6> PMID: [14744958](https://pubmed.ncbi.nlm.nih.gov/14744958/)
81. Zimmet P, Alberti G, Shaw J. A new IDF worldwide definition of the metabolic syndrome: the rationale and the results. *Diabetes Voice.* 2005; 50[3]:31–3.
82. Esposito Vinzi V, Chin WW, Henseler J, Wang H. *Handbook of Partial Least Squares: Concepts, Methods and Applications.* Heidelberg, Dordrecht, London, New York: Springer; 2010.
83. Sanchez G. PLS path modeling with R [Internet]. Online book. 2013. [http://gastonsanchez.com/PLS\\_Path\\_Modeling\\_with\\_R.pdf](http://gastonsanchez.com/PLS_Path_Modeling_with_R.pdf)
84. Wetzels M, Odekerken-Schroder G, Van Oppen C. Using PLS path modeling for assessing hierarchical construct models: guidelines and empirical illustration. *Manag Inf Syst Q.* 2009; 33[1]:11.
85. Larimore (Comanche) J a., McClellan GS. Native American student retention in U.S. postsecondary education. *New Dir Student Serv.* 2005; 2005[109]:17–32.
86. Piano MR, Benowitz NL, Fitzgerald GA, Corbridge S, Heath J, Hahn E, et al. Impact of smokeless tobacco products on cardiovascular disease: Implications for policy, prevention, and treatment: A policy statement from the American Heart Association. *Circulation.* 2010; 122[15]:1520–44. <https://doi.org/10.1161/CIR.0b013e3181f432c3> PMID: [20837898](https://pubmed.ncbi.nlm.nih.gov/20837898/)
87. Pappas RS, Stanfill SB, Watson CH, Ashley DL. Analysis of toxic metals in commercial moist snuff and Alaskan iqmik. *J Anal Toxicol.* 2008; 32[4]:281–91. PMID: [18430295](https://pubmed.ncbi.nlm.nih.gov/18430295/)
88. Blue Bird Jernigan V, Wetherill MS, Hearod J, Jacob T, Salvatore AL, Cannady T, et al. Food Insecurity and Chronic Diseases Among American Indians in Rural Oklahoma: The THRIVE Study. *Am J Public Health.* 2017 Jan; 107[3]:441–6. <https://doi.org/10.2105/AJPH.2016.303605> PMID: [28103070](https://pubmed.ncbi.nlm.nih.gov/28103070/)
89. Bauer KW, Widome R, Himes JH, Smyth M, Rock BH, Hannan PJ, et al. High Food Insecurity and Its Correlates Among Families Living on a Rural American Indian Reservation. *Am J Public Health.* 2012 May; 102[7]:1346–52. <https://doi.org/10.2105/AJPH.2011.300522> PMID: [22594740](https://pubmed.ncbi.nlm.nih.gov/22594740/)
90. Albert MA, Glynn RJ, Buring J, Ridker PM. Impact of traditional and novel risk factors on the relationship between socioeconomic status and incident cardiovascular events. *Circulation.* 2006; 114[24]:2619–26. <https://doi.org/10.1161/CIRCULATIONAHA.106.660043> PMID: [17116764](https://pubmed.ncbi.nlm.nih.gov/17116764/)
91. Mensah GA, Mokdad AH, Ford ES, Greenlund KJ, Croft JB. State of disparities in cardiovascular health in the United States. *Circulation.* 2005; 111[10]:1233–41. <https://doi.org/10.1161/01.CIR.0000158136.76824.04> PMID: [15769763](https://pubmed.ncbi.nlm.nih.gov/15769763/)
92. Omran AR. *The epidemiologic transition: a theory of the epidemiology of population change.* 1971. Milbank Q. Wiley Online Library; 2005; 83[4]:731–57.
93. Chen D, Rivera-Andrade Á, González J, Burt D, Mendoza-Montano C, Patrie J, et al. Prevalence of risk factors for noncommunicable diseases in an indigenous community in Santiago Atitlán, Guatemala. *Rev Panam Salud Publica.* 2017 Apr 20; 41[2]:e7.
94. Dombrowski K, Khan B, Moses J, Channell E, Dombrowski N. Network sampling of social divisions in a rural Inuit community. *Identities.* 2014; 21[2]:134–51.
95. BurnSilver S, Magdanz J, Stotts R, Berman M, Kofinas G. Are Mixed Economies Persistent or Transitional? Evidence Using Social Networks from Arctic Alaska. *Am Anthropol.* 2016 Mar; 118[1]:121–9.
96. U.S. Census Bureau. American FactFinder [Internet]. 2016. <http://factfinder.census.gov>

97. Boyer BB, Mohatt G V, Plaetke R, Herron J, Stanhope KL, Stephensen C, et al. Metabolic syndrome in Yup'ik Eskimos: the Center for Alaska Native Health Research (CANHR) Study. *Obesity* (Silver Spring). 2007 Nov; 15[11]:2535–40.
98. Koller KR, Metzger JS, Jolly SE, Umans JG, Hopkins SE, Kaufmann C, et al. Cardiometabolic correlates of low type 2 diabetes incidence in western Alaska Native people—The WATCH study. *Diabetes Res Clin Pract*. Elsevier; 2015; 108[3]:423–31.
99. Jacobson TA, Glickstein SB, Rowe JD, Soni PN. Effects of eicosapentaenoic acid and docosahexaenoic acid on low-density lipoprotein cholesterol and other lipids: A review. *J Clin Lipidol*. Mosby, Inc; 2012; 6[1]:5–18.
100. Gerhard GT, Patton BD, Lindquist SA, Wander RC. Comparison of three species of dietary fish: effects on serum concentrations of low-density-lipoprotein cholesterol and apolipoprotein in normotriglyceridemic subjects. *Am J Clin Nutr*. Am Soc Nutrition; 1991; 54[2]:334–9.
101. Williams M V, Baker DW, Parker RM, Nurss JR. Relationship of functional health literacy to patients' knowledge of their chronic disease. A study of patients with hypertension and diabetes. *Arch Intern Med*. American Medical Association; 1998 Jan 26; 158[2]:166–72.
102. Beaulieu-Jones BR, O'Brien DM, Hopkins SE, Moore JH, Boyer BB, Gilbert-Diamond D. Sex, adiposity, and hypertension status modify the inverse effect of marine food intake on blood pressure in Alaska Native (Yup'ik) people. *J Nutr*. 2015 May; 145[5]:931–8. <https://doi.org/10.3945/jn.114.209619> PMID: 25788581
103. Mendelsohn ME. Protective effects of estrogen on the cardiovascular system. *Am J Cardiol*. 2002; 89[12]:12–7.
104. O'Brien DM, Thummel KE, Bulkow LR, Wang Z, Corbin B, Klejka J, et al. Declines in traditional marine food intake and vitamin D levels from the 1960s to present in young Alaska Native women. *Public Health Nutr*. 2016;[14]:1–8.
105. Jørgensen ME, Moustgaard H, Bjerregaard P, Borch-Johnsen K. Gender differences in the association between westernization and metabolic risk among Greenland Inuit. *Eur J Epidemiol*. 2006; 21[10]:741–8. <https://doi.org/10.1007/s10654-006-9063-4> PMID: 17077992
106. Kaholokula JKA, Nacapoy AH, Grandinetti A, Chang HK. Association between acculturation modes and type 2 diabetes among native hawaiians. *Diabetes Care*. 2008; 31[4]:698–700. <https://doi.org/10.2337/dc07-1560> PMID: 18202248
107. Lara M, Gamboa C, Kahramanian MI, Morales LS, Hayes Bautista DE. Acculturation and Latino health in the United States: a review of the literature and its sociopolitical context. *Annu Rev Public Health*. 2005; 26:367–97. <https://doi.org/10.1146/annurev.publhealth.26.021304.144615> PMID: 15760294
108. Veenstra G, Burnett PJ. A Relational approach to health practices: towards transcending the agency-structure divide. *From Heal Behav to Heal Pract Crit Perspect*. 2014; 36[2]:31–42.
109. Abel T. Cultural capital and social inequality in health. *J Epidemiol Community Health*. 2008; 62[7]:e13. PMID: 18572429
110. Veenstra G, Abel T. Capital interplays and the self-rated health of young men: results from a cross-sectional study in Switzerland. *Int J Equity Health*. 2015 Dec 18; 14[1]:38.
111. Abel T, Frohlich KL. Capitals and capabilities: Linking structure and agency to reduce health inequalities. *Soc Sci Med*. Elsevier Ltd; 2012 Jan; 74[2]:236–44.
112. McIvor O, Napoleon A. Language and culture as protective factors for at-risk communities. *Int J Indig Heal*. 2009; 5[1]:6.
113. Rasmus S, Charles B, Mohatt G V. Creating Qungasvik (a Yup'ik intervention “toolbox”): case examples from a community-developed and culturally-driven intervention. *Am J Community Psychol*. 2014 Sep; 54[1–2]:140–52. <https://doi.org/10.1007/s10464-014-9651-5> PMID: 24764018
114. Dressler WW, Bindon JR. The Health Consequences of Cultural Consonance: Cultural Dimensions of Lifestyle, Social Support, and Arterial Blood Pressure in an African American Community. *Am Anthropol*. 2000; 102[2]:244–60.
115. Singer MK. Applying the concept of culture to reduce health disparities through health behavior research. *Prev Med (Baltim)*. 2012; 55[5]:356–361.
116. Brabec M, Godoy R, Reyes-García V, Leonard WR. BMI, income, and social capital in a native Amazonian society: interaction between relative and community variables. *Am J Hum Biol*. 2007; 19[4]:459–474. <https://doi.org/10.1002/ajhb.20606> PMID: 17546614
117. Steptoe A, Kivimaki M. Stress and cardiovascular disease: an update on current knowledge. *Annu Rev Public Heal*. 2013; 34:337–54.
118. Kawagley AO, Norris-Tull D, Norris-Tull RA. The indigenous worldview of Yupiaq culture: Its scientific nature and relevance to the practice and teaching of science. *J Res Sci Teach*. Wiley Online Library; 1998; 35[2]:133–44.