

2018

A remarkable teratological specimen of *Trichiotinus rufobrunneus* (Casey) (Coleoptera: Scarabaeidae: Cetoniinae: Trichiini)

Héctor Jaime Gasca-Álvarez

Corporacion Sentido Natural, hjgasca@sentidonatural.org

Paul E. Skelley

Florida Department of Agriculture and Consumer Services, Paul.Skelley@FreshFromFlorida.com

Cuauhtemoc Deloya

Instituto de Ecología, A.C., cuauhtemoc.deloya@inecol.mx

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

Gasca-Álvarez, Héctor Jaime; Skelley, Paul E.; and Deloya, Cuauhtemoc, "A remarkable teratological specimen of *Trichiotinus rufobrunneus* (Casey) (Coleoptera: Scarabaeidae: Cetoniinae: Trichiini)" (2018). *Insecta Mundi*. 1175.

<http://digitalcommons.unl.edu/insectamundi/1175>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0680

A remarkable teratological specimen of
Trichiotinus rufobrunneus (Casey)
(Coleoptera: Scarabaeidae: Cetoniinae: Trichiini)

Héctor Jaime Gasca-Álvarez
Programa de Investigación
Corporación Sentido Natural
Bogotá, Colombia.

Paul Skelley
Florida State Collection of Arthropods
Florida Department of Agriculture and Consumer Services
P. O. Box 147100
Gainesville, FL 32614-7100 USA

Cuauhtémoc Deloya
Red de Interacciones Multitróficas
Instituto de Ecología, A.C.
Carretera antigua a Coatepec 351, El Haya, 91070
Xalapa, Veracruz, México

Date of issue: December 28, 2018

Héctor Jaime Gasca-Álvarez, Paul Skelley and Cuauhtémoc Deloya
A remarkable teratological specimen of *Trichiotinus rufobrunneus* (Casey)
(Coleoptera: Scarabaeidae: Cetoniinae: Trichiini)

Insecta Mundi 0680: 1–5

ZooBank Registered: urn:lsid:zoobank.org:pub:6849188C-81E7-45E8-86C4-ABCB481A61EE

Published in 2018 by

Center for Systematic Entomology, Inc.

P.O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, insectamundi@gmail.com

Head Layout Editor: Robert G. Forsyth

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Robert G. Forsyth

A remarkable teratological specimen of *Trichiotinus rufobrunneus* (Casey) (Coleoptera: Scarabaeidae: Cetoniinae: Trichiini)

Héctor Jaime Gasca-Álvarez

Programa de Investigación
Corporación Sentido Natural
Bogotá, Colombia.
hjasca@sentidonatural.org

Paul Skelley

Florida State Collection of Arthropods
Florida Department of Agriculture and Consumer Services
P. O. Box 147100
Gainesville, FL 32614-7100 USA
Paul.Skelley@freshfromflorida.com

Cuauhtémoc Deloya

Red de Interacciones Multitróficas
Instituto de Ecología, A.C.
Carretera antigua a Coatepec 351, El Haya, 91070
Xalapa, Veracruz, México
cuauhtemoc.deloya@inecol.mx

Abstract. An unusual eye malformation observed in *Trichiotinus rufobrunneus* (Casey) (Coleoptera: Scarabaeidae: Cetoniinae: Trichiini), is described and illustrated. The functionality of the ectopic compound eye is discussed. According to label data, larval association with oak rotten log habitats is reported.

Key words. Teratology, morphology, Scarabaeoidea.

Resumen. Se describe e ilustra una malformación inusual en los ojos de *Trichiotinus rufobrunneus* (Casey, 1914). Se discute la funcionalidad de los ojos compuestos ectópicos. De acuerdo con los datos de etiqueta, se reporta la asociación de la larva con madera de roble en descomposición.

Palabras clave. Teratología, morfología, Scarabaeoidea.

Introduction

Morphological malformations are common in Coleoptera and have been widely documented (Mocquerys 1880; Ellis 1915; Balazuc 1948, 1969; Ortuño and Hernández 1993; Ortuño and Ramos-Abuín 2008; Castro-Tovar et al. 2014). Malformations are frequently reported in Carabidae, Staphylinidae, Cerambycidae, Chrysomelidae (Frank 1981; Gandhi and Herms 2008; Asiain and Márquez 2009; Ortuño and Peláez 2004; Clark and Belo-Neto 2010; Verdugo 2013; Verdugo and del Saz Fucho 2012; Verdugo and Toribio 2015, 2017) and in the superfamily Scarabaeoidea (see Gasca-Álvarez et al. 2017).

Several authors have reiterated the scientific importance of describing and illustrating insect abnormalities. They state that some cases can provide relevant information about environmental influences on developmental processes (Glasgow 1925; Cockayne 1937; Savini and Furth 2004; Clark and Belo-Neto 2010).

The origin of these developmental abnormalities has been difficult to establish. Some can be coded in the DNA and are transferable to subsequent generations. In some cases, malformations can be produced by the effects of a single gene, or are characterized by diverse phenotypic expressions that may involve multiple genes and their interactions with the environment (Clark and Belo-Neto 2010; Palomar-Morales et al. 2016). These abnormalities can be caused by the action of endogenous or exogenous factors that influence embryonic and/or postembryonic development. Virus infections or parasites, can be other factors associated with the occurrence of malformations.

Continuing documentation of the teratology of scarab beetles (see Gasca-Álvarez et al. 2017), we report herein a notable case in a male specimen of *Trichiotinus rufobrunneus* (Casey)

Materials and Methods

The male specimen is housed in the Florida State Collections of Arthropods (Gainesville, Florida, United States of America), and labelled: “FLORIDA: Levy Co. 3.8 mi SW Archer 23-III-1988. P. Skelley: reared from grub in rotten oak log”. Photographs were taken using the AutoMontage system by Syncroscopy located at the Florida State Collections of Arthropods. Terminology follows Balazuc (1948, 1969) and Ortuño and Hernández (1993).

Results and Discussion

The specimen exhibits a well-developed third holoptic eye, smaller than the normal eyes and located in the frons, close to right eye (Fig. 1). The size and shape of the head, and the antennal insertion appears to be typical, compared with other normal specimens. This malformation is a new case reported in Scarabaeoidea. Other different cases involving eye malformations in Coleoptera have been described for two species of Chrysomelidae. In *Donacia bicolora* Zschach, a case of cephalic hemiatrophy produced the complete reduction of the right eye (Balazuc 1948). Clark and Belo-Neto (2010) found a female of *Pseudoluperus longulus* (LeConte) from Utah (USA), with the head surface covered almost entirely by a single holoptic eye.

This finding in *T. rufobrunneus* raises the question of whether this compound eye is a functional structure. Researchers at Indiana University (USA), demonstrated that the down-regulation via RNAi of *orthodenticle*, a single head patterning gene, results in development of functional ectopic compound eyes at the middorsal adult head of five scarab species (Zattara et al. 2016, 2017). The RNA sequencing experiments were performed on *Onthophagus taurus* (Schreber), *O. sagittarius* (Fabricius), *O. binodis* (Thunberg), *Digitonthophagus gazella* (Fabricius) (all Onthophagini) and *Liatongus militaris* (Castelnau) (Oniticellini). According to the authors, the ommatidial organization of these induced structures reveals the presence of rudimentary ommatidial lenses, crystalline cones, and associated neural-like tissue. To establish the functionality, a light-aversion behavioral assay demonstrated that these ectopic compound eyes have the ability to respond to visual stimuli when wild-type eyes are surgically removed (Zattara et al. 2016, 2017). In all scarab species studied, these ectopic eyes are located in the middle of the head.

This contrasts with the observed specimen of *Trichiotinus* Casey, where the third compound eye is not centrally located (Fig. 1b). According to the position of the third compound eye, its inner margin is somewhat in line with the lateral margin of the clypeus, similar to the position of the “normal” eye on the right side of the head. Moreover, the head is asymmetrical, the mouthparts are exposed and the “normal” eye on the right side is placed more lateral in regard to its regular position. Possibly, the apparent third compound eye is in fact a rudiment of the original right eye (Fig. 1c), which for an unknown reason stopped its development and a was replaced with a new one.

Little is known about the natural history of *T. rufobrunneus*. This flower chafer has a limited distribution in Florida and is likely restricted to oak scrub habitats (Hoffmann 1935; Philips et al. 2016). Adults have been reported as floral visitors or pollinators of the saw palmetto *Serenoa repens* (Bartram) Small (Arecaceae) (Deyrup and Deyrup 2012), *Asimina obovata* Nash (Annonaceae) (Gottsberger 2012) and *Opuntia* spp. (Cactaceae) (Eisner 2003). Larval feeding habits were not known until now. According to data label, the larvae of *T. rufobrunneus* fed and developed in rotting oak trunks. In some places in Florida, the populations are relatively abundant in the spring. When many *Trichiotinus* specimens are resting or when feeding on pollen within a flower, they take a position with the pygidium facing outward towards the flower’s opening, thus giving it the appearance of the head of a bee or wasp (Ratcliffe 1991; Eisner 2003).

Acknowledgments

We are grateful to Howard Frank, Petr Šípek and William B. Warner for their constructive comments and suggestions, which helped us to improve the manuscript.

Literature Cited

- Asiain, J., and J. Márquez. 2009.** New teratological examples in Neotropical Staphylinidae (Insecta: Coleoptera), with a compilation of previous teratological records. *Revista Mexicana de Biodiversidad* 80: 129–139.
- Balazuc, J. 1948.** La tératologie des coléoptères et expériences de transplantation sur *Tenebrio molitor* L. *Mémoires du Muséum National d'Histoire Naturelle (N.S.)* 25: 1–293.
- Balazuc, J. 1969.** Supplément à la tératologie des coléoptères. *Redia* 51: 39–111.
- Castro-Tovar, A., M. Baena, and M. A. López Vergara. 2014.** Nuevos casos de teratologías en Coleoptera (Insecta). *Zoologica Baetica* 25: 3–12.
- Cockayne, E. A. 1937.** Insect teratology, reduplication of legs in Coleoptera, Diptera, and Hymenoptera. *Transactions of the Royal Entomological Society of London* 86: 191–200.
- Clark, S., and L. Belo-Neto. 2010.** Remarkable teratological specimen of *Pseudoluperus longulus* (LeConte) (Coleoptera: Chrysomelidae) from Utah, U.S.A. *The Coleopterists Bulletin* 64: 383–385.
- Deyrup, M., and L. Deyrup. 2012.** The diversity of insects visiting flowers of saw palmetto (Arecaceae). *Florida Entomologist* 95(3): 711–730.
- Eisner, T. 2003.** *For Love of Insects*. Belknap Press of Harvard University Press; Cambridge, MA. 464 p.
- Ellis, H. W. 1915.** Teratological specimens of Coleoptera. *Proceedings of the Entomological Society of London* 17: XLVI.
- Frank, J. H. 1981.** A review of teratology in Staphylinidae, with description of a teratological specimen of *Tachinus axillaris* Erichson (Coleoptera, Staphylinidae, Tachyporinae) from Florida. *Florida Entomologist* 64: 337–340.
- Gandhi, K. J. K., and D. Herms. 2008.** Report on the largest known occurrence of morphological anomalies in ground beetles (Coleoptera: Carabidae). *The Coleopterists Bulletin* 62: 104–113.
- Gasca-Álvarez, H. J., C. Deloya, and P. Reyes-Castillo. 2017.** Teratological cases in five species of *Cotinis* Burmeister (Coleoptera: Scarabaeidae: Cetoniinae: Gymnetini), with a compilation of teratologies in Scarabaeoidea. *The Coleopterist Bulletin* 71(2): 329–338.
- Glasgow, R. D. 1925.** A specimen of *Melanoplus diffentialis* Thomas with four ocelli. *Psyche* 32: 285–290.
- Gottsberger, G. 2012.** How diverse are Annonaceae with regard to pollination? *Botanical Journal of the Linnean Society* 169: 245–261.
- Hoffmann, C. H. 1935.** The biology and taxonomy of the genus *Trichiotinus* (Scarabaeidae; Coleoptera). *Entomologica Americana* 15: 133–214.
- Mocquerys, S. 1880.** *Recueil de Coléoptères*. Léon Deshayes; Rouen, France. 142 p.
- Ortuño, V. M., and J. M. Hernández. 1993.** Diversos casos teratológicos en Coleoptera. *Boletín de la Real Sociedad Española de Historia Natural* 89: 163–179.
- Ortuño, V. M., and L. Peláez. 2004.** Nuevos e interesantes casos de carábidos teratomorfos (Coleoptera, Adepaga, Carabidae). *Bulletin de la Société Entomologique de France* 109: 251–256.
- Ortuño, V. M., and J. A. Ramos-Abún. 2008.** Reflexiones sobre la teratología y descripción de cuatro teratosis apendiculares en Coleoptera. *Boletín de la Sociedad Entomológica Aragonesa* 43: 435–439.
- Palomar-Morales, M., G. Chirino-Galindo, C. Álvarez-Rodríguez, and M. G. Villanueva-Santiago. 2016.** Embriología y teratología, manuales de animales en laboratorio. Universidad Nacional Autónoma de México, Facultad de Estudios Superiores Iztacala; Mexico City, Mexico. 115 p.
- Philips, T. K., M. Callahan, J. Orozco, and N. Rowland. 2016.** Phylogenetic analysis of the North American beetle genus *Trichiotinus* (Coleoptera: Scarabaeidae: Trichiinae). *Psyche* 2016: 1–9.
- Ratcliffe, B. C. 1991.** The scarab beetles of Nebraska. *Bulletin of the University of Nebraska State Museum* 12: 1–333.
- Savini, V., and D. Furth. 2004.** Teratología en Coleoptera: un caso en *Gioia bicolor* (Blake 1969) (Chrysomelidae, Alticinae) de Jamaica. *Entomotropica* 19: 165–167.

- Verdugo A. 2013.** A propósito de una hemimeria protorácica asociada a la ausencia de la pata protorácica izquierda en *Iberodorcadion zenete* Anichtchenko & Verdugo, 2004 (Coleoptera: Cerambycidae: Dorcadionini). *Revista Gaditana de Entomología* 4(1):123–127.
- Verdugo, A., and A. Del Saz Fucho. 2012.** A propósito de un caso de esquistomelia binaria heterodinámica de antena derecha en *Iberodorcadion perezii* (Graells, 1849) ssp. *nudipenne* (Escalera, 1908) (Coleoptera: Cerambycidae: Dorcadionini). *Revista Gaditana de Entomología* 3(1–2): 11–16.
- Verdugo, A., and M. Toribio. 2015.** Sobre un caso de malformación del tipo „duplicación del edeago“ en *Acupalpus maculatus* (Schaum, 1860) (Coleoptera: Carabidae: Harpalinae). *Revista Gaditana de Entomología* 6(1): 131–136.
- Verdugo, A., and M. Toribio. 2017.** Un nuevo caso de malformación del tipo esquistomelia ternaria heterodinámica en *Calathus (Calathus) fuscipes graecus* Dejean, 1831 (Coleoptera: Carabidae: Platyninae). *Revista Gaditana de Entomología* 8(1): 39–42.
- Zattara, E. E., H. A. Busey, D. M. Linz, Y. Tomoyasu, and A. P. Moczek. 2016.** Neofunctionalization of embryonic head patterning genes facilitates the positioning of novel traits on the dorsal head of adult beetles. *Proceedings of the Royal Society B* 283(1834): 1–8.
- Zattara, E. E., A. L. M. Macagno, H. A. Busey, and A. P. Moczek. 2017.** Development of functional ectopic compound eyes in scarabaeid beetles by knockdown of orthodenticle. *Proceedings of the National Academy of Sciences* 114(45): 12021–12026.

Received November 21, 2018; accepted November 30, 2018.

Review editor Oliver Keller.




Figure 1. Teratological *Trichiotinus rufobrunneus* (Casey). a) Habitus. b) Head. c) Detail of third eye.

