

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Carro, Lorena and Razmilic, Valeria and Nouioui, Imen and Richardson, Lee and Pan, Che and Golinska, Patrycja and Asenjo, Juan A. and Bull, Alan T. and Klenk, Hans-Peter and Goodfellow, Michael (2018) Hunting for cultivable *Micromonospora* strains in soils of the Atacama Desert. *Antonie van Leeuwenhoek*, 111 (8). pp. 1375-1387. ISSN 0003-6072.

### DOI

<https://doi.org/10.1007/s10482-018-1049-1>

### Link to record in KAR

<https://kar.kent.ac.uk/66520/>

### Document Version

Author's Accepted Manuscript

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

# Hunting for cultivable *Micromonospora* strains in soils of the Atacama Desert

Lorena Carro<sup>1</sup>, Valeria Razmilic<sup>1,2</sup>, Imen Nouioui<sup>1</sup>, Lee Richardson<sup>1</sup>, Che Pan<sup>1</sup>, Patrycja Golinska<sup>3</sup>, Juan A. Asenjo<sup>2</sup>, Alan T. Bull<sup>4</sup>, Hans-Peter Klenk<sup>1</sup>, Michael Goodfellow<sup>1</sup>

<sup>1</sup> School of Natural and Environmental Sciences, Newcastle University, Newcastle-upon-Tyne, UK.

<sup>2</sup> Centre for Biotechnology and Bioengineering (CeBiB), Department of Chemical Engineering and Biotechnology, Universidad de Chile, Beauchef 850, Santiago, Chile.

<sup>3</sup> Department of Microbiology, Nicolaus Copernicus University, Torun, Poland

<sup>4</sup> School of Biosciences, University of Kent, Canterbury, Kent CT2 7NJ, UK

Corresponding author: Lorena Carro [lcg@usal.es](mailto:lcg@usal.es), Telephone: +44 (0)1912084853.

## Abstract (150-250)

Innovative procedures were used to selectively isolate small numbers of *Micromonospora* strains from extreme hyper-arid and high altitude Atacama Desert soils. *Micromonosporae* were recognized on isolation plates by their ability to produce filamentous microcolonies that were strongly attached to the agar. Most of the isolates formed characteristic orange colonies that lacked aerial hyphae and turned black on spore formation whereas those from the high altitude soil were dry, blue-green and covered by white aerial hyphae. The isolates were assigned to seven multi- and eleven single-membered groups based on BOX-PCR profiles. Representatives of the groups were assigned to either multi-membered clades that also contained marker strains or formed distinct phyletic lines in the *Micromonospora* 16S rRNA gene tree, many of the isolates were considered to be putatively novel species of *Micromonospora*. Most of the isolates from the high altitude soils showed activity against wild type strains of *Bacillus subtilis* and *Pseudomonas fluorescens* while those from the rhizosphere of *Parastrephia quadrangulares* and from the Lomas Bayas hyper-arid soil showed resistance to UV radiation.

**Keywords:** *Micromonospora*, Atacama Desert, BOX-PCR, polyphasic taxonomy, UV radiation.

## 37 Introduction

38 Given their importance in biotechnology, bioprospecting and ecology, actinobacteria  
39 remain a source of interest to the microbiological community (Goodfellow and Fiedler  
40 2010; Demain 2014; Barka et al. 2016). Actinobacteria were initially isolated from the  
41 Atacama Desert over fifty years ago (Cameron et al. 1966; Opfell and Zebal 1967)  
42 though the first extensive survey of these organisms in this desert was only reported  
43 recently (Okoro et al. 2009). Okoro and her colleagues isolated small numbers of  
44 filamentous actinobacteria from arid, hyper-arid and extreme hyper-arid Atacama  
45 Desert soils using selective isolation procedures and polyphasic taxonomic methods  
46 which showed that a high proportion of the isolates belonged to the genera  
47 *Amycolatopsis*, *Lechevaleria* and *Streptomyces* many of which were assigned to  
48 putative new species and contained novel non-ribosomal peptide synthase genes.  
49 Subsequent studies have shown that Atacama Desert habitats are not only a rich source  
50 of novel streptomycetes but also of rare and poorly studied taxa, some of which produce  
51 bioactive compounds and have been given validly published names (Bull et al. 2016;  
52 Busarakam et al. 2014, 2016; Goodfellow et al. 2017; Idris et al. 2017a; Idris et al.  
53 2017b; Trujillo et al. 2017). *Streptomyces leewenhoekii* strains (Busarakam et al. 2014),  
54 for instance, synthesize novel macrolactone and polyketide antibiotics (Nachtigall et al.  
55 2011; Rateb et al. 2011a; Rateb et al. 2011b) and chaxapeptin, a new lasso peptide  
56 (Elsayed et al. 2015) while the type strain of *Lentzia chajnantorensis* (Idris et al. 2017b)  
57 produces novel diene and monoene glycosides, several of which show anti-HIV  
58 integrase activity (Wichner et al. 2017). Complementary metagenomic analyses of  
59 Atacama Desert habitats have revealed a remarkable actinobacterial diversity most of  
60 which has not been detected using culture-dependent methods (Bull et al. 2017; Idris  
61 et al. 2017c). Improved selective isolation and cultivation methods are needed to isolate  
62 components of this diversity, not least *Micromonospora* strains which are known to be a  
63 rich source of new specialist metabolites and have the potential to defend plants against  
64 root-infecting fungi (Carro et al. 2018a).

65 *Micromonospora* (Ørskov 1923), the type genus of the family  
66 *Micromonosporaceae* (Krasil'nikov 1938; Zhi et al. 2009), currently encompasses 79  
67 species with validly published names (Parte 2014) many of which have been  
68 circumscribed using polyphasic methods (Genilloud 2012; Carro et al. 2018b) though  
69 the genus remains underspeciated (Carro et al. 2012a; Carro et al. 2013a).  
70 *Micromonosporae* have been isolated from diverse natural habitats (Genilloud 2012),  
71 notably from rhizosphere soil (Carro et al. 2013b; Thawai et al. 2016) from desert  
72 locations in China (Ding et al. 2013) and from tissues of a broad range of plants, such as  
73 *Triticum aestivum* (Coombs and Franco 2003), *Zea mays* (Shen et al. 2014), and  
74 *Parathelypteris beddomei* (Zhao et al. 2017); as well as from nitrogen-fixing root  
75 nodules of actinorhizal plants (Trujillo et al. 2006; Carro et al. 2013a) and legumes  
76 (Trujillo et al. 2007; Garcia et al. 2010; Trujillo et al. 2010; Carro et al. 2018b).

77 It is apparent from culture-independent studies that micromonosporae form a  
78 small, but integral part of actinobacterial communities in Atacama Desert habitats (Bull  
79 et al. 2017; Idris et al. 2017c). The failure to isolate them from such habitats may reflect  
80 the use of isolation media mainly designed to be selective for streptomycetes

81 (Busarakam 2014; Idris 2016). The primary aims of the present study were to isolate  
82 micromonosporae from diverse Atacama Desert soils using procedures designed to be  
83 selective for members of this taxon and to determine whether any such isolates  
84 represented putatively novel taxa.

85

## 86 **Materials and Methods**

### 87 *Sampling sites*

88 Soil samples (Table 1) were collected from several locations in the Atacama Desert  
89 (Fig. 1) between 2010 and 2016 (ATB, MG) and an additional one in 2012 (Professor  
90 Luis Cáceres, University of Antofagasta). The samples were collected aseptically using  
91 spatulas sterilized in the field with ethanol and contained in sterile polycarbonate  
92 bottles. Following transport to the UK the samples were stored at 4°C. Further details of  
93 the sampling sites can be found elsewhere (Bull et al. 2017; Idris et al. 2017c).

### 94 *Selective isolation*

95 Three isolation procedures were used: a) Phenol-heat protocol (P-H): a gram of each of  
96 the soil samples was diluted in 0.85% sodium chloride solution containing 1.5% phenol  
97 (Hayakawa et al. 1991), the preparations shaken on a rotary shaker for an hour,  
98 incubated at 70°C for 40 minutes in a water bath (Nonomura and Ohara, 1969), shaken  
99 again at room temperature for 2 hours, serial dilutions prepared in the saline solution,  
100 and the resultant preparations shaken at room temperature prior to spreading over chitin-  
101 vitamin (CHV) (Zhang 2011), Gause number 1 (G n° 1) (Gause et al. 1983), humic acid-  
102 vitamin (HV) (Hayakawa and Nonomura 1987), M65 (DSMZ medium number 65),  
103 R2A (Reasoner et al. 1979) and Zhang' starch soil extract agar (ZSSE) (Zhang 2011), all  
104 of the media were supplemented with 50µg/ml of cycloheximide and 50µg/ml of  
105 nalidixic acid; b) sprinkle protocol (S): 0.5g of soil particles of individual samples  
106 which had been preheated at 120°C for 15 minutes were sprinkled directly over the  
107 isolation media and c) a standard dilution plate protocol (D): dilutions of each soil  
108 sample in saline were used to inoculate each of the isolation media following the  
109 procedure described by Goodfellow et al. (1967). All of the inoculated plates were  
110 incubated at 28°C and examined weekly for up to 28 days for the presence of  
111 *Micromonospora* colonies that were detected, using a stereoscopic microscope, and then  
112 transferred to the medium from which they were isolated but without the antibiotics, the  
113 resultant plates were incubated at 28°C for 15 days. The 45 isolated strains were  
114 maintained on either M65 or ZSSE agar and in 20% v/v glycerol at -80°C for long-term  
115 preservation.

### 116 *Dereplication of isolates*

117 Genomic DNA for BOX-PCR was extracted from all the isolates using a Bacterial DNA  
118 Extraction Kit (Sigma) following the instructions of the manufacturer and BOX-PCR  
119 fingerprint profiles generated using modifications of the methods described by  
120 Versalovic et al. (1994) and Trujillo et al. (2010). To this end, Bioline 2x MiFi™ mix  
121 was used for PCR amplification in a final volume of 25 µl per reaction following the  
122 manufacturer's recommendations; the thermal cycling parameters were: 7 minutes at  
123 95 °C, 30 cycles of 1 minute at 94 °C, 1 minute at 52 °C and 3 minutes at 72 °C followed  
124 by a final extension at 72 °C for 10 minutes. Five microliters of each PCR product was

125 loaded onto a 2% agarose gel containing 10 µl of GelRed™ (Crisafuli et al. 2015) per  
126 100 ml and electrophoresis run at 70 V for 3 hours in freshly prepared 1x TBE-EDTA  
127 buffer at pH 8.0 using a Bio-Rad Pac 300 power supply; a DNA molecular weight  
128 marker 1kb HyperLadder™ (Bioline) was used as the molecular size standard. After  
129 electrophoresis, gels were photographed, stored on disk as TIFF files, and manually  
130 aligned into 9 multi- and 10 single-membered similarity groups.

### 131 *Phylogeny*

132 PCR-mediated amplification of genomic DNA of 20 representatives of the BOX-PCR  
133 groups was conducted using the universal primers 27F and 1522R (Lane 1991). Bioline  
134 2x MiFi™ mix was used for PCR amplification in a final volume of 50 µl per reaction  
135 following the manufacturer's recommendations. The thermal cycling parameters were:  
136 9 minutes at 95 °C, 30 cycles of 1 minute at 94 °C, 1 minute at 56 °C and 2 minutes at  
137 72 °C followed by a final extension at 72 °C for 10 minutes. The PCR products were  
138 purified using a QIAquick® PCR purification kit, according to the manufacturer's  
139 instructions (Qiagen) and sequenced using the EZ-seq Barcode Service (Macrogen).  
140 The resultant sequences (around 1400 nucleotides) were manually aligned using  
141 MUSCLE (Edgar 2004) and then compared with corresponding sequences of  
142 *Micromonospora* type strains retrieved from the EzBioCloud server (Yoon et al. 2016).  
143 Phylogenetic distances were calculated with the Kimura 2-parameter model (Kimura  
144 1980) and tree topologies inferred using the neighbour-joining (Saitou and Nei 1987)  
145 and maximum-likelihood algorithms (Felsenstein 1981) and one thousand bootstrap  
146 replications. Tree reconstructions were carried out using MEGA 7 software (Kumar et  
147 al. 2016).

### 148 *Stress tests*

149 All of the strains were examined for their ability to grow in the presence of various  
150 concentrations of sodium chloride (1, 3, 5, 7 and 9%, w/v) and at a range of temperature  
151 (4, 10, 20, 28, 37 and 45 °C) and pH values (4.5, 5.5, 6.5, 8.0 and 9.0) using M65 and  
152 ZSSE agar as basal media; pH values were determined using phosphate buffers, as  
153 described previously (Carro et al. 2012b). All of the plates, apart from those from the  
154 temperature tests, were incubated at 28°C for 15 days. The strains were also examined  
155 for their ability to grow following exposure to 100 mJoules for 30 and for 60 minutes in  
156 a UV chamber (Biorad) using *Geodermatophilus poikilotrophi* DSM 44209<sup>T</sup> as the  
157 positive control (Montero-Calasanz et al. 2014)) set at a wavelength of 254 nm (UVC).  
158 The capacity of the strains to grow under anaerobic conditions was tested on M65 and  
159 ZSSE agar plates that were incubated for 28°C for 10 days in anaerobic atmosphere  
160 generation bags (Sigma-Aldrich 68061).

### 161 *Antibiotic sensitivity assays*

162 The isolates were examined for their ability to inhibit the growth of wild type strains of  
163 *Bacillus subtilis*, *Escherichia coli* and *Pseudomonas fluorescens* using a standard plug  
164 assay (Fiedler, 2004). Lawns of each of the isolates were prepared on yeast extract-malt  
165 extract and oatmeal agar (International *Streptomyces* Project [ISP] media 2 and 3;  
166 Shirling and Gottlieb 1966) plates incubated at 28°C for 14 days. Agar plugs taken from  
167 the incubated plates were transferred to sterile square Petri Dishes and overnight Luria  
168 broth cultures [OD<sub>600</sub> of approximately 0.6] of each of the wild type strains in molten  
169 nutrient agar added to a final concentration of OD<sub>600</sub> of 0.0125. The inoculated dishes

170 were incubated at 28°C and examined for zones of inhibition around the agar plugs after  
171 24 and 48 hours.

## 172 **Results and Discussion**

### 173 *Isolation of micromonosporae*

174 The ability to isolate specific fractions of actinobacterial communities from the  
175 Atacama Desert biome is of paramount importance especially since metagenomic  
176 analyses have revealed a staggering degree of actinobacterial diversity therein: the vast  
177 majority of which has gone undetected using culture-dependent procedures (Idris et al.  
178 2017c), this partly reflects the use of isolation media that favour the growth of  
179 streptomycetes (Williams et al. 1984). Isolation procedures designed to recover specific  
180 actinobacterial genera from environmental samples are many and varied but tend to  
181 reflect the biological properties of the target organisms (Goodfellow 2010), as was the  
182 case in the present study. It is, for instance, known that *Micromonospora* spores are  
183 activated by heat pretreatment regimes, are resistant to phenol and germinate on nutrient  
184 rich media such as humic acid-vitamin agar (Hoskisson et al. 2000; Shen et al. 2014;  
185 Carro et al. 2018b). It is, therefore, encouraging that over half of the *Micromonospora*  
186 strains were isolated from the soil samples using the phenol-heat procedure, albeit on  
187 several isolation media (Table 2).

188 Micromonosporae growing on the isolation plates were recognized, under the  
189 stereoscopic microscope, by their ability to produce filamentous microcolonies that  
190 were strongly attached to agar, these colonies were readily distinguished from those of  
191 aerobic, endospore-forming bacilli (data not shown). It was particularly interesting that  
192 most of the *Micromonospora* strains came from the ALMA 4 sample as complementary  
193 culture-independent studies have shown that soil samples from this location contain  
194 micromonosporal propagules (Bull et al., 2017). The strains isolated from this site  
195 proved to be unusual as they formed dry, filamentous blue-green colonies covered by  
196 white aerial hyphae following growth for three weeks on M65 and ZSSE media (Fig. 1).  
197 The ability of these isolates to form aerial hyphae is interesting as some  
198 *Micromonospora* type strains have been shown to contain putative homologous genes to  
199 those associated with aerial hyphae formation and spore maturation in streptomycetes  
200 (Carro et al. 2018a). Strains isolated from soil samples taken from the three remaining  
201 locations showed a typical micromonosporal phenotype, one characterized by the  
202 production of filamentous orange colonies that lack aerial hyphae and turn blue-black  
203 upon spore formation (Genilloud 2012). Five of these strains were isolated from  
204 extreme hyper-arid soil collected from the Lomas Bayas region (Idris et al. 2017c), one  
205 of the driest areas in the Atacama Desert; all of these isolates were recovered by plating  
206 serial dilutions of the soil onto M65 agar. In turn, the 13 strains isolated from the  
207 rhizosphere of *Parastrephia quadrangularis* (*Compositae*, tribe *Asteraceae*) were  
208 obtained either by sprinkling mineral particles or spreading serial dilutions of the  
209 phenol-heat protocol onto HV and ZSSE agar and incubating for four weeks at 28°C.  
210 *Micromonospora* strains were recovered using all three selective isolation procedures  
211 (Table 2).

### 212 *BOX-PCR profiles*

213 The BOX-PCR profiles of the isolates encompass considerable genetic diversity with  
214 fragments ranging from 0.05 to 2.0 kb (Fig. 2) thereby providing further evidence that  
215 this method is effective in distinguishing between *Micromonospora* strains (Maldonado  
216 et al. 2008; Trujillo et al. 2010). The isolates were assigned to seven multi- and 11  
217 single-membered groups defined at the 60% similarity level (Table 3). Five of the  
218 multi-membered groups and five of the singletons were composed of strains isolated  
219 from the ALMA 4 soil samples, the largest of these taxa, group XI, contained seven  
220 isolates. In turn, the strains isolated from *P. quadrangularis* rhizosphere soil were  
221 assigned to a multi-membered group that encompassed 10 strains and to two single  
222 membered groups; the five Lomas Bayas strains formed one multi- and 3 single-  
223 membered taxa while the single isolates from ALMA 5 and Yungay Core Region  
224 samples gave unique profiles (Fig. 2). It is interesting that the multi-membered groups  
225 only contained strains isolated from a single location and that few strains were clones  
226 (Fig. 2; Table 3).

### 227 *Phylogenetic analyses*

228 Almost complete 16S rRNA gene sequences (1372 – 1507 nucleotides [nt]) were  
229 generated for the 20 isolates chosen to represent the BOX-PCR groups (Table 3). The  
230 generic assignment of all of these strains was confirmed as they were recovered in the  
231 *Micromonospora* 16S rRNA gene tree (Fig. 3). The 16S rRNA gene sequence  
232 similarities between these isolates and the type strains of their closest phylogenetic  
233 neighbours fell within the range 98.4 to 99.9 % (Table 3). Several of the isolates were  
234 assigned to well-supported clusters in the *Micromonospora* tree though none of these  
235 taxa included strains isolated from more than one location. Ten out of the twelve  
236 ALMA 4 strains formed a well-supported phyletic line within a weakly supported clade  
237 that encompassed the type strains of *Micromonospora costi* (Thawai 2015), their nearest  
238 neighbour, and *Micromonospora fulviviridis* (Kroppenstedt et al. 2005); it is interesting  
239 that the isolates of this clade were recovered from different samples of ALMA 4 soil  
240 using two of the three isolation protocols and several selective isolation media (Table  
241 2). Isolate 5R2A7, one of the remaining ALMA strains, formed a well-supported clade  
242 together with the type strain of *Micromonospora coriariae* (Trujillo et al. 2006), its  
243 nearest phylogenetic neighbour, and with *Micromonospora cremea* (Carro et al. 2012b)  
244 whereas isolate ATA32, the remaining ALMA 4 strain, formed an unsupported clade  
245 with the type strain of *Micromonospora narathiwatensis* (Thawai et al. 2007) though it  
246 was most closely related to *Micromonospora eburnea* (Thawai et al. 2005).

247 It is interesting that two out of the three strains recovered from the rhizosphere  
248 of *P. quadrangularis*, isolates STR1-7 and STR1s-6, were loosely associated with the  
249 type strains of *Micromonospora lupini* (Trujillo et al. 2007), *Micromonospora taraxaci*  
250 (Zhao et al. 2014) and *Micromonospora violae* (Zhang et al. 2014) which were isolated  
251 from a root nodule of *Lupinus angustifolius* and the roots of *Taraxacum mongolicum*  
252 and *Viola philippica*, respectively (Fig. 3). Moreover, the final strain, isolate STR1s-5,  
253 was most closely related to the type strain of *Micromonospora ureilytica* (Carro et al.  
254 2016b), which was isolated from a root nodule of *Pisum sativum*. In turn, two of the  
255 four strains isolated from the extreme hyper-arid Lomas Bayas soil, isolates LB19 and  
256 LB32, were phylogenetically close to the type strain of *Micromonospora saelicesensis*  
257 (Trujillo et al. 2007), an isolate from a root nodule of *L. angustifolius*. It is also  
258 interesting that isolate LB39 was recovered in the well supported clade that included the

259 type strains of *M. chokoriensis*, *M. taraxaci* and *M. violae* (Fig. 3); the final strain from  
260 the Lomas Bayas soil, isolate LB4, formed a well-supported clade with the type strain  
261 of *Micromonospora chalcea* (Foulerton 1905, Orskov, 1923) which was isolated from  
262 air. It can be seen from the Figure 3 that the sole strain from the extreme hyper-arid  
263 Yungay Core soil, isolate Y6-2, forms a well-supported clade with the type strain of  
264 *Micromonospora pisi* (Garcia et al. 2010), an isolate from a root nodule of *Pisum*  
265 *sativum*.

266 To date, representatives of dereplicated groups of actinobacteria isolated from  
267 Atacama Desert soils which show low pairwise 16S rRNA gene sequence similarities,  
268 ( $\leq 90\%$ ; Meier-Kolthoff et al. 2013) with the type strains of their nearest phylogenetic  
269 neighbour have been invariably assigned to new species when subject to polyphasic  
270 taxonomic analyses, as exemplified by *Lechevalieria*, *Pseudonocardia* and  
271 *Streptomyces* species (Okoro et al. 2010; Busarakam et al. 2014; Trujillo et al. 2017).  
272 This cut-off point has also proved to be a reliable indicator for the presumptive  
273 recognition of novel *Micromonospora* species (Trujillo et al. 2007; Garcia et al. 2010;  
274 Carro et al. 2016a) though pairwise 16S rRNA gene sequence similarities very much  
275 higher than the 99.0% threshold have been found to be indicative of new  
276 *Micromonospora* species, as exemplified by *M. taraxaci* and *M. violae* (Zhang et al.  
277 2014; Zhao et al. 2014). Given these indicators it seems likely that further comparative  
278 taxonomic analyses will show that most, if not all, of the representatives of the BOX-  
279 PCR groups will be found to represent novel *Micromonospora* species. Indeed, isolates  
280 ATA38, ATA39, and ATA43 from ALMA4, isolate STR1-7 from the rhizosphere of *P.*  
281 *quadrangularis*, and isolate Y6\_2 from Yungay are cases in point as they share 16S  
282 rRNA gene sequence similarities with their closest phylogenetic neighbours at or below  
283 the 99.0% threshold (Table 3).

#### 284 *Adaptation to extreme conditions*

285 In general, the pH and temperature profiles of the isolates were typical of  
286 *Micromonospora* strains (Genilloud 2012) as they grew from 20 to 37°C, at pH 6.0 to  
287 8.0, but not at 4, 12 or 45°C or below pH 5.0. None of the isolates grew under anaerobic  
288 conditions, but most of them were able to grow in the presence of 1%, w/v sodium  
289 chloride; six out of the thirteen strains isolated from the rhizosphere of *P.*  
290 *quadrangularis* grew in the presence of 3%, w/v sodium chloride. It was also interesting  
291 that only 30% of the samples from ALMA soil grew on M65 agar following exposure to  
292 UV light (UVC) at 100 mJoules/second for 30 minutes, this number increased to 60%  
293 on ZSSE agar. Similarly, high percentages were observed for strains isolated from the  
294 rhizosphere and Lomas Bayas soils on M65 (64 and 80%) and ZSSE (50 and 60%) agar,  
295 respectively. These results are interesting as it has been shown that the type strain of  
296 *Modestobacter caceresii*, an isolate recovered from a soil sample from the Yungay Core  
297 Region, has the capacity to protect and repair damaged caused by UV radiation  
298 (Busarakam et al. 2016). Interestingly, the sole *Micromonospora* strain obtained from  
299 this area, isolate Y6-2, grew after an hour exposure to UV light at 100 mJoules/second,  
300 only around 20% of the isolates grew under this condition. Although UVC radiation no  
301 longer reaches the Earth's surface it has been used as a selective tool with reference to  
302 early life on the planet and high altitude biology. Recently Paulino-Lima et al. (2016)  
303 screened desert soils, including Atacama soil, for UVC-resistant bacteria and  
304 interestingly over 40% of their isolates were members of the phylum *Actinobacteria*;



305 *Micromonospora* strains were not recovered by these authors. Consequently, the  
306 distinctive radiation resistance of *Micromonospora* strain Y6-2 suggests that it could be  
307 a novel target for detailed radiation-resistant physiological and biochemical researches  
308 that should be extended to include the other radiation stressors naturally present on  
309 the Earth surface. The study of these radiations will be of interest for future studies of  
310 the strains and to determine if key osmoprotectant proteins are implicated in the process  
311 as previously shown in *Rhodobacter* (Pérez et al 2017).

312 Approximately half of the isolates showed weak activity in the plug assays  
313 against the wild type strains of *B. subtilis* and *P. fluorescens* though no activity was  
314 shown against the *E. coli* strain. However, it is important that these preliminary  
315 antibiotic sensitivity studies are extended as the genomes of *Micromonospora* type  
316 strains have much greater potential to synthesize novel specialised metabolites than  
317 previously realised (Carro et al. 2018a). Indeed, novel strains of *Micromonospora* and  
318 corresponding strains of other genera classified in the family *Micromonosporaceae*  
319 should be given greater prominence in the search for new classes of bioactive  
320 compounds, notably antibiotics that are needed to control drug resistant pathogens.

321 It can be concluded that the isolation procedures used in this study provide an  
322 effective way of isolating small numbers of putatively novel *Micromonospora* species  
323 from Atacama Desert soils. Indeed, innovative selective isolation procedures based on  
324 the biological properties of target organisms are needed to cultivate elements of the  
325 extensive actinobacterial dark matter detected in soils of the Atacama Desert landscape  
326 using culture-independent methods (Bull et al. 2017; Idris et al. 2017c)

327

### 328 **Acknowledgments**

329 This project was partly funded by a UK-Newton Project for UK-Chile collaboration  
330 (JIC CA 586) and by support from the Basal Programme of CONICYT for the Centre  
331 for Biotechnology and Bioengineering (CeBiB) (project FB0001). IN and LC thank  
332 Newcastle University for postdoctoral fellowships and ATB and MG are indebted to the  
333 Leverhulme Trust for Emeritus Fellowships.

334

### 335 **Conflict of Interest**

336 The authors declare that they have no conflict of interest.

337

338 This article does not contain any studies with human participants or animals performed  
339 by any of the authors.

340

342 **References**

- 343 Barka EA, Vatsa P, Sanchez L et al. (2016) Taxonomy, physiology, and natural products of  
 344 *Actinobacteria*. *Microbiol Mol Biol Rev* 80:1-43 doi:10.1128/MMBR.00019-15
- 345 Bull AT, Asenjo JA, Goodfellow M, Gomez-Silva B (2016) The Atacama Desert: Technical  
 346 resources and the growing importance of novel microbial diversity. *Annu Rev*  
 347 *Microbiol* 70:215-234 doi:10.1146/annurev-micro-102215-095236
- 348 Bull AT, Idris H, Sanderson R, Asenjo J, Andrews B, Goodfellow M (2017) High altitude, hyper-  
 349 arid soils of the Central-Andes harbor mega-diverse communities of actinobacteria.  
 350 *Extremophiles* doi:10.1007/s00792-017-0976-5
- 351 Busarakam K (2014) Novel Actinobacterial Diversity in Arid Atacama Desert Soils as a Source of  
 352 New Drug Leads. Ph. D. Thesis. Newcastle University, Newcastle upon Tyne
- 353 Busarakam K, Bull AT, Girard G, Labeda DP, van Wezel GP, Goodfellow M (2014) *Streptomyces*  
 354 *leeuwenhoekii* sp. nov., the producer of chaxalactins and chaxamycins, forms a distinct  
 355 branch in *Streptomyces* gene trees. *Antonie van Leeuwenhoek* 105:849-861  
 356 doi:10.1007/s10482-014-0139-y
- 357 Busarakam K, Bull AT, Trujillo ME, Riesco R, Sangal V, van Wezel GP, Goodfellow M (2016)  
 358 *Modestobacter caceresii* sp. nov., novel actinobacteria with an insight into their  
 359 adaptive mechanisms for survival in extreme hyper-arid Atacama Desert soils. *Syst*  
 360 *Appl Microbiol* 39:243-251 doi:10.1016/j.syapm.2016.03.007
- 361 Cameron RE, Gensel DR, Blank CB (1966) Soil studies -desert microflora XII. Abundance of  
 362 microflora in soil samples from the Chile Atacama Desert. space programs summary.  
 363 In: *Supporting Research and Advanced Developments, Space Programs Summary* No.  
 364 37–38, vol IV. Jet Propulsion Lab, Pasadena, CA, pp 140–147
- 365 Carro L, Spröer C, Alonso P, Trujillo ME (2012a) Diversity of *Micromonospora* strains isolated  
 366 from nitrogen fixing nodules and rhizosphere of *Pisum sativum* analyzed by multilocus  
 367 sequence analysis. *Syst Appl Microbiol* 35:73-80 doi:10.1016/j.syapm.2011.11.003
- 368 Carro L, Pukall R, Spröer C, Kroppenstedt RM, Trujillo ME (2012b) *Micromonospora cremea* sp.  
 369 nov. and *Micromonospora zamorensis* sp. nov., isolated from the rhizosphere of *Pisum*  
 370 *sativum*. *Int J Syst Evol Microbiol* 62:2971-2977 doi:10.1099/ijs.0.038695-0
- 371 Carro L, Pujic P, Trujillo ME, Normand P (2013a) *Micromonospora* is a normal occupant of  
 372 actinorhizal nodules. *J Biosci* 38:685-693
- 373 Carro L, Pukall R, Spröer C, Kroppenstedt RM, Trujillo ME (2013b) *Micromonospora*  
 374 *halotolerans* sp. nov., isolated from the rhizosphere of a *Pisum sativum* plant. *Antonie*  
 375 *van Leeuwenhoek* 103:1245-1254 doi:10.1007/s10482-013-9903-7
- 376 Carro L, Riesco R, Spröer C, Trujillo ME (2016a) *Micromonospora luteifusca* sp. nov. isolated  
 377 from cultivated *Pisum sativum*. *Syst Appl Microbiol* 39:237-242  
 378 doi:10.1016/j.syapm.2016.04.003
- 379 Carro L, Riesco R, Spröer C, Trujillo ME (2016b) *Micromonospora ureilytica* sp. nov.,  
 380 *Micromonospora noduli* sp. nov. and *Micromonospora vinacea* sp. nov., isolated from  
 381 *Pisum sativum* nodules. *Int J Syst Evol Microbiol* 66:3509-3514  
 382 doi:10.1099/ijsem.0.001231
- 383 Carro L et al. (2018a) Genome-based classification of micromonosporae with a focus on their  
 384 biotechnological and ecological potential. *Sci Rep* 8(1):525 doi: 10.1038/s41598-017-  
 385 17392-0
- 386 Carro L, Veyisoglu A, Riesco R, Spröer C, Klenk HP, Sahin N, Trujillo ME (2018b)  
 387 *Micromonospora phytophila* sp. nov. and *Micromonospora luteiviridis* sp. nov., isolated  
 388 as natural inhabitants of plant nodules. *Int J Syst Evol Microbiol* 68:248-253  
 389 doi:10.1099/ijsem.0.002490

390 Coombs JT, Franco CM (2003) Isolation and identification of actinobacteria from surface-  
391 sterilized wheat roots. *Appl Environ Microbiol* 69:5603-5608

392 Crisafuli FA, Ramos EB, Rocha MS (2015) Characterizing the interaction between DNA and  
393 GelRed fluorescent stain. *Eur Biophys J* 44:1-7 doi:10.1007/s00249-014-0995-4

394 Demain AL (2014) Importance of microbial natural products and the need to revitalize their  
395 discovery. *J Ind Microbiol Biotechnol* 41:185-201 doi:10.1007/s10295-013-1325-z

396 Ding D, Chen G, Wang B, Wang Q, Liu D, Peng M, Shi P (2013) Cultivable actinomycetes from  
397 desert ecosystem in northeast of Qinghai Plateau. *Ann Microbiol* 63:259-266

398 Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high  
399 throughput. *Nucleic Acids Res* 32:1792-1797 doi:10.1093/nar/gkh340

400 Elsayed SS, Trusch F, Deng H et al. (2015) Chaxapeptin, a lasso peptide from extremotolerant  
401 *Streptomyces leeuwenhoekii* strain C58 from the hyperarid Atacama Desert. *J Org*  
402 *Chem* 80:10252-10260 doi:10.1021/acs.joc.5b01878

403 Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach.  
404 *J Mol Evol* 17:368-376

405 Fiedler HP (2004) Screening for biodiversity. In: Bull AT (eds) *Microbial Diversity and*  
406 *Bioprospecting*. American Society of Microbiology, Washington DC, pp. 324-335

407 Foulerton A (1905). New species of *Streptothrix* isolated from the air. *Lancet* 1199-1200

408 Garcia LC, Martinez-Molina E, Trujillo ME (2010) *Micromonospora pisi* sp. nov., isolated from  
409 root nodules of *Pisum sativum*. *Int J Syst Evol Microbiol* 60:331-337  
410 doi:10.1099/ijs.0.012708-0

411 Gause GF, Preobrazhenskaya TP, Sveshnikova GV, Terekhova LP, Maksimova TS (1983) A Guide  
412 for Determination of Actinomycetes. Nauka, Moscow, Russia

413 Genilloud O (2012) Genus I. *Micromonospora* Ørskov 1923, 156<sup>AL</sup>. In: Goodfellow M, Kämpfer  
414 P, Busse HJ, Trujillo ME, Suzuki KI, Ludwig W, Whitman WB (eds) *Bergey's Manual of*  
415 *Systematic Bacteriology*, 2<sup>nd</sup> edn., vol 5. The *Actinobacteria*, Part B. Springer, New York,  
416 pp 1039–1057

417 Goodfellow M, Hill IR, Gray TRG (1967) Bacteria in a pine forest soil. In: Gray TRG, Parkinson D  
418 (eds) *The Ecology of Soil Bacteria*. University Press, Liverpool, pp 500-515.

419 Goodfellow M (2010) Selective isolation of actinobacteria. In: Baltz RH et al. (eds) *Manual of*  
420 *Industrial Microbiology and Biotechnology*. 3rd edn. American Society of Microbiology,  
421 pp 13-27

422 Goodfellow M, Fiedler HP (2010) A guide to successful bioprospecting: informed by  
423 actinobacterial systematics. *Antonie van Leeuwenhoek* 98:119-142  
424 doi:10.1007/s10482-010-9460-2

425 Goodfellow M, Busarakam K, Idris H et al. (2017) *Streptomyces asenjonii* sp. nov., isolated from  
426 hyper-arid Atacama Desert soils and emended description of *Streptomyces*  
427 *viridosporus* Pridham et al. 1958. *Antonie van Leeuwenhoek* 110:1133-1148  
428 doi:10.1007/s10482-017-0886-7

429 Hayakawa M, Nonomura H (1987) Humic acid vitamin agar, a new medium for the selective  
430 isolation of soil actinomycetes. *J Ferment Technol* 65:501-509

431 Hayakawa M, Sadakata T, Kajiura T, Nonomura H. (1991) New methods for the highly selective  
432 isolation of *Micromonospora* and *Microbispora* from soil. *J Ferment Bioengin* 72:320–  
433 326

434 Hoskisson PA, Hobbs G, Sharples GP (2000) Response of *Micromonospora echinospora* (NCIMB  
435 12744) spores to heat treatment with evidence of a heat activation phenomenon. *Lett*  
436 *Appl Microbiol* 30:114-117

437 Idris H (2016) Actinobacterial Diversity in Atacama Deset habitats as a road map to  
438 biodiscovering. Ph. D. Thesis. Newcastle University, Newcastle upon Tyne

439 Idris H, Labeda DP, Nouioui I et al. (2017a) *Streptomyces aridus* sp. nov., isolated from a high  
440 altitude Atacama Desert soil and emended description of *Streptomyces noboritoensis*

441 Isono et al. 1957. *Antonie van Leeuwenhoek* 110:705-717 doi:10.1007/s10482-017-  
442 0838-2

443 Idris H, Nouioui I, Asenjo JA, Bull AT, Goodfellow M (2017b) *Lentzea chajnantorensis* sp. nov.,  
444 an actinobacterium from a very high altitude Cerro Chajnantor gravel soil in northern  
445 Chile. *Antonie van Leeuwenhoek* 110:795-802 doi:10.1007/s10482-017-0851-5

446 Idris H, Goodfellow M, Sanderson R, Asenjo JA, Bull AT (2017c) Actinobacterial rare biospheres  
447 and dark matter revealed in habitats of the Chilean Atacama Desert. *Sci Rep* 7:8373  
448 doi:10.1038/s41598-017-08937-4

449 Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions  
450 through comparative studies of nucleotide sequences. *J Mol Evol* 16:111-120

451 Krasil'nikov NA (1938) *Ray Fungi and Related Organisms. Actinomycetales*. Akad Nauk, Moscow

452 Kroppenstedt RM, Mayilraj S, Wink JM, Kallow W, Schumann P, Secondini C, Stackebrandt E  
453 (2005) Eight new species of the genus *Micromonospora*, *Micromonospora citrea* sp.  
454 nov., *Micromonospora echinaurantiaca* sp. nov., *Micromonospora echinofusca* sp. nov.,  
455 *Micromonospora fulviviridis* sp. nov., *Micromonospora inyonensis* sp. nov.,  
456 *Micromonospora peucetia* sp. nov., *Micromonospora sagamiensis* sp. nov., and  
457 *Micromonospora viridifaciens* sp. nov. *Syst Appl Microbiol* 28:328-339  
458 doi:10.1016/j.syapm.2004.12.011

459 Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis  
460 Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33:1870-1874  
461 doi:10.1093/molbev/msw054

462 Lane DJ (1991) 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M (eds) *Nucleic Acid*  
463 *Techniques in Bacterial Systematics*. Wiley, New York, pp 115-175

464 Maldonado LA, Stach JE, Ward AC, Bull AT, Goodfellow M (2008) Characterisation of  
465 micromonosporae from aquatic environments using molecular taxonomic methods.  
466 *Antonie van Leeuwenhoek* 94:289-298 doi:10.1007/s10482-008-9244-0

467 Meier-Kolthoff JP, Goker M, Spröer C, Klenk HP (2013) When should a DDH experiment be  
468 mandatory in microbial taxonomy?. *Arch Microbiol* 195:413-418 doi:10.1007/s00203-  
469 013-0888-4

470 Montero-Calasanz MC, Hofner B, Göker M et al. (2014) *Geodermatophilus poikilotropi* sp. nov.:  
471 a multitolerant actinomycete isolated from dolomitic marble. *Biomed Res Int* 9:14767

472 Nachtigall J, Kulik A, Helaly S et al. (2011) Atacamycins A-C, 22-membered antitumor  
473 macrolactones produced by *Streptomyces* sp. C38. *J Antibiot (Tokyo)* 64:775-780  
474 doi:10.1038/ja.2011.96

475 Nonomura H, Ohara Y (1969) Distribution of actinomycetes in soil. VI. A culture method  
476 effective for both preferential isolation and enumeration of *Microbispora* and  
477 *Streptosporangium* strains in soil (Part 1) *J Ferment Technol* 47:463-469

478 Okoro CK, Brown R, Jones AL, Andrews BA, Asenjo JA, Goodfellow M, Bull AT (2009) Diversity  
479 of culturable actinomycetes in hyper-arid soils of the Atacama Desert, Chile. *Antonie*  
480 *van Leeuwenhoek* 95:121-133 doi:10.1007/s10482-008-9295-2

481 Okoro CK, Bull AT, Mutreja A, Rong X, Huang Y, Goodfellow M (2010) *Lechevalieria*  
482 *atacamensis* sp. nov., *Lechevalieria deserti* sp. nov. and *Lechevalieria roselyniae* sp.  
483 nov., isolated from hyperarid soils. *Int J Syst Evol Microbiol* 60:296-300  
484 doi:10.1099/ijs.0.009985-0

485 Opfell JB, Zebal GP (1967) Ecological patterns of micro-organisms in desert soils. *Life Sci Space*  
486 *Res* 5:187-203

487 Ørskov J (1923) *Investigations into the Morphology of the Ray Fungi*. Levin and Munksgaard,  
488 Copenhagen

489 Parte AC (2014) LPSN—list of prokaryotic names with standing in nomenclature. *Nucleic Acids*  
490 *Research* 42:D613-D616 doi:10.1093/nar/gkt1111

491 Paulino-Lima IG, Fujishima K, Navarrete JU et al. (2016) Extremely high UV-C radiation resistant  
492 microorganisms from desert environments with different manganese concentrations. J  
493 Photochem Photobiol B-Biol 163: 327-336

494 Pérez V, Hengst M, Kurte L et al. (2017) Bacterial survival under extreme UV radiation : a  
495 comparative proteomics study of *Rhodobacter* sp., isolated from high altitude  
496 wetlands in Chile Front Microbiol 8:1173 doi:10.3389/fmicb.2017.01173

497 Rateb ME, Houssen WE, Arnold M et al. (2011a) Chaxamycins A-D, bioactive ansamycins from a  
498 hyper-arid desert *Streptomyces* sp. J Nat Prod 74:1491-1499 doi:10.1021/np200320u

499 Rateb ME, Houssen WE, Harrison WT et al. (2011b) Diverse metabolic profiles of a  
500 *Streptomyces* strain isolated from a hyper-arid environment. J Nat Prod 74:1965-1971  
501 doi:10.1021/np200470u

502 Reasoner DJ, Blannon JC, Geldreich EE (1979) Rapid seven-hour fecal coliform test. Appl  
503 Environ Microbiol 38:229-236

504 Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing  
505 phylogenetic trees. Mol Biol Evol 4:406-425

506 Shen Y, Zhang Y, Liu C et al. (2014) *Micromonospora zae* sp. nov., a novel endophytic  
507 actinomycete isolated from corn root (*Zea mays* L.). J Antibiot (Tokyo) 67:739-743  
508 doi:10.1038/ja.2014.54

509 Shirling EB, Gottlieb D (1966) Methods for characterization of *Streptomyces* species. Int J Syst  
510 Evol Microbiol 16:313-340 doi:doi:10.1099/00207713-16-3-313

511 Thawai C (2015) *Micromonospora costi* sp. nov., isolated from a leaf of *Costus speciosus*. Int J  
512 Syst Evol Microbiol 65:1456-1461 doi:10.1099/ijms.0.000120

513 Thawai C, Kittiwongwattana C, Thanaboripat D, Laosinwattana C, Koohakan P, Parinthawong N  
514 (2016) *Micromonospora soli* sp. nov., isolated from rice rhizosphere soil. Antonie van  
515 Leeuwenhoek 109:449-456 doi:10.1007/s10482-016-0651-3

516 Thawai C, Tanasupawat S, Itoh T, Suwanborirux K, Suzuki K, Kudo T (2005) *Micromonospora*  
517 *eburnea* sp. nov., isolated from a Thai peat swamp forest. Int J Syst Evol Microbiol  
518 55:417-422 doi:10.1099/ijms.0.63217-0

519 Thawai C, Tanasupawat S, Suwanborirux K, Itoh T, Kudo T (2007) *Micromonospora*  
520 *narathiwatensis* sp. nov., from Thai peat swamp forest soils. J Gen Appl Microbiol  
521 53:287-293

522 Trujillo ME, Kroppenstedt RM, Schumann P, Carro L, Martinez-Molina E (2006)  
523 *Micromonospora coriariae* sp. nov., isolated from root nodules of *Coriaria myrtifolia*.  
524 Int J Syst Evol Microbiol 56:2381-2385 doi:10.1099/ijms.0.64449-0

525 Trujillo ME, Kroppenstedt RM, Fernandez-Molinero C, Schumann P, Martinez-Molina E (2007)  
526 *Micromonospora lupini* sp. nov. and *Micromonospora saelicesensis* sp. nov., isolated  
527 from root nodules of *Lupinus angustifolius*. Int J Syst Evol Microbiol 57:2799-2804  
528 doi:10.1099/ijms.0.65192-0

529 Trujillo ME, Alonso-Vega P, Rodriguez R, Carro L, Cerda E, Alonso P, Martinez-Molina E (2010)  
530 The genus *Micromonospora* is widespread in legume root nodules: the example of  
531 *Lupinus angustifolius*. ISME J 4:1265-1281 doi:10.1038/ismej.2010.55

532 Trujillo ME, Idris H, Riesco R, Nouioui I, Igual JM, Bull AT, Goodfellow M (2017) *Pseudonocardia*  
533 *nigra* sp. nov., isolated from Atacama Desert rock Int J Syst Evol Microbiol 67:2980-  
534 2985 doi:10.1099/ijsem.0.002063

535 Versalovic J, Schneider M, De Bruijn FJ, Lupski JR (1994) Genomic fingerprinting of bacteria  
536 using repetitive sequence-based polymerase chain reaction. Meth Mol Cell Biol 5:25-  
537 40

538 Wichner D, Idris H, Houssen WE et al. (2017) Isolation and anti-HIV-1 integrase activity of  
539 lentzeosides A-F from extremotolerant *Lentzea* sp. H45, a strain isolated from a high-  
540 altitude Atacama Desert soil. J Antibiot 70:448-453 doi:10.1038/ja.2016.78

541 Williams ST, Goodfellow M, Vickers JC (1984) New microbes from old habitats? In: Carr WG,  
542 Kelly D (eds) *The Microbe 1984. II Prokaryotes and Eukaryotes*. Cambridge University  
543 Press, Cambridge, pp 219-256

544 Yoon S-H, Ha S-M, Kwon S, Lim J, Kim Y, Seo H, Chun J (2016) Introducing EzBioCloud: A  
545 taxonomically united database of 16S rRNA and whole genome assemblies. *Int J Syst*  
546 *Evol Microbiol* doi:doi:10.1099/ijsem.0.001755

547 Zhang J (2011) Improvement of an isolation medium for actinomycetes. *Modern Appl Sci*  
548 5:124-127

549 Zhang Y, Liu H, Zhang X et al. (2014) *Micromonospora violae* sp. nov., isolated from a root of  
550 *Viola philippica* Car. *Antonie van Leeuwenhoek* 106:219-225 doi:10.1007/s10482-014-  
551 0184-6

552 Zhao J, Guo L, He H et al. (2014) *Micromonospora taraxaci* sp. nov., a novel endophytic  
553 actinomycete isolated from dandelion root (*Taraxacum mongolicum* Hand.-Mazz.).  
554 *Antonie van Leeuwenhoek* 106:667-674 doi:10.1007/s10482-014-0237-x

555 Zhao S, Liu C, Zheng W et al. (2017) *Micromonospora parathelypteridis* sp. nov., an endophytic  
556 actinomycete with antifungal activity isolated from the root of *Parathelypteris*  
557 *beddomei* (Bak.) Ching. *Int J Syst Evol Microbiol* 67:268-274 doi:  
558 10.1099/ijsem.0.001614

559 Zhi XY, Li WJ, Stackebrandt E (2009) An update of the structure and 16S rRNA gene sequence-  
560 based definition of higher ranks of the class *Actinobacteria*, with the proposal of two  
561 new suborders and four new families and emended descriptions of the existing higher  
562 taxa. *Int J Syst Evol Microbiol* 59:589-608 doi:10.1099/ijms.0.65780-0

563

564

565

566 **Figure legends**

567

568 **Figure 1.** A. Sampling sites: Lomas Bayas (LB), Salar de Tara (ST), Atacama Large  
569 Millimeter Array observatory (ALMA) and Yungay (Y6); B. Salar de Tara site; C.  
570 *Parastrephia. quadrangularis* plant growing at the Salar the Tara; D. Typical orange  
571 colony of a *Micromonospora* isolate; E. Typical black *Micromonospora* colony  
572 following spore production; F-H. Morphology observed for some Atacama  
573 micromonosporae; F. Sporulating *Micromonospora* colonies; G. Mountain-shaped  
574 *Micromonospora* isolates; H. Colony from ALMA4 isolate covered by white aerial  
575 hyphae.

576

577 **Figure 2.** BOX–PCR fingerprints showing the genetic diversity of *Micromonospora*  
578 strains isolated from Atacama Desert soils.

579

580 **Figure 3.** Neighbour-joining phylogenetic tree based on almost complete 16S rRNA  
581 gene sequences showing relationships between the *Micromonospora* strains and  
582 between them and *Micromonospora* type strains. Numbers at the nodes indicate the  
583 levels of bootstrap support (%), only values above 50% are shown. Asterisks indicate  
584 branches of the tree that were also recovered in the maximum-likelihood tree.  
585 *Catellatospora citrea* was used as the outgroup. Bar, 0.005 substitutions per nucleotide  
586 position.

587

589 Table 1. Location, sampling sites and dates of collection of soil samples from the  
590 Atacama Desert.

Location	Sampling site and code	Collection date	Altitude (masl)	Latitude (°S)	Longitude (°W)	Habitat	N° of isolates
Cerro Chajnantor+	ALMA 4 ATA, 4G	20.10.12	4000	23°03'31''	67°52'27''	Subsurface soil (30 cm)	25
	ALMA 5 5R2A	20.10.12	5046	23°00'49''	67°45'31''	Surface soil (2 cm)	1
Lomas Bayas	LB 3	24.02.14	1500	23°24'27''	69°31'03''	Extreme hyper-arid surface soil (2 cm)	5
Yungay Core Region	Cerro Aguas Blancas, Y6	13.11.10	1047	24°06'18''	70°01'15''	Extreme hyper-arid subsurface soil (30 cm)	1
Salar de Tara	STR1	05.10.16	4174	23°03'97''	67°18'87''	Rhizosphere of <i>Parastrephia quadrangularis</i>	13

591 +The Atacama Large Millimeter Array (ALMA) Observatory is situated on Cerro  
592 Chajnantor: ALMA is operated as an international partnership which includes the  
593 European Southern Observatory (ESO). Permission to collect soil samples from Cerro  
594 Chajnantor was given by the Director of the ESO.

595 Table 2. Source and substrate mycelial colour of the isolated *Micromonospora* strains.

	Isolates	Isolation protocol	Isolation media	Colour of substrate mycelium
	4G51, 4G53, 4G55, 4G57	D	G nº 1	green
ALMA 4	ATA32, ATA34, ATA47	P-H	ZSSE	orange
	ATA48	P-H	CHV	orange
	ATA31, ATA33, ATA35, ATA36a, ATA36b, ATA37, ATA38, ATA39, ATA40, ATA42, ATA43, ATA44, ATA45	P-H	ZSSE	green
	ATA50	P-H	CHV	green
	ATA46, ATA51	P-H	HV	green
	ATA52	P-H	HV	black
ALMA 5	5R2A7	D	R2A	orange
Lomas Bayas	LB 4, LB 19, LB 32, LB 39, LB 41	D	M65	orange
Yungay core region	Y6_2	P-H	HV	orange



	STR1-41, STR1-74, STR1-85	P-H	HV	orange
Salar de Tara	STR1s-5, STR1s-6, STR1s-7, STR1s-11	S	HV	orange
(rhizosphere)	STR1-7, STR1-71, STR1-72	P-H	ZSSE	orange
	STR1s-13a, STR1s-14, STR1s-16	S	ZSSE	orange

596

597

598 Table 3. Assignment of isolates to BOX groups and determination of their nearest  
599 phylogenetic neighbours.

<b>BOX group</b>	<b>Isolates</b>	<b>Representative isolates</b>	<b>% Similarity to closest phylogenetic type strain</b>	
<b>I</b>	ATA31, ATA33, ATA45, ATA47, ATA48, ATA52	ATA 45, ATA52	<i>M. costi</i>	99.1
<b>II</b>	ATA35, ATA37, ATA40	ATA40	<i>M. costi</i>	99.1
<b>III</b>	ATA38	ATA38	<i>M. costi</i>	98.4
<b>IV</b>	ATA39	ATA39	<i>M. costi</i>	98.4
<b>V</b>	ATA42	ATA42	<i>M. costi</i>	99.1
<b>VI</b>	ATA43	ATA43	<i>M. terminaliae</i>	98.8
<b>VII</b>	ATA32, ATA34	ATA32	<i>M. eburnea</i>	99.4
<b>VIII</b>	LB4, LB41	LB4	<i>M. chalcea</i>	99.6
<b>IX</b>	LB19	LB19	<i>M. saelicesensis</i>	99.8
<b>X</b>	LB32	LB32	<i>M. saelicesensis</i>	99.8
<b>XI</b>	LB39	LB39	<i>M. chokoriensis</i>	99.9
<b>XII</b>	ATA36, ATA44, 4G51, ATA 46, 4G55, 4G53, 4G57	4G51, 4G57	<i>M. costi</i>	99.1
<b>XIII</b>	ATA51a, ATA51b	ATA51b	<i>M. costi</i>	99.1
<b>XIV</b>	5R2A7	5R2A7	<i>M. coriariae</i>	99.8
<b>XV</b>	STR1-7, STR1-41, STR1-72, STR1-74, STR1-85, STR1s-7, STR1s-11, STR1s-13A, STR1s-14, STR1s-16	STR1-7	<i>M. chokoriensis</i>	99
<b>XVI</b>	STR1s-5	STR1s-5	<i>M. chokoriensis</i>	99.4
<b>XVII</b>	STR1s-6	STR1s-6	<i>M. ureilytica</i>	99.6
<b>XVIII</b>	Y6-2	Y6-2	<i>M. pisi</i>	98.8

600

601