

## Research Article

# On the Generalized Weighted Lebesgue Spaces of Locally Compact Groups

**I. Akbarbaglu and S. Maghsoudi**

*Department of Mathematics, Zanjan University, Zanjan 45195-313, Iran*

Correspondence should be addressed to S. Maghsoudi, s\_maghsodi@znu.ac.ir

Received 11 June 2011; Accepted 16 August 2011

Academic Editor: Narcisa C. Apreutesei

Copyright © 2011 I. Akbarbaglu and S. Maghsoudi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let  $G$  be a locally compact group with a fixed left Haar measure  $\lambda$  and  $\Omega$  be a system of weights on  $G$ . In this paper, we deal with locally convex space  $L^p(G, \Omega)$  equipped with the locally convex topology generated by the family of norms  $(\|\cdot\|_{p, \omega})_{\omega \in \Omega}$ . We study various algebraic and topological properties of the locally convex space  $L^p(G, \Omega)$ . In particular, we characterize its dual space and show that it is a semireflexive space. Finally, we give some conditions under which  $L^p(G, \Omega)$  with the convolution multiplication is a topological algebra and then characterize its closed ideals and its spectrum.

## 1. Introduction

Throughout this paper, let  $G$  denote a locally compact Hausdorff group with a fixed left Haar measure  $\lambda$ . By a weight function on  $G$ , we mean an arbitrary strictly positive measurable function on  $G$ , and, by a system of weights on  $G$ , a set of weight functions  $\Omega$  such that given  $\omega_1, \omega_2$  in  $\Omega$  and  $c > 0$ , there is an  $\nu \in \Omega$  such that  $c\omega_i(x) \leq \nu(x)$  ( $i = 1, 2$ ) for locally almost all  $x \in G$ .

For a weight function  $\omega$  and  $1 \leq p < \infty$ , let  $L^p(G, \omega)$  denote the space of all complex-valued measurable functions  $f$  on  $G$  such that  $f\omega \in L^p(G)$ , the usual Lebesgue space on  $G$  with respect to  $\lambda$ ; see [1] for more details. Then,  $L^p(G, \omega)$  with the norm  $\|\cdot\|_{p, \omega}$  defined by  $\|f\|_{p, \omega} := \|f\omega\|_p$  for all  $f \in L^p(G, \omega)$  is a Banach space. We also denote by  $L^\infty(G, 1/\omega)$  the space of all measurable complex-valued functions  $f$  on  $G$  such that  $f/\omega \in L^\infty(G)$ , the space defined in [1]. Then,  $L^\infty(G, 1/\omega)$  with the norm  $\|\cdot\|_{\infty, \omega}$  defined by  $\|f\|_{\infty, \omega} := \|f/\omega\|_\infty$  for all  $f \in L^\infty(G, 1/\omega)$  is a Banach space. Furthermore, for  $1 \leq p < \infty$ , the topological dual of  $L^p(G, \omega)$  coincides with  $L^q(G, 1/\omega)$ , where  $q$  is the exponential conjugate

to  $p$  defined by  $1/p + 1/q = 1$ . In fact, the mapping  $T$  from  $L^q(G, 1/\omega)$  to  $L^p(G, \omega)$  defined by

$$\langle T(f), g \rangle = \int_G f(x)g(x)d\lambda(x) \quad (1.1)$$

is an isometric isomorphism; see for example [2]. For measurable functions  $f$  and  $g$  on  $G$ , the convolution multiplication

$$(f * g)(x) = \int_G f(y)g(y^{-1}x)d\lambda(y) \quad (1.2)$$

is defined at each point  $x \in G$  for which this makes sense. The algebraic and topological properties of weighted  $L^p$ -spaces have been studied extensively; see for example [2–5].

Let  $1 \leq p < \infty$  and  $\Omega$  be a system of weights on  $G$ , we set

$$L^p(G, \Omega) = \bigcap_{\omega \in \Omega} L^p(G, \omega). \quad (1.3)$$

In this paper, we equip the space  $L^p(G, \Omega)$  with the natural locally convex topology generated by the family of norms  $\|\cdot\|_{p, \omega}$ , where  $\omega$  runs through  $\Omega$ . For a similar study in other contexts, see [6–8]. We investigate certain algebraic and topological properties of the locally convex space  $L^p(G, \Omega)$ . Our results generalize and improve some interesting results of [5] and partially answer a question raised in [3].

## 2. Preliminaries and Some Basic Results

Let  $G$  be a locally compact Hausdorff group with a fixed left Haar measure  $\lambda$  and  $\Omega$  be a system of weights on  $G$ . We equip  $L^p(G, \Omega)$  with the locally convex topology generated by the family of norms  $(\|\cdot\|_{p, \omega})_{\omega \in \Omega}$  and denote this topology by  $\tau_\Omega$ . So  $(L^p(G, \Omega), \tau_\Omega)$  has a basis of closed absolutely convex neighbourhoods at the origin of the form

$$V_{p, \omega} = \left\{ f \in L^p(G, \Omega) : \|f\|_{p, \omega} \leq 1 \right\}, \quad (\omega \in \Omega). \quad (2.1)$$

Note that the topology  $\tau_\Omega$  is Hausdorff, because if  $f \in L^p(G, \Omega)$  and  $f \neq 0$ , we have  $\lambda(\{x \in G : f(x) \neq 0\}) > 0$ . Put  $E = \{x \in G : f(x) \neq 0\}$  and fix  $\omega \in \Omega$ . Then,

$$\|f\|_{p, \omega} = \left( \int_G (|f|\omega)^p d\lambda \right)^{1/p} \geq \left( \int_E (|f|\omega)^p d\lambda \right)^{1/p} > 0, \quad (2.2)$$

and thus  $\tau_\Omega$  is Hausdorff.

If  $\Omega$  and  $\Gamma$  are two systems of weights on  $G$  and for every  $\omega \in \Omega$ , there is a  $\nu \in \Gamma$  such that  $\omega \leq \nu$  (pointwise locally almost everywhere on  $G$ ), then we write  $\Omega \leq \Gamma$ . In the case which  $\Gamma \leq \Omega$  and  $\Omega \leq \Gamma$ , we write  $\Omega \sim \Gamma$ .

**Proposition 2.1.** *Let  $\Omega$  and  $\Gamma$  be two systems of weights on  $G$  and  $T : G \rightarrow G$  be a measurable mapping such that  $\Omega \leq \Gamma \circ T := \{\nu \circ T : \nu \in \Gamma\}$ . If the Radon-Nikodym function  $h = d(\lambda \circ T^{-1})/d\lambda$  belongs to  $L^\infty(G)$ , then the mapping  $f \mapsto f \circ T$  is a continuous linear map from  $(L^p(G, \Gamma), \tau_\Gamma)$  into  $(L^p(G, \Omega), \tau_\Omega)$ .*

*Proof.* Given  $f \in L^p(G, \Gamma)$  and  $\omega \in \Omega$ , choose  $\nu \in \Gamma$  such that  $\omega \leq \nu \circ T$ . Then we have

$$\begin{aligned} \|f \circ T\|_{p, \omega} &= \left( \int_G (|f \circ T(x)| \omega(x))^p d\lambda(x) \right)^{1/p} \leq \left( \int_G (|f \circ T(x)| (\nu \circ T)(x))^p d\lambda(x) \right)^{1/p} \\ &= \left( \int_G (|f(x)| \nu(x))^p d(\lambda \circ T^{-1})(x) \right)^{1/p} \leq \left( \int_G (|f(x)| \nu(x))^p h(x) d\lambda(x) \right)^{1/p} \\ &\leq \|h\|_\infty \|f\|_{p, \nu} < \infty. \end{aligned} \quad (2.3)$$

Hence,  $\omega(f \circ T) \in L^p(G)$ . Since  $\omega \in \Omega$  was arbitrary,  $f \circ T \in L^p(G, \Omega)$ . Continuity also follows from the above relations.  $\square$

The space of all bounded Borel measurable functions on  $G$  with compact support will be denoted by  $B_c(G)$ . Let us remark that if  $B_c(G) \subseteq L^p(G, \Omega)$ , then  $B_c(G)$  is norm dense in  $L^p(G, \omega)$  for any weight  $\omega$  on  $G$ ; see for example [9].

**Corollary 2.2.** *Let  $\Omega$  and  $\Gamma$  be two systems of weights on  $G$ . Then,*

- (i) *If  $\Omega \leq \Gamma$ , then the induced topology  $\tau_\Omega$  on  $L^p(G, \Gamma)$  is weaker than  $\tau_\Gamma$ .*
- (ii) *If  $B_c(G) \subseteq L^p(G, \Gamma) \subseteq L^p(G, \Omega)$  and  $\tau_\Omega \subseteq \tau_\Gamma$ , then  $\Omega \leq \Gamma$ . In particular,  $\Omega \sim \Gamma$  if and only if  $L^p(G, \Gamma) = L^p(G, \Omega)$ .*

*Proof.* (i) is trivial. For (ii), we observe that for any  $\omega \in \Omega$ , there is a  $\nu \in \Gamma$  such that  $V_{p, \nu} \subseteq V_{p, \omega} \cap L^p(G, \Gamma)$ . So the identity map  $I$  from  $(L^p(G, \Omega), \|\cdot\|_{p, \nu})$  into  $(L^p(G, \omega), \|\cdot\|_{p, \omega})$  is continuous. Since  $L^p(G, \Gamma)$  is dense in  $(L^p(G, \nu), \|\cdot\|_{p, \nu})$ ,  $I$  can be extended continuously to a continuous linear mapping on  $L^p(G, \nu)$ . The extension map is again the identity map. So  $L^p(G, \nu) \subseteq L^p(G, \omega)$ . Hence, there exists a constant  $c > 0$  such that  $\omega(x) \leq c \nu(x)$  locally almost everywhere; see Lemma 2.1 in [10]. This proves that  $\Omega \leq \Gamma$ .  $\square$

Let us recall the definition of the projective limit of a family of locally convex spaces. Let  $(\Lambda, \leq)$  be a partially ordered set and  $\{X_\alpha : \alpha \in \Lambda\}$  be a family of locally convex spaces, and for  $\alpha \leq \beta$ ,  $f_{\alpha, \beta}$  be a linear map from  $X_\beta$  into  $X_\alpha$ . Suppose that  $f_{\alpha\gamma} = f_{\alpha\beta} \circ f_{\beta\gamma}$  for all  $\alpha \leq \beta \leq \gamma$  and  $f_{\alpha\alpha}$  be the identity map on  $X_\alpha$  for all  $\alpha \in \Lambda$ . Then, the projective limit of the family  $(X_\alpha, f_{\alpha, \beta})$  is defined as

$$\lim_{\alpha} (X_\alpha, f_{\alpha, \beta}) = \left\{ (x_\alpha) \in \prod_{\alpha \in \Lambda} X_\alpha : x_\alpha = f_{\alpha, \beta}(x_\beta), \text{ whenever } \alpha \leq \beta \right\}; \quad (2.4)$$

for more details see for example [11].

**Proposition 2.3.** *Let  $\Omega$  be a system of weights on  $G$ . Then  $(L^p(G, \Omega), \tau_\Omega)$  is a complete space.*

*Proof.* We note that for any two weights  $\omega, \nu \in \Omega$  with  $\omega \leq \nu$ ,  $L^p(G, \nu) \subseteq L^p(G, \omega)$ . Let the mapping  $I_{\omega, \nu} : L^p(G, \nu) \rightarrow L^p(G, \omega)$  be the canonical injection. Then, it is clear that  $(L^p(G, \Omega), \tau_\Omega)$  is isomorphic to the projective limit system  $\lim_{\omega} (L^p(G, \omega), I_{\omega, \nu})$  of the Banach spaces  $(L^p(G, \omega), \|\cdot\|_{p, \omega})$ ,  $\omega \in \Omega$ , and, hence, is complete; see Lemma 3.2.1 in [12].  $\square$

**Proposition 2.4.** *The locally convex space  $(L^p(G, \Omega), \tau_\Omega)$  is normable if and only if the topology  $\tau_\Omega$  is generated by  $\|\cdot\|_{p, \omega}$  for some  $\omega \in \Omega$ .*

*Proof.* If  $L^p(G, \Omega)$  is normable, then it has a neighbourhood  $V$  of zero that is norm bounded with respect to  $\|\cdot\|_{p, \omega}$  for every  $\omega \in \Omega$ . Hence, there is  $\omega' \in \Omega$  so that  $V_{p, \omega'} = \{f \in L^p(G, \Omega) : \|f\|_{p, \omega'} \leq 1\}$  is norm bounded in the space  $(L^p(G, \nu), \|\cdot\|_{p, \nu})$  for every  $\nu \in \Omega$ . This implies that there is a positive constant  $c_\nu$  so that  $V_{p, \omega'} \subseteq c_\nu V_{p, \nu}$ , and our claim is proved. The converse is clear.  $\square$

### 3. The Dual and Bidual of $L^p(G, \Omega)$ , $1 \leq p < \infty$

In this section we deal with the dual space of  $(L^p(G, \Omega), \tau_\Omega)$  and, among other things, characterize its equicontinuous subsets.

**Theorem 3.1.** *If  $1 \leq p < \infty$  and  $B_c(G) \subseteq L^p(G, \Omega)$ , then the dual space of  $(L^p(G, \Omega), \tau_\Omega)$  is  $\Omega \cdot L^q(G) := \{\omega f : \omega \in \Omega, f \in L^q(G)\}$  with  $1/p + 1/q = 1$ .*

*Proof.* Let  $h \in L^q(G, 1/\omega)$ . We define the linear functional  $F : L^p(G, \Omega) \rightarrow \mathbb{C}$  by  $F(f) = \int_G fh \, d\lambda$ , then  $F \in (L^p(G, \Omega), \tau_\Omega)^*$ .

Conversely, let  $F \in (L^p(G, \Omega), \tau_\Omega)^*$ . First, we know that  $B_c(G) \subseteq L^p(G, \Omega) \subseteq L^p(G, \omega)$  for every  $\omega \in \Omega$ . So there is a  $\nu \in \Omega$  such that  $|F(f)| \leq 1$  whenever  $f \in \{h \in L^p(G, \Omega) : \|h\|_{p, \nu} \leq 1\}$ . As  $F$  is bounded in the intersection of the unit ball of  $(L^p(G, \nu), \|\cdot\|_{p, \nu})$  with  $(L^p(G, \Omega), \|\cdot\|_{p, \nu})$ ,  $F$  is continuous on  $L^p(G, \Omega)$  with the topology induced by the norm  $\|\cdot\|_{p, \nu}$ . Since  $L^p(G, \Omega)$  is dense in  $(L^p(G, \nu), \|\cdot\|_{p, \nu})$ ,  $F$  can be extended continuously to a continuous linear form on  $L^p(G, \nu)$  which we denote by  $\tilde{F}$ . Then, we have  $\tilde{F} \in (L^p(G, \nu), \|\cdot\|_{p, \nu})^*$ , and hence there is a unique  $h \in L^q(G, 1/\nu)$  so that

$$\tilde{F}(f) = \int_G fh \, d\lambda \quad (f \in L^p(G, \nu)); \quad (3.1)$$

therefore, we obtain the following isomorphism:

$$\Phi : \bigcup_{\omega \in \Omega} L^q\left(G, \frac{1}{\omega}\right) \longrightarrow (L^p(G, \Omega), \tau_\Omega)^*, \quad (3.2)$$

defined by  $\Phi(h) = F_h$ , where  $F_h(f) = \int_G fh \, d\lambda$  for all  $f \in L^p(G, \Omega)$ .  $\square$

**Lemma 3.2.** *Let  $\Omega$  be a system of weights on  $G$ . For every  $\omega \in \Omega$ , define the mapping  $T_\omega : L^p(G, \Omega) \rightarrow L^p(G)$  by  $T_\omega(f) = f\omega$ . Then,  $V_{p, \omega}^\circ = T_\omega^*(B^\circ)$  for  $\omega \in \Omega$ , where  $B$  is the closed unit ball of  $L^p(G)$  and  $B^\circ$  denotes its polar.*

*Proof.* It is clear that  $T_\omega$  is a well-defined continuous linear map. Also,  $T_\omega(L^p(G, \Omega))$  is dense in  $(L^p(G), \|\cdot\|_p)$ . Therefore  $T_\omega^*$  (the adjoint of  $T_\omega$ ) is weak\* continuous and one to one linear map

from  $L^q(G)$  into  $\Omega \cdot L^q(G)$ , where  $1/p + 1/q = 1$ . Now, since  $B^\circ$  is  $\sigma(L^q(G), L^p(G))$ -compact by the Alaoglu theorem and so  $T_\omega^*(B^\circ)$  is  $\sigma(\Omega \cdot L^q(G), L^p(G, \Omega))$ -compact, while  $T_\omega^*(B^\circ)$  is obviously convex. So we find that

$$\begin{aligned} V_{p,\omega} &= \{f \in L^p(G, \Omega) : T_\omega(f) \in B\} = T_\omega^{-1}(B) = \{f \in L^p(G, \Omega) : T_\omega(f) \in B^{\circ\circ}\} \\ &= \{f \in L^p(G, \Omega) : |T_\omega^*(g)(f)| \leq 1, \text{ for every } g \in B^\circ\} = T_\omega^*(B^\circ)^\circ. \end{aligned} \quad (3.3)$$

From which it follows that

$$V_{p,\omega}^\circ = T_\omega^*(B^\circ)^{\circ\circ} = T_\omega^*(B^\circ). \quad (3.4)$$

□

We have the following characterization of the equicontinuous subsets of  $\Omega \cdot L^q(G)$ .

**Theorem 3.3.** *Let  $1 \leq p < \infty$  and  $M$  be a subset of  $(L^p(G, \Omega), \tau_\Omega)^* = \Omega \cdot L^q(G)$ . The following are equivalent.*

- (a)  $M$  is  $\tau_\Omega$ -equicontinuous.
- (b) There are  $\omega \in \Omega$  and an equicontinuous subset  $M'$  of  $(L^p(G), \|\cdot\|_p)^* = L^q(G)$  so that  $M \subseteq \omega \cdot M'$ .
- (c) There are  $\omega \in \Omega$  and  $\alpha > 0$  such that  $\sup\{\|f/\omega\|_q : f \in M\} \leq \alpha < \infty$  whenever  $1/p + 1/q = 1$ .

*Proof.* (a  $\Rightarrow$  b) By (a), there is  $\omega \in \Omega$  so that  $M \subseteq V_{p,\omega}^\circ$ , where  $V_{p,\omega} = \{f \in L^p(G, \Omega) : \|f\|_{p,\omega} \leq 1\}$ . According to Lemma 3.2, we have  $V_{p,\omega}^\circ = T_\omega^*(B^\circ)$ , where  $B$  is the closed unit ball of  $L^p(G)$ . Hence  $M \subseteq \omega B^\circ$ .

- (b  $\Rightarrow$  c) There is  $\alpha > 0$  so that  $M' \subseteq \alpha B^\circ$  by (b). So  $M \subseteq \alpha \omega B^\circ$ , and  $\sup_{f \in M} \|f/\omega\|_q \leq \alpha$ .
- (c  $\Rightarrow$  a) If  $p = 1$ , it is clear that

$$M \subseteq \left\{ f \in L^p(G, \Omega) : \int_G |f(x)|\omega(x)d\lambda(x) \leq \frac{1}{\alpha} \right\}^\circ, \quad (3.5)$$

and if  $1 < p < \infty$ , by Hölder's inequality, for  $h \in M$  and

$$f \in W = \left\{ f \in L^p(G, \Omega) : \|f\|_{p,\omega} \leq \frac{1}{\alpha} \right\}, \quad (3.6)$$

we have

$$\left| \int_G hf \, d\lambda \right| \leq \int_G \left| \frac{h}{\omega} \right| |f\omega| \, d\lambda \leq \left\| \frac{h}{\omega} \right\|_q \|f\omega\|_p \leq 1. \quad (3.7)$$

Hence,  $M \subseteq W^\circ$ , and this guarantees that  $M$  is  $\tau_\Omega$ -equicontinuous in both cases. □

**Proposition 3.4.** *Let  $\Omega$  be a system of weights on  $G$ . Then, the set of extreme points of  $V_{p,\omega}^\circ$  is the set  $\{\omega f : f \in L^q(G), \|f\|_q = 1\}$  for  $1 < p < \infty$ , and  $\{f \in L^\infty(G) : |f| = 1 \text{ l.a.e.}\}$  for  $p = \infty$ .*

*Proof.* Fix  $\omega \in \Omega$  and let  $T_\omega : L^p(G, \Omega) \rightarrow L^p(G)$  be the map defined in Lemma 3.2. From Lemma 3.2, it follows that for any extreme point  $h$  of  $V_{p,\omega}^\circ$ , there is an extreme point  $f$  of  $B^\circ$  so that  $h = T_\omega^*(f) = f\omega$ .

Conversely, let  $\omega \in \Omega$  be arbitrary and let  $h = \omega f$ , where  $f$  is an extreme point of  $B^\circ$ . Clearly,  $h \in V_{p,\omega}^\circ$ , and if  $h = cg + (1-c)k$  for some  $g, k \in V_{p,\omega}^\circ$  and  $0 < c < 1$ , then there are  $m, n \in B^\circ$  such that  $T_\omega^*(m) = g$  and  $T_\omega^*(n) = k$ . Thus,  $T_\omega^*(f) = h = T_\omega^*(cm + (1-c)n)$  and since  $T_\omega^*$  is one to one,  $f = cm + (1-c)n$ . However  $f$  is an extreme point of  $B^\circ$ , which implies that  $f = m = n$ , and hence  $h = g = k$ , that is,  $h$  is an extreme point of  $V_{p,\omega}^\circ$ . Now the rest of the proof is easy to complete; see for example Section 2.14 in [13].  $\square$

Let us recall that a locally convex space  $(E, \tau)$  is called semireflexive if  $(E, \tau)^{**} = E$ .

**Theorem 3.5.** *Let  $\Omega$  be a system of weights on  $G$ . Then  $(L^p(G, \Omega), \tau_\Omega)$  is semireflexive.*

*Proof.* If  $F \in (L^p(G, \Omega), \tau_\Omega)^{**}$ , then the restriction of  $F$  to  $L^q(G, 1/\omega)$ , for every  $\omega \in \Omega$ , belongs to  $L^q(G, 1/\omega)^*$ , where  $L^q(G, 1/\omega)$  was considered with the induced strong topology on  $(L^p(G, \Omega), \tau_\Omega)^*$ . Now if  $\{h_\alpha\}_{\alpha \in I} \subseteq L^q(G, 1/\omega)$  and  $h_\alpha \rightarrow h$  for some  $h \in L^q(G, 1/\omega)$  in the norm  $\|\cdot\|_{q, 1/\omega}$ , then for every weakly bounded set  $A$  in  $L^p(G, \Omega)$ ,

$$\int_G f(h_\alpha - h) d\lambda \rightarrow 0 \quad \text{uniformly on } A. \quad (3.8)$$

This means that  $h_\alpha \rightarrow h$  in the strong topology of  $(L^p(G, \Omega), \tau_\Omega)^*$ . Hence, for every  $\omega \in \Omega$ , there is a unique  $f_\omega \in L^p(G, \omega)$  so that

$$F(h) = \int_G f_\omega h d\lambda \quad \text{on } L^q\left(G, \frac{1}{\omega}\right). \quad (3.9)$$

Now note that if  $\omega, \nu \in \Omega$  with  $\omega \leq \nu$ , then  $L^p(G, \nu) \subseteq L^p(G, \omega)$  and  $L^q(G, 1/\omega) \subseteq L^q(G, 1/\nu)$ . Therefore for every  $h \in L^q(G, 1/\omega)$ ,

$$\int_G f_\omega h d\lambda = \int_G f_\nu h d\lambda, \quad (3.10)$$

and hence  $f_\omega = f_\nu$  almost everywhere. This implies that

$$F \in \lim_{\omega} (L^p(G, \omega), I_{\omega, \nu}) = L^p(G, \Omega). \quad (3.11)$$

Conversely, if  $f \in L^p(G, \Omega)$ , then it is obvious that the linear form

$$F(h) = \int_G f h d\lambda \quad (h \in (L^p(G, \Omega), \tau_\Omega)^*) \quad (3.12)$$

is continuous with respect to the strong topology on  $(L^p(G, \Omega), \tau_\Omega)^*$ . So the canonical imbedding  $J : L^p(G, \Omega) \rightarrow (L^p(G, \Omega), \tau_\Omega)^{**}$  is onto. Hence  $L^p(G, \Omega)$  is semireflexive.  $\square$

#### 4. $L^p(G, \Omega)$ As a Topological Algebra

In this section, we study conditions on a system of weights  $\Omega$  for that  $L^p(G, \Omega)$  with the convolution multiplication to be a topological algebra. We commence with some definitions.

If  $f$  is a function on  $G$ , the left translate of  $f$  by  $x \in G$  is the function given by  $L_x f(y) = f(x^{-1}y)$ . A subset  $\mathcal{F}$  of functions on  $G$  is called left translation invariant if  $L_x f \in \mathcal{F}$  for all  $f \in \mathcal{F}$  and  $x \in G$ .

A weight function  $\omega$  on a locally compact group  $G$  is called left moderate if

$$\ell(s) := \operatorname{ess\,sup}_t \frac{\omega(st)}{\omega(t)} < \infty, \tag{4.1}$$

for all  $s \in G$ . It is easy to see that  $\ell(s) > 0$ ,  $\ell(st) \leq \ell(s)\ell(t)$ ; see [4] or [9]. Let us remark that any submultiplicative and any locally integrable left moderate measurable function is bounded and bounded away from zero on any compact subset of  $G$ ; see Theorem 2.7 in [10]. In particular,  $\ell$  is bounded on compact sets. The condition that  $\omega$  is left moderate is equivalent to that the space  $L^p(G, \omega)$  (for  $1 \leq p \leq \infty$ ) being translation invariant; see for more details [4]. Observe that for  $f \in L^p(G, \omega)$  and  $x \in G$ ,

$$\begin{aligned} \|L_x f\|_{p, \omega} &= \left( \int_G (|f(x^{-1}t)|\omega(t))^p d\lambda(t) \right)^{1/p} \\ &= \left( \int_G (|f(t)|\omega(xt))^p d\lambda(t) \right)^{1/p} \\ &\leq \left( \int_G (|f(t)|\ell(x)\omega(t))^p d\lambda(t) \right)^{1/p} \\ &= \ell(x) \|f\|_{p, \omega}. \end{aligned} \tag{4.2}$$

**Lemma 4.1.** *Let  $\Omega$  be a system of weights on  $G$ . Then  $L^p(G, \Omega)$  is left translation invariant if and only if every element of  $\Omega$  is left moderate.*

*Proof.* The “if” part is clear by the remarks above. For the converse, we need only to note that  $L^p(G, \Omega)$  is dense in  $(L^p(G, \omega), \|\cdot\|_{p, \omega})$  for  $\omega \in \Omega$ .  $\square$

**Theorem 4.2.** *Let  $\Omega$  be a system of locally integrable left moderate weights on  $G$  and  $f \in L^p(G, \Omega)$ . Then, the map  $x \mapsto L_x f$  from  $G$  into  $(L^p(G, \Omega), \tau_\Omega)$  is continuous.*

*Proof.* Assume first that  $f \in B_c(G)$  with  $K = \operatorname{supp}(f)$ . Let  $x \in G$ ,  $\omega \in \Omega$ , and  $(x_\alpha)$  be a net in  $G$  convergent to  $x$ . Choose a compact neighbourhood  $U$  of  $x$ , then  $\operatorname{supp}(L_x f) \subseteq UK$  whenever  $x \in U$ . Let

$$k = \sup\{\omega(s) : s \in UK\} < \infty. \tag{4.3}$$

Choose  $\alpha_0$  such that  $x_\alpha \in U$  for all  $\alpha \leq \alpha_0$  and  $\|L_{x_\alpha}f - L_x f\|_p \leq \epsilon/k$ . Then

$$\begin{aligned} \|L_{x_\alpha}f - L_x f\|_{p,\omega} &= \left( \int_{UF} \left( |f(x_\alpha^{-1}t) - f(x^{-1}t)| \omega(t) \right)^p d\lambda(t) \right)^{1/p} \\ &\leq k \left( \int_{UF} \left( |f(x_\alpha^{-1}t) - f(x^{-1}t)| \right)^p d\lambda(t) \right)^{1/p} \\ &= k \|L_{x_\alpha}f - L_x f\|_p \\ &\leq \epsilon, \end{aligned} \quad (4.4)$$

for all  $\alpha \geq \alpha_0$ .

Finally, let  $f$  be an arbitrary element of  $L^p(G, \Omega)$  and  $\epsilon > 0$ . Let  $M$  be an upper bound for the function  $\ell$  on the compact neighbourhood  $U$  of  $x$ ; recall that  $\ell$  is submultiplicative. For every  $\omega \in \Omega$ , there exists  $g_\omega \in B_c(G)$  such that  $\|f - g_\omega\|_{p,\omega} \leq \epsilon/3M$ . By the first part, we can choose  $\alpha_0$  such that

$$\|L_{x_\alpha}g_\omega - L_x g_\omega\|_{p,\omega} \leq \frac{\epsilon}{3}, \quad x_\alpha \in U, \quad (4.5)$$

for all  $\alpha \geq \alpha_0$ . One can conclude that

$$\begin{aligned} \|L_{x_\alpha}f - L_x f\|_{p,\omega} &\leq \|L_{x_\alpha}f - L_{x_\alpha}g_\omega\|_{p,\omega} + \|L_{x_\alpha}g_\omega - L_x g_\omega\|_{p,\omega} + \|L_x g_\omega - L_x f\|_{p,\omega} \\ &\leq \ell(x_\alpha)\epsilon + \epsilon/3 + \ell(x)\epsilon \\ &\leq \frac{M\epsilon}{3M} + \frac{\epsilon}{3} + \frac{M\epsilon}{3M} = \epsilon, \end{aligned} \quad (4.6)$$

for all  $\alpha \geq \alpha_0$ . This finishes the proof.  $\square$

We now focus on some systems of weights for that  $L^p(G, \Omega)$  to be an algebra under usual convolution

$$f * g(t) = \int_G f(s)g(s^{-1}t)d\lambda(s) \quad (f, g \in L^p(G, \Omega)), \quad (4.7)$$

whenever this integral makes sense. For  $p = 1$ , it is well known that  $L^1(G, \omega)$  is a convolution algebra if and only if  $\omega$  is weakly submultiplicative; that is, for all  $x, y \in G$ ,

$$\omega(st) \leq c\omega(s)\omega(t), \quad (4.8)$$

for some  $c > 0$ .



For any two weight functions  $\omega$  and  $\nu$  on  $G$ , we set

$$\Phi_{[\omega,\nu]}(x) = \int_G \left( \frac{\omega(x)}{\nu(y)\nu(y^{-1}x)} \right)^q d\lambda(y). \quad (4.9)$$

In the case where  $\omega = \nu$ , we simply write  $\Phi_\omega = \Phi_{[\omega,\omega]}$ .

The following lemma is similar to Lemma 2.2 in [9].

**Lemma 4.3.** *Let  $1 < p < \infty$  and  $\Omega$  be a system of weights on  $G$ . If  $L^p(G, \Omega)$  is a convolution algebra, then  $\omega^p$  is locally integrable for each  $\omega \in \Omega$ .*

The next result gives a sufficient condition for that  $L^p(G, \Omega)$  to be a convolution algebra.

**Theorem 4.4.** *Let  $1 < p < \infty$  and  $\Omega$  be a system of weights on  $G$ . If for every  $\omega \in \Omega$ , there is a  $\nu \in \Omega$  such that  $\Phi_{[\omega,\nu]} \in L^\infty(G)$ , where  $q$  is the conjugate exponent to  $p$ , then the space  $(L^p(G, \Omega), \tau_\Omega)$  is a complete locally convex algebra with continuous multiplication.*

*Proof.* We must show that

$$\|f * g\|_{p,\omega} \leq \|f\|_{p,\nu} \|g\|_{p,\nu} \quad (4.10)$$

for all  $f, g \in L^p(G, \Omega)$ . By Lemma 4.3,  $B_c(G)$  is dense in  $(L^p(G, \Omega), \|\cdot\|_{p,\omega})$ , thus for any  $\omega \in \Omega$ , it suffices to show that

$$\|f * g\|_{p,\omega} \leq \|f\|_{p,\nu} \|g\|_{p,\nu'} \quad (4.11)$$

for all  $f, g \in B_c(G)$ . For this, let  $f, g \in B_c(G)$ . Writing

$$f * g(x) = \int_G f(y)g(y^{-1}x) \frac{\nu(y)\nu(y^{-1}x)}{\nu(y)\nu(y^{-1}x)} d\lambda(y), \quad (4.12)$$

and using Hölder's inequality, we obtain

$$\begin{aligned} & |f * g(x)| \\ & \leq \left( \int_G (|f(y)|\nu(y))^p (|g(y^{-1}x)|\nu(y^{-1}x))^p d\lambda(y) \right)^{1/p} \left( \int_G \left( \frac{1}{\nu(y)\nu(y^{-1}x)} \right)^q d\lambda(y) \right)^{1/q}. \end{aligned} \quad (4.13)$$

This shows that

$$\begin{aligned} & \left| \int_G (|f * g(x)|\omega(x))^p d\lambda(x) \right| \\ & \leq \int_G \left( \int_G (|f(y)|v(y))^p (|g(y^{-1}x)|v(y^{-1}x))^p d\lambda(y) \right) \Phi_{[\omega,v]}(x)^{p/q} d\lambda(x) \quad (4.14) \\ & \leq \|f\|_{p,v}^p \|g\|_{p,v}^p \|\Phi_{[\omega,v]}\|_\infty^{p/q}. \end{aligned}$$

Whence

$$\|f * g\|_{p,\omega} \leq \|\Phi_{[\omega,v]}\|_\infty^{1/q} \|f\|_{p,v} \|g\|_{p,v}. \quad (4.15)$$

This completes the proof.  $\square$

The following corollary is a direct consequence of Theorem 4.4.

**Corollary 4.5.** *Let  $1 < p < \infty$  and  $\Omega$  be a system of weights on  $G$  such that for every  $\omega \in \Omega$ ,  $\Phi_\omega \in L^\infty(G)$ . Then  $L^p(G, \Omega)$  is a complete locally multiplicative convex algebra.*

The next result provides us with a class of weights  $\omega$  on the additive group  $\mathbb{R}^n$  for which the usual weighted Lebesgue space  $L^p(\mathbb{R}^n, \omega)$  becomes a Banach algebra.

**Proposition 4.6.** *Let  $1 < p < \infty$  and  $n$  be a natural number. Let  $\varpi : \mathbb{R}^n \rightarrow (0, +\infty)$  be a function such that*

- (i) *If  $\|x\| \leq \|y\|$ , then  $\varpi(x) \leq \varpi(y)$ .*
- (ii)  *$\varpi^{-1} \in L^1(\mathbb{R}^n)$ .*
- (iii) *There exists a positive number  $M$  such that  $\varpi(2x) \leq M\varpi(x)$  for all  $x \in \mathbb{R}^n$ .*

*Then  $L^p(\mathbb{R}^n, \sqrt[q]{\varpi})$  is a Banach algebra, where  $q$  is the conjugate exponent to  $p$ .*

*Proof.* For any  $x \in \mathbb{R}^n$ , let  $A_x = \{y \in \mathbb{R}^n : 2\|y\| \geq \|x\|\}$  and observe that

$$\begin{aligned} \varpi(y) & \geq \varpi\left(\frac{x}{2}\right), \quad \text{if } y \in A_x, \\ \varpi(x-y) & \geq \varpi\left(\frac{x}{2}\right), \quad \text{if } y \in \mathbb{R}^n \setminus A_x. \end{aligned} \quad (4.16)$$

Hence, for  $x \in \mathbb{R}^n$ ,

$$\begin{aligned} \Phi_{\sqrt[q]{\varpi}}(x) & = \int_{\mathbb{R}^n} \frac{\varpi(x)}{\varpi(y)\varpi(x-y)} dy \\ & = \int_{\mathbb{R}^n \setminus A_x} \frac{\varpi(x)}{\varpi(y)\varpi(x-y)} dy + \int_{A_x} \frac{\varpi(x)}{\varpi(y)\varpi(x-y)} dy \end{aligned}$$

$$\begin{aligned} &\leq \left( \frac{\varpi(x)}{\varpi(x/2)} \right) \left( \int_{\mathbb{R}^n \setminus A_x} \frac{1}{\varpi(y)} dy + \int_{A_x} \frac{1}{\varpi(x-y)} dy \right) \\ &\leq 2M \|\varpi\|_1. \end{aligned} \tag{4.17}$$

Thus,  $\Phi_{\varpi} \in L^\infty(\mathbb{R}^n)$ , and now the result follows from Corollary 4.5. □

*Example 4.7.* Let  $1 \leq p < \infty$ ,  $q$  be the conjugate exponent to  $p$ , and  $n \in \mathbb{N}$ . Set

$$\omega(x) = (a + \|x\|^r)^{s/q} b^{(1/q) \ln(c + \|x\|^t)} \quad (x \in \mathbb{R}^n), \tag{4.18}$$

where  $sr > n$ ,  $b > 1$  and  $a, c, t > 0$ . Then  $L^p(\mathbb{R}^n, \omega)$  is a Banach algebra.

We are going to prove the converse of Theorem 4.4. For this, we fix some notation. If  $f, g$  be two complex-valued functions on  $G$ , then  $f \otimes g$  denotes the function on  $G \times G$  given by  $f \otimes g(x, y) = f(x)g(y)$  for all  $x, y \in G$ . Also for any two sets  $\mathcal{F}$  and  $\mathcal{K}$  of functions on  $G$  we set  $\mathcal{F} \otimes \mathcal{K} = \{f \otimes g : f \in \mathcal{F}, g \in \mathcal{K}\}$ . For a locally compact group  $G$ , note that the cartesian product  $G \times G$  is a locally compact group by defining the product  $(x, y)(s, t) = (xs, yt)$  for all  $x, y, s, t \in G$ .

We need the following easy lemma in the sequel.

**Lemma 4.8.** *Let  $1 < p < \infty$  and  $\Omega$  be a system of weights on  $G$  such that  $B_c(G) \subseteq L^p(G, \Omega)$ . Then  $B_c(G) \otimes B_c(G)$  is dense in  $(L^p(G \times G, \omega \otimes \omega), \|\cdot\|_{p, \omega \otimes \omega})$ .*

*Proof.* Since  $B_c(G)$  is norm dense in  $L^p(G, \omega)$ , then  $B_c(G) \otimes B_c(G)$  is projective tensor norm dense in  $L^p(G, \omega) \widehat{\otimes}_\pi L^p(G, \omega)$ , where  $\widehat{\otimes}$  is the projective tensor product. Hence  $L^p(G, \omega) \otimes L^p(G, \omega)$  is  $\pi$ -dense in  $L^p(G, \omega) \widehat{\otimes}_\pi L^p(G, \omega)$ . On the other hand, it is known that  $L^p(G, \omega) \widehat{\otimes}_\pi L^p(G, \omega)$  is isometric with  $(L^p(G \times G, \omega \otimes \omega), \|\cdot\|_{p, \omega \otimes \omega})$ . In fact, the linear map

$$\varrho : L^p(G, \omega) \widehat{\otimes}_\pi L^p(G, \omega) \longrightarrow L^p(G \times G, \omega \otimes \omega), \quad \varrho(f \otimes g)(x, y) = f(x)g(y) \tag{4.19}$$

for all  $f, g \in L^p(G, \omega)$  and  $x, y \in G$ , can be extended to a surjective isometry; for more details, see for example [14]. Now we conclude that  $B_c(G) \otimes B_c(G)$  is  $\|\cdot\|_{p, \omega \otimes \omega}$ -dense in  $L^p(G \times G, \omega \otimes \omega)$ . □

The next theorem is our main result in this section.

**Theorem 4.9.** *Let  $1 < p < \infty$ ,  $G$  be  $\sigma$ -compact, and  $\Omega$  be a system of weights on  $G$ . If the space  $(L^p(G, \Omega), \tau_\Omega)$  is an algebra with continuous multiplication, then for every  $\omega \in \Omega$  there exists a  $\nu \in \Omega$  such that  $\Phi_{[\omega, \nu]} \in L^\infty(G)$ .*

*Proof.* Choose an arbitrary  $\omega \in \Omega$ . Then, by assumption, there exists some  $\nu \in \Omega$  such that for every  $f, g \in L^p(G, \Omega)$ ,  $\|f * g\|_{p, \omega} \leq \|f\|_{p, \nu} \|g\|_{p, \nu}$ . Now for every  $h \in L^q(G, 1/\omega)$ ,

$$F(f) = \int_G f(x)h(x)d\lambda(x) \quad (f \in B_c(G)) \tag{4.20}$$

defines a continuous linear functional on  $B_c(G)$  with the norm  $\|F\| = \|h\|_{q,1/\omega}$ . Also for every  $f, g \in B_c(G)$ ,  $f * g \in B_c(G)$ , and we have

$$\begin{aligned} F(f * g) &= \int_G f * g(x)h(x)d\lambda(x) = \int_G \left( \int_G f(y)g(y^{-1}x)d\lambda(y) \right) h(x)d\lambda(x) \\ &= \int_G \int_G f(y)g(x)h(yx)d\lambda(x)d\lambda(y) < \infty. \end{aligned} \quad (4.21)$$

Set  $F(f \otimes g) = F(f * g) = \int_{G \times G} f(y)g(x)h(yx) d\lambda \times \lambda(x, y)$  for  $f, g \in B_c(G)$ . By Lemma 4.8,  $F$  can be extended to a  $\|\cdot\|_{p,\nu \otimes \nu}$ -continuous functional on  $L^p(G \times G, \nu \otimes \nu)$ . Since  $G$  is  $\sigma$ -compact, by Exercise 15.14 in [1], it follows that the function  $(x, y) \mapsto h(yx)$  belongs to  $L^q(G \times G, 1/(\nu \otimes \nu))$ . But

$$\begin{aligned} \int_G \int_G \left( \frac{h(yx)}{\nu(x)\nu(y)} \right)^q d\lambda(x)d\lambda(y) &= \int_G \int_G \left( \frac{h(x)}{\nu(y)\nu(y^{-1}x)} \right)^q d\lambda(x)d\lambda(y) \\ &= \int_G \left( \frac{h(x)}{\omega(x)} \right)^q \Phi_{[\omega,\nu]}(x)d\lambda(x) < \infty. \end{aligned} \quad (4.22)$$

Since  $(h/\omega)^q \in L^1(G)$  is arbitrary, we conclude that  $\Phi_{[\omega,\nu]} \in L^\infty(G)$ ; see Section 14 in [15] or Theorem 20.15 in [1].  $\square$

As an immediate consequence of Theorem 4.9, we obtain the following corollary that partially answers a question raised in [3].

**Corollary 4.10.** *Let  $\omega$  be a weight on  $\sigma$ -compact group  $G$  and  $1 < p < \infty$ . Then  $L^p(G, \omega)$  is a convolution algebra if and only if  $\Phi_\omega \in L^\infty(G)$ .*

## 5. Ideals and the Spectrum of the Algebra $L^p(G, \Omega)$

We commence this section with the following proposition.

**Proposition 5.1.** *Let  $1 \leq p < \infty$  and let  $L^p(G, \Omega)$  be a translation invariant algebra. Then*

- (i)  $(L^p(G, \Omega), \tau_\Omega)$  has an approximate identity.
- (ii)  $(L^p(G, \Omega), \tau_\Omega)$  has a bounded approximate identity or an identity if and only if  $G$  is discrete.

*Proof.* (i) Let  $U$  be a fixed relatively compact neighbourhood of the identity element  $e$ , and let  $\mathcal{U}$  be the family of all neighbourhoods of  $e$  contained in  $U$  directed by reverse inclusion. Set  $e_V := \chi_V / \lambda(V)$ , and note that since elements of  $\Omega$  are locally integrable,  $e_V \in L^p(G, \Omega)$ . Given

$\epsilon > 0$  and  $\omega \in \Omega$ , then by Theorem 4.2, there exists a neighbourhood  $W$  of the identity such that  $\|f - L_t f\|_{p,\omega} < \epsilon$  for  $t \in W$ . Now, for  $V \in \mathcal{U}$  with  $V \subseteq W$ , and  $g \in L^q(G, 1/\omega)$ , we have

$$\begin{aligned} |\langle e_V * f - f, g \rangle| &= \left| \int_G (e_V * f - f)(x)g(x)d\lambda(x) \right| \\ &\leq \int_G \int_V \frac{|f(t^{-1}x) - f(x)|}{\lambda(V)} d\lambda(t) |g(x)| d\lambda(x) \leq \frac{1}{\lambda(V)} \int_V \langle |L_t f - f|, |g| \rangle d\lambda(t) \\ &\leq \sup_{t \in V} \|L_t f - f\|_{p,\omega} \|g\|_{q,1/\omega} < \epsilon \|g\|_{q,1/\omega}. \end{aligned} \tag{5.1}$$

Hence,  $\|e_V * f - f\|_{p,\omega} \leq \epsilon$  for all neighborhoods  $V \subseteq W$ , from which it follows that  $e_V * f \rightarrow f$  in  $\tau_\Omega$ -topology.

(ii) Let  $(e_\alpha)_\alpha$  be a bounded left approximate identity for  $L^p(G, \Omega)$ . Fix an  $\omega \in \Omega$ , then  $\|e_\alpha\|_{p,\omega} \leq M$  for some positive number  $M$ . Let  $f \in L^p(G, \omega)$ . Since  $L^p(G, \Omega)$  is dense in  $L^p(G, \omega)$  with the norm  $\|\cdot\|_{p,\omega}$ , then given  $\epsilon > 0$ , there exists  $g \in L^p(G, \Omega)$  such that  $\|f - g\|_{p,\omega} \leq \epsilon/3(M + 1)$ . Choose  $\alpha_0$  such that  $\|e_\alpha * g - g\|_{p,\omega} \leq \epsilon/3$  for all  $\alpha \geq \alpha_0$ . Then it follows that

$$\begin{aligned} \|e_\alpha * f - f\|_{p,\omega} &\leq \|e_\alpha * f - e_\alpha * g\|_{p,\omega} + \|e_\alpha * g - g\|_{p,\omega} + \|f - g\|_{p,\omega} \\ &\leq M \frac{\epsilon}{3(M + 1)} + \frac{\epsilon}{3} + \frac{\epsilon}{3(M + 1)} < \epsilon, \end{aligned} \tag{5.2}$$

for all  $\alpha \geq \alpha_0$ . This means that  $(L^p(G, \omega), \|\cdot\|_{p,\omega})$  has a bounded left approximate identity. But according to Theorem 4.2 in [9], this is equivalent to that  $G$  is discrete.  $\square$

The next theorem shows that closed ideals of the algebra  $(L^p(G, \Omega), \tau_\Omega)$  are exactly translation invariant subspaces.

**Theorem 5.2.** *Let  $1 \leq p < \infty$  and  $L^p(G, \Omega)$  be a translation invariant algebra. Then a closed linear subspace of  $L^p(G, \Omega)$  is an ideal in  $L^p(G, \Omega)$  if and only if it is two-sided translation invariant.*

*Proof.* Suppose that  $I$  is a  $\tau_\Omega$ -closed two-sided translation invariant subspace of  $L^p(G, \Omega)$ . We have to show that  $g * f \in I$  and  $f * g \in I$  for all  $f \in I$  and  $g \in L^p(G, \Omega)$ . Let  $h \in L^q(G, 1/\omega)$ , for some  $\omega \in G$ , such that  $\int_G f(x)h(x)d\lambda(x) = 0$  for all  $f \in I$ . Then, for  $f \in I$  and any  $g \in L^p(G, \Omega)$ ,

$$\begin{aligned} \int_G (g * f)(x)h(x)d\lambda(x) &= \int_G h(x) \left( \int_G g(y)f(y^{-1}x)d\lambda(y) \right) d\lambda(x) \\ &= \int_G g(y) \left( \int_G L_y f(x)h(x)d\lambda(x) \right) d\lambda(y) \\ &= 0. \end{aligned} \tag{5.3}$$

Since  $(L^p(G, \Omega), \tau_\Omega)^* = \Omega \cdot L^q(G)$ , the Hahn-Banach theorem implies that  $g * f \in I$  for all  $f \in I$  and  $g \in L^p(G, \Omega)$ . Thus  $I$  is a left ideal, and using the right translation invariance of  $I$ , it is readily seen, in the same way, that  $I$  is also a right ideal.

Conversely, let  $I$  be a closed ideal of  $(L^p(G, \Omega), \tau_\Omega)$ , and  $x \in G$ . Let  $(e_\alpha)$  be an approximate identity for  $L^p(G, \Omega)$ . Then for each  $f \in L^p(G, \Omega)$ , we have

$$\|L_x(e_\alpha) * f - L_x f\|_{p, \omega} \leq \ell(x) \|e_\alpha * f - f\|_{p, \omega} \longrightarrow 0. \quad (5.4)$$

Hence,  $L_x(e_\alpha) * f \rightarrow L_x f$  in  $\tau_\Omega$ -topology. As  $I$  is a  $\tau_\Omega$ -closed left ideal, it follows that  $L_x f \in I$ ; that is,  $I$  is left translation invariant. Similarly, it is shown that  $I$  is also right translation invariant.  $\square$

We denote by  $\Delta(L^p(G, \Omega))$  the spectrum of  $(L^p(G, \Omega), \tau_\Omega)$  consisting of all  $\tau_\Omega$ -continuous nonzero linear functionals  $\Phi$  on  $L^p(G, \Omega)$  which are multiplicative; that is,

$$\Phi(f * g) = \Phi(f)\Phi(g) \quad (f, g \in L^p(G, \Omega)). \quad (5.5)$$

We conclude this work with the following result which is a characterization of the spectrum of  $(L^p(G, \Omega), \tau_\Omega)$ .

**Proposition 5.3.** *Let  $\Omega$  be a system of weights on  $\delta$ -compact group  $G$ . Then*

$$\Delta(L^p(G, \Omega)) = \left\{ \Phi_\rho : \rho \in L^q\left(G, \frac{1}{\omega}\right), \omega \in \Omega, \rho(xy) = \rho(x)\rho(y) \right\}, \quad (5.6)$$

where

$$\Phi_\rho(f) = \int_G f(x)\rho(x)d\lambda(x) \quad (f \in L^p(G, \Omega)). \quad (5.7)$$

*Proof.* Let  $\rho \in L^q(G, 1/\omega)$  for some  $\omega \in \Omega$  such that  $\rho(xy) = \rho(x)\rho(y)$  for almost all  $x, y \in G$ . Then,  $\Phi_\rho$  is  $\|\cdot\|_{p, \omega}$ -continuous and so  $\tau_\Omega$ -continuous. Moreover, for  $f, g \in L^p(G, \Omega)$ ,

$$\begin{aligned} \Phi_\rho(f * g) &= \int_G \int_G f(x)g(y)\rho(xy)d\lambda(x)d\lambda(y) \\ &= \int_G \int_G f(x)g(y)\rho(x)\rho(y)d\lambda(x)d\lambda(y) \\ &= \Phi_\rho(f)\Phi_\rho(g). \end{aligned} \quad (5.8)$$

That is,  $\Phi_\rho \in \Delta(L^p(G, \Omega))$ .

Conversely, let  $\Phi \in \Delta(L^p(G, \Omega))$ . Then  $\Phi$  is bounded on a  $\tau_\Omega$ -neighbourhood of zero. Thus  $\Phi$  is bounded on the set  $\{f \in L^p(G, \Omega) : \|f\|_{p, \omega} < 1\} \cap L^p(G, \omega)$  for some  $\omega \in \Omega$ . Therefore  $\Phi$  can be extended to an element  $\bar{\Phi}$  in  $(L^p(G, \omega), \|\cdot\|_{p, \omega})^*$ . It follows that there exists a function  $\rho \in L^q(G, 1/\omega)$  such that

$$\bar{\Phi}(f) = \int_G f\rho d\lambda, \quad (5.9)$$

for all  $f \in L^p(G, \omega)$ . Since for  $f, g \in B_c(G)$ ,  $\Phi(f)\Phi(g) = \Phi(f * g)$ , we infer that

$$\begin{aligned} \int_{G \times G} f(y)g(x)\rho(y)\rho(x)d\lambda \times \lambda(x, y) &= \int_G \int_G f(y)g(x)\rho(y)\rho(x)d\lambda(y)d\lambda(x) \\ &= \int_G \int_G f(y)g(x)\rho(yx)d\lambda(y)d\lambda(x) \quad (5.10) \\ &= \int_{G \times G} f(y)g(x)\rho(yx)d\lambda \times \lambda(x, y). \end{aligned}$$

By an argument similar to the proof of Theorem 4.9, we deduce that  $\rho(xy) = \rho(x)\rho(y)$  for almost all  $x, y \in G$ .  $\square$

## References

- [1] E. Hewitt and K. Stromberg, *Real and Abstract Analysis*, Springer, New York, NY, USA, 1975.
- [2] Yu. N. Kuznetsova, "Weighted  $L_p$ -algebras on groups," *Functional Analysis and Its Applications*, vol. 40, no. 3, pp. 234–236, 2006.
- [3] F. Abtahi, R. Nasr-Isfahani, and A. Rejali, "Weighted  $L^p$ -conjecture for locally compact groups," *Periodica Mathematica Hungarica*, vol. 60, no. 1, pp. 1–11, 2010.
- [4] R. E. Edwards, "The stability of weighted Lebesgue spaces," *Transactions of the American Mathematical Society*, vol. 93, pp. 369–394, 1959.
- [5] A. El Kinani and A. Benazzouz, "m-convex structure in the weighted space  $L^p_\Omega(\mathbb{R}^n)$ ," *Bulletin of the Belgian Mathematical Society*, vol. 10, no. 1, pp. 49–57, 2003.
- [6] J. O. Olaleru, "Weighted locally convex spaces of measurable functions," *Revista Colombiana de Matemáticas*, vol. 38, no. 1, pp. 7–15, 2004.
- [7] K. Reiher, "Weighted inductive and projective limits of normed Köthe function spaces," *Results in Mathematics*, vol. 13, no. 1-2, pp. 147–161, 1988.
- [8] J. Wengenroth, "Weighted (LF)-spaces of measurable functions," *Results in Mathematics*, vol. 32, no. 1-2, pp. 169–178, 1997.
- [9] Yu. N. Kuznetsova, "The invariant weighted algebras  $L^p(G, \omega)$ ," *Mathematical Notes*, vol. 84, no. 4, pp. 529–537, 2008.
- [10] H. G. Feichtinger, "Gewichtsfunktionen auf lokalkompakten Gruppen," *Österreichische Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Klasse. Sitzungsberichte*, vol. 188, no. 8–10, pp. 451–471, 1979.
- [11] H. H. Schaefer, *Topological Vector Spaces*, Springer, New York, NY, USA, 1971.
- [12] A. Mallios, *Topological Algebras. Selected Topics*, vol. 124 of *North-Holland Mathematics Studies*, North-Holland Publishing, Amsterdam, The Netherlands, 1986.
- [13] R. B. Holmes, *Geometric Functional Analysis and Its Applications*, Springer, New York, NY, USA, 1975.
- [14] E. Kaniuth, *A Course in Commutative Banach Algebras*, vol. 246 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2009.
- [15] K. Floret, *Maß- und Integrationstheorie*, B. G. Teubner, Stuttgart, Germany, 1981.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

