

# Identification of a nutrient sensing transcriptional network in monocytes by using inbred rat models of cafeteria diet

Martínez-Micaelo N<sup>1</sup>, González-Abuín N<sup>1</sup>, Terra X<sup>1</sup>, Ardévol A<sup>1</sup>, Pinent M<sup>1</sup>, Petretto E<sup>3,4</sup>, Behmoaras J<sup>2</sup>, Blay M<sup>1</sup>

<sup>1</sup>Mobiofood Research Group. Department of Biochemistry and Biotechnology, Universitat Rovira i Virgili, Tarragona, Spain.

<sup>2</sup>Centre of Complement and Inflammation Research, Imperial College London, Du Cane Road, London, W12 0NN, UK

<sup>3</sup>MRC Clinical Sciences Centre, Imperial College London, Hammersmith Hospital, Du Cane Road, London, W12 0NN, UK

<sup>4</sup>Duke-NUS Graduate Medical School Singapore, 8 College Road, 169857 Singapore, Republic of Singapore

## **RUNNING TITTLE:**

Differential metabolic response and circulating monocytes transcriptome of WKY and LEW rat to cafeteria diet

**KEY WORDS:** Cafeteria diet, inbred rats, monocytes' transcriptomics, LEW, WKY

1,2 Corresponding authors: mteresa.blay@urv.cat and Jacques.behmoaras@imperial.ac.uk

## ABSTRACT

Obesity has reached pandemic levels worldwide. The current models of diet-induced obesity in rodents use predominantly high-fat based diets that do not take into account the consumption of variety of highly palatable, energy dense foods that are prevalent in Western society. We and others have shown that cafeteria diet (CAF) is a robust and reproducible model of human metabolic syndrome with tissue inflammation in the rat. We have previously shown that inbred rat strains such as Wistar Kyoto (WKY) and Lewis (LEW) show different susceptibilities to CAF diets with distinct metabolic and morphometric profiles. Here we show a difference in plasma MCP-1 levels and investigate the effect of CAF diet on peripheral blood monocyte transcriptome as powerful stress-sensing immune cells in WKY and LEW rats. We found that 75.5% of the differentially expressed transcripts under CAF diet were up-regulated in WKY rats and were functionally related to the activation of the immune response. Using a gene co-expression network constructed from the genes differentially expressed between CAF diet-fed LEW and WKY rats, we identified the Acyl-CoA synthetase short-chain family member 2 (*Acss2*) as a *hub* gene for a nutrient sensing cluster of transcripts in monocytes. *Acss2* genomic region is significantly enriched for previously established metabolism quantitative trait loci in the rat. Notably, monocyte expression levels of *Acss2* significantly correlated with plasma glucose, triglyceride, leptin and NEFA levels as well as morphometric measurements such as body weight and the total fat following CAF in the rat. These results show the importance of the genetic background in nutritional genomics and identify inbred rat strains as potential models for CAF-induced obesity.

# 1. INTRODUCTION

Obesity represents an important health problem in human populations and there is need of new animal models that mimics optimally the main characteristics of human obesity.

Susceptibility to obesity has a large underlying genetic component with modulatory environmental factors such as diet (Barsh et al., 2000; Bouchard, 2008; Hetherington and Cecil, 2009; Ordovas, 2008; Pérusse and Bouchard, 2000; Speakman, 2004). The administration of cafeteria (CAF) diet to rats is considered as a robust model of the human metabolic syndrome and its related pathologies, and has been widely used to understand the genetic factors that underlie obesity. Feeding with a cafeteria diet, obesity is due to the continuous hyperphagia as result of the voluntary intake of highly palatable and energy-dense cafeteria-style foods, which are typically found in western-type diets (Sampey et al., 2011).

Importantly, the genetic background of the selected strain is an important determinant of the phenotypic diversity observed in diet-induced obesity models. Hence, by comparing Lewis (LEW) and Wistar Kyoto (WKY) rats responses to an obesogenic diet, we recently showed significant differences in metabolic and morphometric parameters. (Martínez-Micaelo et al., 2016). Specifically, our results suggested the leptin-signalling pathway as a divergent point between the strain-specific adaptations to CAF diet. LEW rats display a typical obesity phenotype characterized by an increase in body weight, adiposity and dislipemic and hyperglucemic profiles, accompanied by a preferential metabolization of carbohydrates as opposed to WKY rats, which metabolized preferentially lipids. Despite their body weight and adiposity gain, WKY rats maintained metabolic homeostasis as a result of the obesogenic diet showing a differential regulation in the leptin axis.

In obesity, the prolonged over-nutrition is characterised by the restrained ability of the adipose tissue to store energy as fat. In parallel, a cytokine-driven response characterised by the secretion of pro-inflammatory chemokines underlie obesity-induced inflammatory reactions. Among others,

monocyte chemoattractant protein-1 (MCP-1) is one of the most important cytokines due to its ability to recruit peripheral monocytes to the adipose tissue (Iyer et al., 2010; Osborn and Olefsky, 2012). The continuous activation of the local inflammatory response underlies the chronic low-grade systemic inflammation that results in the widespread activation of the immune system (Lumeng and Saltiel, 2011). Furthermore, although the induced inflammatory response is triggered and takes place primarily in adipose tissue, other organs are also involved.

Peripheral blood monocytes are powerful stress-sensing immune cells that play central roles in the regulation of innate immune responses through the release of inflammatory cytokines and the activation of the adaptive immunity (Parihar et al., 2010). Monocytes are of particular interest because, as circulating immune cells, they are exposed to the systemic environment, including the metabolic factors and pro-inflammatory cytokines produced and secreted by organs and tissues. Thus, the regulation of monocyte expression profile may reflect the physiological state of the whole organism.

Here, we studied the CAF diet-induced modulation of peripheral monocyte transcriptome from inbred LEW and WKY rats. Considering the isogenic and homozygous nature of these inbred strains and using the CAF diet as induced-obesity, both the genetic background and environmental factors can be controlled, providing a powerful tool for understanding the “gene-diet” interactions that underlie the development of complex traits such as obesity. These inbred strains have been widely studied for their differential susceptibility to experimental glomerulonephritis through macrophage activation (Behmoaras et al., 2008; Behmoaras et al., 2010) and present a markedly distinct macrophage transcriptome, which result in differential macrophage activation (D’Souza et al., 2013; Maratou et al., 2011). Here, we show that the peripheral monocyte transcriptome of LEW and WKY rats is modulated by the CAF diet differentially. Our results suggest that monocyte transcriptome differences are caused by the contrasting metabolic and morphometric responses of LEW and WKY rats to the diet induced-obesity.

These results establish new models of nutritional genomics in the CAF-induced obesity in the rat.

## **2. MATERIALS AND METHODS**

### **Animals and experimental design**

Male LEW (LEW/Crl) and WKY (WKY/NCrl) (Charles River, Margate, UK) rats, weighing 190 g and 130 g, respectively, were housed in a 22 °C temperature-controlled room with a 12-h light/dark cycle. After an adaptation period, rats from each strain were randomly distributed into two experimental groups (n=5) and fed with either a standard chow diet (STD) (Panlab, Barcelona, Spain) or a cafeteria diet (CAF), generating a hypercaloric diet distributed in a 10.4%, 38.6% and 50.8% of daily total energy intake from protein, carbohydrates and lipids respectively. After seven weeks of the indicated diet, the rats were fasted for 9 h and euthanised by exsanguination under anaesthesia. Blood was collected from abdominal aorta, and circulating monocytes were immediately isolated. See Figure 1 and (Martínez-Micaelo et al., 2016) for more detailed experimental design. Heparinised plasma was obtained by centrifugation from a blood sample and stored at -80 °C until analysis. All the procedures were performed with the approval of the Animal Ethics Committee of the Universitat Rovira i Virgili (Tarragona, Spain).

### **Measurement of inflammatory parameters in plasma**

MCP-1 was measured by an enzymatic colorimetric test kits provided by (ABCAM, Cambridge, USA).

### **Isolation of circulating monocytes, RNA extraction and microarray preparation**

Peripheral blood mononuclear cells (PBMCs) were isolated by density-gradient centrifugation using HISTOPAQUE-1083 solution (SIGMA, Madrid). Monocytes were purified based on their adherence to plastic in serum-free RPMI-supplemented medium. Thus, after 90 min, the non-adherent cells were removed by several washes with warm PBS, and the total RNA from the

adherent monocytes was extracted using the TRIzol reagent according to the manufacturer's instructions (Invitrogen, Barcelona, Spain) and purified using RNeasy microkits (Qiagen). The quality of total RNA isolated was determined using the Agilent 2100 Bioanalyser. The RNA Integrity Number (RIN) of RNA ranged from 9.10-10. One hundred and fifty nanograms of total RNA were amplified, labelled and hybridised to Rat Gene 1.0 ST arrays (Affymetrix, Santa Clara, CA, USA) using the Ambion WT Expression kit (Life Technologies) according to manufacturer's instructions.

### **Microarray analysis of differential gene expression**

The data from microarrays were normalised using the robust multi-array average (RMA) method (Irizarry et al., 2003), implemented in the Affymetrix Bioconductor package, and the differentially expressed genes were determined using the linear model implemented in the limma Bioconductor package. The Benjamini and Hochberg method were used to adjust P-values for multiple testing and control of the false discovery rate. Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways analysis was performed using the GOstats Bioconductor package (Falcon and Gentleman, 2007).

### **Gene co-expression networks**

A gene co-expression network was constructed from the gene expression profiles of the transcripts whose expression was differentially modulated (absolute fold change >1.5) when comparing the expression profiles of circulating monocytes from CAF diet-fed WKY and LEW rats. The co-expression network was inferred using a graphical Gaussian model (GGM) implemented in the R package GeneNet (Schäfer and Strimmer, 2005). Briefly, a partial-correlation matrix was estimated by computing the partial correlation between the expression profiles of each gene pair. Bayesian posterior edge probability >0.95 (corresponding to a local false discovery rate <5%) was used to determine the significance of the resulting pairwise partial correlations. In the resulting co-expression network the nodes represent the set of genes that were differentially expressed and co-expressed between WKY and LEW rats, and the edges link the pairs of genes whose expression

is not conditionally independent, defined as the pairwise partial correlation once the common effects of the other genes in the subset are removed (Opgen-Rhein and Strimmer, 2007). To identify sub-clusters (modules) within the network we used the MCODE algorithm implemented in the clusterMaker plugin (Morris et al., 2011) of Cytoscape software (Cline et al., 2007), and the modules were functionally annotated with GO terms and KEGG pathways using the GOstats Bioconductor package (Falcon and Gentleman, 2007). The selected clusters were analysed using the cyto-Hubba plugin (Lin et al., 2008) to determine the hubs and bottlenecks that represent the key regulatory genes within the network.

### **Statistical analysis**

The results are expressed as the mean  $\pm$  SEM. Strain and diet-specific effects (represented by a hash tag or an asterisk, respectively) were assessed using two-way ANOVA. Differences were considered to be significant when the  $p$ -values were  $<0.05$ . These calculations were performed using SPSS 17.0 software.

## **3. RESULTS**

### **3.1 The metabolic response of LEW and WKY rats to CAF diet is strain-dependent.**

We have previously shown in detail strain-specific phenotypic differences in CAF diet-induced obesity in LEW and WKY rats (Martínez-Micaelo et al., 2016).

Strain-specific differences were also observed in the modulation of the circulating levels of inflammatory biomarkers. Although CAF diet provoked a significant enhance in the levels of MCP-1 in LEW rats ( $241.29 \pm 13.0$  vs.  $297.97 \pm 11.9$ ,  $P$ -value = 0.017), no dietary effects were observed in WKY rats' circulating MCP-1 levels ( $273.57 \pm 22.0$  vs.  $305.10 \pm 10.9$ ,  $P$ -value = 0.263).

### **3.2 CAF diet differentially modulates gene expression profiles in circulating monocytes of LEW and WKY rats.**

To study genetic factors underlying the metabolic-immunological crosstalk, gene expression profiles of circulating monocytes from STD- and CAF-fed LEW and WKY rats were determined. Microarray data were analysed in two ways: 1) comparing the monocyte expression profiles of CAF-fed WKY and LEW rats in order to identify genes that underlie genotype-dependent adaptation to diet-induced obesity and 2) analysing the strain-specific modulation of the transcriptome of circulating monocytes by the CAF diet. The study design is described in Figure 1.

#### **3.2.1 Strain-dependent adaptation of LEW and WKY rats to CAF diet.**

The expression profiles of circulating monocytes from WKY and LEW CAF-fed rats were first compared in order to determine the genetic basis of the strain-dependent response to obesity. This microarray data analysis identified 8,479 transcripts that were differentially expressed ( $p < 0.01$ ) between the circulating monocytes from CAF-fed WKY and LEW rats; 75.5% of these transcripts were significantly up-regulated in WKY rats. We prioritized a set of 29 transcripts that were significantly differentially expressed after correction for multiple testing ( $FDR < 5\%$ ) and with fold change ( $fc$ )  $> 5$  when comparing the expression profiles from WKY and LEW rats (Figure 2A). Notably, this subset of highly differentially expressed transcripts included macrophage activation 2 like (*Mpa2l*), a target gene of nuclear factor- $\kappa$ B (NF- $\kappa$ B) (Rothgiesser et al., 2010) and lactate dehydrogenase A (*Ldha*), the gene encoding for the enzyme that catalyses the conversion of lactate to pyruvate during the glycolysis under anaerobic conditions. *Mpa2l* was 26 fold over-expressed in WKY rats whereas *Ldha* was markedly down-regulated ( $fc = -60.95$ ) in these rats. Furthermore, 6 of the 29 transcripts with high differential expression between WKY and LEW rats (*Klra7*, *Irg1*, *Ly49si1*, *Plk2*, *Rnd3*, *Lilrb3l*) were also previously identified as differentially expressed transcripts in bone marrow-derived macrophages from WKY and LEW rats (Maratou et al., 2011).



To gain insights into the molecular processes and pathways underlying these transcriptional responses, we analysed the set of 954 transcripts with  $fc > 1.5$  (up- and down-regulated) between CAF-fed WKY and LEW rats for enrichment for gene ontology (GO) terms. The GO terms were manually curated to remove redundant categories resulting from overlapping gene sets (Supplementary Table S1), and the most relevant terms that were significantly over-represented are shown in Figure 2A and B. The overexpressed genes in CAF-fed WKY rats were enriched for terms related to the activation of the inflammatory and immune responses resulting from the up-regulation of classical inflammatory mediators such as interferon gamma (*Ifng*,  $fc = 2.43$ ), tumour necrosis factor (*Tnf*,  $fc = 2.11$ ), nitric oxide synthase 2 (*Nos2*,  $fc = 1.68$ ), interleukins (IL) including *IL1a* ( $fc = 2.66$ ), *IL6* ( $fc = 2.41$ ), *IL12a* ( $fc = 2.34$ ), *IL18* ( $fc = 1.96$ ) and *IL23a* ( $fc = 3.45$ ) as well as genes involved in the regulation of eicosanoid production such as the gene encoding for cyclooxygenase 2 (prostaglandin-endoperoxide synthase 2 or *Ptgs2*,  $fc = 2.71$ ) and the gene encoding the 5-lipoxygenase (arachidonate 5-lipoxygenase or *Alox*,  $fc = 2.70$ ). Moreover, the over-expressed genes were also enriched for terms related to the positive regulation of the JAK-STAT cascade, specifically through the involvement of the tyrosine phosphorylation of STAT3. Nevertheless, the genes that were down-regulated in CAF-fed WKY rats compared to CAF-fed LEW rats were significantly enriched for very few gene ontologies (Figure 2C), suggesting the prominent significant functional gene enrichment in the immune system GO category.

We then investigated gene regulatory networks by constructing a gene co-expression network based on the analysis of the microarray-derived expression profiles assessed in circulating monocytes. In order to identify the individual genes and the gene clusters whose co-regulation underlie the strain-specific responses to diet, a gene co-expression network was inferred based on the differentially expressed transcripts that showed a  $fc > 1.5$  (up- and down-regulated) between the circulating monocytes from CAF-fed WKY and LEW rats (Figure 3A). The resulting network is composed of a single connected component formed by 742 nodes and 8,215 edges. The molecular complex detection (MCODE) cluster algorithm was employed to extract

functional modules of densely interconnected genes (Bader and Hogue, 2003). The biological relevance of the genes included in each of the 16 MCODE-derived clusters was tested with GO and KEGG analysis of the functional significance of biochemical pathways (Figure 3B). Although not all of the modules were enriched significantly for GO terms or KEGG pathways, the transcripts in cluster 2, forming 40 nodes and 205 edges (representing 40 annotated protein-coding genes), were functionally enriched for nutrient sensors (P-value =  $2.94 \times 10^{-4}$ ), responders to hormone stimuli (P-value =  $1.90 \times 10^{-3}$ ), and components of the NOD-like receptor signalling pathway (P-value =  $6.84 \times 10^{-3}$ ). Remarkably, the topology analysis of the module structure of cluster 2 identified the Acyl-CoA synthetase short-chain family member 2 (*Acss2*) as the most essential *hub* within the cluster in terms of the node with the largest degree and the highest betweenness score. In addition, the functional relevance of *Acss2* as regulator of nutrient sensing was determined by analysing the overlap of quantitative trait loci (QTLs) within the *Acss2* genomic region based on previously established QTL mapping results obtained from the Rat Genome Database (de la Cruz et al., 2005). This analysis revealed that a large majority of the 30 overlapping QTLs are related to traits associated with the regulation of metabolism such as body weight QTL 94, glucose level QTL 39 and serum leptin concentration QTL 7 (Supplementary Table S2).

In order to link the monocyte transcriptome to the obesity phenotypes observed in CAF diet, we investigated the association between the expression of *Acss2* in peripheral monocytes and metabolism-related traits (Table 1, Figure S2). Notably, significant and strong correlations were found between the *Acss2* expression in monocytes and (i) plasma levels of glucose, triglycerides, leptin and NEFAs and (ii) total fat and body weight as morphometric parameters (Table 1, Figure S2).

The topological analysis inside this nutrient sensing cluster also revealed a functional sub-module consisting of 9 highly co-expressed nodes; all of these nodes were overexpressed in CAF-fed WKY rats compared to the LEW-fed

rats and significantly enriched for inflammatory response genes and genes that function in response to lipid and glucocorticoid stimuli.

### **3.2.3 Strain-specific modulation of the transcriptome by CAF diet in circulating monocytes**

We also determined the effects of CAF diet on monocyte gene expression profile from LEW and WKY rats (Figure 4A and 4B). We show that the phenotypic response of LEW rats to CAF diet was translated in significant changes in monocyte gene expression, and although 228 and 195 transcripts were significantly up- and down-regulated respectively, the magnitude of fc by CAF diet was relatively moderate. Furthermore, gene ontology and KEGG analysis of the differentially expressed transcripts did not reveal any significant enrichment for any obvious obesity-related pathways when CAF and STD diet was compared in LEW rats. Interestingly, the modulation of the circulating monocyte expression profile by CAF diet in the WKY rats resulted in more significant changes in monocyte gene expression levels (483 up-regulated and 449 down-regulated transcripts). Gene ontology (GO) and KEGG analyses showed an enrichment for Wnt and peroxisome proliferator-activated receptor (PPAR) signalling pathways for up-regulated and down-regulated genes, respectively.

## 4. DISCUSSION

In the present study, we show differential monocyte transcriptome responses in inbred rat strains subjected to a metabolic challenge based on dietary energy density. Based on our previous results (Martínez-Micaelo et al., 2016), although both strains showed increased weight gain and adiposity in response to the CAF diet, plasma metabolic profiling evidenced contrasting phenotypic responses. Indeed, although CAF diet challenge induced hyperglycaemia, hypertriglyceridaemia and higher levels of NEFAs in LEW rats, this was not the case for WKY rats. These strains are therefore distinct models of diet induced obesity based on their biochemical response to CAF diet. Furthermore, increased leptin circulating levels was observed independently of the genetic background. In our previous work, using the same model of diet induced obesity, we reported a strong relationship between plasma leptin levels and the nutritional status-associated traits, such as the food intake, adiposity or plasma triglycerides and glucose levels in LEW rats. The metabolic phenotype resulting from central leptin resistance is considered as a pivotal event underlying the development of obesity and its associated dysfunctions (Zhang and Scarpace, 2006). In contrast, the diet-induced hyperleptinaemia was not translated into the impairment in lipid and glucose metabolism in WKY rats. Hence our previous data suggested that the regulation of the leptin-signalling axis was, at least partly, the underlying cause of the observed strain-specific responses to CAF diet.

We show that the diet-induced modulation of the inflammatory response is strain-specific. Although CAF diet promoted inflammation measured by increased fat-derived MCP-1 circulating levels in LEW rats, this was not the case in WKY rats, suggesting the genetic control of CAF-induced inflammatory markers in plasma.

The genetic determinants of the phenotypic response to diet is well-established (Pérusse and Bouchard, 2000; Speakman, 2004). Our results obtained in WKY and LEW rats suggest that the factors underlying the differential response to the dietary energy density are genetically determined

in these strains. The role of peripheral monocytes as professional stress-sensing circulating cells and the pivotal role of leptin in the regulation of energy metabolism, suggest that monocytes and the regulation of their gene expression profiles are valuable for the determination of the host genetic factors influencing the plasticity of the response to diet-induced obesity. Given the contrasting metabolic profiles of WKY and LEW rats, we hypothesised that their monocyte transcriptome will be an intermediate phenotype reflecting the genetic control of CAF-diet in the rat. The differentially expressed genes underlying the strain-dependent response to CAF diet were determined by comparing the expression profiles of monocytes from CAF-fed WKY and LEW rats. Notably, the genes that were significantly overexpressed in WKY rats were significantly enriched for the activation of inflammatory and immune response GO categories. Indeed, genes encoding classical inflammatory mediators, including *Ifng*, *Tnf*, *Nos2*, several interleukins and regulators of eicosanoid production were significantly over-expressed in CAF-fed WKY rats.

The measurement of a broader range of inflammatory plasma biomarkers (not only MCP-1 levels) in the two strains will allow more conclusive correlations between the monocyte transcriptome and systemic markers of obesity-induced inflammation. Off note, the expression of *Mcp1* is not differentially expressed in response to cafeteria diet in monocytes from WKY and LEW strains, so the levels of this inflammatory biomarker in plasma is not explained by the peripheral monocyte transcriptome, but more likely is a result of adiposity gained during the CAF diet.

A co-expression network of the differentially expressed genes was inferred in order to explore transcriptional changes in terms of gene-gene interactions. The integration of the co-expression network inferred from the significant pairwise partial correlations between the strain-specific differential expressed genes and the MCODE-based clustering of nodes lead to biological function-enriched clusters of highly co-expressed genes, based on the assumption that functionally related genes are frequently co-expressed across organisms, constituting conserved transcription modules. Thus, by extracting the cluster

of genes differentially co-expressed, we identified a module where highly co-expressed genes within the cluster were significantly enriched for nutrient sensing, response to hormone stimuli and the NOD-like receptor signalling pathways. Furthermore, *Acss2* was identified as the most highly connected hub gene and also the gene with the highest betweenness centrality score, two topological measures that highlight the impact of this node within the cluster. Based on its position within the cluster, *Acss2* expression changes may have a large impact on the underlying biological functions (Carter et al., 2004). *Acss2* encodes a member of the Acyl-CoA synthetase short-chain family, an enzyme that catalyses the ligation of the acetate derived from the  $\beta$ -oxidation of fatty acids to CoA to produce acetyl-CoA for oxidation through the citric acid cycle (TCA). *Acss2* is therefore an essential enzyme for energy expenditure under ketogenic conditions (Luong et al., 2000). QTLs related to complex obesity-related traits such as body weight, glucose and leptin levels were found to overlap within *Acss2* genomic region. Interestingly, in a previous study investigating glucose intolerance biomarkers in circulating white blood cells (WBC) in high-fat diet (Wopereis et al., 2012), *Acss2* gene expression was proposed as an accurate diagnostic biomarker for glucose intolerance. In addition, *Acss2* gene expression in WBC has been linked with hepatic concentrations of TCA-involved metabolites, including citrate, malate, succinate and fumarate, whose production depends directly on the ACSS2 protein activity (Wopereis et al., 2012). Consistent with these results, we found that, in addition to the overexpression of *Acss2* in CAF-fed WKY rat monocytes, which showed significantly lower glucose levels than CAF-fed LEW rats, there is a significant and strong correlation between *Acss2* gene expression in circulating monocytes and the corresponding glucose plasma levels. *Acss2* levels also significantly correlate with plasma leptin levels and the body weight, traits related to the overlapping QTLs, and with the plasma levels of triglycerides and NEFAs. Therefore, our results confirm that a low expression of *Acss2* in circulating monocytes is an indicator of impaired glucose metabolism. Furthermore, *Acss2* is involved in *de novo* lipogenesis, and in the regulation of the triglyceride storage capacity of adipose tissue, playing a key role as a nutrient sensor. In addition, the expression of *Acss2* is regulated by sterol regulatory element-binding proteins (SREBPs),

transcription factors, which are key regulators of nutritional homeostasis (Osborne, 2000). SREBPs, especially SREBP-1, which is abundantly expressed in monocytes, have also been described as a link between the lipid metabolism and the innate immune response through its direct regulation of core genes involved in the inflammasome activation (Im et al., 2011), supporting the significant over-representation of the NOD-like receptor signalling pathway in the co-expressed genes within the *Acss2*-regulated nutrient sensing cluster. In summary, our results strongly suggested that *Acss2*, a nutrient sensing protein and a diagnostic biomarker of glucose resistance, is a key regulator gene for the differential responsiveness to dietary energy density and a pivotal genetic link for the integration of metabolic and immune homeostasis.

When CAF-diet effects were compared to STD diet, this translated into a moderate change in gene expression profile in circulating monocytes of the LEW rat whereas a substantial set of genes was differentially modulated by the diet in WKY rats including microRNA 223 (*Mir223*) expression. miR-223, the mature form of *Mir223*, is a microRNA that is highly expressed in monocytes and directly targets STAT3 to regulate its activation; the down-regulation of miR-223 expression, such as was induced by the CAF diet in monocytes from WKY rats, is associated with the activation of STAT3 (Chen et al., 2012).

In conclusion, the differential phenotypic and transcriptomics response of LEW and WKY rats to a CAF diet highlighted the prone and resistant responses of these inbred strains to diet-induced obesity. These results suggest that inbred rat strains could be used in nutritional genomics to establish the complex intermediate role of innate immune cells such as monocytes in CAF-induced obesity.

## **ACKNOWLEDGEMENTS**

This study was supported by a grant (AGL2014-55347R) from the Ministerio de Economía y Competitividad of the Spanish Government. J.B. and E.P. acknowledges funding from the Medical Research Council (MR/M004716/1). N. M. holds a PhD grant from the Universitat Rovira i Virgili and mobility grant from the Ministerio de Educación of the Spanish Government.

The authors declare no competing financial interest.



## REFERENCES

- Bader, G. D. and Hogue, C. W. V** (2003). An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* **4**, 2.
- Barsh, G. S., Farooqi, I. S. and O’Rahilly, S.** (2000). Genetics of body-weight regulation. *Nature* **404**, 644–651.
- Behmoaras, J., Bhargal, G., Smith, J., McDonald, K., Mutch, B., Lai, P. C., Domin, J., Game, L., Salama, A., Foxwell, B. M., et al.** (2008). Jund is a determinant of macrophage activation and is associated with glomerulonephritis susceptibility. *Nat. Genet.* **40**, 553–9.
- Behmoaras, J., Smith, J., D’Souza, Z., Bhargal, G., Chawanasuntoropoj, R., Tam, F. W. K., Pusey, C. D., Aitman, T. J. and Cook, H. T.** (2010). Genetic loci modulate macrophage activity and glomerular damage in experimental glomerulonephritis. *J. Am. Soc. Nephrol.* **21**, 1136–44.
- Bouchard, C.** (2008). Gene–environment interactions in the etiology of obesity: defining the fundamentals. *Obesity* **16**, S5–S10.
- Carter, S. L., Brechbühler, C. M., Griffin, M. and Bond, A. T.** (2004). Gene co-expression network topology provides a framework for molecular characterization of cellular state. *Bioinformatics* **20**, 2242–50.
- Chen, Q., Wang, H., Liu, Y., Song, Y., Lai, L., Han, Q., Cao, X. and Wang, Q.** (2012). Inducible microRNA-223 down-regulation promotes TLR-triggered IL-6 and IL-1 $\beta$  production in macrophages by targeting STAT3. *PLoS One* **7**, e42971.
- Cline, M. S., Smoot, M., Cerami, E., Kuchinsky, A., Landys, N., Workman, C., Christmas, R., Avila-Campilo, I., Creech, M., Gross, B., et al.** (2007). Integration of biological networks and gene expression data using Cytoscape. *Nat. Protoc.* **2**, 2366–82.
- D’Souza, Z., McAdoo, S. P., Smith, J., Pusey, C. D., Cook, H. T., Behmoaras, J. and Aitman, T. J.** (2013). Experimental crescentic glomerulonephritis: a new biconogenic rat model. *Dis. Model. Mech.* **6**, 1477–86.

- de la Cruz, N., Bromberg, S., Pasko, D., Shimoyama, M., Twigger, S., Chen, J., Chen, C.-F., Fan, C., Foote, C., Gopinath, G. R., et al.** (2005). The Rat Genome Database (RGD): developments towards a phenome database. *Nucleic Acids Res.* **33**, D485–91.
- Falcon, S. and Gentleman, R.** (2007). Using GOstats to test gene lists for GO term association. *Bioinformatics* **23**, 257–8.
- Hetherington, M. M. and Cecil, J. E.** (2009). Gene-environment interactions in obesity.
- Im, S.-S., Yousef, L., Blaschitz, C., Liu, J. Z., Edwards, R. A., Young, S. G., Raffatellu, M. and Osborne, T. F.** (2011). Linking lipid metabolism to the innate immune response in macrophages through sterol regulatory element binding protein-1a. *Cell Metab.* **13**, 540–9.
- Irizarry, R. A., Hobbs, B., Collin, F., Beazer-Barclay, Y. D., Antonellis, K. J., Scherf, U. and Speed, T. P.** (2003). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* **4**, 249–64.
- Iyer, A., Fairlie, D. P., Prins, J. B., Hammock, B. D. and Brown, L.** (2010). Inflammatory lipid mediators in adipocyte function and obesity. *Nat. Rev. Endocrinol.* **6**, 71–82.
- Lin, C.-Y., Chin, C.-H., Wu, H.-H., Chen, S.-H., Ho, C.-W. and Ko, M.-T.** (2008). Hubba: hub objects analyzer--a framework of interactome hubs identification for network biology. *Nucleic Acids Res.* **36**, W438–43.
- Lumeng, C. N. and Saltiel, A. R.** (2011). Inflammatory links between obesity and metabolic disease. *J. Clin. Invest.* **121**, 2111.
- Luong, A., Hannah, V. C., Brown, M. S. and Goldstein, J. L.** (2000). Molecular characterization of human acetyl-CoA synthetase, an enzyme regulated by sterol regulatory element-binding proteins. *J. Biol. Chem.* **275**, 26458–66.
- Maratou, K., Behmoaras, J., Fewings, C., Srivastava, P., D'Souza, Z., Smith, J., Game, L., Cook, T. and Aitman, T.** (2011). Characterization of the macrophage transcriptome in glomerulonephritis-susceptible and -

resistant rat strains. *Genes Immun.* **12**, 78–89.

**Martínez-Micaelo, N., González-Abuín, N., Ardévol, A., Pinent, M., Petretto, E., Behmoaras, J. and Blay, M.** (2016). Leptin signal transduction underlies the differential metabolic response of LEW and WKY rats to cafeteria diet. *J. Mol. Endocrinol.* **56**, 1–10.

**Morris, J. H., Apeltsin, L., Newman, A. M., Baumbach, J., Wittkop, T., Su, G., Bader, G. D. and Ferrin, T. E.** (2011). clusterMaker: a multi-algorithm clustering plugin for Cytoscape. *BMC Bioinformatics* **12**, 436.

**Opgen-Rhein, R. and Strimmer, K.** (2007). From correlation to causation networks: a simple approximate learning algorithm and its application to high-dimensional plant gene expression data. *BMC Syst. Biol.* **1**, 37.

**Ordovas, J. M.** (2008). Genotype–phenotype associations: modulation by diet and obesity. *Obesity* **16**, S40–S46.

**Osborn, O. and Olefsky, J. M.** (2012). The cellular and signaling networks linking the immune system and metabolism in disease. *Nat. Med.* **18**, 363–374.

**Osborne, T. F.** (2000). Sterol regulatory element-binding proteins (SREBPs): key regulators of nutritional homeostasis and insulin action. *J. Biol. Chem.* **275**, 32379–82.

**Parihar, A., Eubank, T. D. and Doseff, A. I.** (2010). Monocytes and macrophages regulate immunity through dynamic networks of survival and cell death. *J. Innate Immun.* **2**, 204–215.

**Pérusse, L. and Bouchard, C.** (2000). Gene-diet interactions in obesity. *Am. J. Clin. Nutr.* **72**, 1285s–1290s.

**Rothgiesser, K. M., Erener, S., Waibel, S., Lüscher, B. and Hottiger, M. O.** (2010). SIRT2 regulates NF-κB dependent gene expression through deacetylation of p65 Lys310. *J. Cell Sci.* **123**, 4251–8.

**Sampey, B. P., Vanhoose, A. M., Winfield, H. M., Freerman, A. J., Muehlbauer, M. J., Fueger, P. T., Newgard, C. B. and Makowski, L.** (2011). Cafeteria Diet Is a Robust Model of Human Metabolic Syndrome With Liver and Adipose Inflammation: Comparison to High -Fat Diet

*Obesity* **19**, 1109–1117.

**Schäfer, J. and Strimmer, K.** (2005). An empirical Bayes approach to inferring large-scale gene association networks. *Bioinformatics* **21**, 754–64.

**Speakman, J. R.** (2004). Obesity: the integrated roles of environment and genetics. *J. Nutr.* **134**, 2090S–2105S.

**Wopereis, S., Radonjic, M., Rubingh, C., Erk, M. van, Smilde, A., Duyvenvoorde, W. van, Cnubben, N., Kooistra, T., Ommen, B. van and Kleemann, R.** (2012). Identification of prognostic and diagnostic biomarkers of glucose intolerance in ApoE3Leiden mice. *Physiol. Genomics* **44**, 293–304.

**Zhang, Y. and Scarpace, P. J.** (2006). The role of leptin in leptin resistance and obesity. *Physiol. Behav.* **88**, 249–56.

## Figure legends

**Figure 1. Experimental design used to study the role of inbred rat models in nutritional genomics.** The phenotypic response (described previously in (Martínez-Micaelo et al., 2016)) showed a differential susceptibility of LEW and WKY rats to CAF diet, here we studied the differential modulation of the peripheral monocyte transcriptome of LEW and WKY rats by the CAF diet, as consequence of the prone and resistant responses of LEW and WKY rats to the diet induced-obesity, respectively.

**Figure 2. Gene expression profile in circulating monocytes from CAF-fed LEW and WKY rats.** (A) Strain-specific differentially expressed genes. Transcripts with more than 5-fold differences between CAF-fed WKY and LEW rats are shown; the red bars correspond to the transcripts that are overexpressed in WKY rats, and the green bars indicate the transcripts that are overexpressed in LEW rats. (B and C) A summary of the over-represented Gene Ontology Categories based on a Gene Set Enrichment analysis of the differentially expressed genes between CAF-fed WKY and LEW rats.

**Figure 3. Gene co-expression networks for differentially expressed transcripts in CAF-fed WKY and LEW rats.** (A) A simplified representation of the network and the MCODE-based clusters. Cluster 2, significantly enriched as a nutrient sensor, is highlighted, and the significantly over-represented Gene Ontology categories from the selected sub-cluster are shown. The edge shape is related to the direction of the partial correlation; the continuous lines represent partial positive correlations, and dotted lines refer to partial negative correlation. The node colour represents the fold change between the expression of the corresponding transcript in CAF-fed WKY and LEW rats; red nodes correspond to genes that are over-expressed in WKY rats, and green nodes correspond to genes that are over-expressed in LEW rats. The highlighted node within cluster 2 represents a high-scoring differentially expressed gene. (B) Functional enrichment analysis of the MCODE-based co-expressed gene clusters. Significantly over-represented GO terms and KEEG pathways within each cluster are represented.

**Figure 4. CAF diet effect on the expression profile of circulating monocytes in WKY and LEW rats.** (A) CAF diet effect on the transcriptome of circulating monocytes of LEW rats when compared to STD diet. (B) Effect of the CAF diet on the gene expression profile of circulating monocytes of WKY rats.

**Table 1.** Significant Pearson correlations between the log2 expression values of Acyl-CoA synthetase short-chain 2 (Acss2) in circulating monocytes of STD diet- and CAF diet-fed LEW and WKY rats with metabolic-related phenotypes.


<b>Phenotypes</b>	<b>Pearson's r</b>	<b>P-value</b>
Glucose	-0.714	$1.28 \times 10^{-3}$
Triglycerides	-0.683	$2.52 \times 10^{-3}$
Leptin	-0.612	$8.97 \times 10^{-3}$
NEFAs	-0.612	$9.02 \times 10^{-3}$
Total fat weight	-0.592	$1.22 \times 10^{-2}$
Body weight	-0.590	$1.26 \times 10^{-2}$




**Peripheral blood monocyte transcriptome**

**1) Genotype-dependent adaptation to diet-induced obesity**

Differentially expressed genes and functional enrichment analysis




Gene co-expression network



**2) Strain-specific modulation of transcriptomics profile of circulating monocytes by the diet**


Effects of the diet on gene expression profile from LEW and WKY rats




**FIGURE 1**




A




B



C



# A Gene co-expression network



# Functional enrichment of Nutrient-sensing sub-cluster



# B Functional enrichment of co-expressed gene clusters







FIGURE 3

**A**

Diet-induced differentially expressed genes in LEW



**B**

Diet-induced differentially expressed genes in WKY

