

Stochastic Discrete Hamiltonian Variational Integrators

Darryl D. Holm* Tomasz M. Tyranowski†

Mathematics Department
Imperial College London
London SW7 2AZ, UK

Abstract

Variational integrators are derived for structure-preserving simulation of stochastic Hamiltonian systems with a certain type of multiplicative noise arising in geometric mechanics. The derivation is based on a stochastic discrete Hamiltonian which approximates a type-II stochastic generating function for the stochastic flow of the Hamiltonian system. The generating function is obtained by introducing an appropriate stochastic action functional and its corresponding variational principle. The discrete Hamiltonian is then constructed using this stochastic variational principle, in combination with a Galerkin type of discretization based on a certain space of polynomials for approximating the paths of the system, together with quadrature rules to approximate the relevant integrals in the action functional. The resulting integrators are symplectic; they preserve integrals of motion related to Lie group symmetries; and they include stochastic symplectic Runge-Kutta methods as a special case. Several new low-stage stochastic symplectic methods of strong order 1.0 derived using this approach are presented and tested numerically to demonstrate their superior long-time stability and energy behavior compared to nonsymplectic methods.

1 Introduction

Stochastic differential equations (SDEs) play an important role in modeling dynamical systems subject to internal or external random fluctuations. Standard references include [4], [20], [21], [22], [33], [40]. Within this class of problems, we are interested in stochastic Hamiltonian systems, which take the form (see [5], [23], [34])

$$\begin{aligned} dq &= \frac{\partial H}{\partial p} dt + \frac{\partial h}{\partial p} \circ dW(t), \\ dp &= -\frac{\partial H}{\partial q} dt - \frac{\partial h}{\partial q} \circ dW(t), \end{aligned} \tag{1.1}$$

where $H = H(q, p)$ is the Hamiltonian function, $h = h(q, p)$ is the stochastic Hamiltonian, $W(t)$ is the standard one-dimensional Wiener process, and \circ denotes Stratonovich integration. The system

*d.holm@imperial.ac.uk

†t.tyranowski@imperial.ac.uk

(1.1) can be formally regarded as a classical Hamiltonian system with the randomized Hamiltonian given by $\widehat{H}(q, p) = H(q, p) + h(q, p) \circ \dot{W}$, where $H(q, p)$ is the deterministic Hamiltonian and $h(q, p)$ is *another* Hamiltonian, to be specified, which multiplies (in the Stratonovich sense, denoted as \circ) a one-dimensional Gaussian white noise, \dot{W} . Such systems can be used to model, e.g., mechanical systems with uncertainty, or error, assumed to arise from random forcing, limited precision of experimental measurements, or unresolved physical processes on which the deterministic Hamiltonian might otherwise depend. Particular examples include modeling synchrotron oscillations of particles in particle storage rings (see [43], [13]) and stochastic dynamics of the interactions of singular solutions of the EPDiff basic fluids equation (see [19]).

As occurs for other SDEs, most Hamiltonian SDEs cannot be solved analytically and one must resort to numerical simulations to obtain approximate solutions. In principle, general purpose stochastic numerical schemes for SDEs can be applied to stochastic Hamiltonian systems. However, as for their deterministic counterparts, stochastic Hamiltonian systems possess several important geometric features. In particular, their phase space flows preserve the symplectic structure. When simulating these systems numerically, it is therefore advisable that the numerical scheme also preserves such geometric features. Geometric integration of deterministic Hamiltonian systems has been thoroughly studied (see [14], [32], [42] and the references therein) and symplectic integrators have been shown to demonstrate superior performance in long-time simulations of Hamiltonian systems, compared to non-symplectic methods; so it is natural to pursue a similar approach for stochastic Hamiltonian systems. This is a relatively recent pursuit. Stochastic symplectic integrators were first proposed in [34] and [35]. Stochastic generalizations of symplectic partitioned Runge-Kutta methods were analyzed in [27] and [28]. A stochastic generating function approach to constructing stochastic symplectic methods, based on approximately solving a corresponding stochastic Hamilton-Jacobi equation satisfied by the generating function, was proposed in [49] and [50], and this idea was further pursued in [1], [3], [12]. Stochastic symplectic integrators constructed via composition methods were proposed and analyzed in [36]. A first order weak symplectic numerical scheme and an extrapolation method were proposed and their global error was analyzed in [2]. More recently, an approach based on Padé approximations has been used to construct stochastic symplectic methods for linear stochastic Hamiltonian systems (see [45]).

An important class of geometric integrators are *variational integrators*. This type of numerical schemes is based on discrete variational principles and provides a natural framework for the discretization of Lagrangian systems, including forced, dissipative, or constrained ones. These methods have the advantage that they are symplectic, and in the presence of a symmetry, satisfy a discrete version of Noether's theorem. For an overview of variational integration for deterministic systems see [31]; see also [17], [24], [25], [37], [38], [41], [47], [48]. Variational integrators were introduced in the context of finite-dimensional mechanical systems, but were later generalized to Lagrangian field theories (see [30]) and applied in many computations, for example in elasticity, electrodynamics, or fluid dynamics; see [26], [39], [44], [46].

Stochastic variational integrators were first introduced in [7] and further studied in [6]. However, those integrators were restricted to the special case when the stochastic Hamiltonian $h = h(q)$ was independent of p , and only low-order Runge-Kutta types of discretization were considered. In the present work we extend the idea of stochastic variational integration to general stochastic Hamiltonian systems by generalizing the variational principle introduced in [25] and applying a Galerkin type of discretization (see [31], [24], [25], [38], [37]), which leads to a more general class of stochastic symplectic integrators than those presented in [6], [7], [27], and [28]. Our approach consists in approximating a generating function for the stochastic flow of the Hamiltonian system,

but unlike in [49] and [50], we do make this discrete approximation by exploiting its variational characterization, rather than solving the corresponding Hamilton-Jacobi equation.

Main content The main content of the remainder of this paper is, as follows.

In Section 2 we introduce a stochastic variational principle and a stochastic generating function suitable for considering stochastic Hamiltonian systems, and we discuss their properties.

In Section 3 we present a general framework for constructing stochastic Galerkin variational integrators, prove the symplecticity and conservation properties of such integrators, show they contain the stochastic symplectic Runge-Kutta methods of [27], [28] as a special case, and finally present several particularly interesting examples of new low-stage stochastic symplectic integrators of strong order 1.0 derived with our general methodology.

In Section 4 we present the results of our numerical tests, which verify the theoretical convergence rates and the excellent long-time performance of our integrators compared to some popular non-symplectic methods.

Section 5 contains the summary of our work.

2 Variational principle for stochastic Hamiltonian systems

The stochastic variational integrators proposed in [7] and [6] were formulated for dynamical systems which are described by a Lagrangian and which are subject to noise whose magnitude depends only on the position q . Therefore, these integrators are applicable to (1.1) only when the stochastic Hamiltonian $h = h(q)$ is independent of p and the Hamiltonian H is non-degenerate (i.e., the associated Legendre transform is invertible). However, in the case of general $h = h(q, p)$ the paths $q(t)$ of the system become almost surely nowhere differentiable, which poses a difficulty in interpreting the meaning of the corresponding Lagrangian. Therefore, we need a different sort of action functional and variational principle to construct stochastic symplectic integrators for (1.1). To this end, we will generalize the approach taken in [25]. To begin, in the next section, we will introduce an appropriate stochastic action functional and show that it can be used to define a type-II generating function for the stochastic flow of the system (1.1).

2.1 Stochastic variational principle

Let the Hamiltonian $H : T^*Q \rightarrow \mathbb{R}$ and the stochastic Hamiltonian $h : T^*Q \rightarrow \mathbb{R}$ be defined on the cotangent bundle T^*Q of the configuration manifold Q , and let (q, p) denote the canonical coordinates on T^*Q . For simplicity, in this work we assume that the configuration manifold has a vector space structure, $Q \cong \mathbb{R}^N$, so that $T^*Q = Q \times Q^* \cong \mathbb{R}^N \times \mathbb{R}^N$ and $TQ = Q \times Q \cong \mathbb{R}^N \times \mathbb{R}^N$. In this case, the natural pairing between one-forms and vectors can be identified with the scalar product on \mathbb{R}^N , that is, $\langle (q, p), (q, \dot{q}) \rangle = p \cdot \dot{q}$, where (q, \dot{q}) denotes the coordinates on TQ . Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space with the filtration $\{\mathcal{F}_t\}_{t \geq 0}$, and let $W(t)$ denote a standard one-dimensional Wiener process on that probability space (such that $W(t)$ is \mathcal{F}_t -measurable). For a given time interval $[t_a, t_b]$ with $t_b > t_a \geq 0$, define the space

$$C([t_a, t_b]) = \{(q, p) : \Omega \times [t_a, t_b] \longrightarrow T^*Q \mid q, p \text{ are almost surely continuous } \mathcal{F}_t\text{-adapted semimartingales}\}. \quad (2.1)$$

Since we assume $T^*Q \cong \mathbb{R}^N \times \mathbb{R}^N$, the space $C([t_a, t_b])$ is a vector space (see [40]). Therefore, we can identify the tangent space $TC([t_a, t_b]) \cong C([t_a, t_b]) \times C([t_a, t_b])$. We can now define the following stochastic action functional, $\mathcal{B} : \Omega \times C([t_a, t_b]) \longrightarrow \mathbb{R}$,

$$\mathcal{B}[q(\cdot), p(\cdot)] = p(t_b)q(t_b) - \int_{t_a}^{t_b} [p \circ dq - H(q(t), p(t)) dt - h(q(t), p(t)) \circ dW(t)], \quad (2.2)$$

where \circ denotes Stratonovich integration, and we have omitted writing the elementary events $\omega \in \Omega$ as arguments of functions, following the standard convention in stochastic analysis.

Theorem 2.1 (Stochastic Variational Principle in Phase Space). *Suppose that $H(q, p)$ and $h(q, p)$ are C^2 functions of their arguments with globally Lipschitz derivatives. If the curve $(q(t), p(t))$ in T^*Q satisfies the stochastic Hamiltonian system (1.1) for $t \in [t_a, t_b]$, where $t_b \geq t_a > 0$, then the pair $(q(\cdot), p(\cdot))$ is a critical point of the stochastic action functional (2.2), that is,*

$$\delta\mathcal{B}[q(\cdot), p(\cdot)] \equiv \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \mathcal{B}[q(\cdot) + \epsilon\delta q(\cdot), p(\cdot) + \epsilon\delta p(\cdot)] = 0, \quad (2.3)$$

almost surely for all variations $(\delta q(\cdot), \delta p(\cdot)) \in C([t_a, t_b])$ such that almost surely $\delta q(t_a) = 0$ and $\delta p(t_b) = 0$.

Proof. Let the curve $(q(t), p(t))$ in T^*Q satisfy (1.1) for $t \in [t_a, t_b]$. It then follows that the stochastic processes $q(t)$ and $p(t)$ are almost surely continuous, \mathcal{F}_t -adapted semimartingales, that is, $(q(\cdot), p(\cdot)) \in C([t_a, t_b])$ (see [4], [40]). We calculate the variation (2.3) as

$$\begin{aligned} \delta\mathcal{B}[q(\cdot), p(\cdot)] &= p(t_b)\delta q(t_b) - \int_{t_a}^{t_b} p(t) \circ d\delta q(t) - \int_{t_a}^{t_b} \delta p(t) \circ dq(t) \\ &\quad + \int_{t_a}^{t_b} \left[\frac{\partial H}{\partial q}(q(t), p(t)) \delta q(t) + \frac{\partial H}{\partial p}(q(t), p(t)) \delta p(t) \right] dt \\ &\quad + \int_{t_a}^{t_b} \left[\frac{\partial h}{\partial q}(q(t), p(t)) \delta q(t) + \frac{\partial h}{\partial p}(q(t), p(t)) \delta p(t) \right] \circ dW(t), \end{aligned} \quad (2.4)$$

where we have used the end point condition, $\delta p(t_b) = 0$. Since the Hamiltonians are C^2 and the processes $q(t)$, $p(t)$ are almost surely continuous, in the last two lines we have used a dominated convergence argument to interchange differentiation with respect to ϵ and integration with respect to t and $W(t)$. Upon applying the integration by parts formula for semimartingales (see [40]), we find

$$\int_{t_a}^{t_b} p(t) \circ d\delta q(t) = p(t_b)\delta q(t_b) - p(t_a)\delta q(t_a) - \int_{t_a}^{t_b} \delta q(t) \circ dp(t). \quad (2.5)$$

Substituting and rearranging terms produces,

$$\begin{aligned} \delta\mathcal{B}[q(\cdot), p(\cdot)] &= \int_{t_a}^{t_b} \delta q(t) \left[\circ dp(t) + \frac{\partial H}{\partial q}(q(t), p(t)) dt + \frac{\partial h}{\partial q}(q(t), p(t)) \circ dW(t) \right] \\ &\quad - \int_{t_a}^{t_b} \delta p(t) \left[\circ dq(t) - \frac{\partial H}{\partial p}(q(t), p(t)) dt - \frac{\partial h}{\partial p}(q(t), p(t)) \circ dW(t) \right], \end{aligned} \quad (2.6)$$

where we have used $\delta q(t_a) = 0$. The terms in the brackets vanish almost surely, since $(q(t), p(t))$ satisfy (1.1). Therefore, $\delta\mathcal{B} = 0$, almost surely. \square

Remark: The converse theorem, that is, if $(q(\cdot), p(\cdot))$ is a critical point of the stochastic action functional (2.2), then the curve $(q(t), p(t))$ satisfies (1.1), was proved (for Lagrangian systems) by Bou-Rabee & Owhadi [7] in the special case when $h = h(q)$ is independent of p . In that case, one can assume that $q(t)$ is continuously differentiable. In the general case, however, $q(t)$ is not differentiable, and the ideas of [7] cannot be applied directly. In fact, the converse theorem may not hold, at least not without more stringent assumptions; see the discussion in [23]. We leave this as an open question. Here, we will use the action functional (2.2) and the variational principle (2.3) to construct numerical schemes, and we will directly verify that these numerical schemes converge to solutions of (1.1).

2.2 Stochastic type-II generating function

When the Hamiltonian functions $H(q, p)$ and $h(q, p)$ satisfy standard measurability and regularity conditions (e.g., as in Theorem 2.1), then the system (1.1) possesses a pathwise unique stochastic flow $F_{t, t_0} : \Omega \times T^*Q \rightarrow T^*Q$. It can be proved that for fixed t, t_0 this flow is mean-square differentiable with respect to the q, p arguments, and is also almost surely a diffeomorphism (see [4], [20], [21], [22]). Moreover, F_{t, t_0} almost surely preserves the canonical symplectic form $\Omega_{T^*Q} = \sum_{i=1}^N dq^i \wedge dp^i$ (see [35], [5], [23]), that is,

$$F_{t, t_0}^* \Omega_{T^*Q} = \Omega_{T^*Q}, \quad (2.7)$$

where F_{t, t_0}^* denotes the pull-back by the flow F_{t, t_0} . We will show below that the action functional (2.2) can be used to construct a type II generating function for F_{t, t_0} . Let $(\bar{q}(t), \bar{p}(t))$ be a particular solution of (1.1) on $[t_a, t_b]$. Suppose that for almost all $\omega \in \Omega$ there is an open neighborhood $\mathcal{U}(\omega) \subset Q$ of $\bar{q}(\omega, t_a)$, an open neighborhood $\mathcal{V}(\omega) \subset Q^*$ of $\bar{p}(\omega, t_b)$, and an open neighborhood $\mathcal{W}(\omega) \subset T^*Q$ of the curve $(\bar{q}(\omega, t), \bar{p}(\omega, t))$ such that for all $q_a \in \mathcal{U}(\omega)$ and $p_b \in \mathcal{V}(\omega)$ there exists a pathwise unique solution $(\bar{q}(\omega, t; q_a, p_b), \bar{p}(\omega, t; q_a, p_b))$ of (1.1) which satisfies $\bar{q}(\omega, t_a; q_a, p_b) = q_a$, $\bar{p}(\omega, t_b; q_a, p_b) = p_b$, and $(\bar{q}(\omega, t; q_a, p_b), \bar{p}(\omega, t; q_a, p_b)) \in \mathcal{W}(\omega)$ for $t_a \leq t \leq t_b$. (As in the deterministic case, for t_b sufficiently close to t_a one can argue that such neighborhoods exist; see [29].) Define the function $S : \mathcal{Y} \rightarrow \mathbb{R}$ as

$$S(q_a, p_b) = \mathcal{B}[\bar{q}(\cdot; q_a, p_b), \bar{p}(\cdot; q_a, p_b)], \quad (2.8)$$

where the domain $\mathcal{Y} \subset \Omega \times T^*Q$ is given by $\mathcal{Y} = \bigcup_{\omega \in \Omega} \{\omega\} \times \mathcal{U}(\omega) \times \mathcal{V}(\omega)$.

Theorem 2.2. *The function $S(q_a, p_b)$ is a type-II stochastic generating function for the stochastic mapping F_{t_b, t_a} , that is, $F_{t_b, t_a} : (q_a, p_a) \rightarrow (q_b, p_b)$ is implicitly given by the equations*

$$q_b = D_2 S(q_a, p_b), \quad p_a = D_1 S(q_a, p_b), \quad (2.9)$$

where the derivatives are understood in the mean-square sense.

Proof. Under appropriate regularity assumptions on the Hamiltonians, the solutions $\bar{q}(t; q_a, p_b)$ and $\bar{p}(t; q_a, p_b)$ are mean-square differentiable with respect to the parameters q_a and p_b , and the partial derivatives are semimartingales (see [4]). We calculate the derivative of S as

$$\begin{aligned} \frac{\partial S}{\partial q_a}(q_a, p_b) &= p_b \frac{\partial \bar{q}(t_b)}{\partial q_a} - \int_{t_a}^{t_b} \frac{\partial \bar{p}(t)}{\partial q_a} \circ d\bar{q}(t) - \int_{t_a}^{t_b} \bar{p}(t) \circ d \frac{\partial \bar{q}(t)}{\partial q_a} \\ &+ \int_{t_a}^{t_b} \left[\frac{\partial \bar{q}(t)}{\partial q_a} \frac{\partial H}{\partial q}(\bar{q}(t), \bar{p}(t)) + \frac{\partial \bar{p}(t)}{\partial q_a} \frac{\partial H}{\partial p}(\bar{q}(t), \bar{p}(t)) \right] dt \\ &+ \int_{t_a}^{t_b} \left[\frac{\partial \bar{q}(t)}{\partial q_a} \frac{\partial h}{\partial q}(\bar{q}(t), \bar{p}(t)) + \frac{\partial \bar{p}(t)}{\partial q_a} \frac{\partial h}{\partial p}(\bar{q}(t), \bar{p}(t)) \right] \circ dW(t), \end{aligned} \quad (2.10)$$

where for notational convenience we have omitted writing q_a and p_b explicitly as arguments of $\bar{q}(t)$ and $\bar{p}(t)$. Applying the integration by parts formula for semimartingales (see [40]), we find

$$\int_{t_a}^{t_b} \bar{p}(t) \circ d \frac{\partial \bar{q}(t)}{\partial q_a} = p_b \frac{\partial \bar{q}(t_b)}{\partial q_a} - \bar{p}(t_a) - \int_{t_a}^{t_b} \frac{\partial \bar{q}(t)}{\partial q_a} \circ d\bar{p}(t). \quad (2.11)$$

Substituting and rearranging terms, we obtain the result,

$$\begin{aligned} \frac{\partial S}{\partial q_a}(q_a, p_b) &= \bar{p}(t_a) + \int_{t_a}^{t_b} \frac{\partial \bar{q}(t)}{\partial q_a} \left[\circ d\bar{p} + \frac{\partial H}{\partial q}(\bar{q}(t), \bar{p}(t)) dt + \frac{\partial h}{\partial q}(\bar{q}(t), \bar{p}(t)) \circ dW(t) \right] \\ &+ \int_{t_a}^{t_b} \frac{\partial \bar{p}(t)}{\partial q_a} \left[\circ d\bar{q} - \frac{\partial H}{\partial p}(\bar{q}(t), \bar{p}(t)) dt - \frac{\partial h}{\partial p}(\bar{q}(t), \bar{p}(t)) \circ dW(t) \right] \\ &= \bar{p}(t_a), \end{aligned} \quad (2.12)$$

since $(\bar{q}(t), \bar{p}(t))$ is a solution of (1.1). Similarly we show $\partial S / \partial p_b(q_a, p_b) = \bar{q}(t_b)$. By definition of the flow, then $F_{t_b, t_a}(q_a, \bar{p}(t_a)) = (\bar{q}(t_b), p_b)$. □

We can consider $S(q_a, p_b)$ as a function of time if we let t_b vary. Let us denote this function as $S_t(q_a, p)$. Below we show that $S_t(q_a, p)$ satisfies a certain stochastic partial differential equation, which is a stochastic generalization of the Hamilton-Jacobi equation considered in [25].

Proposition 2.3 (Type II Stochastic Hamilton-Jacobi Equation). *Let the time-dependent type-II generating function be defined as*

$$S_2(q_a, p, t) \equiv S_t(q_a, p) = p\bar{q}(t) - \int_{t_a}^t \left[\bar{p}(\tau) \circ d\bar{q}(\tau) - H(\bar{q}(\tau), \bar{p}(\tau)) d\tau - h(\bar{q}(\tau), \bar{p}(\tau)) \circ dW(\tau) \right], \quad (2.13)$$

where $\bar{q}(\tau) \equiv \bar{q}(\tau; q_a, p)$ and $\bar{p}(\tau) \equiv \bar{p}(\tau; q_a, p)$ as before. Then the function $S_2(q_a, p, t)$ satisfies the following stochastic partial differential equation

$$dS_2 = H\left(\frac{\partial S_2}{\partial p}, p\right) dt + h\left(\frac{\partial S_2}{\partial p}, p\right) \circ dW(t), \quad (2.14)$$

where dS_2 denotes the stochastic differential of S_2 with respect to the t variable.

Proof. Choose an arbitrary pair $(q_a, p_a) \in T^*Q$ and define the particular solution $(\bar{q}(\tau), \bar{p}(\tau)) = F_{\tau, t_a}(q_a, p_a)$. Form the function $S_2(q_a, \bar{p}(t), t)$ and consider its total stochastic differential $\bar{d}S_2(q_a, \bar{p}(t), t)$ with respect to time. On one hand, the rules of Stratonovich calculus imply

$$\bar{d}S_2(q_a, \bar{p}(t), t) = dS_2(q_a, \bar{p}(t), t) + \frac{\partial S_2}{\partial p}(q_a, \bar{p}(t), t) \circ d\bar{p}(t), \quad (2.15)$$

where dS_2 denotes the partial stochastic differential of S_2 with respect to the t argument. On the other hand, integration by parts in (2.13) implies

$$\bar{d}S_2(q_a, \bar{p}(t), t) = \bar{q}(t) \circ d\bar{p}(t) + H(\bar{q}(t), \bar{p}(t)) dt + h(\bar{q}(t), \bar{p}(t)) \circ dW(t). \quad (2.16)$$

Comparing (2.15) and (2.16), and using (2.9), we obtain

$$dS_2(q_a, \bar{p}(t), t) = H\left(\frac{\partial S_2}{\partial p}(q_a, \bar{p}(t), t), \bar{p}(t)\right) dt + h\left(\frac{\partial S_2}{\partial p}(q_a, \bar{p}(t), t), \bar{p}(t)\right) \circ dW(t). \quad (2.17)$$

This equation is satisfied along a particular path $\bar{p}(t)$, however, as in the discussion preceding Theorem 2.2, we can argue, slightly informally, that for almost all $\omega \in \Omega$, and for t sufficiently close to t_a , one can find open neighborhoods $\mathcal{U}(\omega) \subset Q$ and $\mathcal{V}(\omega) \subset Q^*$ which can be connected by the flow F_{t, t_a} , i.e., given $q_a \in \mathcal{U}(\omega)$ and $p \in \mathcal{V}(\omega)$, there exists a pathwise unique solution $(\bar{q}(\omega, \tau), \bar{p}(\omega, \tau))$ such that $\bar{q}(\omega, t_a) = q_a$ and $\bar{p}(\omega, t) = p$. With this assumption, (2.17) can be reformulated as the full-blown stochastic PDE (2.14). □

Remark: Similar stochastic Hamilton-Jacobi equations were introduced in [49], [50], where they were used for constructing stochastic symplectic integrators by considering series expansions of generating functions in terms of multiple Stratonovich integrals. This was a direct generalization of a similar technique for deterministic Hamiltonian systems (see [14]). In this work we explore the generalized Galerkin framework for constructing approximations of the generating function $S(q_a, p_b)$ in (2.9) by using its variational characterization (2.8). Our approach is a stochastic generalization of the techniques proposed in [25] and [38] for deterministic Hamiltonian and Lagrangian systems.

2.3 Stochastic Noether's theorem

Let a Lie group G act on Q by the left action $\Phi : G \times Q \rightarrow Q$. The Lie group G then acts on TQ and T^*Q by the tangent $\Phi^{TQ} : G \times TQ \rightarrow TQ$ and cotangent $\Phi^{T^*Q} : G \times T^*Q \rightarrow T^*Q$ lift actions, respectively, given in coordinates by the formulas (see [18], [29])

$$\begin{aligned}\Phi_g^{TQ}(q, \dot{q}) &\equiv \Phi^{TQ}(g, (q, \dot{q})) = \left(\Phi_g^i(q), \frac{\partial \Phi_g^i}{\partial q^j}(q) \dot{q}^j \right), \\ \Phi_g^{T^*Q}(q, p) &\equiv \Phi^{T^*Q}(g, (q, p)) = \left(\Phi_g^i(q), p_j \frac{\partial \Phi_{g^{-1}}^j}{\partial q^i}(\Phi_g(q)) \right),\end{aligned}\tag{2.18}$$

where $i, j = 1, \dots, N$ and summation is implied over repeated indices. Let \mathfrak{g} denote the Lie algebra of G and $\exp : \mathfrak{g} \rightarrow G$ the exponential map (see [18], [29]). Each element $\xi \in \mathfrak{g}$ defines the infinitesimal generators ξ_Q , ξ_{TQ} , and ξ_{T^*Q} , which are vector fields on Q , TQ , and T^*Q , respectively, given by

$$\xi_Q(q) = \left. \frac{d}{d\lambda} \right|_{\lambda=0} \Phi_{\exp \lambda \xi}(q), \quad \xi_{TQ}(q, \dot{q}) = \left. \frac{d}{d\lambda} \right|_{\lambda=0} \Phi_{\exp \lambda \xi}^{TQ}(q, \dot{q}), \quad \xi_{T^*Q}(q, p) = \left. \frac{d}{d\lambda} \right|_{\lambda=0} \Phi_{\exp \lambda \xi}^{T^*Q}(q, p).\tag{2.19}$$

The momentum map $J : T^*Q \rightarrow \mathfrak{g}^*$ associated with the action Φ^{T^*Q} is defined as the mapping such that for all $\xi \in \mathfrak{g}$ the function $J_\xi : T^*Q \ni (q, p) \rightarrow \langle J(q, p), \xi \rangle \in \mathbb{R}$ is the Hamiltonian for the infinitesimal generator ξ_{T^*Q} , i.e.,

$$\xi_{T^*Q}^q = \frac{\partial J_\xi}{\partial p}, \quad \xi_{T^*Q}^p = -\frac{\partial J_\xi}{\partial q},\tag{2.20}$$

where $\xi_{T^*Q}(q, p) = (q, p, \xi_{T^*Q}^q(q, p), \xi_{T^*Q}^p(q, p))$. The momentum map J can be explicitly expressed as (see [18], [29])

$$J_\xi(q, p) = p \cdot \xi_Q(q).\tag{2.21}$$

Noether's theorem for deterministic Hamiltonian systems relates symmetries of the Hamiltonian to quantities preserved by the flow of the system. It turns out that this result carries over to the stochastic case, as well. A stochastic version of Noether's theorem was proved in [5] and [23]. For completeness, and for the benefit of the reader, below we state and provide a less involved proof of Noether's theorem for stochastic Hamiltonian systems.

Theorem 2.4 (Stochastic Noether's theorem). *Suppose that the Hamiltonians $H : T^*Q \rightarrow \mathbb{R}$ and $h : T^*Q \rightarrow \mathbb{R}$ are invariant with respect to the cotangent lift action $\Phi^{T^*Q} : G \times T^*Q \rightarrow T^*Q$ of the Lie group G , that is,*

$$H \circ \Phi_g^{T^*Q} = H, \quad h \circ \Phi_g^{T^*Q} = h,\tag{2.22}$$

for all $g \in G$. Then the cotangent lift momentum map $J : T^*Q \rightarrow \mathfrak{g}^*$ associated with Φ^{T^*Q} is almost surely preserved along the solutions of the stochastic Hamiltonian system (1.1).

Proof. Equation (2.22) implies that the Hamiltonians are infinitesimally invariant with respect to the action of G , that is, for all $\xi \in \mathfrak{g}$ we have

$$dH \cdot \xi_{T^*Q} = 0, \quad dh \cdot \xi_{T^*Q} = 0, \quad (2.23)$$

where dH and dh denote differentials with respect to the variables q and p . Let $(q(t), p(t))$ be a solution of (1.1) and consider the stochastic process $J_\xi(q(t), p(t))$, where $\xi \in \mathfrak{g}$ is arbitrary. Using the rules of Stratonovich calculus we can calculate the stochastic differential

$$\begin{aligned} dJ_\xi(q(t), p(t)) &= \frac{\partial J_\xi}{\partial q}(q(t), p(t)) \circ dq + \frac{\partial J_\xi}{\partial p}(q(t), p(t)) \circ dp \\ &= - \left(\frac{\partial H}{\partial q} \xi_{T^*Q}^q + \frac{\partial H}{\partial p} \xi_{T^*Q}^p \right) dt - \left(\frac{\partial h}{\partial q} \xi_{T^*Q}^q + \frac{\partial h}{\partial p} \xi_{T^*Q}^p \right) \circ dW \\ &= -(dH \cdot \xi_{T^*Q}) dt - (dh \cdot \xi_{T^*Q}) \circ dW = 0, \end{aligned} \quad (2.24)$$

where we used (1.1), (2.20), and (2.23). Therefore, $J_\xi(q(t), p(t)) = \text{const}$ almost surely for all $\xi \in \mathfrak{g}$, which completes the proof. \square

3 Stochastic Galerkin Hamiltonian Variational Integrators

If the converse to Theorem 2.1 holds, then the generating function $S(q_a, p_b)$ defined in (2.8) could be equivalently characterized by

$$S(q_a, p_b) = \underset{\substack{(q,p) \in C([t_a, t_b]) \\ q(t_a)=q_a, p(t_b)=p_b}}{\text{ext}} \mathcal{B}[q(\cdot), p(\cdot)], \quad (3.1)$$

where the extremum is taken pointwise in the probability space Ω . This characterization allows us to construct stochastic Galerkin variational integrators by choosing a finite dimensional subspace of $C([t_a, t_b])$ and a quadrature rule for approximating the integrals in the action functional \mathcal{B} . Galerkin variational integrators for deterministic systems were first introduced in [31], and further developed in [17], [24], [25], [37], and [38]. In the remainder of the paper, we will generalize these ideas to the stochastic case.

3.1 Construction of the integrator

Suppose we would like to solve (1.1) on the interval $[0, T]$ with the initial conditions $(q_0, p_0) \in T^*Q$. Consider the discrete set of times $t_k = k \cdot \Delta t$ for $k = 0, 1, \dots, K$, where $\Delta t = T/K$ is the time step. In order to determine the discrete curve $\{(q_k, p_k)\}_{k=0, \dots, K}$ that approximates the exact solution of (1.1) at times t_k we need to construct an approximation of the exact stochastic flow F_{t_{k+1}, t_k} on each interval $[t_k, t_{k+1}]$, so that $(q_{k+1}, p_{k+1}) \approx F_{t_{k+1}, t_k}(q_k, p_k)$. Let us consider the space $C^s([t_k, t_{k+1}]) \subset C([t_k, t_{k+1}])$ defined as

$$C^s([t_k, t_{k+1}]) = \{(q, p) \in C([t_k, t_{k+1}]) \mid q \text{ is a polynomial of degree } s\}. \quad (3.2)$$

For convenience, we will express $q(t)$ in terms of Lagrange polynomials. Consider the control points $0 = d_0 < d_1 < \dots < d_s = 1$ and let the corresponding Lagrange polynomials of degree s be denoted by $l_{\mu,s}(\tau)$, that is, $l_{\mu,s}(d_\nu) = \delta_{\mu\nu}$. A polynomial trajectory $q_d(t; q^\mu)$ can then be expressed as

$$q_d(t_k + \tau \Delta t; q^\mu) = \sum_{\mu=0}^s q^\mu l_{\mu,s}(\tau), \quad \dot{q}_d(t_k + \tau \Delta t; q^\mu) = \frac{1}{\Delta t} \sum_{\mu=0}^s q^\mu \dot{l}_{\mu,s}(\tau), \quad (3.3)$$

where $q^\nu = q_d(t_k + d_\nu \Delta t; q^\mu)$ for $\nu = 0, \dots, s$ are the control values, \dot{q}_d denotes the time derivative of q_d , and $\dot{l}_{\mu,s}$ denotes the derivative of the Lagrange polynomial $l_{\mu,s}$ with respect to its argument. The restriction of the action functional (2.2) to the space $C^s([t_k, t_{k+1}])$ takes the form

$$\mathcal{B}^s[q_d(\cdot; q^\mu), p(\cdot)] = p(t_{k+1})q^s - \int_{t_k}^{t_{k+1}} [p(t)\dot{q}_d(t) - H(q_d(t), p(t))] dt + \int_{t_k}^{t_{k+1}} h(q_d(t), p(t)) \circ dW(t), \quad (3.4)$$

since for differentiable functions $dq_d(t) = \dot{q}_d(t) dt$, where for brevity $q_d(t) \equiv q_d(t; q^\mu)$. Next we approximate the integrals in (3.4) using numerical quadrature rules $(\alpha_i, c_i)_{i=1}^r$ and $(\beta_i, c_i)_{i=1}^r$, where $0 \leq c_1 < \dots < c_r \leq 1$ are the quadrature points, and α_i, β_i are the corresponding weights. At this point we only make a general assumption that for each i we have $\alpha_i \neq 0$ or $\beta_i \neq 0$. More specific examples will be presented in Section 3.4. The approximate action functional takes the form

$$\begin{aligned} \bar{\mathcal{B}}^s[q_d(\cdot; q^\mu), p(\cdot)] &= p(t_{k+1})q^s - \Delta t \sum_{i=1}^r \alpha_i [p(t_k + c_i \Delta t) \dot{q}_d(t_k + c_i \Delta t) - H(q_d(t_k + c_i \Delta t), p(t_k + c_i \Delta t))] \\ &\quad + \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t), p(t_k + c_i \Delta t)), \end{aligned} \quad (3.5)$$

where $\Delta W = W(t_{k+1}) - W(t_k)$ is the increment of the Wiener process over the considered time interval and is a Gaussian random variable with zero mean and variance Δt . The way of approximating the Stratonovich integral above was inspired by the ideas presented in [7], [11], [28], [34], and [35]. Note that since we only used $\Delta W = \int_{t_k}^{t_{k+1}} dW(t)$ in the above approximation, we can in general expect strong convergence of order 1.0 only. To obtain strong convergence of higher order we would also need to include higher-order multiple Stratonovich integrals, e.g., to achieve convergence of order 1.5 we would need to include terms involving $\Delta Z = \int_{t_k}^{t_{k+1}} \int_{t_k}^t dW(\xi) dt$ (see [11], [34], [35]). We can now approximate the generating function $S(q_k, p_{k+1})$ with the discrete Hamiltonian function $H_d^+(q_k, p_{k+1})$ defined as

$$\begin{aligned} H_d^+(q_k, p_{k+1}) &= \underset{\substack{q^1, \dots, q^s \in Q \\ P_1, \dots, P_r \in Q^* \\ q^0 = q_k}}{\text{ext}} \left\{ p_{k+1} q^s - \Delta t \sum_{i=1}^r \alpha_i [P_i \dot{q}_d(t_k + c_i \Delta t) - H(q_d(t_k + c_i \Delta t), P_i)] \right. \\ &\quad \left. + \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t), P_i) \right\}, \end{aligned} \quad (3.6)$$

where we denoted $P_i \equiv p(t_k + c_i \Delta t)$. The numerical scheme $(q_k, p_k) \longrightarrow (q_{k+1}, p_{k+1})$ is now implicitly generated by $H_d^+(q_k, p_{k+1})$ as in (2.9):

$$q_{k+1} = D_2 H_d^+(q_k, p_{k+1}), \quad p_k = D_1 H_d^+(q_k, p_{k+1}). \quad (3.7)$$

Equations (3.6) and (3.7) can be written together as the following system:

$$-p_k = \sum_{i=1}^r \alpha_i \left[P_i \dot{l}_{0,s}(c_i) - \Delta t \frac{\partial H}{\partial q}(t_k + c_i \Delta t) l_{0,s}(c_i) \right] - \Delta W \sum_{i=1}^r \beta_i \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{0,s}(c_i), \quad (3.8a)$$

$$0 = \sum_{i=1}^r \alpha_i \left[P_i \dot{l}_{\mu,s}(c_i) - \Delta t \frac{\partial H}{\partial q}(t_k + c_i \Delta t) l_{\mu,s}(c_i) \right] - \Delta W \sum_{i=1}^r \beta_i \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{\mu,s}(c_i), \quad (3.8b)$$

$$p_{k+1} = \sum_{i=1}^r \alpha_i \left[P_i \dot{l}_{s,s}(c_i) - \Delta t \frac{\partial H}{\partial q}(t_k + c_i \Delta t) l_{s,s}(c_i) \right] - \Delta W \sum_{i=1}^r \beta_i \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{s,s}(c_i), \quad (3.8c)$$

$$\alpha_i \dot{q}_d(t_k + c_i \Delta t) = \alpha_i \frac{\partial H}{\partial p}(t_k + c_i \Delta t) + \beta_i \frac{\Delta W}{\Delta t} \frac{\partial h}{\partial p}(t_k + c_i \Delta t), \quad (3.8d)$$

$$q_{k+1} = q^s, \quad (3.8e)$$

where $\mu = 1, \dots, s-1$ in (3.8b), $i = 1, \dots, r$ in (3.8d), and for brevity we have introduced the notation

$$H(t_k + c_i \Delta t) \equiv H(q_d(t_k + c_i \Delta t), p(t_k + c_i \Delta t)) \quad (\text{similarly for } h).$$

Equation (3.8a) corresponds to the second equation in (3.7), equations (3.8b), (3.8c) and (3.8d) correspond to extremizing (3.6) with respect to q^1, \dots, q^s , and P_1, \dots, P_r , respectively, and finally (3.8e) is the first equation in (3.7). Knowing (q_k, p_k) , the system (3.8) allows us to solve for (q_{k+1}, p_{k+1}) : we first simultaneously solve (3.8a), (3.8b) and (3.8d) ($(s+r)N$ equations) for q^1, \dots, q^s and P_1, \dots, P_r ($(s+r)N$ unknowns); then $q_{k+1} = q^s$ and (3.8c) is an explicit update rule for p_{k+1} . When $h \equiv 0$, then (3.8) reduces to the deterministic Galerkin variational integrator discussed in [38].

3.2 Properties of stochastic Galerkin variational integrators

The key features of variational integrators are their symplecticity and exact preservation of the discrete counterparts of conserved quantities (momentum maps) related to the symmetries of the Lagrangian or Hamiltonian (see [31]). These properties carry over to the stochastic case, as was first demonstrated in [7] for Lagrangian systems. In what follows, we will show that the stochastic Galerkin Hamiltonian variational integrators constructed in Section 3.1 also possess these properties.

Theorem 3.1 (Symplecticity of the discrete flow). *Let $F_{t_{k+1}, t_k}^+ : \Omega \times T^*Q \rightarrow T^*Q$ be the discrete stochastic flow implicitly defined by the discrete Hamiltonian H_d^+ as in (3.7). Then F_{t_{k+1}, t_k}^+ is almost surely symplectic, that is,*

$$(F_{t_{k+1}, t_k}^+)^* \Omega_{T^*Q} = \Omega_{T^*Q}, \quad (3.9)$$

where $\Omega_{T^*Q} = \sum_{i=1}^N dq^i \wedge dp^i$ is the canonical symplectic form on T^*Q .

Proof. The proof follows immediately by observing that (see [25])

$$0 = ddH^+(q_k, p_{k+1}) = \sum_{i=1}^N dq_{k+1}^i \wedge dp_{k+1}^i - \sum_{i=1}^N dq_k^i \wedge dp_k^i = (F_{t_{k+1}, t_k}^+)^* \Omega_{T^*Q} - \Omega_{T^*Q}, \quad (3.10)$$

where d in the above formula denotes the differential operator with respect to the variables q and p and is understood in the mean-square sense. The result holds almost surely, because equation (3.7) holds almost surely. \square

The discrete counterpart of stochastic Noether's theorem readily generalizes from the corresponding theorem in the deterministic case.

Theorem 3.2 (Discrete stochastic Noether's theorem). *Let Φ^{T^*Q} be the cotangent lift action of the action Φ of the Lie group G on the configuration space Q . If the generalized discrete stochastic Lagrangian $R_d(q_k, p_{k+1}) = p_{k+1}q_{k+1} - H_d^+(q_k, p_{k+1})$, where $q_{k+1} = D_2H_d^+(q_k, p_{k+1})$, is invariant under the action of G , that is,*

$$R_d(\Phi_g(q_k), \pi_{Q^*} \circ \Phi_g^{T^*Q}(q_{k+1}, p_{k+1})) = R_d(q_k, p_{k+1}), \quad \text{for all } g \in G, \quad (3.11)$$

where $\pi_{Q^*} : Q \times Q^* \rightarrow Q^*$ is the projection onto Q^* , then the cotangent lift momentum map J associated with Φ^{T^*Q} is almost surely preserved, i.e., a.s. $J(q_{k+1}, p_{k+1}) = J(q_k, p_k)$.

Proof. See the proof of Theorem 4 in [25]. In our case the result holds almost surely, because equation (3.7) holds almost surely. \square

For applications, it is useful to know under what conditions the discrete Hamiltonian (3.6) inherits the symmetry properties of the Hamiltonians H and h . Not unexpectedly, this depends on the behavior of the interpolating polynomial (3.3) under the group action. We say that the polynomial $q_d(t; q^\mu)$ is equivariant with respect to G if for all $g \in G$ we have

$$\Phi_g^{TQ}(q_d(t; q^\mu), \dot{q}_d(t; q^\mu)) = (q_d(t; \Phi_g(q^\mu)), \dot{q}_d(t; \Phi_g(q^\mu))). \quad (3.12)$$

Theorem 3.3. *Suppose that the Hamiltonians $H : T^*Q \rightarrow \mathbb{R}$ and $h : T^*Q \rightarrow \mathbb{R}$ are invariant with respect to the cotangent lift action $\Phi^{T^*Q} : G \times T^*Q \rightarrow T^*Q$ of the Lie group G , that is,*

$$H \circ \Phi_g^{T^*Q} = H, \quad h \circ \Phi_g^{T^*Q} = h, \quad (3.13)$$

for all $g \in G$, and suppose the interpolating polynomial $q_d(t; q^\mu)$ is equivariant with respect to G . Then the generalized discrete stochastic Lagrangian $R_d(q_k, p_{k+1}) = p_{k+1}q_{k+1} - H_d^+(q_k, p_{k+1})$ corresponding to the discrete Hamiltonian (3.6), where $q_{k+1} = D_2H_d^+(q_k, p_{k+1})$, is invariant with respect to the action of G .

Proof. The proof is similar to the proofs of Lemma 3 in [25] and Theorem 3 in [38]. Let us, however, carefully examine the actions of G on Q , TQ , and T^*Q . Let $q_k \in Q$ and $p_{k+1} \in Q^*$, and let $q_{k+1} = D_2H_d^+(q_k, p_{k+1})$. First, note that for the stochastic discrete Hamiltonian (3.6), we have

$$R(q_k, p_{k+1}) = \underset{\substack{q^1, \dots, q^s \in Q \\ P_1, \dots, P_r \in Q^* \\ q^0 = q_k}}{\text{ext}} \left\{ \Delta t \sum_{i=1}^r \alpha_i \left[P_i \dot{q}_d(t_k + c_i \Delta t; q^\mu) - H(q_d(t_k + c_i \Delta t; q^\mu), P_i) \right] \right. \\ \left. - \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t; q^\mu), P_i) \right\}, \quad (3.14)$$

where we used (3.8e). Consider $\tilde{q}_k = \Phi_g(q_k)$ and $(\tilde{q}_{k+1}, \tilde{p}_{k+1}) = \Phi_g^{T^*Q}(q_{k+1}, p_{k+1})$ for $g \in G$, and calculate (3.14) for the transformed values \tilde{q}_k and \tilde{p}_{k+1} :

$$R(\tilde{q}_k, \tilde{p}_{k+1}) = \underset{\substack{\tilde{q}^1, \dots, \tilde{q}^s \in Q \\ \tilde{P}_1, \dots, \tilde{P}_r \in Q^* \\ \tilde{q}^0 = \tilde{q}_k}}{\text{ext}} \left\{ \Delta t \sum_{i=1}^r \alpha_i \left[\tilde{P}_i \dot{q}_d(t_k + c_i \Delta t; \tilde{q}^\mu) - H(q_d(t_k + c_i \Delta t; \tilde{q}^\mu), \tilde{P}_i) \right] \right. \\ \left. - \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t; \tilde{q}^\mu), \tilde{P}_i) \right\}. \quad (3.15)$$

Let us perform a change of variables with respect to which we extremize. First, define $q^\mu = \Phi_{g^{-1}}(\tilde{q}^\mu)$, so that $\tilde{q}^\mu = \Phi_g(q^\mu)$ for $\mu = 0, \dots, s$. From (3.12) we have $q_d(t_k + c_i \Delta t; \tilde{q}^\mu) = \Phi_g(q_d(t_k + c_i \Delta t; q^\mu))$, which we use to define P_i by $(q_d(t_k + c_i \Delta t; \tilde{q}^\mu), \tilde{P}_i) = \Phi_g^{T^*Q}(q_d(t_k + c_i \Delta t; q^\mu), P_i)$ for $i = 1, \dots, r$. Since Φ_g and $\Phi_g^{T^*Q}$ are bijective, extremization with respect to q^μ and P_i is equivalent to extremization with respect to \tilde{q}^μ and \tilde{P}_i , and $\tilde{q}^0 = \tilde{q}_k$ implies $q^0 = q_k$. Moreover, from (3.12) and (2.18) we have that $\tilde{P}_i \dot{q}_d(t_k + c_i \Delta t; \tilde{q}^\mu) = P_i \dot{q}_d(t_k + c_i \Delta t; q^\mu)$. Finally, the invariance of the Hamiltonians implies

$$R(\tilde{q}_k, \tilde{p}_{k+1}) = \underset{\substack{q^1, \dots, q^s \in Q \\ P_1, \dots, P_r \in Q^* \\ q^0 = q_k}}{\text{ext}} \left\{ \Delta t \sum_{i=1}^r \alpha_i \left[P_i \dot{q}_d(t_k + c_i \Delta t; q^\mu) - H(q_d(t_k + c_i \Delta t; q^\mu), P_i) \right] \right. \\ \left. - \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t; q^\mu), P_i) \right\} = R(q_k, p_{k+1}), \quad (3.16)$$

which completes the proof. □

Remark: One can easily verify that the interpolating polynomial (3.3) is in particular equivariant with respect to linear actions (translations, rotations, etc.), therefore the stochastic Galerkin variational integrator (3.8) preserves quadratic momentum maps (such as linear and angular momentum) related to linear symmetries of the Hamiltonians H and h .

3.3 Stochastic symplectic partitioned Runge-Kutta methods

A general class of stochastic Runge-Kutta methods for Stratonovich ordinary differential equations was introduced and analyzed in [9], [10], and [11]. These ideas were later used by Ma & Ding & Ding [27] and Ma & Ding [28] to construct symplectic Runge-Kutta methods for stochastic Hamiltonian

systems. An s -stage stochastic symplectic partitioned Runge-Kutta method for (1.1) is defined in [28] by the following system:

$$Q_i = q_k + \Delta t \sum_{j=1}^s a_{ij} \frac{\partial H}{\partial p}(Q_j, P_j) + \Delta W \sum_{j=1}^s b_{ij} \frac{\partial h}{\partial p}(Q_j, P_j), \quad i = 1, \dots, s, \quad (3.17a)$$

$$P_i = p_k - \Delta t \sum_{j=1}^s \bar{a}_{ij} \frac{\partial H}{\partial q}(Q_j, P_j) - \Delta W \sum_{j=1}^s \bar{b}_{ij} \frac{\partial h}{\partial q}(Q_j, P_j), \quad i = 1, \dots, s, \quad (3.17b)$$

$$q_{k+1} = q_k + \Delta t \sum_{i=1}^s \alpha_i \frac{\partial H}{\partial p}(Q_i, P_i) + \Delta W \sum_{i=1}^s \beta_i \frac{\partial h}{\partial p}(Q_i, P_i), \quad (3.17c)$$

$$p_{k+1} = p_k - \Delta t \sum_{i=1}^s \alpha_i \frac{\partial H}{\partial q}(Q_i, P_i) - \Delta W \sum_{i=1}^s \beta_i \frac{\partial h}{\partial q}(Q_i, P_i), \quad (3.17d)$$

where Q_i and P_i for $i = 1, \dots, s$ are the position and momentum internal stages, respectively, and the coefficients of the method a_{ij} , \bar{a}_{ij} , b_{ij} , \bar{b}_{ij} , α_i , β_i satisfy the symplectic conditions

$$\alpha_i \bar{a}_{ij} + \alpha_j a_{ji} = \alpha_i \alpha_j, \quad (3.18a)$$

$$\beta_i \bar{a}_{ij} + \alpha_j b_{ji} = \beta_i \alpha_j, \quad (3.18b)$$

$$\alpha_i \bar{b}_{ij} + \beta_j a_{ji} = \alpha_i \beta_j, \quad (3.18c)$$

$$\beta_i \bar{b}_{ij} + \beta_j b_{ji} = \beta_i \beta_j, \quad (3.18d)$$

for $i, j = 1, \dots, s$. We now prove that in the special case when $r = s$, the stochastic Galerkin variational integrator (3.8) is equivalent to the stochastic symplectic partitioned Runge-Kutta method (3.17).

Theorem 3.4. *Let $r = s$ and let $\bar{l}_{i,s-1}(\tau)$ for $i = 1, \dots, s$ denote the Lagrange polynomials of degree $s-1$ associated with the quadrature points $0 \leq c_1 < \dots < c_s \leq 1$. Moreover, let the weights α_i be given by*

$$\alpha_i = \int_0^1 \bar{l}_{i,s-1}(\tau) d\tau, \quad (3.19)$$

and assume $\alpha_i \neq 0$ for $i = 1, \dots, s$. Then the stochastic Galerkin Hamiltonian variational integrator (3.8) is equivalent to the stochastic partitioned Runge-Kutta method (3.17) with the coefficients

$$a_{ij} = \int_0^{c_i} \bar{l}_{j,s-1}(\tau) d\tau, \quad (3.20a)$$

$$\bar{a}_{ij} = \frac{\alpha_j(\alpha_i - a_{ji})}{\alpha_i}, \quad (3.20b)$$

$$b_{ij} = \frac{\beta_j a_{ij}}{\alpha_j}, \quad (3.20c)$$

$$\bar{b}_{ij} = \frac{\beta_j(\alpha_i - a_{ji})}{\alpha_i}, \quad (3.20d)$$

for $i, j = 1, \dots, s$.

Proof. The proof follows the main steps of the proof of Theorem 2.6.2 in [31]. The time derivative \dot{q}_d is a polynomial of degree $s-1$. Therefore, it can be uniquely expressed in terms of the Lagrange polynomials $\bar{l}_{j,s-1}(\tau)$ as

$$\dot{q}_d(t_k + \tau \Delta t) = \sum_{j=1}^s \dot{q}_d(t_k + c_j \Delta t) \bar{l}_{j,s-1}(\tau). \quad (3.21)$$

Upon integrating with respect to time, we find

$$q_d(t_k + \tau \Delta t) = q_k + \Delta t \sum_{j=1}^s \dot{q}_d(t_k + c_j \Delta t) \int_0^\tau \bar{l}_{j,s-1}(\xi) d\xi, \quad (3.22)$$

where we have used $q^0 = q_k$. For $\tau = 1$ this gives

$$q_{k+1} = q_k + \Delta t \sum_{j=1}^s \alpha_j \dot{q}_d(t_k + c_j \Delta t), \quad (3.23)$$

where we have used $q^s = q_{k+1}$ and (3.19). Define the internal stages $Q_j \equiv q_d(t_k + c_j \Delta t)$. Then, upon using (3.8d), equation (3.23) becomes (3.17c). For $\tau = c_i$ equation (3.22) gives

$$Q_i = q_k + \Delta t \sum_{j=1}^s a_{ij} \dot{q}_d(t_k + c_j \Delta t), \quad (3.24)$$

where a_{ij} is defined by (3.20a). Upon substituting (3.8d), equation (3.24) becomes (3.17a), where b_{ij} is defined by (3.20c). Next, sum equations (3.8a), (3.8b), and (3.8c). Noting that $\sum_{\mu=0}^s l_{\mu,s}(\tau) = 1$, this gives equation (3.17d). Finally, we note that for each $i = 1, \dots, s$ we have a unique decomposition

$$\int_0^\tau \bar{l}_{i,s-1}(\xi) d\xi - \alpha_i = \sum_{\mu=0}^s m_{i\mu} l_{\mu,s}(\tau), \quad (3.25)$$

since the left-hand side is a polynomial of degree s , and therefore it can be uniquely expressed as a linear combination of the Lagrange polynomials $l_{\mu,s}(\tau)$ with the coefficients $m_{i\mu}$. Evaluating this identity at $\tau = d_0 = 0$, $\tau = d_s = 1$, and differentiating it with respect to τ yield the following three equations, respectively,

$$\begin{aligned} -\alpha_i &= \sum_{\mu=0}^s m_{i\mu} l_{\mu,s}(0) = m_{i0}, \\ 0 &= \sum_{\mu=0}^s m_{i\mu} l_{\mu,s}(1) = m_{is}, \\ \bar{l}_{i,s-1}(\tau) &= \sum_{\mu=0}^s m_{i\mu} \dot{l}_{\mu,s}(\tau). \end{aligned} \quad (3.26)$$

We form a linear combination of equations (3.8a), (3.8b), and (3.8c) with the coefficients m_{j0} , $m_{j\mu}$, and m_{js} , respectively. By using the identities (3.26) and rearranging the terms, we obtain (3.17d), where \bar{a}_{ij} and \bar{b}_{ij} are defined by (3.20b) and (3.20d), respectively. One can easily verify that the coefficients (3.20) satisfy the conditions (3.18). □

3.4 Examples

In the construction of the integrator (3.8) we may choose the degree s of the approximating polynomials and the quadrature rules $(\alpha_i, c_i)_{i=1}^r$ and $(\beta_i, c_i)_{i=1}^r$. In the deterministic case, the higher the degree of the polynomials and the higher the order of the quadrature rule, then the higher the order of convergence of the resulting integrator (see [38]). In our case, however, as explained in Section 3.1, we cannot in general achieve strong order of convergence higher than 1.0, because we only used ΔW in (3.5). Since the system (3.8) requires solving $(s+r)N$ equations for $(s+r)N$ variables, from the computational point of view it makes sense to only consider methods with low values of s and r . In this work we focus on the following classical numerical integration formulas (see [14], [15], [16]):

- Gauss-Legendre quadratures (Gau): midpoint rule ($r = 1$), etc.
- Lobatto quadratures (Lob): trapezoidal rule ($r = 2$), Simpson’s rule ($r = 3$), etc.
- Open trapezoidal rule (Otr; $r = 2$)
- Milne’s rule (Mil; $r = 3$)
- Rectangle rule (Rec; $r = 1$)—only in the case when $h = h(q)$.

In [38] notation $PsNrQu$ was proposed to denote a Galerkin variational integrator based on polynomials of degree s and a quadrature rule of order u with r quadrature points. We adopt a similar notation, keeping in mind that u denotes the *classical* order of the used quadrature rule—when the rule is applied to a stochastic integral, as in (3.5), its classical order is not attained in general. We also use a three-letter code to identify which integration formula is used. For example, $P2N2Q4Gau$ denotes the integrator defined by (3.8) using polynomials of degree 2 and the Gauss-Legendre quadrature formula of classical order 4 with 2 quadrature points for both the Lebesgue and Stratonovich integrals in (3.5). If two different quadrature rules are used, we first write the rule applied to the Lebesgue integral, followed by the rule applied to the Stratonovich integral, e.g., $P1N1Q2GauN2Q2Lob$. Below we give several examples of integrators obtained by using polynomials of degree $s = 1, 2$ and the quadrature rules listed above.

3.4.1 General stochastic Hamiltonian

For a general stochastic potential $h = h(q, p)$, equation (3.8d), which represents the discretization of the Legendre transform, needs to contain both $\partial H/\partial p$ and $\partial h/\partial p$ terms to correctly approximate the continuous system. Therefore, we only consider methods with $\alpha_i = \beta_i \neq 0$ for all $i = 1, \dots, r$. A few examples of interest are listed below.

1. $P1N1Q2Gau$ (*Stochastic midpoint method*)

Using the midpoint rule ($r = 1$, $c_1 = 1/2$, $\alpha_1 = \beta_1 = 1$) together with polynomials of degree $s = 1$ gives a stochastic Runge-Kutta method (3.17) with $a_{11} = \bar{a}_{11} = b_{11} = \bar{b}_{11} = 1/2$. Noting that $Q_1 = (q_k + q_{k+1})/2$ and $P_1 = (p_k + p_{k+1})/2$, this method can be written as

$$\begin{aligned}
q_{k+1} &= q_k + \frac{\partial H}{\partial p} \left(\frac{q_k + q_{k+1}}{2}, \frac{p_k + p_{k+1}}{2} \right) \Delta t + \frac{\partial h}{\partial p} \left(\frac{q_k + q_{k+1}}{2}, \frac{p_k + p_{k+1}}{2} \right) \Delta W, \\
p_{k+1} &= p_k - \frac{\partial H}{\partial q} \left(\frac{q_k + q_{k+1}}{2}, \frac{p_k + p_{k+1}}{2} \right) \Delta t - \frac{\partial h}{\partial q} \left(\frac{q_k + q_{k+1}}{2}, \frac{p_k + p_{k+1}}{2} \right) \Delta W.
\end{aligned} \tag{3.27}$$

The stochastic midpoint method was considered in [28] and [35].

2. *P2N2Q2Lob (Stochastic Störmer-Verlet method)*

If the trapezoidal rule ($r = 2$, $c_1 = 0$, $c_2 = 1$, $\alpha_1 = \beta_1 = 1/2$, $\alpha_2 = \beta_2 = 1/2$) is used with polynomials of degree $s = 2$, we obtain another partitioned Runge-Kutta method (3.17) with $a_{11} = a_{12} = 0$, $a_{21} = a_{22} = 1/2$, $\bar{a}_{11} = \bar{a}_{21} = 1/2$, $\bar{a}_{12} = \bar{a}_{22} = 0$, $(b_{ij}) = (a_{ij})$, $(\bar{b}_{ij}) = (\bar{a}_{ij})$. Noting that $Q_1 = q_k$, $Q_2 = q_{k+1}$, and $P_1 = P_2$, this method can be more efficiently written as

$$\begin{aligned}
P_1 &= p_k - \frac{1}{2} \frac{\partial H}{\partial q} (q_k, P_1) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q} (q_k, P_1) \Delta W, \\
q_{k+1} &= q_k + \frac{1}{2} \frac{\partial H}{\partial p} (q_k, P_1) \Delta t + \frac{1}{2} \frac{\partial H}{\partial p} (q_{k+1}, P_1) \Delta t + \frac{1}{2} \frac{\partial h}{\partial p} (q_k, P_1) \Delta W + \frac{1}{2} \frac{\partial h}{\partial p} (q_{k+1}, P_1) \Delta W, \\
p_{k+1} &= P_1 - \frac{1}{2} \frac{\partial H}{\partial q} (q_{k+1}, P_1) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q} (q_{k+1}, P_1) \Delta W.
\end{aligned} \tag{3.28}$$

This method is a stochastic generalization of the Störmer-Verlet method (see [14]) and was considered in [28]. This method is particularly efficient, because the first equation can be solved separately from the second one, and the last equation is an explicit update.

3. *P1N2Q2Lob (Stochastic trapezoidal method)*

This integrator is based on polynomials of degree $s = 1$ with control points $d_0 = 0$, $d_1 = 1$, and the trapezoidal rule. Equations (3.8) take the form

$$\begin{aligned}
p_k &= \frac{1}{2} (P_1 + P_2) + \frac{1}{2} \frac{\partial H}{\partial q} (q_k, P_1) \Delta t + \frac{1}{2} \frac{\partial h}{\partial q} (q_k, P_1) \Delta W, \\
p_{k+1} &= \frac{1}{2} (P_1 + P_2) - \frac{1}{2} \frac{\partial H}{\partial q} (q_{k+1}, P_2) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q} (q_{k+1}, P_2) \Delta W, \\
q_{k+1} &= q_k + \frac{\partial H}{\partial p} (q_k, P_1) \Delta t + \frac{\partial h}{\partial p} (q_k, P_1) \Delta W, \\
q_{k+1} &= q_k + \frac{\partial H}{\partial p} (q_{k+1}, P_2) \Delta t + \frac{\partial h}{\partial p} (q_{k+1}, P_2) \Delta W.
\end{aligned} \tag{3.29}$$

This integrator is a stochastic generalization of the trapezoidal method for deterministic systems (see [31]). One may easily verify that if the Hamiltonians are separable, that is, $H(q, p) = T_0(p) + U_0(q)$ and $h(q, p) = T_1(p) + U_1(q)$, then $P_1 = P_2$ and (3.29) is equivalent to the Störmer-Verlet method (3.28).

4. *P1N3Q4Lob*

If we use Simpson's rule ($r = 3$, $c_1 = 0$, $c_2 = 1/2$, $c_3 = 1$, $\alpha_1 = 1/6$, $\alpha_2 = 2/3$, $\alpha_3 = 1/6$, $\beta_i = \alpha_i$), the resulting integrator (3.8) requires solving simultaneously $4N$ nonlinear equations, so it is computationally expensive in general. However, if the Hamiltonians H and h are separable, then (3.8d) implies $P_1 = P_2 = P_3$, and the integrator can be rewritten as

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial T_0}{\partial p}(P_1)\Delta t + \frac{\partial T_1}{\partial p}(P_1)\Delta W, \\ p_{k+1} &= P_1 - \frac{1}{3} \frac{\partial U_0}{\partial q}\left(\frac{q_k + q_{k+1}}{2}\right)\Delta t - \frac{1}{6} \frac{\partial U_0}{\partial q}(q_{k+1})\Delta t - \frac{1}{3} \frac{\partial U_1}{\partial q}\left(\frac{q_k + q_{k+1}}{2}\right)\Delta W - \frac{1}{6} \frac{\partial U_1}{\partial q}(q_{k+1})\Delta W, \end{aligned} \quad (3.30)$$

where

$$P_1 = p_k - \frac{1}{6} \frac{\partial U_0}{\partial q}(q_k)\Delta t - \frac{1}{3} \frac{\partial U_0}{\partial q}\left(\frac{q_k + q_{k+1}}{2}\right)\Delta t - \frac{1}{6} \frac{\partial U_1}{\partial q}(q_k)\Delta W - \frac{1}{3} \frac{\partial U_1}{\partial q}\left(\frac{q_k + q_{k+1}}{2}\right)\Delta W, \quad (3.31)$$

and $H(q, p) = T_0(p) + U_0(q)$ and $h(q, p) = T_1(p) + U_1(q)$. In this case only the first equation needs to be solved for q_{k+1} , and then the second equation is an explicit update.

5. *P1N2Q2Otr*

Like the method (3.29), this integrator is based on polynomials of degree $s = 1$ with control points $d_0 = 0$, $d_1 = 1$, but uses the open trapezoidal rule ($r = 2$, $c_1 = 1/3$, $c_2 = 2/3$, $\alpha_1 = 1/2$, $\alpha_2 = 1/2$, $\beta_i = \alpha_i$). Equations (3.8) take the form

$$\begin{aligned} p_k &= \frac{1}{2}(P_1 + P_2) + \frac{1}{3} \frac{\partial H}{\partial q}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta t + \frac{1}{6} \frac{\partial H}{\partial q}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta t \\ &\quad + \frac{1}{3} \frac{\partial h}{\partial q}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta W + \frac{1}{6} \frac{\partial h}{\partial q}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta W, \\ p_{k+1} &= \frac{1}{2}(P_1 + P_2) - \frac{1}{6} \frac{\partial H}{\partial q}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta t - \frac{1}{3} \frac{\partial H}{\partial q}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta t \\ &\quad - \frac{1}{6} \frac{\partial h}{\partial q}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta W - \frac{1}{3} \frac{\partial h}{\partial q}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta W, \\ q_{k+1} &= q_k + \frac{\partial H}{\partial p}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta t + \frac{\partial h}{\partial p}\left(\frac{q_{k+1} + 2q_k}{3}, P_1\right)\Delta W, \\ q_{k+1} &= q_k + \frac{\partial H}{\partial p}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta t + \frac{\partial h}{\partial p}\left(\frac{2q_{k+1} + q_k}{3}, P_2\right)\Delta W. \end{aligned} \quad (3.32)$$

In general one has to solve the first, third, and fourth equation simultaneously ($3N$ equations for $3N$ variables). In case of separable Hamiltonians we have $P_1 = P_2$ and one only needs to solve N nonlinear equations: P_1 can be explicitly calculated from the third equation and substituted into the first one, and the resulting nonlinear equation then has to be solved for q_{k+1} .

6. $P2N2Q2Otr$

If the open trapezoidal rule is used with polynomials of degree $s = 2$, we obtain yet another partitioned Runge-Kutta method (3.17) with $a_{11} = \bar{a}_{22} = 1/2$, $a_{12} = \bar{a}_{12} = -1/6$, $a_{21} = \bar{a}_{21} = 2/3$, $a_{22} = \bar{a}_{11} = 0$, $(b_{ij}) = (a_{ij})$, $(\bar{b}_{ij}) = (\bar{a}_{ij})$. Inspecting equations (3.17) we see that, for example, Q_2 is explicitly given in terms of Q_1 and P_1 , therefore one only needs to solve $3N$ equations for the $3N$ variables Q_1 , P_1 , P_2 , and the remaining equations are explicit updates. This method further simplifies for separable Hamiltonians H and h .

7. $P1N3Q4Mil$

A method similar to 4.3 is obtained if we use Milne's rule ($r = 3$, $c_1 = 1/4$, $c_2 = 1/2$, $c_3 = 3/4$, $\alpha_1 = 2/3$, $\alpha_2 = -1/3$, $\alpha_3 = 2/3$, $\beta_i = \alpha_i$) instead of Simpson's rule. The resulting integrator is also computationally expensive in general, but if the Hamiltonians H and h are separable, then (3.8d) implies $P_1 = P_2 = P_3$, and the integrator can be rewritten as

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial T_0}{\partial p}(P_1)\Delta t + \frac{\partial T_1}{\partial p}(P_1)\Delta W, \\ p_{k+1} &= p_k - \frac{2}{3} \frac{\partial U_0}{\partial q} \left(\frac{3q_k + q_{k+1}}{4} \right) \Delta t + \frac{1}{3} \frac{\partial U_0}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta t - \frac{2}{3} \frac{\partial U_0}{\partial q} \left(\frac{q_k + 3q_{k+1}}{4} \right) \Delta t \\ &\quad - \frac{2}{3} \frac{\partial U_1}{\partial q} \left(\frac{3q_k + q_{k+1}}{4} \right) \Delta W + \frac{1}{3} \frac{\partial U_1}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta W - \frac{2}{3} \frac{\partial U_1}{\partial q} \left(\frac{q_k + 3q_{k+1}}{4} \right) \Delta W, \end{aligned} \quad (3.33)$$

where

$$\begin{aligned} P_1 &= p_k - \frac{1}{2} \frac{\partial U_0}{\partial q} \left(\frac{3q_k + q_{k+1}}{4} \right) \Delta t + \frac{1}{6} \frac{\partial U_0}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta t - \frac{1}{6} \frac{\partial U_0}{\partial q} \left(\frac{q_k + 3q_{k+1}}{4} \right) \Delta t \\ &\quad - \frac{1}{2} \frac{\partial U_1}{\partial q} \left(\frac{3q_k + q_{k+1}}{4} \right) \Delta W + \frac{1}{6} \frac{\partial U_1}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta W - \frac{1}{6} \frac{\partial U_1}{\partial q} \left(\frac{q_k + 3q_{k+1}}{4} \right) \Delta W, \end{aligned} \quad (3.34)$$

and $H(q, p) = T_0(p) + U_0(q)$ and $h(q, p) = T_1(p) + U_1(q)$. In this case only the first equation needs to be solved for q_{k+1} , and then the second equation is an explicit update.

3.4.2 Stochastic Hamiltonian independent of momentum

In case the stochastic Hamiltonian $h = h(q)$ is independent of the momentum variable p , the term $\partial h/\partial p$ does not enter equation (3.8d), and therefore we can allow a choice of quadrature rules such that $\alpha_i = 0$ or $\beta_i = 0$ for some i . If $\alpha_i = 0$, however, the system (3.8) becomes underdetermined, but at the same time the corresponding P_i does not enter any of the remaining equations, therefore we can simply ignore it. To simplify the notation, let $i_1 < \dots < i_{\bar{r}}$ be the set of indices such that $\alpha_{i_m} \neq 0$, and denote $\bar{\alpha}_m \equiv \alpha_{i_m}$, $\bar{c}_m \equiv c_{i_m}$ for $m = 1, \dots, \bar{r}$. Similarly, let $j_1 < \dots < j_{\tilde{r}}$ be the set of indices such that $\beta_{j_m} \neq 0$, and denote $\tilde{\beta}_m \equiv \beta_{j_m}$, $\tilde{c}_m \equiv c_{j_m}$ for $m = 1, \dots, \tilde{r}$. In (3.8) leave out the terms and equations corresponding to $\alpha_i = 0$ or $\beta_i = 0$, and replace α_i , β_i , c_i and r by $\bar{\alpha}_i$, $\tilde{\beta}_i$, \bar{c}_i , \tilde{c}_i , \bar{r} and \tilde{r} , accordingly. In other words, this is equivalent to using the quadrature rules $(\bar{\alpha}_i, \bar{c}_i)_{i=1}^{\bar{r}}$ and $(\tilde{\beta}_i, \tilde{c}_i)_{i=1}^{\tilde{r}}$ in (3.6). We then simultaneously solve (3.8a), (3.8b) and (3.8d) ($(s + \bar{r})N$ equations) for q^1, \dots, q^s and $P_1, \dots, P_{\bar{r}}$ ($(s + \bar{r})N$ unknowns). A few examples of such integrators are listed below.

1. *P1N1Q1Rec (Stochastic symplectic Euler method)*

The rectangle rule ($\bar{r} = 1$, $\bar{c}_1 = 1$, $\bar{\alpha}_1 = 1$) does not yield a convergent numerical scheme in the general case, but when $h = h(q)$, the Itô and Stratonovich interpretations of (1.1) are equivalent, and the rectangle rule can be used to construct efficient integrators. In fact, applying the rectangle rule to both the Lebesgue and Stratonovich integrals and using polynomials of degree $s = 1$ yield a method which can be written as

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial H}{\partial p}(q_{k+1}, p_k) \Delta t, \\ p_{k+1} &= p_k - \frac{\partial H}{\partial q}(q_{k+1}, p_k) \Delta t - \frac{\partial h}{\partial q}(q_{k+1}) \Delta W. \end{aligned} \quad (3.35)$$

This method is a straightforward generalization of the symplectic Euler scheme (see [14], [31]) and is particularly computationally efficient, as only the first equation needs to be solved for q_{k+1} , and then the second equation is an explicit update. Moreover, in case the Hamiltonian H is separable, the method becomes explicit.

2. *P1N1Q1RecN2Q2Lob*

The accuracy of the stochastic symplectic Euler scheme above can be improved by applying the trapezoidal rule to the Stratonovich integral instead of the rectangle rule. The resulting integrator takes the form

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial H}{\partial p}(q_{k+1}, P_1) \Delta t, \\ p_{k+1} &= p_k - \frac{\partial H}{\partial q}(q_{k+1}, P_1) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q}(q_k) \Delta W - \frac{1}{2} \frac{\partial h}{\partial q}(q_{k+1}) \Delta W, \end{aligned} \quad (3.36)$$

where

$$P_1 = p_k - \frac{1}{2} \frac{\partial h}{\partial q}(q_k) \Delta W. \quad (3.37)$$

While having a similar computational cost, this method yields a more accurate solution than (3.35) (see Section 4 for numerical tests).

3. *P1N1Q1RecN1Q2Gau*

Similarly, if we apply the midpoint rule instead of the trapezoidal rule, we obtain the following modification of the stochastic symplectic Euler method:

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial H}{\partial p}(q_{k+1}, P_1) \Delta t, \\ p_{k+1} &= p_k - \frac{\partial H}{\partial q}(q_{k+1}, P_1) \Delta t - \frac{\partial h}{\partial q}\left(\frac{q_k + q_{k+1}}{2}\right) \Delta W, \end{aligned} \quad (3.38)$$

where

$$P_1 = p_k - \frac{1}{2} \frac{\partial h}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta W. \quad (3.39)$$

This method demonstrates a similar performance as (3.36) (see Section 4 for numerical tests).

4. *P2N2Q2LobN1Q1Rec*

A modification of the stochastic Störmer-Verlet method (3.28) is obtained if we use the rectangle rule to approximate the Stratonovich integral:

$$\begin{aligned} P_1 &= p_k - \frac{1}{2} \frac{\partial H}{\partial q}(q_k, P_1) \Delta t, \\ q_{k+1} &= q_k + \frac{1}{2} \frac{\partial H}{\partial p}(q_k, P_1) \Delta t + \frac{1}{2} \frac{\partial H}{\partial p}(q_{k+1}, P_1) \Delta t, \\ p_{k+1} &= P_1 - \frac{1}{2} \frac{\partial H}{\partial q}(q_{k+1}, P_1) \Delta t - \frac{\partial h}{\partial q}(q_{k+1}) \Delta W. \end{aligned} \quad (3.40)$$

This integrator has a similar computational cost as the stochastic Störmer-Verlet method (see Section 4), but it yields a slightly less accurate solution (see Section 4).

5. *P1N1Q2GauN2Q2Lob*

This integrator is a modification of the stochastic midpoint method (3.27). We apply the midpoint rule ($\tilde{r} = 1$, $\tilde{c}_1 = 1/2$, $\tilde{\alpha}_1 = 1$) to the Lebesgue integral in (3.4), and the trapezoidal rule ($\tilde{r} = 2$, $\tilde{c}_1 = 0$, $\tilde{c}_2 = 1$, $\tilde{\beta}_1 = 1/2$, $\tilde{\beta}_2 = 1/2$) to the Stratonovich integral. The resulting numerical scheme can be written as

$$\begin{aligned} q_{k+1} &= q_k + \frac{\partial H}{\partial p} \left(\frac{q_k + q_{k+1}}{2}, P_1 \right) \Delta t, \\ p_{k+1} &= p_k - \frac{\partial H}{\partial q} \left(\frac{q_k + q_{k+1}}{2}, P_1 \right) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q}(q_k) \Delta W - \frac{1}{2} \frac{\partial h}{\partial q}(q_{k+1}) \Delta W, \end{aligned} \quad (3.41)$$

where

$$P_1 = \frac{p_k + p_{k+1}}{2} + \frac{1}{4} \Delta W \left[\frac{\partial h}{\partial q}(q_{k+1}) - \frac{\partial h}{\partial q}(q_k) \right]. \quad (3.42)$$

6. *P1N2Q2LobN1Q2Gau*

If instead we apply the trapezoidal rule to the Lebesgue integral and the midpoint rule to the Stratonovich integral in (3.4), we obtain a modification of the stochastic trapezoidal rule (3.29):

$$\begin{aligned}
p_k &= \frac{1}{2}(P_1 + P_2) + \frac{1}{2} \frac{\partial H}{\partial q}(q_k, P_1) \Delta t + \frac{1}{2} \frac{\partial h}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta W, \\
p_{k+1} &= \frac{1}{2}(P_1 + P_2) - \frac{1}{2} \frac{\partial H}{\partial q}(q_{k+1}, P_2) \Delta t - \frac{1}{2} \frac{\partial h}{\partial q} \left(\frac{q_k + q_{k+1}}{2} \right) \Delta W, \\
q_{k+1} &= q_k + \frac{\partial H}{\partial p}(q_k, P_1) \Delta t, \\
q_{k+1} &= q_k + \frac{\partial H}{\partial p}(q_{k+1}, P_2) \Delta t.
\end{aligned} \tag{3.43}$$

3.5 Convergence

There are two ways of measuring accuracy of a stochastic numerical integrator like (3.8), namely strong convergence and weak convergence (see [21], [33]). Let $\bar{z}(t) = (\bar{q}(t), \bar{p}(t))$ be the exact solution to (1.1) with the initial conditions q_0 and p_0 , and let $z_k = (q_k, p_k)$ denote the numerical solution at time t_k obtained by applying (3.8) iteratively k times with the constant time step Δt . The numerical solution is said to converge strongly with strong global order r if there exist $\delta > 0$ and a constant $C > 0$ such that for all $\Delta t \in (0, \delta)$ we have

$$E(\|z_K - \bar{z}(T)\|) \leq C \Delta t^r, \tag{3.44}$$

where $T = K \Delta t$, as defined before, and E denotes the expected value. On the contrary, the numerical solution is said to converge weakly with weak global order r if there exist $\delta > 0$ and a constant $C > 0$ such that for all $\Delta t \in (0, \delta)$ and f from a sufficiently large class of functions (usually polynomials) we have

$$\|E(f(z_K)) - E(f(\bar{z}(T)))\| \leq C \Delta t^r. \tag{3.45}$$

We are here primarily interested in strong convergence. In principle, in order to determine the strong order of convergence of the Galerkin variational integrator (3.8) we need to calculate the power series expansions of q_{k+1} and p_{k+1} in terms of the powers of Δt and ΔW , and compare them to the Stratonovich-Taylor expansions for the exact solution $\bar{q}(t_k + \Delta t)$ and $\bar{p}(t_k + \Delta t)$ (see [11], [21], [33]). It is quite a tedious task to do in the general case, therefore we only discuss the examples presented in Section 3.4. For instance, in case of the stochastic trapezoidal method (3.29) we plug in series expansions for P_1 , P_2 , q_{k+1} and p_{k+1} , and determine their coefficients by expanding the derivatives of the Hamiltonians into Taylor series around (q_k, p_k) and comparing the terms corresponding to the like powers of Δt and ΔW . We find that

$$\begin{aligned}
q_{k+1} &= q_k + \frac{\partial H}{\partial p} \Delta t + \frac{\partial h}{\partial p} \Delta W + \frac{1}{2} \left(\frac{\partial h}{\partial p} \frac{\partial^2 h}{\partial p \partial q} - \frac{\partial h}{\partial q} \frac{\partial^2 h}{\partial p^2} \right) \Delta W^2 + \dots, \\
p_{k+1} &= p_k - \frac{\partial H}{\partial q} \Delta t - \frac{\partial h}{\partial q} \Delta W - \frac{1}{2} \left(\frac{\partial h}{\partial p} \frac{\partial^2 h}{\partial q^2} - \frac{\partial h}{\partial q} \frac{\partial^2 h}{\partial p \partial q} \right) \Delta W^2 + \dots,
\end{aligned} \tag{3.46}$$

where the derivatives of the Hamiltonians are evaluated at (q_k, p_k) . Let $\bar{q}(t; q_k, p_k)$ and $\bar{p}(t; q_k, p_k)$ denote the exact solution of (1.1) such that $\bar{q}(t_k; q_k, p_k) = q_k$ and $\bar{p}(t_k; q_k, p_k) = p_k$. Using (1.1) we

calculate the Stratonovich-Taylor expansions for $\bar{q}(t_{k+1}; q_k, p_k)$ and $\bar{p}(t_{k+1}; q_k, p_k)$, and comparing them to (3.46) we find that

$$\begin{aligned} E(q_{k+1} - \bar{q}(t_{k+1}; q_k, p_k)) &= O(\Delta t^2), & \sqrt{E(\|q_{k+1} - \bar{q}(t_{k+1}; q_k, p_k)\|^2)} &= O(\Delta t^{\frac{3}{2}}), \\ E(p_{k+1} - \bar{p}(t_{k+1}; q_k, p_k)) &= O(\Delta t^2), & \sqrt{E(\|p_{k+1} - \bar{p}(t_{k+1}; q_k, p_k)\|^2)} &= O(\Delta t^{\frac{3}{2}}). \end{aligned} \quad (3.47)$$

Using Theorem 1.1 from [33], we conclude that the stochastic trapezoidal method (3.29) has strong order of convergence $r = 1$. In a similar fashion we prove that all methods presented in Section 3.4 are strongly convergent with strong order 1. We further verify these results numerically in Section 4.1.

3.6 Methods of order 3/2

In order to construct stochastic Galerkin variational integrators of higher order one needs to include higher order terms in the discretization of the Stratonovich integral in (3.5). For example, a method of strong order 3/2 must include terms involving $\Delta Z = \int_{t_k}^{t_{k+1}} \int_{t_k}^t dW(\xi) dt$ (see [11], [34], [35]). Inspired by the theory presented in [11], we can add extra terms to the discrete Hamiltonian (3.6) and write it as

$$\begin{aligned} H_d^+(q_k, p_{k+1}) &= \underset{\substack{q^1, \dots, q^s \in Q \\ P_1, \dots, P_r \in Q^* \\ q^0 = q_k}}{\text{ext}} \left\{ p_{k+1} q^s - \Delta t \sum_{i=1}^r \alpha_i \left[P_i \dot{q}_d(t_k + c_i \Delta t) - H(q_d(t_k + c_i \Delta t), P_i) \right] \right. \\ &\quad \left. + \Delta W \sum_{i=1}^r \beta_i h(q_d(t_k + c_i \Delta t), P_i) + \frac{\Delta Z}{\Delta t} \sum_{i=1}^r \gamma_i h(q_d(t_k + c_i \Delta t), P_i) \right\}. \end{aligned} \quad (3.48)$$

The random variables ΔW and ΔZ have a Gaussian joint distribution (see [21], [35]), and at each time step they can be simulated by two independent $\mathcal{N}(0, 1)$ -distributed random variables χ and η as

$$\Delta W = \chi \sqrt{\Delta t}, \quad \Delta Z = \frac{1}{2} \Delta t^{\frac{3}{2}} \left(\chi + \frac{1}{\sqrt{3}} \eta \right). \quad (3.49)$$

In order to achieve strong convergence of order 3/2 one needs to determine appropriate values for the parameters s , r , α_i , β_i , γ_i , and c_i . However, we will not attempt to achieve this in the present work. Instead, we will show that some known stochastic symplectic integrators can be derived as stochastic Galerkin variational integrators.

Suppose the Hamiltonian is separable, i.e., $H(q, p) = T(p) + U(q)$, and the stochastic Hamiltonian $h = h(q)$ does not depend on momentum. Consider the discrete Hamiltonian

$$\begin{aligned} H_d^+(q_k, p_{k+1}) &= \underset{\substack{q^1, \dots, q^s \in Q \\ P_1, \dots, P_r \in Q^* \\ q^0 = q_k}}{\text{ext}} \left\{ p_{k+1} q^s - \Delta t \sum_{i=1}^r \left[\bar{\alpha}_i P_i \dot{q}_d(t_k + c_i \Delta t) - \bar{\alpha}_i U(q_d(t_k + c_i \Delta t)) - \alpha_i T(P_i) \right] \right. \\ &\quad \left. + \Delta W \sum_{i=1}^r \bar{\beta}_i h(q_d(t_k + c_i \Delta t)) + \frac{\Delta Z}{\Delta t} \sum_{i=1}^r \bar{\gamma}_i h(q_d(t_k + c_i \Delta t)) \right\}, \end{aligned} \quad (3.50)$$

where different weights $\bar{\alpha}_i$ and α_i were applied to the potential $U(q)$ and kinetic $T(p)$ terms, respectively. Similar to (3.8), the corresponding stochastic variational integrator takes the form

$$\begin{aligned}
-p_k &= \sum_{i=1}^r \bar{\alpha}_i \left[P_i \dot{l}_{0,s}(c_i) - \Delta t \frac{\partial U}{\partial q}(t_k + c_i \Delta t) l_{0,s}(c_i) \right] - \sum_{i=1}^r \left(\bar{\beta}_i \Delta W + \bar{\gamma}_i \frac{\Delta Z}{\Delta t} \right) \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{0,s}(c_i), \\
0 &= \sum_{i=1}^r \bar{\alpha}_i \left[P_i \dot{l}_{\mu,s}(c_i) - \Delta t \frac{\partial U}{\partial q}(t_k + c_i \Delta t) l_{\mu,s}(c_i) \right] - \sum_{i=1}^r \left(\bar{\beta}_i \Delta W + \bar{\gamma}_i \frac{\Delta Z}{\Delta t} \right) \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{\mu,s}(c_i), \\
p_{k+1} &= \sum_{i=1}^r \bar{\alpha}_i \left[P_i \dot{l}_{s,s}(c_i) - \Delta t \frac{\partial U}{\partial q}(t_k + c_i \Delta t) l_{s,s}(c_i) \right] - \sum_{i=1}^r \left(\bar{\beta}_i \Delta W + \bar{\gamma}_i \frac{\Delta Z}{\Delta t} \right) \frac{\partial h}{\partial q}(t_k + c_i \Delta t) l_{s,s}(c_i), \\
\bar{\alpha}_i \dot{q}_d(t_k + c_i \Delta t) &= \alpha_i \frac{\partial T}{\partial p}(P_i), \\
q_{k+1} &= q^s,
\end{aligned} \tag{3.51}$$

where $\mu = 1, \dots, s-1$ in the second equation, and $i = 1, \dots, r$ in the fourth equation. In the special case when $r = s$ and

$$\bar{\alpha}_i = \int_0^1 \bar{l}_{i,s-1}(\tau) d\tau, \quad i = 1, \dots, s, \tag{3.52}$$

we can show, similar to Theorem 3.4, that the stochastic Galerkin variational integrator (3.51) is equivalent to the stochastic partitioned Runge-Kutta method

$$\begin{aligned}
Q_i &= q_k + \Delta t \sum_{j=1}^s a_{ij} \frac{\partial T}{\partial p}(P_j), & i = 1, \dots, s, \\
P_i &= p_k - \Delta t \sum_{j=1}^s \bar{a}_{ij} \frac{\partial U}{\partial q}(Q_j) - \sum_{j=1}^s \left(\bar{b}_{ij} \Delta W + \bar{\lambda}_{ij} \frac{\Delta Z}{\Delta t} \right) \frac{\partial h}{\partial q}(Q_j), & i = 1, \dots, s, \\
q_{k+1} &= q_k + \Delta t \sum_{i=1}^s \alpha_i \frac{\partial T}{\partial p}(P_i), \\
p_{k+1} &= p_k - \Delta t \sum_{i=1}^s \bar{\alpha}_i \frac{\partial U}{\partial q}(Q_i) - \sum_{i=1}^s \left(\bar{\beta}_i \Delta W + \bar{\gamma}_i \frac{\Delta Z}{\Delta t} \right) \frac{\partial h}{\partial q}(Q_i),
\end{aligned} \tag{3.53}$$

with the coefficients

$$\begin{aligned}
a_{ij} &= \frac{\alpha_j}{\bar{\alpha}_j} \int_0^{c_i} \bar{l}_{j,s-1}(\tau) d\tau, & \bar{a}_{ij} &= \frac{\bar{\alpha}_j(\alpha_i - a_{ji})}{\alpha_i}, \\
\bar{b}_{ij} &= \frac{\bar{\beta}_j(\alpha_i - a_{ji})}{\alpha_i}, & \bar{\lambda}_{ij} &= \frac{\bar{\gamma}_j(\alpha_i - a_{ji})}{\alpha_i}, & i, j &= 1, \dots, s,
\end{aligned} \tag{3.54}$$

where we assume $\alpha_i \neq 0$ and $\bar{\alpha}_i \neq 0$ for all i . Partitioned Runge-Kutta methods of type (3.53) were considered in [35]. In particular, it was shown that for $s = 2$ the choice of the coefficients

$$\begin{aligned}
\alpha_1 = 2/3, \quad \alpha_2 = 1/3, \quad \bar{\alpha}_1 = 1/4, \quad \bar{\alpha}_2 = 3/4, \quad \bar{\beta}_1 = -1/2, \quad \bar{\beta}_2 = 3/2, \quad \bar{\gamma}_1 = 3/2, \quad \bar{\gamma}_2 = -3/2, \\
a_{11} = 0, \quad a_{12} = 0, \quad \bar{a}_{11} = 1/4, \quad \bar{a}_{12} = 0, \quad \bar{b}_{11} = -1/2, \quad \bar{b}_{12} = 0, \quad \bar{\lambda}_{11} = 3/2, \quad \bar{\lambda}_{12} = 0, \\
a_{21} = 2/3, \quad a_{22} = 0, \quad \bar{a}_{21} = 1/4, \quad \bar{a}_{22} = 3/4, \quad \bar{b}_{21} = -1/2, \quad \bar{b}_{22} = 3/2, \quad \bar{\lambda}_{21} = 3/2, \quad \bar{\lambda}_{22} = -3/2,
\end{aligned} \tag{3.55}$$

gives a method of strong order 3/2 (see Theorem 4.3 in [35]).

4 Numerical experiments

In this section we present the results of our numerical experiments. We verify numerically the convergence results from Section 3.5 and investigate the conservation properties of our integrators. In particular, we show that our stochastic variational integrators demonstrate superior behavior in long-time simulations compared to some popular general purpose non-symplectic stochastic algorithms.

4.1 Numerical convergence analysis

4.1.1 Kubo oscillator

In order to test the convergence of the numerical algorithms from Section 3.4.1 we performed computations for the Kubo oscillator, which is defined by $H(q, p) = p^2/2 + q^2/2$ and $h(q, p) = \beta(p^2/2 + q^2/2)$, where β is the noise intensity (see [35]). The Kubo oscillator is used in the theory of magnetic resonance and laser physics. The exact solution is given by

$$\bar{q}(t) = p_0 \sin(t + \beta W(t)) + q_0 \cos(t + \beta W(t)), \quad \bar{p}(t) = p_0 \cos(t + \beta W(t)) - q_0 \sin(t + \beta W(t)), \tag{4.1}$$

where q_0 and p_0 are the initial conditions. Simulations with the initial conditions $q_0 = 0$, $p_0 = 1$ and the noise intensity $\beta = 0.1$ were carried out until the time $T = 3.2$ for a number of decreasing time steps Δt . In each case 2000 sample paths were generated. Let $z_{\Delta t}(t) = (q_{\Delta t}(t), p_{\Delta t}(t))$ denote the numerical solution. We used the exact solution (4.1) as a reference for computing the absolute error $E(|z_{\Delta t}(T) - \bar{z}(T)|)$ and the mean error $|E(z_{\Delta t}(T)) - E(\bar{z}(T))|$, where $\bar{z}(t) = (\bar{q}(t), \bar{p}(t))$. The dependence of these errors on the time step Δt is depicted in Figure 4.1. We verified that our algorithms have both strong and weak order of convergence 1.0. The integrators *P1N3Q4Lob*, *P1N3Q4Mil*, *P1N2Q2Lob* (stochastic trapezoidal method), and *P2N2Q2Lob* (stochastic Störmer-Verlet method) turned out to be the most accurate, with the latter two having least computational cost.

4.1.2 Synchrotron oscillations of particles in storage rings

We carried out a similar test for the numerical schemes from Section 3.4.2. We performed computations for the stochastic Hamiltonian system defined by $H(q, p) = p^2/2 - \cos q$ and $h(q) = \beta \sin q$, where β is the noise intensity. Systems of this type are used for modeling synchrotron oscillations of




Figure 4.1: The absolute (*left*) and mean (*right*) errors at the time $T = 3.2$ as a function of the time step Δt for the simulations of the Kubo oscillator with the initial conditions $q_0 = 0$, $p_0 = 1$ and the noise intensity $\beta = 0.1$. In each case 2000 sample paths were generated. The tested integrators proved to be both strongly and weakly convergent of order 1.0.

a particle in a storage ring. Due to fluctuating electromagnetic fields, a particle performs stochastically perturbed oscillations with respect to a reference particle which travels with fixed energy along the design orbit of the accelerator; in this description p corresponds to the energy deviation of the particle from the reference particle, and q measures the longitudinal phase difference of both particles (see [13], [43] for more details). Simulations with the initial conditions $q_0 = 0$, $p_0 = 1$ and the noise intensity $\beta = 0.1$ were carried out until the time $T = 3.2$ for a number of decreasing time steps Δt . In each case 2000 sample paths were generated. The absolute and mean errors were calculated with respect to a high-precision reference solution generated using the order 3/2 strong Taylor scheme (see [21], Chapter 10.4) with a very fine time step $\Delta t = 2 \cdot 10^{-6}$. The dependence of these errors on the time step Δt is depicted in Figure 4.2. We verified that our algorithms have strong order of convergence 1.0. The simulations also suggest that the weak order of convergence of the tested algorithms is 3/2 (perhaps except for $P1N1Q2GauN2Q2Lob$, as in that case the result is not entirely conclusive). It is worth noting that the integrators $P1N1Q1RecN2Q2Lob$ and $P1N1Q1RecN1Q2Gau$ turned out to be as weakly accurate as the stochastic Störmer-Verlet method ($P2N2Q2Lob$), while being less computationally expensive.

4.2 Long-time energy behavior

4.2.1 Kubo oscillator

One can easily check that in the case of the Kubo oscillator the Hamiltonian $H(q, p)$ stays constant for almost all sample paths, i.e., $H(\bar{q}(t), \bar{p}(t)) = H(q_0, p_0)$ almost surely. We used this example to test the performance of the integrators from Section 3.4.1. Simulations with the initial conditions $q_0 = 0$, $p_0 = 1$, the noise intensity $\beta = 0.1$, and the relatively large time step $\Delta t = 0.25$ were carried out until the time $T = 1000$ (approximately 160 periods of the oscillator in the absence




Figure 4.2: The absolute (*left*) and mean (*right*) errors at the time $T = 3.2$ as a function of the time step Δt for the simulations of the synchrotron oscillations of a particle in a storage ring with the initial conditions $q_0 = 0$, $p_0 = 1$ and the noise intensity $\beta = 0.1$. In each case 2000 sample paths were generated. The tested integrators proved to be strongly convergent of order 1.0. The simulations also suggest the tested integrators have weak order of convergence $3/2$ (although it is not conclusive for $P1N1Q2GauN2Q2Lob$). Note that the weak error plots for $P1N1Q1RecN1Q2Gau$, $P1N1Q1RecN2Q2Lob$, $P1N2Q2LobN1Q2Gau$, and $P2N2Q2Lob$ overlap very closely.

of noise) for a single realization of the Wiener process. For comparison, similar simulations were carried out using non-symplectic explicit methods like Milstein’s scheme and the order 3/2 strong Taylor scheme (see [21]). The numerical value of the Hamiltonian $H(q, p)$ as a function of time for each of the integrators is depicted in Figure 4.3. We find that the non-symplectic schemes do not preserve the Hamiltonian well, even if small time steps are used. For example, we find that Milstein’s scheme does not give a satisfactory solution even with $\Delta t = 0.001$, and though the Taylor scheme with $\Delta t = 0.05$ yields a result comparable to the variational integrators, the growing trend of the numerical Hamiltonian is evident. On the other hand, the variational integrators give numerical solutions for which the Hamiltonian oscillates around the true value (one can check via a direct calculation that the stochastic midpoint method (3.27) in this case preserves the Hamiltonian exactly; of course this does not necessarily hold in the general case).

4.2.2 Anharmonic oscillator

In general the Hamiltonian $H(q, p)$ does not stay constant for stochastic Hamilton equations. To determine how well our integrators perform in such cases we considered the anharmonic oscillator defined by $H(q, p) = p^2/2 + \gamma q^4$ and $h(q) = \beta q$, where β is the noise intensity and γ is a parameter. One can calculate the expected value of the Hamiltonian analytically as

$$E\left(H(q(t), p(t))\right) = H(q_0, p_0) + \frac{\beta^2}{2}t, \quad (4.2)$$

that is, the mean value of the Hamiltonian grows linearly in time (see [43]). Simulations with the initial conditions $q_0 = 0$, $p_0 = 1$, the parameter $\gamma = 0.1$, and the noise intensity $\beta = 0.1$ were carried out until the time $T = 784$ (approximately 100 periods of the oscillator in the absence of noise). In each case 10,000 sample paths were generated. The numerical value of the mean Hamiltonian $E(H)$ as a function of time for each of the integrators is depicted in Figure 4.4. We see that the variational integrators accurately capture the linear growth of $E(H)$, whereas the Taylor scheme fails to reproduce that behavior even when a smaller time step is used. It is worth noting that the integrators *P1N1Q1RecN2Q2Lob* and *P1N1Q1RecN1Q2Gau* yield a very accurate solution, while being computationally efficient, as discussed in Section 3.4.2.

5 Summary

In this paper we have presented a general framework for constructing a new class of stochastic symplectic integrators for stochastic Hamiltonian systems. We generalized the approach of Galerkin variational integrators introduced in [25], [31], [38] to the stochastic case, following the ideas underlying the stochastic variational integrators introduced in [7]. The solution of the stochastic Hamiltonian system was approximated by a polynomial of degree s , and the action functional was approximated by a quadrature formula based on r quadrature points. We showed that the resulting integrators are symplectic, preserve integrals of motion related to Lie group symmetries, and include stochastic symplectic Runge-Kutta methods introduced in [27], [28], [35] as a special case when $r = s$. We pointed out several new low-stage stochastic symplectic methods of strong order 1.0, both for the case of a general stochastic Hamiltonian $h = h(q, p)$ and a stochastic Hamiltonian $h = h(q)$ independent of p , and demonstrated their superior long-time stability and energy behavior via numerical experiments.




Figure 4.3: *Top*: The numerical Hamiltonian for the solutions obtained with Milstein's scheme and the order 3/2 strong Taylor scheme. We see that the Hamiltonian tends to blow up despite using small time steps. *Bottom*: The numerical Hamiltonian for the solutions obtained with the integrators derived in Section 3.4.1. For comparison, the solution obtained with the Taylor scheme for $\Delta t = 0.05$ is also included. Note that for clarity the same color code is applied when the plots for some integrators overlap very closely.




Figure 4.4: *Top:* The numerical value of the mean Hamiltonian $E(H)$ is shown for the solutions computed with the order 3/2 strong Taylor scheme using the time step $\Delta t = 0.05$ and the variational integrators derived in Section 3.4.1 using the time step $\Delta t = 0.25$ or $\Delta t = 0.5$. The variational integrators accurately capture the linear growth of $E(H)$, whereas the Taylor scheme fails to reproduce that behavior. *Middle:* The difference between the numerical value of the mean Hamiltonian $E(H)$ and the exact value (4.2) is shown for the integrators derived in Section 3.4.1. *Bottom:* Same for the integrators derived in Section 3.4.2. The integrators $P1N1Q1RecN2Q2Lob$ and $P1N1Q1RecN1Q2Gau$ prove to be particularly accurate, while having a low computational cost.

Our work can be extended in several ways. In Section 3.6 we indicated how higher-order stochastic variational integrators can be constructed and showed that a type of stochastic symplectic partitioned Runge-Kutta methods of strong order $3/2$ considered in [35] can be recast in that formalism. It would be interesting to derive new stochastic integrators of order $3/2$ by choosing appropriate values for the parameters in (3.48) or (3.50). It would also be interesting to apply the Galerkin approach to construct stochastic variational integrators for constrained (see [6]) and dissipative (see [8]) stochastic Hamiltonian systems, and systems defined on Lie groups (see [24]). Another important problem would be stochastic variational error analysis. That is, rather than considering how closely the numerical solution follows the exact trajectory of the system, one could investigate how closely the discrete Hamiltonian matches the exact generating function. In the deterministic setting these two notions of the order of convergence are equivalent (see [31]). It would be instructive to know if a similar result holds in the stochastic case. Finally, it would be interesting to develop higher-order weakly convergent stochastic variational integrators. As mentioned in Section 3.1 and Section 3.6, higher-order strongly convergent methods require inclusion of higher-order multiple Stratonovich integrals, which are cumbersome to simulate in practice. In many cases, though, one is only interested in calculating the probability distribution of the solution rather than precisely approximating each sample path. In such cases weakly convergent methods are much easier to use (see [21], [33]).

Acknowledgments

We would like to thank Nawaf Bou-Rabee, Mickael Chekroun, Dan Crisan, Nader Ganaba, Melvin Leok, Houman Owhadi, and Wei Pan for useful comments and references. This work was partially supported by the European Research Council Advanced Grant 267382 FCCA. Parts of this project were completed while the authors were visiting the Institute for Mathematical Sciences, National University of Singapore in 2016.

References

- [1] C. Anton, J. Deng, and Y. S. Wong. Weak symplectic schemes for stochastic Hamiltonian equations. *Electronic Transactions on Numerical Analysis*, 43:1–20, 2014.
- [2] C. Anton, Y. S. Wong, and J. Deng. On global error of symplectic schemes for stochastic Hamiltonian systems. *International Journal of Numerical Analysis and Modeling, Series B*, 4(1):80–93, 2013.
- [3] C. Anton, Y. S. Wong, and J. Deng. Symplectic numerical schemes for stochastic systems preserving Hamiltonian functions. In I. Dimov, I. Faragó, and L. Vulkov, editors, *Numerical Analysis and Its Applications: 5th International Conference, NAA 2012, Lozenetz, Bulgaria, June 15-20, 2012, Revised Selected Papers*, pages 166–173. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
- [4] L. Arnold. *Stochastic Differential Equations: Theory and Applications*. Dover Books on Mathematics. Dover Publications, 2013.

- [5] J. Bismut. Mécanique aléatoire. In P. Hennequin, editor, *Ecole d'Eté de Probabilités de Saint-Flour X - 1980*, volume 929 of *Lecture Notes in Mathematics*, pages 1–100. Springer Berlin Heidelberg, 1982.
- [6] N. Bou-Rabee and H. Owhadi. Stochastic variational partitioned Runge-Kutta integrators for constrained systems. Unpublished, arXiv:0709.2222, 2007.
- [7] N. Bou-Rabee and H. Owhadi. Stochastic variational integrators. *IMA Journal of Numerical Analysis*, 29(2):421–443, 2009.
- [8] N. Bou-Rabee and H. Owhadi. Long-run accuracy of variational integrators in the stochastic context. *SIAM J. Numer. Anal.*, 48(1):278–297, 2010.
- [9] K. Burrage and P. M. Burrage. High strong order explicit Runge-Kutta methods for stochastic ordinary differential equations. *Appl. Numer. Math.*, 22:81–101, 1996.
- [10] K. Burrage and P. M. Burrage. General order conditions for stochastic Runge-Kutta methods for both commuting and non-commuting stochastic ordinary differential equation systems. *Appl. Numer. Math.*, 28:161–177, 1998.
- [11] K. Burrage and P. M. Burrage. Order Conditions of Stochastic Runge-Kutta Methods by B-Series. *SIAM Journal on Numerical Analysis*, 38(5):1626–1646, 2000.
- [12] J. Deng, C. Anton, and Y. S. Wong. High-order symplectic schemes for stochastic Hamiltonian systems. *Communications in Computational Physics*, 16:169–200, 2014.
- [13] G. Dôme. Theory of RF acceleration. In S. Turner, editor, *Proceedings of CERN Accelerator School, Oxford, September 1985*, volume 1, pages 110–158. CERN European Organization for Nuclear Research, 1987.
- [14] E. Hairer, C. Lubich, and G. Wanner. *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. Springer Series in Computational Mathematics. Springer, New York, 2002.
- [15] E. Hairer, S. Nørsett, and G. Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*, volume 8 of *Springer Series in Computational Mathematics*. Springer, 2nd edition, 1993.
- [16] E. Hairer and G. Wanner. *Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems*, volume 14 of *Springer Series in Computational Mathematics*. Springer, 2nd edition, 1996.
- [17] J. Hall and M. Leok. Spectral variational integrators. *Numer. Math.*, 130(4):681–740, Aug. 2015.
- [18] D. D. Holm, T. Schmah, and C. Stoica. *Geometric Mechanics and Symmetry: From Finite to Infinite Dimensions*. Oxford Texts in Applied and Engineering Mathematics. Oxford University Press, Oxford, 2009.
- [19] D. D. Holm and T. M. Tyrannowski. Variational principles for stochastic soliton dynamics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 472(2187), 2016.

- [20] N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. Kodansha scientific books. North-Holland, 1989.
- [21] P. Kloeden and E. Platen. *Numerical Solution of Stochastic Differential Equations*. Applications of Mathematics : Stochastic Modelling and Applied Probability. Springer, 1995.
- [22] H. Kunita. *Stochastic Flows and Stochastic Differential Equations*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1997.
- [23] J. A. Lázaro-Camí and J. P. Ortega. Stochastic Hamiltonian Dynamical Systems. *Reports on Mathematical Physics*, 61(1):65 – 122, 2008.
- [24] M. Leok and T. Shingel. General techniques for constructing variational integrators. *Frontiers of Mathematics in China*, 7(2):273–303, 2012.
- [25] M. Leok and J. Zhang. Discrete Hamiltonian variational integrators. *IMA Journal of Numerical Analysis*, 31(4):1497–1532, 2011.
- [26] A. Lew, J. E. Marsden, M. Ortiz, and M. West. Asynchronous variational integrators. *Archive for Rational Mechanics and Analysis*, 167(2):85–146, 2003.
- [27] Q. Ma, D. Ding, and X. Ding. Symplectic conditions and stochastic generating functions of stochastic Runge-Kutta methods for stochastic Hamiltonian systems with multiplicative noise. *Applied Mathematics and Computation*, 219(2):635–643, 2012.
- [28] Q. Ma and X. Ding. Stochastic symplectic partitioned Runge-Kutta methods for stochastic Hamiltonian systems with multiplicative noise. *Appl. Math. Comput.*, 252(C):520–534, Feb. 2015.
- [29] J. Marsden and T. Ratiu. *Introduction to Mechanics and Symmetry*, volume 17 of *Texts in Applied Mathematics*. Springer Verlag, 1994.
- [30] J. E. Marsden, G. W. Patrick, and S. Shkoller. Multisymplectic geometry, variational integrators, and nonlinear PDEs. *Communications in Mathematical Physics*, 199(2):351–395, 1998.
- [31] J. E. Marsden and M. West. Discrete mechanics and variational integrators. *Acta Numerica*, 10(1):357–514, 2001.
- [32] R. I. McLachlan and G. R. W. Quispel. Geometric integrators for ODEs. *Journal of Physics A: Mathematical and General*, 39(19):5251–5285, 2006.
- [33] G. Milstein. *Numerical Integration of Stochastic Differential Equations*. Mathematics and Its Applications. Springer Netherlands, 1995.
- [34] G. N. Milstein, Y. M. Repin, and M. V. Tretyakov. Symplectic integration of Hamiltonian systems with additive noise. *SIAM J. Numer. Anal.*, 39(6):2066–2088, June 2001.
- [35] G. N. Milstein, Y. M. Repin, and M. V. Tretyakov. Numerical methods for stochastic systems preserving symplectic structures. *SIAM J. Numer. Anal.*, 40(4):1583 – 1604, 2002.
- [36] T. Misawa. Symplectic integrators to stochastic Hamiltonian dynamical systems derived from composition methods. *Mathematical Problems in Engineering*. Vol. 2010, Article ID 384937, 12 pages, 2010.

- [37] S. Ober-Blöbaum. Galerkin variational integrators and modified symplectic Runge–Kutta methods. *IMA Journal of Numerical Analysis*, 2016.
- [38] S. Ober-Blöbaum and N. Saake. Construction and analysis of higher order Galerkin variational integrators. *Advances in Computational Mathematics*, 41(6):955–986, 2015.
- [39] D. Pavlov, P. Mullen, Y. Tong, E. Kanso, J. E. Marsden, and M. Desbrun. Structure-preserving discretization of incompressible fluids. *Physica D: Nonlinear Phenomena*, 240(6):443–458, 2011.
- [40] P. Protter. *Stochastic Integration and Differential Equations*. Stochastic Modelling and Applied Probability. Springer Berlin Heidelberg, 2005.
- [41] C. W. Rowley and J. E. Marsden. Variational integrators for degenerate Lagrangians, with application to point vortices. In *Decision and Control, 2002, Proceedings of the 41st IEEE Conference on*, volume 2, pages 1521–1527. IEEE, 2002.
- [42] J. M. Sanz-Serna. Symplectic integrators for Hamiltonian problems: an overview. *Acta Numerica*, 1:243–286, 1992.
- [43] M. Seeßelberg, H. P. Breuer, H. Mais, F. Petruccione, and J. Honerkamp. Simulation of one-dimensional noisy Hamiltonian systems and their application to particle storage rings. *Zeitschrift für Physik C Particles and Fields*, 62(1):63–73, 1994.
- [44] A. Stern, Y. Tong, M. Desbrun, and J. E. Marsden. Variational integrators for Maxwell’s equations with sources. *PIERS Online*, 4(7):711–715, 2008.
- [45] L. Sun and L. Wang. Stochastic symplectic methods based on the Padé approximations for linear stochastic Hamiltonian systems. *Journal of Computational and Applied Mathematics*, 2016. <http://dx.doi.org/10.1016/j.cam.2016.08.011>.
- [46] T. M. Tyranowski and M. Desbrun. R-adaptive multisymplectic and variational integrators. *IMA Journal of Numerical Analysis*, 2013. (submitted, arXiv:1303.6796).
- [47] T. M. Tyranowski and M. Desbrun. Variational partitioned Runge-Kutta methods for Lagrangians linear in velocities. Preprint arXiv:1401.7904, 2013.
- [48] J. Vankerschaver and M. Leok. A novel formulation of point vortex dynamics on the sphere: geometrical and numerical aspects. *J. Nonlin. Sci.*, 24(1):1–37, 2014.
- [49] L. Wang. *Variational Integrators and Generating Functions for Stochastic Hamiltonian Systems*. PhD thesis, Karlsruhe Institute of Technology, 2007.
- [50] L. Wang and J. Hong. Generating functions for stochastic symplectic methods. *Discrete and Continuous Dynamical Systems*, 34(3):1211–1228, 2014.