

RESEARCH ARTICLE


Is systematic training in opioid overdose prevention effective?

Albert Espelt^{1,2,3,4*}, Marina Bosque-Prous¹, Cinta Folch⁵, Ana Sarasa-Renedo^{1,6}, Xavier Majó⁷, Jordi Casabona⁷, M. Teresa Brugals¹, REDAN Group¹

1 Agència de Salut Pública de Barcelona, Plaça Lesseps, Barcelona, Spain, **2** Centros de Investigación Biomédica en Red. Epidemiología y Salud Pública (CIBERESP), Calle Melchor Fernández Almagro, Madrid, Spain, **3** Facultat de Ciències de la Salut de Manresa, Universitat de Vic Universitat Central de Catalunya (UVicUCC), Av. Universitària, Manresa, Spain, **4** Departament de Psicobiologia i Metodologia en Ciències de la Salut, Universitat Autònoma de Barcelona, Campus UAB, Bellaterra, Spain, **5** Centre d'Estudis Epidemiològics sobre les Infeccions de Transmissió Sexual i Sida de Catalunya (CEEISCAT), Agència de Salut Pública de Catalunya (ASPC), Generalitat de Catalunya, Carretera Canyet s/n, Badalona, Spain, **6** Programa de Epidemiologia Aplicada y de Campo (PEAC), Instituto de Salud Carlos III, Calle Sinesio Delgado, Madrid, Spain, **7** Subdirecció General de Drogodependències, Departament de Salut de la Generalitat de Catalunya, Carrer de Roc Boronat, Barcelona, Spain

† A full list of members of the REDAN Group is provided in the Acknowledgments.

* aespelt@umanresa.cat



OPEN ACCESS

Citation: Espelt A, Bosque-Prous M, Folch C, Sarasa-Renedo A, Majó X, Casabona J, et al. (2017) Is systematic training in opioid overdose prevention effective? PLoS ONE 12(10): e0186833. <https://doi.org/10.1371/journal.pone.0186833>

Editor: Georgios Nikolopoulos, University of Cyprus, CYPRUS

Received: December 15, 2016

Accepted: October 9, 2017

Published: October 31, 2017

Copyright: © 2017 Espelt et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The study consists of two databases: the database of the Harm Reduction survey and the mortality database. The data from the mortality database are from the Institute of Legal Medicine and include personal data of the participants, which are very sensitive data. These data can also be made accessible for research purposes upon request to the Drug Information System of Barcelona (sicas@aspb.cat). All other data are available in the manuscript and Supporting Information files.

Abstract

The objectives were to analyze the knowledge about overdose prevention, the use of naloxone, and the number of fatal overdoses after the implementation of Systematic Training in Overdose Prevention (STOOP) program. We conducted a quasi-experimental study, and held face-to-face interviews before ($n = 725$) and after ($n = 722$) implementation of systematic training in two different samples of people who injected opioids attending harm reduction centers. We asked participants to list the main causes of overdose and the main actions that should be taken when witnessing an overdose. We created two dependent variables, the number of (a) correct and (b) incorrect answers. The main independent variable was *Study Group*: Intervention Group (IG), Comparison Group (CG), Pre-Intervention Group With Sporadic Training in Overdose Prevention (PREIGS), or Pre-Intervention Group Without Training in Overdose Prevention (PREIGW). The relationship between the dependent and independent variables was assessed using a multivariate Poisson regression analysis. Finally, we conducted an interrupted time series analysis of monthly fatal overdoses before and after the implementation of systematic program during the period 2006–2015. Knowledge of overdose prevention increased after implementing systematic training program. Compared to the PREIGW, the IG gave more correct answers (IRR = 1.40;95%CI: 1.33–1.47), and fewer incorrect answers (IRR = 0.33;95%CI: 0.25–0.44). Forty percent of people who injected opioids who received a naloxone kit had used the kit in response to an overdose they witnessed. These courses increase knowledge of overdose prevention in people who use opioids, give them the necessary skills to use naloxone, and slightly diminish the number of fatal opioid overdoses in the city of Barcelona.

Funding: This work was supported by the Spanish Network on Addictive Disorders [grant numbers RD06/0001/1018, and RD12/0028/0018] (<http://www.eng.isciii.es/ISCIII/es/contenidos/fd-investigacion/fd-ejecucion/fd-centros-participados/centros-participados-redes-retics.shtml>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Most deaths due to illicit drugs are caused by heroin and illicit opioids, with overdose being a leading cause of death among people who use opioids. In 2010, the estimated average EU mortality rate due to overdose among 15-64-year-olds was 18.3 deaths per million inhabitants (7,000–8,000 deaths per year) [1]. Opioid overdose can be fatal or non-fatal, and while fatal overdoses are an important public health problem worldwide, non-fatal overdoses are also important because they cause significant morbidity among victims [2]. While 3–5% of overdoses result in death [3–6], the annual incidence of non-fatal overdose ranges from 9 to 22% [4,7,8]. Opioid overdose can be prevented by taking certain risk factors and risky behaviors into account (e.g. controlling heroin administration route, and avoiding the concomitant use of other drugs) [9], and overdoses can be reversed using simple measures. However, three out of ten people who use opioids in Spain have insufficient knowledge of overdose risk factors or actions to take when witnessing an overdose [10,11].

Training programs in the prevention and management of opioid overdose have proven effective in increasing the relevant knowledge among people who inject Opioids (PWIO) in various settings [12–15], and these programs may be driving the ongoing decrease in overdose mortality [16,17]. In 2009, we designed a generalized Systematic Training in Opioid Overdose Prevention (STOOP) program to be implemented in Catalonia. The program consists in systematic training courses that started in all harm reduction centers in 2009, and was gradually extended to therapeutic communities and treatment centers. By 2013, the STOOP program had already been implemented in all harm reduction centers, treatment centers and therapeutic communities of Catalonia [18]. A specific manual, created to educate and assist in overdose prevention, is the basis for the implementation of STOOP program throughout the territory [18]. The STOOP program is addressed to groups of PWIO and people who use psycho-stimulants, and explain the risks, signs and symptoms of an overdose, and the differences between overdoses caused by opioids and those caused by other psycho-stimulants (see S1 and S2 Tables). This program addresses common myths about dealing with overdoses, and users are instructed on the correct actions to take (i.e. management) when an overdose occurs. Users who have acquired sufficient knowledge (assessed using a test after completing the full program) are given a naloxone kit (two 1 ml bottles of Naloxone (0.4mg/ml), 1 retractable syringe, 1 mask, 2 alcohol wipes, and a brochure with additional information) [18].

To inform future management and policymaking, it is necessary to evaluate the Systematic Training in Opioid Overdose Prevention (STOOP) program. The objectives of this study were to evaluate knowledge about overdose prevention, the use of naloxone, and the number of fatal overdoses following implementation of the STOOP program in Catalonia.

Material and methods

Study design and subjects

We used a quasi-experimental pre-post study design, including a comparison group [19]. The study sample consisted of people who injected opioids and who were attending any of the 18 existing harm reduction centers in Catalonia. Harm reduction centers included needle exchange programs, outreach programs, and supervised injecting facilities. The inclusion criteria were: having injected opioids during the 6 months prior to the interview, and having given written informed consent. Participants were recruited in two distinct periods (each sample was selected independently), before (from October 2008 to March 2009) and after implementation of STOOP program (from October 2010 to April 2011). An independent sample was selected for each period ($n_1 = 725$; $n_2 = 722$, respectively) [20] (Fig 1). Subjects were




Fig 1. Design of the quasi-experimental study.

<https://doi.org/10.1371/journal.pone.0186833.g001>

assigned to strata in proportion to the volume of visits in each center and the percentage of individuals in each center by country of birth. In centers with less than 5% of foreign-born users, we recruited only native participants. Participants were randomly selected within harm reduction centers.

Finally, we designed an interrupted time series analysis of monthly fatal overdoses in the city of Barcelona (Catalonia) before and after implementation of the STOOP program, including data between 2006 and 2015. The cut-off point was set in January 2013 when the STOOP program had already been implemented in all harm reduction centers, treatment centers and therapeutic communities.

Data collection

Before and after implementing the STOOP program, trained interviewers conducted face-to-face interviews in each center using an anonymous structured questionnaire adapted from the Itinere project [21] and the World Health Organization [22]. We collected and analyzed data from the following variables associated with knowledge of overdose prevention: age, sex, educational level, irregular income, regular residence, country of birth, lifetime treatment for drug dependency (including drug-free residential treatment or admission to therapeutic communities, in-hospital detoxification, out-patient drug-free treatment, methadone maintenance, and other medication or other treatments), frequency of drug injection, poly-drug use and lifetime history of overdose.

To encourage participation, those respondents participating in the interview before the implementation of STOOP program received 12 Euros and those participating after the implementation of STOOP program received 24 Euros. All the participants provided their written informed consent to participate in this study. The study protocol was approved by a Clinical Ethics Review Board (Hospital Universitari Germans Trias I Pujol, Badalona, Spain).

Variables

Dependent variables. *Knowledge about overdose prevention:* To assess this, we used two open questions, participants were asked to list the main causes of overdose and the main

actions that should be taken when witnessing an overdose, as has been done elsewhere [10]. An opioid overdose was defined as an episode that occurred following use of heroin, methadone or other opioids, and is characterized by extreme difficulty in breathing, loss of consciousness and problems waking up or recovering consciousness, and sometimes bluish skin or lips [10]. As described elsewhere [11], the responses to each open question were collected, transcribed verbatim and coded. Responses were then classified in 11 answer categories for causes of overdose (9 correct and 2 incorrect answers) and 15 actions to take when witnessing an opioid overdose (9 correct and 6 incorrect answers) (Table 1). This classification was reviewed and agreed upon separately by three experienced researchers (co-authors MTB, ASR, AE), and a regional working document was used to resolve inconsistencies [23]. From this information, we built two dependent variables based on the number of (1) correct or (2) incorrect responses, revealing participants' level of knowledge about the causes of overdose, and actions to take when witnessing an overdose.

Number of fatal opioid overdoses per month. Data on monthly fatal overdoses between 2006 and 2014 came from the register of the Legal Medicine Institute from the city of Barcelona (Catalonia). Overdoses due to other substances or that were suspected to be intentional (suicides) were excluded from the analysis.

Independent variables. The main independent variable was *Study Group*: Pre-Implementation Group Without any Courses (PREIGW), Pre-Implementation Group With Sporadic Courses (PREIGS), Comparison Group (CG) or Intervention Group (IG). The PREIGW consisted of all people who injected opioids interviewed between October 2008 and March 2009 who reported that they had not attended any STOOOP course ($n = 529$). The PREIGS consisted of all people who injected opioids interviewed between October 2008 and March 2009 who reported that they had attended some Opioid Prevention Training course ($n = 196$). The IG ($n = 220$) consisted of people who injected opioids interviewed between October 2010 and April 2011, when STOOOP program had already been implemented and reported that they had attended to an Opioid Overdose Prevention Training course at least once in the previous 2 years. Individuals interviewed after the implementation of the STOOOP program who self-reported that they had not participated in any course were included in the CG ($n = 502$). Before the implementation of the STOOOP program there were only sporadic courses with this objective in Catalonia. Thus, existing sporadic courses focused on training a few key PWIO on how to administer naloxone injections to their peers, and were less well prepared and more heterogeneous than the STOOOP program.

Finally, to describe and evaluate the effectiveness of naloxone distribution, we separately analyzed the following 4 variables from the IG: having received a naloxone kit in the previous 12 months, having witnessed any overdose in the previous 12 months, having helped a peer suffering an overdose, and having administered the naloxone kit to the peer suffering an overdose.

Statistical analyses

To perform an initial pre-post evaluation in terms of curve shifting, for each study group (PREIGW, PREIGS, GC or IG) we plotted the distribution of the number of correct and incorrect answers about causes or actions in overdose prevention and management (Fig 2). Similarly, we assessed users' level of knowledge by calculating the mean number of correct and incorrect answers about causes or actions, and the corresponding 95% Confidence Intervals (95%CI). To further assess whether the STOOOP program resulted in greater knowledge among people who injected opioids, we fit a multivariate Poisson regression model to obtain the adjusted Incidence Rate Ratio (IRR) and 95%CI [STATA syntax: poisson DependentVariable

Table 1. Knowledge about overdose prevention: cited causes of overdose, and actions to take when this happens, according to the pre-implementation, comparison, and intervention groups.

	PREIGW	PREIGS	CG	IG
	%	%	%	%
Causes of opioid overdose				
Correct answers				
Use of heroin together with other drugs	45.0	66.3	50.8	69.1
Amount injected	66.4	59.2	66.2	59.1
Stronger or purer than usual	15.7	18.4	28.7	32.7
Lower tolerance to heroin	11.7	20.4	16.8	27.7
Change of drug supplier	1.7	1.5	3.6	8.2
Health causes (weakness, predisposition, low defenses. . .)	9.8	10.7	11.1	5.5
Psychological problems / suicide attempt	2.6	2.0	2.0	1.4
Injecting whole dose at once or very quickly	4.3	1.0	1.8	0.9
Intravenous route	5.5	4.1	0.4	0.5
Incorrect answers				
Adulterated or cut heroin	23.8	25.5	16.6	13.2
Meaningless and false causes	5.9	4.1	3.6	2.3
Action to take when witnessing an opioid overdose				
Correct answers				
First aid	47.8	76.0	80.0	89.0
Call emergency services	59.7	63.8	69.5	72.6
Use of naloxone	0.0	0.0	9.9	43.8
Check consciousness	7.0	15.3	3.1	16.0
Wake up/keep the person awake	12.9	6.6	16.7	7.3
Call police/call for help	6.4	3.1	2.7	3.2
Remove syringe	0.2	1.0	0.6	2.3
Observation	1.9	1.5	3.3	0.9
Facilitate breathing	0.9	0.5	0.6	0.0
Incorrect answers				
Shower the person	18.5	17.3	16.0	4.1
Inject substance other than naloxone	11.0	1.5	8.4	2.7
Hit/shake the person	7.6	8.7	6.8	2.3
Meaningless actions	6.0	11.7	5.6	2.3
Make the person move/stand up	14.7	9.2	9.1	1.8
Abandon him/her	0.4	0.0	1.0	0.5


PREIGW: Pre-Implementation Group Without Training in Overdose Prevention; PREIGS: Pre-Implementation Group With Sporadic Training in Overdose Prevention; CG: Comparison Group; IG: Intervention Group

<https://doi.org/10.1371/journal.pone.0186833.t001>

IndependentVariables, vce(robust) irr] [24]. The main independent variable was *Study Group*. The model was adjusted for variables associated with overdose prevention knowledge, as described above [11].

In people from the IG we created a flow chart to summarize the use of naloxone during the year after attending the overdose prevention course.

Finally, we performed a descriptive analysis of the number of fatal opioid overdoses for each month since 2006. To evaluate changes in the number of fatal opioid overdoses after implementation of systematic overdose prevention training courses, we performed an interrupted time-series analysis using quasi-poisson regression models for overdispersed count data, as previously suggested [25]. We compared the number of fatal opioid overdoses per



PREIGW, Pre-Implementation Group Without Training in Overdose Prevention;

PREIGS, Pre-Implementation Group With Sporadic Training in Overdose Prevention;

CG, Comparison Group; IG, Intervention Group

Fig 2. Distribution of people who injected opioids recruited through harm reduction centers in Catalonia according to the number of a) correct and b) incorrect answers about overdose risk factors, and adequate/inadequate actions for reversing or minimizing the effects of an overdose.

<https://doi.org/10.1371/journal.pone.0186833.g002>

month throughout the time series, controlling for time trend and seasonal patterns using linear trend and including Fourier series terms in the model [26]. [STATA syntax: `glm Overdose-Deaths Intervention trend sin1 cos1 sin2 cos2 sin3 cos3 sin4 cos4, f(poisson) l(log) scale(x2) eform`]. We selected 2013 as the intervention year because the STOOOP program had not been widely implemented in therapeutic communities and treatment centers before that year [18].

All statistical analyses were performed using STATA 13.0.

Results

General characteristics of the sample

Overall, the study included 1,447 people who injected opioids who were attending harm reduction centers in Catalonia. Table 2 shows the between-group differences in the independent variables. These differences were observed for the following variables: age, lifetime drug treatment history, previous overdose history, poly-drug use, irregular income, regular residence at time of interview and country of birth.

Effect of attending systematic training in opioid overdose prevention program in knowledge acquisition

In the second round of recruitment, 722 participants were interviewed, of which 30% reported having attended at least one of the STOOOP courses (IG). Knowledge about overdose prevention was greater after the implementation of STOOOP program. Comparing the number of responses cited by participants (overdose risk factors and the correct actions to take when witnessing an overdose), we found, both in IG and CG, that the distribution curve shifted towards higher scores for correct answers and lower scores for incorrect answers, with the IG showing

Table 2. General characteristics of people who injected opioids recruited through harm reduction centers in Catalonia.

	PREIGW (n = 529)	PREIGS (n = 196)	CG (n = 502)	IG (n = 220)	p-value
	%	%	%	%	
Sex					
Men	83.7	79.1	84.5	79.5	= 0.19
Age					
>30 years	73.1	88.6	75.6	82.3	<0.01
Educational level					
Secondary or higher	26.2	20.5	23.1	24.7	= 0.39
Age at first injection					
>20	44.5	41.3	44.7	42.2	= 0.81
Residence¹					
Institution or homeless	39.3	41.0	32.5	47.7	<0.01
Irregular income¹					
Yes	60.3	66.2	42.9	51.4	<0.01
Self-perceived health²					
Poor	38.0	39.0	45.2	39.4	= 0.11
Time since last drug injection					
≥30 days	93.6	89.8	91.0	87.3	= 0.04
Use of Supervised Injecting Facility¹					
Less than half of injection days	47.0	50.3	47.6	48.4	= 0.89
Lifetime drug treatment					
No	17.6	2.6	21.9	9.1	<0.01
Poly-drug use¹					
≥3 drugs	94.3	98	80.7	83.2	<0.01
Previous overdose history					
No	50.9	30.1	44.6	34.5	<0.01
Country of birth					
Native (Spain)	53.7	70.9	61.3	61.9	
Eastern Europe	29.3	10.2	25.9	15.1	
Other countries	17.0	18.9	12.8	22.9	<0.01

¹Previous 6 months

²At time of interview; PREIGW, Pre-Implementation Group Without Training in Overdose Prevention; PREIGS, Pre-Implementation Group With Sporadic Training in Overdose Prevention; CG, Comparison Group; IG, Intervention Group.

p-value compares the values for each variable between study groups.

<https://doi.org/10.1371/journal.pone.0186833.t002>

the most marked curve displacement. In other words, the population who injected opioids as a whole gained knowledge on overdose prevention (Fig 2).

The IG showed the highest levels of knowledge, with a mean number of correct and incorrect answers of 4.4 (95%CI: 4.2–4.5) and 0.3 (95%CI: 0.2–0.4), respectively (Table 3). Table 3 also shows the results of the association between study group and the number of correct and incorrect answers about causes and actions in overdose prevention after adjusting for confounding variables related to knowledge of overdose prevention. Individuals in the IG, CG and PREIGS were more likely to give correct answers reflecting adequate knowledge [IRR 1.40 (95%CI: 1.33–1.47), 1.17 (95%CI: 1.12–1.23) and 1.09 (95%CI: 1.04–1.16), respectively] and less likely to give incorrect answers [IRR 0.33 (95%CI: 0.25–0.44), 0.74 (95%CI: 0.64–0.85) and 0.85 (95%CI: 0.71–1.02)] than those in the PREIGW. The IRR of 1.40 indicates that individuals in the IG gave 40% more correct answers than those in the PREIGW.

Table 3. Correct or incorrect answers about causes or actions in overdose prevention among people who injected opioids recruited at harm reduction centers in Catalonia before and after the implementation of Systematic Training in Opioid Overdose Prevention in Catalonia.

	Correct answers				Incorrect answers			
	mean	(95%CI)	aIRR	(95%CI)	mean	(95%CI)	aIRR	(95%CI)
PREIGW	3.1	(3.0–3.2)	1		0.9	(0.8–1.0)	1	
PREIGS	3.6	(3.5–3.7)	1.09	(1.04–1.16)	0.8	(0.7–0.9)	0.85	(0.71–1.02)
CG	3.6	(3.5–3.7)	1.17	(1.12–1.23)	0.7	(0.6–0.7)	0.74	(0.64–0.85)
IG	4.4	(4.2–4.5)	1.40	(1.33–1.47)	0.3	(0.2–0.4)	0.33	(0.25–0.44)

PREIGW: Pre-Implementation Group Without Training in Overdose Prevention; PREIGS: Pre-Implementation Group With Sporadic Training in Overdose Prevention; CG: Comparison Group; IG: Intervention Group. aIRR: Incidence Rate Ratio of correct and incorrect answers before and after the implementation of the STOOOP program; aIRR were adjusted for sex, age, educational level, age at first injection, residence, income, self-perceived health, time since last injection in a harm reduction facility, previous treatment for drug dependency, poly-drug use, previous overdose, previous overdose prevention training, and country of birth.

<https://doi.org/10.1371/journal.pone.0186833.t003>

Table 1 shows that people who injected opioids in the IG generally gave more correct answers than those in the CG, PREIGS and PREIGW (e.g. heroin use together with other drugs, use of stronger or purer heroin than usual, reduced tolerance to heroin, and change of drug supplier), and fewer incorrect answers (e.g. taking adulterated or cut heroin). This difference was even greater for actions to take when witnessing an overdose, especially for giving first aid, using naloxone, and checking consciousness. For example, the percentage of responses such as, showering the individual, make them move/stand up, and injecting substances other than naloxone, was lower in the IG than in the other groups (Table 1).

Implementation of systematic training in opioid overdose prevention and use of naloxone

One hundred fifty-eight participants in the IG (72%) received naloxone, of whom 94 (59%) reported having witnessed ≥ 1 overdoses in the 12 months prior to the interview, 68% of whom (n = 64) had helped the sufferer (59% of these administered naloxone, Fig 3). Thus, 40.4% of people who injected opioids who had received a naloxone kit had used it when witnessing an overdose (Fig 3).

Implementation of systematic training in opioid overdose prevention courses and fatal opioid overdoses

Fig 4 shows the observed distribution of the number of fatal opioid overdoses since 2006, as well as the distribution (and 95%CI) that would be expected if the STOOOP program had not been implemented before and after 2013. The gap between the number of expected and observed fatal opioid overdoses increased over time. In the years 2013 and 2014 there were 27 fewer fatal opioid overdoses than expected if the STOOOP program had not been implemented.

Discussion

Our data indicate that 1) people who use opioids and attended the Systematic Training in Opioid Overdose Prevention (STOOOP) program have greater knowledge of overdose prevention than those who did not receive systematic training courses. Our results suggest that these programs improve the general understanding of the population who use opioids as a whole. Individuals who did not attend the STOOOP program in 2010–2011 had the same level of knowledge as those who attended sporadic courses in 2008–2009. 2) 59% of trained users that




Fig 3. Use of naloxone when witnessing an overdose in the intervention group.

<https://doi.org/10.1371/journal.pone.0186833.g003>




Fig 4. Trend of fatal overdoses (a) and (b) Observed and expected number of fatal opioid overdose per month, Barcelona, 2006–2015.

<https://doi.org/10.1371/journal.pone.0186833.g004>

received naloxone had witnessed an overdose in the previous 12 months, and 40% of them had used naloxone during the most recently witnessed overdose. 3) After 2013, there was a decline in the number of fatal overdoses in Barcelona, with 27 fewer fatal opioid overdoses in 2013–2014 than expected if the STOOOP program had not been implemented.

We observed greater knowledge in the IG than in the PREIGW, consistent with other studies that found an increase in knowledge [12–14] and better knowledge retention in trained PWID [27,28]. The greater overdose prevention knowledge increased gradually between groups ($PREIGW < PREIGC \leq CG < IG$). Thus, STOOOP program could result in an overall increase in knowledge among people who use opioids. The increase among untrained users may be due to greater availability of information at harm reduction centers (training courses were held regularly at each center), and greater awareness among staff who had received training on this topic after introduction of STOOOP program. This general increase in knowledge could also be due to peer diffusion, in that course participants are known to share what they have learned through their actions and their conversations, both within and beyond the context of overdose events [29]. Peer diffusion of this information is particularly important in this community because of the difficulty in recruiting people who use opioids for training, and the stigma attached to illicit drug use [30]. For this reason, the implementation of the STOOOP program was accompanied by specific training on overdose prevention in NGOs, the police department, users' associations, and social educators involved in municipal plans.

Some external factors independent of the STOOOP program (e.g. other campaigns) could have boosted the level of knowledge. However, this is unlikely because the interventions conducted in the population attending harm reduction centers are informed to our organization or supervised by. In addition, the chain of causality between the intervention and the expected outcome was very direct (few intermediate factors) and the study period was relatively short [19,27]. Alternatively, the improvement in knowledge could be due to individual variables related to drug consumption or individual maturation. However, we think this is unlikely because we adjusted the final regression models by drug consumption and individual maturation variables associated with knowledge of opioid overdose prevention. In addition, improvements in knowledge about some causes or actions in overdose prevention were clearly related to attendance at the courses, e.g. an increased number of correct answers about first aid, calling the emergency services and using naloxone, and a decreased number of incorrect answers such as making the victim take a shower or move/stand up (Table 1).

We found that 40.4% of people who use opioids who attended a STOOOP program and received a naloxone kit had used it during the last overdose they witnessed, a similar proportion to that observed in San Francisco (40%) [13] and in New York (58%) [31]. A meta-analysis done in 2015 [32], found that 9% of naloxone kits distributed to trained users will have been administered to a peer within the three months of supply. This is consistent with effective knowledge retention [27], and greater ability to recognize an overdose and act appropriately after training [33]. For example, it has been shown that PWIO with increased knowledge can administer naloxone to an overdose victim and prevent a fatal overdose [34]. However, some trained PWIO did not use their naloxone kit during the last overdose they witnessed. Previous studies have not been able to clarify the reasons for this because of small sample sizes, although the reasons given include loss of the naloxone kit, that the witness was no longer using drugs, and that the victim was already dead when found [27,35]. Tobin and colleagues [13] suggested that naloxone kits were generally not lost, stolen or confiscated, although in our study 16% of participants in the overdose prevention program who received a naloxone kit reported that they were not carrying it when they witnessed an overdose (results not shown). Another important issue is what could happen in countries with a different justice approach to drug use. Thus, people who use opioids need police permission to carry naloxone in public, which

in turn requires fluid communication and agreement between the police and the public administration. Therefore, after the STOOOP program, participants receive a card identifying them as experts in overdose prevention.

Finally, we observed a slight decline in the number of overdose deaths in Barcelona after 2013, which could be related to the STOOOP program. In June of 2012 only 43.5% of PWIO in Catalonia had participated in an overdose prevention course, with different percentages among individuals undergoing treatment (32.1%), those in therapeutic communities (35.2%), and those using harm reduction centers (66.7%) [36]. While we cannot directly attribute this reduction to the implementation of STOOOP program, our results are consistent those reported in Scotland [37] and the USA [17]. Implementation of Scotland's National Naloxone Programme was associated with a 36% decrease in the proportion of opioid-related deaths during the four weeks after release from prison. The results on greater knowledge of overdose prevention are encouraging because greater knowledge could be related to reduced overdose risk [38,39]. Moreover, a systematic review found that the Take-Home-Naloxone provision reduced fatal overdoses among program participants themselves, and also among fellow people who use opioids and the wider community, and significantly reduced overdose mortality with respect to communities without implementation [34].

Strengths and limitations

Finally, we note some strengths and weaknesses of our study. This is a study that uses an extended and representative sample of people who inject opioids and use harm reduction centers. The dependent variable, knowledge of overdose prevention, while not as straightforward as others [40], was asked in exactly the same way in each wave of this study, and the answers were classified in the same way, with broad consensus among the study researchers, as explained elsewhere [11].

In terms of limitations, we could not differentiate between knowledge acquisition, retention and application. Although we could not control for the time between attending the systematic courses and the study interview, we know that they were not done at the same time. The STOOOP program started in the third trimester of 2009 and the first surveys started in October 2010. Moreover, these courses are systematically addressed to all users, and reminders are issued and re-training offered a year after the first course. In this sense, the course remains active all year. Finally, we could not observe the direct impact of knowledge acquisition on some preventive practices, although we observed a slightly decline in the number of fatal overdoses in the city of Barcelona.

Conclusions

We found that Systematic Training in Opioid Overdose Prevention (STOOOP) program increases knowledge among people who injected opioids. In addition, a high percentage of trained people who injected opioids (40%) used naloxone during the last overdose they witnessed. The STOOOP program should also be deployed in prisons, since PWID have a higher risk of a drug-related death once they are released [41,42]. However, further research is required to investigate and develop new strategies to increase the use of naloxone where necessary.

Supporting information

S1 Table. Short Guide on counselling for heroin and cocaine overdose.
(DOCX)

S2 Table. Key points for training users in the prevention and treatment of overdoses.
(DOCX)

S1 File. This is the S1 File Dataset for Tables 1, 2 and 3.
(DTA)

S2 File. This is the S2 File Dataset for Fig 3.
(DTA)

Acknowledgments

The members of the REDAN Group are: Jordi Casabona, Cinta Folch, Anna Esteve, Alexandra Montoliu, Rafa Muñoz, Victoria González, Vicenç Ausina, M. Teresa Brugal, Albert Espelt, Carmen Vecino, Xavier Majó, Joan Colom, Mercè Meroño, Anna Altabas.

The authors would like to thank all interviewers, participants and collaborating centers: Ambit Prevenió; SAPS; Baluard, CAS Vall d'Hebrón; "El Local" Sant Adrià; AEC-Gris; Asaupam Santa Coloma; CAS Reus, AIDE Terrassa, Alba Terrassa, Arrels Lleida; Mercè Fontanilles Lleida; Creu Roja Constantí, IAS Girona.

Author Contributions

Conceptualization: Albert Espelt, Cinta Folch, Xavier Majó, Jordi Casabona.

Data curation: Marina Bosque-Prous.

Formal analysis: Albert Espelt, Ana Sarasa-Renedo.

Investigation: Albert Espelt, Marina Bosque-Prous, Cinta Folch, Ana Sarasa-Renedo, M. Teresa Brugal.

Methodology: Albert Espelt, Marina Bosque-Prous, Xavier Majó, Jordi Casabona, M. Teresa Brugal.

Supervision: M. Teresa Brugal.

References

1. EMCDDA. 2016 Annual report on the state of the drugs problem in Europe. Lisbon: European Monitoring Centre for Drugs and Drug Addiction (EMCDDA); 2016.
2. Warner-Smith M, Darke S, Lynskey M, Hall W. Heroin overdose: causes and consequences. *Addiction*. 2001; 96: 1113–1125. <https://doi.org/10.1080/09652140120060716> PMID: 11487418
3. Darke S, Mattick RP, Degenhardt L. The ratio of non-fatal to fatal heroin overdose. *Addiction*. 2003; 98: 1169–1171.
4. Darke S, Ross J, Hall W. Overdose among heroin users in Sydney, Australia: II. responses to overdose. *Addiction*. 1996; 91: 413–417. PMID: 8867203
5. Espelt A, Barrio G, Álamo-Junquera D, Bravo M, Sarasa-Renedo A, Vallejo F, et al. Lethality of opioid overdose in a community cohort of young heroin users. *Eur Addict Res*. 2015; 21: 300–306. <https://doi.org/10.1159/000377626> PMID: 26022713
6. Neale J. A response to Darke et al., "The ratio of non-fatal to fatal heroin overdose." *Addiction*. 2003; 98: 1171.
7. Brugal MT, Barrio G, De LF, Regidor E, Royuela L, Suelves JM. Factors associated with non-fatal heroin overdose: assessing the effect of frequency and route of heroin administration. *Addiction*. 2002; 97: 319–327. PMID: 11964108
8. Gossop M, Griffiths P, Powis B, Williamson S, Strang J. Frequency of non-fatal heroin overdose: survey of heroin users recruited in non-clinical settings. *BMJ*. 1996; 313: 402. PMID: 8761230
9. Neira-Leon M, Barrio G, Bravo MJ, Brugal MT, de la FL, Domingo-Salvany A, et al. Infrequent opioid overdose risk reduction behaviours among young adult heroin users in cities with wide coverage of HIV

- prevention programmes. *Int J Drug Policy*. 2011; 22: 16–25. <https://doi.org/10.1016/j.drugpo.2010.06.003> PMID: 20800462
10. Neira-Leon M, Barrio G, Brugal MT, de la FL, Ballesta R, Bravo MJ, et al. Do young heroin users in Madrid, Barcelona and Seville have sufficient knowledge of the risk factors for unintentional opioid overdose? *J Urban Health*. 2006; 83: 477–496. <https://doi.org/10.1007/s11524-006-9054-5> PMID: 16739049
 11. Sarasa-Renedo A, Espelt A, Folch C, Vecino C, Majó X, Castellano Y, et al. Overdose prevention in injecting opioid users: the role of substance abuse treatment and training programs. *Gac Sanit*. 2014; 28: 146–54. <https://doi.org/10.1016/j.gaceta.2013.10.012> PMID: 24418018
 12. Seal KH, Thawley R, Gee L, Bamberger J, Kral AH, Ciccarone D, et al. Naloxone distribution and cardiopulmonary resuscitation training for injection drug users to prevent heroin overdose death: a pilot intervention study. *J Urban Health*. 2005; 82: 303–311. <https://doi.org/10.1093/jurban/jti053> PMID: 15872192
 13. Tobin KE, Sherman SG, Beilenson P, Welsh C, Latkin CA. Evaluation of the Staying Alive programme: training injection drug users to properly administer naloxone and save lives. *Int J Drug Policy*. 2009; 20: 131–136. <https://doi.org/10.1016/j.drugpo.2008.03.002> PMID: 18434126
 14. Wagner KD, Valente TW, Casanova M, Partovi SM, Mendenhall BM, Hundley JH, et al. Evaluation of an overdose prevention and response training programme for injection drug users in the Skid Row area of Los Angeles, CA. *Int J Drug Policy*. 2010; 21: 186–193. <https://doi.org/10.1016/j.drugpo.2009.01.003> PMID: 19268564
 15. Maldjian L, Siegler A, Kunins HV. Evaluation of overdose prevention trainings in New York City: Knowledge and self-efficacy among participants 12 months after training. *Subst Abus*. 2016; 37: 459–465. <https://doi.org/10.1080/08897077.2015.1135850> PMID: 26731134
 16. Clark AK, Wilder CM, Winstanley EL. A systematic review of community opioid overdose prevention and naloxone distribution programs. *J Addict Med*. 2014; 8: 153–163. <https://doi.org/10.1097/ADM.000000000000034> PMID: 24874759
 17. Walley AY, Xuan Z, Hackman HH, Quinn E, Doe-Simkins M, Sorensen-Alawad A, et al. Opioid overdose rates and implementation of overdose education and nasal naloxone distribution in Massachusetts: interrupted time series analysis. *BMJ*. 2013; 346: f174. <https://doi.org/10.1136/bmj.f174> PMID: 23372174
 18. Espelt A, Major X, Parés-Badell O, Carvajal S, Gasulla L, Bosque-Prous M, et al. Implementation of Systematic Programs of Overdose Training at Drug Treatment and Prevention Centres in Catalonia, 2008–2013. In: Dichtl A, Heino S, editors. *Naloxon—Überlebenshilfe im Drogennotfall*. Frankfurt: Fachhochschulverlag; 2015. pp. 83–95.
 19. López MJ, Mari-Dell’Olmo M, Pérez-Giménez A, Nebot M. [Evaluative designs in public health: methodological considerations]. *Gac Sanit*. 2011; 25 Suppl 1: 9–16. [https://doi.org/10.1016/S0213-9111\(11\)70003-6](https://doi.org/10.1016/S0213-9111(11)70003-6)
 20. CEESCAT. SIVES 2012: Integrated Surveillance System for STI and HIV in Catalonia [Internet]. Barcelona: Generalitat de Catalunya, Departament de Salut; 2012. Available: http://www.ceeiscat.cat/documents/SIVES_2012_eng.pdf
 21. de la Fuente L, Brugal Puig MT, Ballesta GR, Bravo Poetela MJ, Barrio AG, Domingo SA, et al. [Cohort study methodology of the ITINERE Project on heroin users in three Spanish cities and main characteristics of the participants]. *Rev Esp Salud Publica*. 2005; 79: 475–491. PMID: 16465964
 22. WHO. Multy-city study on drug injecting and risk of HIV infection. Programme on substance abuse—final report. Geneva, Switzerland: World Health Organization; 1994.
 23. Majó X, Illundain E, Valverde C. Manual per educar en prevenció i assistència a les sobredosis. Adreçat als professionals de les drogodependències. Generalitat de Catalunya. Departament de Salut; 2009.
 24. Hilbe JM. *Modeling Count Data*. New York, NY: Cambridge University Press; 2014.
 25. Ver Hoef JM, Boveng PL. Quasi-Poisson vs. negative binomial regression: how should we model overdispersed count data? *Ecology*. 2007; 88: 2766–2772. PMID: 18051645
 26. Bhaskaran K, Gasparrini A, Hajat S, Smeeth L, Armstrong B. Time series regression studies in environmental epidemiology. *Int J Epidemiol*. 2013; 42: 1187–1195. <https://doi.org/10.1093/ije/dyt092> PMID: 23760528
 27. Gaston RL, Best D, Manning V, Day E. Can we prevent drug related deaths by training opioid users to recognise and manage overdoses? *Harm Reduct J*. 2009; 6: 26. <https://doi.org/10.1186/1477-7517-6-26> PMID: 19781073
 28. Strang J. Death matters: understanding heroin/opiate overdose risk and testing potential to prevent deaths. *Addiction*. 2015; 110 Suppl S2: 27–35. <https://doi.org/10.1111/add.12904> PMID: 26042565

29. Sherman SG, Gann DS, Tobin KE, Latkin CA, Welsh C, Bielensohn P. "The life they save may be mine": diffusion of overdose prevention information from a city sponsored programme. *Int J Drug Policy*. 2009; 20: 137–142. <https://doi.org/10.1016/j.drugpo.2008.02.004> PMID: 18502635
30. Rossi C. Monitoring drug control strategies: hidden phenomena, observable events, observable times. *Int J Drug Policy*. 1999; 10: 131–144.
31. Galea S, Worthington N, Piper TM, Nandi VV, Curtis M, Rosenthal DM. Provision of naloxone to injection drug users as an overdose prevention strategy: early evidence from a pilot study in New York City. *Addict Behav*. 2006; 31: 907–912. <https://doi.org/10.1016/j.addbeh.2005.07.020> PMID: 16139434
32. McAuley A, Aucott L, Matheson C. Exploring the life-saving potential of naloxone: A systematic review and descriptive meta-analysis of take home naloxone (THN) programmes for opioid users. *Int J Drug Policy*. 2015; 26: 1183–1188. <https://doi.org/10.1016/j.drugpo.2015.09.011> PMID: 26508033
33. Green TC, Heimer R, Grau LE. Distinguishing signs of opioid overdose and indication for naloxone: an evaluation of six overdose training and naloxone distribution programs in the United States. *Addiction*. 2008; 103: 979–989. <https://doi.org/10.1111/j.1360-0443.2008.02182.x> PMID: 18422830
34. McDonald R, Strang J. Are take-home naloxone programmes effective? Systematic review utilizing application of the Bradford Hill criteria. *Addiction*. 2016; 111: 1177–1187. <https://doi.org/10.1111/add.13326> PMID: 27028542
35. Strang J, Manning V, Mayet S, Best D, Titherington E, Santana L, et al. Overdose training and take-home naloxone for opiate users: prospective cohort study of impact on knowledge and attitudes and subsequent management of overdoses. *Addiction*. 2008; 103: 1648–1657. <https://doi.org/10.1111/j.1360-0443.2008.02314.x> PMID: 18821875
36. Arribas-Ibar E, Sánchez-Niubò A, Majó X, Domingo-Salvany A, Brugal MT. Coverage of overdose prevention programs for opiate users and injectors: a cross-sectional study. *Harm Reduct J*. 2014; 11: 33. <https://doi.org/10.1186/1477-7517-11-33> PMID: 25416534
37. Bird SM, McAuley A, Perry S, Hunter C. Effectiveness of Scotland's National Naloxone Programme for reducing opioid-related deaths: a before (2006–10) versus after (2011–13) comparison. *Addiction*. 2016; 111: 883–891. <https://doi.org/10.1111/add.13265> PMID: 26642424
38. EMCDDA. Preventing overdose deaths in Europe [Internet]. Lisbon, Portugal; 2013. Available: http://www.salledeconsoommation.fr/_media/emcdda-2013-preventing-overdose-deaths.pdf
39. Frisher M, Baldacchino A, Crome I, Bloor R. Preventing opioid overdoses in Europe: A critical assessment of known risk factors and preventative measures. *European Monitoring Centre for Drugs and Drug Addiction (EMCDDA)*; 2012.
40. Williams AV, Strang J, Marsden J. Development of Opioid Overdose Knowledge (OOKS) and Attitudes (OOAS) Scales for take-home naloxone training evaluation. *Drug Alcohol Depend*. 2013; 132: 383–386. <https://doi.org/10.1016/j.drugalcdep.2013.02.007> PMID: 23453260
41. Merrall ELC, Kariminia A, Binswanger IA, Hobbs MS, Farrell M, Marsden J, et al. Meta-analysis of drug-related deaths soon after release from prison. *Addiction*. 2010; 105: 1545–1554. <https://doi.org/10.1111/j.1360-0443.2010.02990.x> PMID: 20579009
42. Binswanger IA, Stern MF, Yamashita TE, Mueller SR, Baggett TP, Blatchford PJ. Clinical risk factors for death after release from prison in Washington State: a nested case-control study. *Addiction*. 2016; 111: 499–510. <https://doi.org/10.1111/add.13200> PMID: 26476210