

On uniqueness of limit cycles in general Bogdanov-Takens bifurcation

Maoan Han^{1*}, Jaume Llibre^{2†} and Junmin Yang^{3‡}

¹ Department of Mathematics, Shanghai Normal University, 100 Guilin Road

Shanghai 200234, China

² Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra

Barcelona, Catalonia, Spain

³ College of Mathematics and Information Science, Hebei Normal University

Shijiazhuang, Hebei 050024, China

Abstract In this paper we present a complete study to the well-known Bogdanov-Takens bifurcation and give a rigorous proof for the uniqueness of limit cycles.

1 Introduction

General theory of Bogdanov-Takens bifurcation is well-known. It was established by Bogdanov in [1] and then introduced in many books, see [2, 3, 4, 5, 13] and [9, 10, 20, 21]. However, to our knowledge, we have not found a really satisfactory and complete study in current works in English. For example, the uniqueness of limit cycles in homoclinic bifurcation was not considered in all of these works in English. The problem was first found and solved in the paper [14]. One can find a relatively complete study on it in books [9, 10, 20, 21] with different details. These books are all in Chinese. The aim of

*Corresponding author. mahan@shnu.edu.cn

†jllibre@mat.uab.cat

‡jmyang@hebtu.edu.cn

this paper is to provide a complete study with all details on the well-known Bogdanov-Takens bifurcation.

We organize the paper as follows. In section 2 we state some preliminary theorems which are useful in studying Bogdanov-Takens bifurcation. The preliminary theorems on Hopf and homoclinic bifurcations were obtained after the paper of Bogdanov [1]. In section 3 we study Bogdanov-Takens bifurcation in detail. In section 4 we provide some examples to show that our proof to Bogdanov-Takens bifurcation, especially the proof to the uniqueness of limit cycles in Hopf and homoclinic bifurcations, is really necessary.

2 Preliminary theorems: a short survey

Consider a C^∞ system on the plane of the form

$$\begin{aligned}\dot{x} &= f(x, y) + \varepsilon f_0(x, y, \varepsilon, \delta), \\ \dot{y} &= g(x, y) + \varepsilon g_0(x, y, \varepsilon, \delta),\end{aligned}\tag{2.1}$$

where f, g, f_0, g_0 are C^∞ functions, ε is small, and $\delta \in G \subset \mathbb{R}^n$ with G compact and $n \geq 1$. Suppose that for $\varepsilon = 0$ system (2.1) has a C^∞ first integral $H(x, y)$, such that the equation $H(x, y) = h$ defines a closed orbit L_h for $h \in J$, where $J = (\alpha, \beta)$ is an open interval. Here, we give a short survey on general theory of limit cycle bifurcation occurred in system (2.1). For simplicity, we suppose that the limit of L_h as $h \rightarrow \alpha$ exists and is an elementary center of $(2.1)|_{\varepsilon=0}$, denoted by C , while the limit of L_h as $h \rightarrow \beta$ exists and is a homoclinic loop of the unperturbed system, denoted by L . The loop L passes through a hyperbolic saddle S . Then C and L together form the boundary of the family $\{L_h\}$. To state some results on the number of limit cycles bifurcated from the family and its boundary, introduce the function below

$$M(h, \delta) = \oint_{L_h} (f g_0 - g f_0) e^{-\int_0^t (f_x + g_y) d\tau} |_{\varepsilon=0} dt, \quad h \in J.\tag{2.2}$$

First, for $h \in J$, we have the following result on the limit cycle bifurcation from the period annulus defined by $\{L_h\}$ which can be found in [18, 12].

Theorem 2.1. *Let $M(h, \delta)$ be the function defined by (2.2). Then*

(i) *For $h_0 \in J$, a necessary condition for L_{h_0} to generate a limit cycle is $M(h_0, \delta_0) = 0$ for some $\delta_0 \in G$.*

(ii) *If*

$$M(h_0, \delta_0) = 0, \quad \frac{\partial M}{\partial h}(h_0, \delta_0) \neq 0,$$

then there exists a constant $\varepsilon_0 > 0$ such that for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$ system (2.1) has a unique limit cycle near L_{h_0} .

(iii) If

$$\frac{\partial^k M}{\partial h^k}(h_0, \delta_0) \neq 0, \quad \frac{\partial^j M}{\partial h^j}(h_0, \delta_0) = 0, \quad j = 0, 1, \dots, k-1$$

for some integer $k \geq 1$, then there exists a constant $\varepsilon_0 > 0$ such that for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$ system (2.1) has at most k limit cycles near L_{h_0} .

According to [18], the above theorem was proved by L. Pontryagin in 1934 in the case $f_x + g_y \equiv 0$. The above general case was obtained by Urabe in 1954, and X. Chen in 1963 respectively.

Now we consider a Hopf bifurcation near the center C . Without loss of generality, we assume that $H(0, 0) = 0$, and that for $x^2 + y^2$ small we have

$$\begin{aligned} f(x, y) + \varepsilon f_0(x, y, \varepsilon, \delta) &= a(\varepsilon, \delta)x - b(\varepsilon, \delta)y + O(x^2 + y^2), \\ g(x, y) + \varepsilon g_0(x, y, \varepsilon, \delta) &= b(\varepsilon, \delta)x + a(\varepsilon, \delta)y + O(x^2 + y^2). \end{aligned} \quad (2.3)$$

In this case the center C has been moved to the origin. By introducing the polar coordinates $x = r \cos \theta$, $y = r \sin \theta$ one can obtain from (2.1) a C^∞ periodic equation of the form

$$\frac{dr}{d\theta} = R(\theta, r, \varepsilon, \delta).$$

Let $r(\theta, x_0, \varepsilon, \delta)$ denote the solution of this equation satisfying $r(0) = x_0$. Define

$$\Delta(x_0, \varepsilon, \delta) = r(2\pi, x_0, \varepsilon, \delta) - x_0.$$

Note that $\Delta(x_0, 0, \delta) = 0$ for $|x_0|$ small. We can write for $|x_0|$ small

$$\Delta(x_0, \varepsilon, \delta) = \varepsilon \sum_{i \geq 1} \Delta_i(\varepsilon, \delta) x_0^i. \quad (2.4)$$

It is known that the right-lower index i of the first nonzero coefficient Δ_i must be odd.

For (2.3) we have

$$\Delta_1(\varepsilon, \delta) = \frac{1}{\varepsilon} [e^{\frac{2\pi a}{b}} - 1], \quad \Delta_1(0, \delta) = \frac{2\pi}{b_0} \frac{\partial a}{\partial \varepsilon}(0, \delta),$$

where $\bar{b}_0 = b(0, \delta)$.

The following theorem was proved in [10, 6].

Theorem 2.2 (Hopf bifurcation) *Let (2.4) hold. Then*

(1) *If*

$$\Delta_{2k+1}(0, \delta_0) \neq 0, \quad \Delta_i(0, \delta_0) = 0, \quad i = 1, \dots, 2k$$

for some $\delta_0 \in G$, $k \geq 0$, then there exist $\varepsilon_0 > 0$ and a neighborhood U of the origin such that system (2.1) has at most k limit cycles in U for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$.

(2) Let $Q(h) = (q(h), 0)$ denote the intersection point of L_h with the positive x -axis with $q(h) > 0$, then

$$\Delta_3(0, \delta) = -\text{sgn}(b_0) \lim_{h \rightarrow 0} \frac{M(h, \delta)}{q^3 \sqrt{f^2(Q) + g^2(Q)}}$$

provided $\Delta_1(0, \delta) = 0$. In general, we have

$$\Delta_{2k+1}(0, \delta) = -\text{sgn}(b_0) \lim_{h \rightarrow 0} \frac{M(h, \delta)}{q^{2k+1} \sqrt{f^2(Q) + g^2(Q)}},$$

provided $\Delta_i(0, \delta) = 0$, $i = 1, \dots, 2k - 1$.

(3) If system (2.1) is analytic, $(f, g) = \pm(-H_y, H_x)$ and $M(h, \delta_0) \neq 0$ for some $\delta_0 \in G$, then for some $k \geq 0$ and $N_k \neq 0$

$$M(h, \delta_0) = N_k h^{k+1} + O(h^{k+\frac{3}{2}}).$$

In this case, $\Delta_i(0, \delta_0) = 0$, $i = 1, \dots, 2k - 1$, and $\Delta_{2k+1}(0, \delta_0) = K_k N_k$ for some constant $K_k \neq 0$. For example, when $H(x, y) = K(x^2 + y^2) + o(x^2 + y^2)$ we have $\Delta_3(0, \delta_0) = \pm \frac{K}{2} N_1$.

On the uniqueness of limit cycles in a Hopf bifurcation we have the following corollary coming from the above theorem.

Corollary 2.1. *If $\frac{\partial a}{\partial \varepsilon}(0, \delta_0) = 0$ and $\Delta_3(0, \delta_0) \neq 0$, then there exists $\varepsilon_0 > 0$ and a neighborhood U of the origin such that for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$, system (2.1) has a unique limit cycle in U if and only if $\mu \varepsilon a(\varepsilon, \delta) < 0$ and $\mu = \text{sgn}(\bar{b}_0 \Delta_3(0, \delta_0))$.*

As usual, the set defined by the equation $a(\varepsilon, \delta) = 0$ on the (ε, δ) space is called a Hopf bifurcation curve.

It was proved in [7] that as $f_x + g_y = 0$, then M is C^∞ at $h = 0$. Moreover, M is analytic at $h = 0$ if system (2.1) is analytic. Therefore in the case of $f_x + g_y = 0$ for h small we can write

$$M(h, \delta) = b_0(\delta)h + b_1(\delta)h^2 + b_2(\delta)h^3 + \dots$$

The formulas for b_0 , b_1 , b_2 can be found in [8]. For example, if

$$(f(x, y), g(x, y)) = (H_y(x, y), -H_x(x, y)),$$

$$H(x, y) = h_c + \frac{1}{2}((x - x_c)^2 + (y - y_c)^2) + \sum_{i+j \geq 3} h_{ij}(x - x_c)^i (y - y_c)^j,$$

$$(f_{0x} + g_{0y})|_{\varepsilon=0} = \sum_{i+j \geq 0} c_{ij}(x - x_c)^i (y - y_c)^j,$$

then for $0 < h - h_c \ll 1$ we have

$$M(h, \delta) = \oint_{L_h} (g_0 dx - f_0 dy)|_{\varepsilon=0} = b_0(\delta)(h - h_c) + b_1(\delta)(h - h_c)^2 + \dots, \quad (2.5)$$

where

$$b_0 = 2\pi c_{00}, \quad b_1 = -c_{10}\pi(h_{12} + 3h_{30}) - c_{01}\pi(h_{21} + 3h_{03}) + c_{20}\pi + c_{02}\pi.$$

Hence, by Theorem 2.1 a necessary condition in order that the center $C(x_c, y_c)$ generates a limit cycle is $c_{00}(\delta_0) = 0$ for some $\delta_0 \in G$. If $c_{00}(\delta_0) = 0$ and $b_1(\delta_0) \neq 0$ for some $\delta_0 \in G$, then there exist a constant $\varepsilon_0 > 0$ and a neighborhood U of the center C such that system (2.1) has at most one limit cycle in U for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$. Further, note that in this case $\bar{b}_0 = -1$, $\Delta_3(0, \delta_0) = -b_1/4$ and $a(\varepsilon, \delta) = \varepsilon c_{00}/2 + O(\varepsilon^2)$. If $\delta \in \mathbb{R}$ and $c'_{00}(\delta_0) = \frac{dc_{00}}{d\delta}(\delta_0) \neq 0$, then there exists a function $\varphi_c(\varepsilon) = \delta_0 + O(\varepsilon)$, corresponding to the Hopf bifurcation, such that for $0 < |\varepsilon| < \varepsilon_0$ and $|\delta - \delta_0| < \varepsilon_0$, system (2.1) has a unique limit cycle near the center C if and only if

$$b_1(\delta_0)c'_{00}(\delta_0)(\delta - \varphi_c(\varepsilon)) < 0.$$

Next, we consider the homoclinic bifurcation. Let L be a homoclinic loop defined by the equation $H(x, y) = \beta$. The loop intersects the x -axis transversally at a point A_0 , which is not singular. Then for h near β the periodic orbit L_h intersects the x -axis at a point $Q(h) = (q(h), 0)$ satisfying $Q(h) \rightarrow A_0$ as $h \rightarrow \beta$. For $|\varepsilon|$ small system (2.1) has a unique saddle $\tilde{S}(x_s(\varepsilon, \delta), y_s(\varepsilon, \delta))$ near the saddle S on L . By the normal form theory there exists a polynomial change of variables which carries system (2.1), near \tilde{S} , into a system of the form

$$\begin{aligned} \dot{u} &= \lambda_1(\varepsilon, \delta)u + a_1(\varepsilon, \delta)u^2v + O(u^3v^2), \\ \dot{v} &= -\lambda_2(\varepsilon, \delta)v - b_1(\varepsilon, \delta)uv^2 + O(u^2v^3), \end{aligned}$$

where $(u, v) = (x - x_s, y - y_s)$.

Introduce the following four quantities

$$\begin{aligned} c_0(\delta) &= M(\beta, \delta), \\ c_1(\delta) &= \frac{\partial}{\partial \varepsilon}(\lambda_1 - \lambda_2)|_{\varepsilon=0} = (f_{0x} + g_{0y})|_{\varepsilon=0, (x,y)=S}, \\ c_2(\delta) &= -\frac{1}{\sqrt{f^2(A_0) + g^2(A_0)}} \lim_{h \rightarrow \beta} \frac{M(h, \delta)}{d(Q(h), A_0)}, \\ c_3(\delta) &= \frac{\partial}{\partial \varepsilon}(a_1 - b_1)|_{\varepsilon=0}. \end{aligned}$$

On the existence of at most one, two, or three limit cycles we have the following theorem which was proved in [10].

Theorem 2.3 (Homoclinic bifurcation). *Let $M(h, \delta)$ be the function defined in (2.2). Then the following statements hold.*

(1) *For $|\varepsilon|$ small a necessary condition in order that system (2.1) has a limit cycle near L is $c_0(\delta_0) = 0$ for some $\delta_0 \in G$.*

(2) *If $c_0(\delta_0) = 0$ and $c_1(\delta_0) \neq 0$, then system (2.1) has at most one limit cycle near L for $|\varepsilon|$ and $|\delta - \delta_0|$ both sufficiently small.*

(3) *If $c_0(\delta_0) = c_1(\delta_0) = 0$, then $c_2(\delta_0)$ is finite, and when $c_2(\delta_0) \neq 0$ system (2.1) has at most two limit cycles near L for $|\varepsilon|$ and $|\delta - \delta_0|$ both sufficiently small.*

(4) *If $c_0(\delta_0) = c_1(\delta_0) = c_2(\delta_0) = 0$ and $c_3(\delta_0) \neq 0$, then system (2.1) has at most three limit cycles near L for $|\varepsilon|$ and $|\delta - \delta_0|$ both sufficiently small.*

(5) *If $f_x + g_y \equiv 0$ and $c_0(\delta_0) = c_1(\delta_0) = 0$, then*

$$c_2(\delta_0) = \oint_L (f_{0x} + g_{0y})|_{(\varepsilon, \delta) = (0, \delta_0)} dt.$$

For the existence of a unique limit cycle the authors of [14] obtained the result below. A proof can be found in [10, 8] also.

Theorem 2.4. *Let $\delta \in \mathbb{R}$ and let $c_0(\delta_0) = 0$ for some $\delta_0 \in G$.*

(1) *If $\frac{\partial c_0}{\partial \delta}(\delta_0) \neq 0$, then there exists a unique differentiable function $\delta^*(\varepsilon) = \delta_0 + O(\varepsilon)$ such that for $|\varepsilon| \neq 0$ and $|\delta - \delta_0|$ both small system (2.1) has a homoclinic loop near L if and only if $\delta = \delta^*(\varepsilon)$.*

(2) *If either $c_1(\delta_0) \neq 0$, or $\lambda_1(\varepsilon, \delta) \equiv \lambda_2(\varepsilon, \delta)$ and $c_2(\delta_0) \neq 0$, then for $|\varepsilon| \neq 0$ and $|\delta - \delta_0|$ both small system (2.1) has at most one limit cycle near L . If additionally $\frac{\partial c_0}{\partial \delta}(\delta_0) \neq 0$, then the limit cycle exists if and only if $\mu(\delta - \delta^*(\varepsilon)) > 0$, where*

$$\mu = \begin{cases} \operatorname{sgn}\left(\frac{\partial c_0}{\partial \delta}(\delta_0)c_1(\delta_0)\right), & \text{if } c_1(\delta_0) \neq 0, \\ \operatorname{sgn}\left(\frac{\partial c_0}{\partial \delta}(\delta_0)c_2(\delta_0)\right), & \text{if } c_1(\delta_0) = 0, c_2(\delta_0) \neq 0. \end{cases}$$

As usual, the curve defined by $\delta = \delta^*(\varepsilon)$ is called a homoclinic bifurcation curve of system (2.1).

An useful tool in the proof of the above theorem is a result on the measure of the distance between the two separatrice near L obtained in 1963 by V. Melnikov in [15].

If $f_x + g_y \equiv 0$ and system $(2.1)_{\varepsilon=0}$ is Hamiltonian, then we have for the function $M(h, \delta)$ the following expansion

$$M(h, \delta) = \sum_{k \geq 0} [c_{2k}(\delta)(h - \beta)^k + c_{2k+1}(\delta)(h - \beta)^{k+1} \ln |h - \beta|].$$

It was proved in [16] that system (2.1) has at most k limit cycles near L for $|\varepsilon| \neq 0$ and $|\delta - \delta_0|$ both small if $c_j(\delta_0) = 0$, $j = 0, \dots, k - 1$ and $c_k(\delta_0) \neq 0$.

For the formulas of c_0, c_1, c_2 and c_3 see [11]. For the formulas with more coefficients see the recent work [17].

3 Bogdanov-Takens bifurcation

In this section we use the theorems listed in section 2 to study the Bogdanov-Takens bifurcation. The problem was studied by Bogdanov in 1976 and Takens in 1974 separately. Consider a C^∞ planar differential system of the form

$$\dot{x} = P(x, y, a), \quad \dot{y} = Q(x, y, a), \tag{3.1}$$

where $a \in \mathbb{R}^2$ for simplicity. Let system (3.1) have a nilpotent singular point (x_0, y_0) for a value of a , say a_0 . Without loss of generality we can suppose that $(x_0, y_0) = (0, 0)$, $a_0 = 0$ and that for $a = 0$ we have

$$\frac{\partial(P, Q)}{\partial(x, y)} \Big|_{(x, y) = (0, 0)} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Then it is easy to see that for $|a|$ small system (3.1) can be transformed into

$$\begin{aligned} \dot{x} &= y, \\ \dot{y} &= g(x, a) + f(x, a)y + O(y^2), \end{aligned} \tag{3.2}$$

near the origin, where

$$\begin{aligned} g(x, a) &= g_0(a) + g_1(a)x + g_2(a)x^2 + O(x^3), \\ f(x, a) &= f_0(a) + f_1(a)x + O(x^2), \\ g_0(0) &= g_1(0) = f_0(0) = 0. \end{aligned}$$

For the sake of simplicity assume $f_1(0)g_2(0) \neq 0$. Since $g_2(0) \neq 0$ there exists a unique function $x_0(a) = O(a)$ such that $g_x(x_0(a), a) = 0$. Then for $|x - x_0(a)|$ small we have

$$g(x, a) = g(x_0(a), a) + (g_2(0) + O(a))(x - x_0(a))^2 + O(|x - x_0(a)|^3),$$

$$f(x, a) = \bar{f}_0(a) + (f_1(0) + O(a))(x - x_0(a)) + O(|x - x_0(a)|^2),$$

where

$$g(x_0(a), a) = g_0(a) + O(|a|^2), \quad \bar{f}_0(a) = f(x_0(a), a).$$

Furthermore suppose

$$a = (a_1, a_2) \in R^2, \quad \det \frac{\partial(g_0, \bar{f}_0)}{\partial(a_1, a_2)} \Big|_{a=0} \neq 0. \quad (3.3)$$

This nondegenerate condition ensures that the change of parameters

$$\mu_1 = g(x_0(a), a), \quad \mu_2 = f(x_0(a), a)$$

has a unique inverse $a = a^*(\mu_1, \mu_2)$. Then setting $u = x - x_0(a)$ and using x again for u from (3.2) under the assumption (3.3) we have

$$\begin{aligned} \dot{x} &= y, \\ \dot{y} &= \mu_1 + \mu_2 y + g_2^*(\mu) x^2 + f_1^*(\mu) x y + O(x^3 + x^2 y + y^2), \end{aligned} \quad (3.4)$$

where $\mu = (\mu_1, \mu_2)$ and

$$g_2^*(0) = g_2(0) \neq 0, \quad f_1^*(0) = f_1(0) \neq 0.$$

By introducing suitable rescaling of the variables x , y and t , from system (3.4) we obtain

$$\begin{aligned} \dot{x} &= y \equiv P(x, y, \mu_1, \mu_2), \\ \dot{y} &= \mu_1 + \mu_2 y + x^2 + xy + O(x^3 + x^2 y + y^2) \\ &\equiv Q(x, y, \mu_1, \mu_2). \end{aligned} \quad (3.5)$$

Our aim next is to study the local behavior of system (3.5) near the origin for small values of the parameters μ_1 and μ_2 .

First, noting that

$$\begin{aligned} Q(x, 0, \mu_1, \mu_2) &= \mu_1 + x^2 + O(x^3), \\ \frac{\partial(P, Q)}{\partial(x, y)} &= \begin{pmatrix} 0 & 1 \\ 2x + y + O(x^2 + xy) & \mu_2 + x + O(x^2 + y) \end{pmatrix}, \end{aligned}$$

one can see that system (3.5) has no singular points for $\mu_1 > 0$, and has the origin as a saddle-node for $\mu_1 = 0$ and $\mu_2 \neq 0$. For $\mu_1 < 0$ system (3.5) has a saddle $(\sqrt{-\mu_1} + O(\mu_1), 0)$ and a singular point $A(\mu_1) = (-\sqrt{-\mu_1} + O(\mu_1), 0)$ of index +1. That is, for each $\mu_2 \neq 0$, system (3.5) undergoes a saddle-node bifurcation at $\mu_1 = 0$.

The eigen-equation of system (3.5) at $A(\mu_1)$ has the form

$$\lambda^2 + \lambda(\mu_2 - \sqrt{-\mu_1} + O(\mu_1)) + 2\sqrt{-\mu_1} + O(\mu_1) = 0.$$

Hence there exist two functions

$$\begin{aligned}\varphi_1(\mu_1) &= 2\sqrt{2}(-\mu_1)^{\frac{1}{4}} + (-\mu_1)^{\frac{1}{2}} + O(|\mu_1|^{\frac{3}{4}}), \\ \varphi_2(\mu_1) &= -2\sqrt{2}(-\mu_1)^{\frac{1}{4}} + (-\mu_1)^{\frac{1}{2}} + O(|\mu_1|^{\frac{3}{4}})\end{aligned}$$

such that $A(\mu_1)$ is a focus for $\varphi_2(\mu_1) < \mu_2 < \varphi_1(\mu_1)$, a node for $\mu_2 > \varphi_1(\mu_1)$ or $\mu_2 < \varphi_2(\mu_1)$, a degenerate node for $\mu_2 = \varphi_1(\mu_1)$ or $\mu_2 = \varphi_2(\mu_1)$. Obviously, there is a function $\varphi(\mu_1) = \sqrt{-\mu_1} + O(|\mu_1|)$, such that $A(\mu_1)$ is stable for $\mu_2 < \varphi(\mu_1)$, unstable for $\mu_2 > \varphi(\mu_1)$. This implies that a possible Hopf bifurcation would occur at $A(\mu_1)$ when μ_2 varies near $\varphi(\mu_1)$.

To study the problem in detail we introduce the rescaling change of variables

$$\mu_2 = -\delta|\mu_1|^{\frac{1}{2}}, \quad x = -u|\mu_1|^{\frac{1}{2}}, \quad y = v|\mu_1|^{\frac{3}{4}}, \quad t = -\tau|\mu_1|^{-\frac{1}{4}},$$

and then using (x, y, t) for (u, v, τ) from system (3.5) we obtain

$$\begin{aligned}\dot{x} &= y, \\ \dot{y} &= 1 - x^2 + \tilde{\varepsilon}(\delta + x)y + O(\tilde{\varepsilon}^2),\end{aligned}$$

where $\tilde{\varepsilon} = |\mu_1|^{\frac{1}{4}}$. Furthermore letting

$$u = \frac{1}{2}(x + 1), \quad v = \frac{1}{2\sqrt{2}}y, \quad \tau = \sqrt{2}t,$$

and also using (x, y, t) for (u, v, τ) , we get the system

$$\begin{aligned}\dot{x} &= y, \\ \dot{y} &= x(1 - x) + \varepsilon(\lambda + x)y + O(\varepsilon^2),\end{aligned}\tag{3.6}$$

where $\lambda = \frac{1}{2}(\delta - 1)$ and $\varepsilon = \sqrt{2}\tilde{\varepsilon}$. The unperturbed system of system (3.6) is Hamiltonian with Hamiltonian

$$H(x, y) = \frac{1}{2}y^2 - \frac{1}{2}x^2 + \frac{1}{3}x^3,$$

whose level curves define a family of closed orbits L_h for $h \in (-\frac{1}{6}, 0)$. The boundary of this family consists of a center at $(1, 0)$ and a homoclinic loop L_0 passing through a saddle at the origin. The first order Melnikov function of system (3.6) has the form

$$M(h, \lambda) = \oint_{L_h} (\lambda + x)y dx = \lambda I_0(h) + I_1(h), \quad h \in (-\frac{1}{6}, 0),$$

where

$$I_j(h) = \oint_{L_h} x^j y dx, \quad j = 0, 1.$$

Direct computations show that $I_0(0) = 6/5$ and $I_1(0) = 36/35$. Thus

$$M(0, \lambda) = \frac{6}{5}\lambda + \frac{36}{35} = \frac{6}{5} \left(\lambda + \frac{6}{7} \right).$$

Then by Theorem 2.4 we immediately have:

Lemma 3.1 *There exist a constant $\epsilon_0 > 0$ and a C^1 function $\lambda^*(\epsilon) = \lambda_0 + O(\epsilon)$ with $\lambda_0 = -6/7$, such that for $0 < \epsilon \leq \epsilon_0$ and $|\lambda - \lambda_0| \leq \epsilon_0$, system (3.6) has a unique limit cycle (no limit cycle, respectively) near L_0 if $\lambda < \lambda^*(\epsilon)$ ($\lambda > \lambda^*(\epsilon)$, respectively), and has a homoclinic loop near L_0 if and only if $\lambda = \lambda^*(\epsilon)$. Moreover, the limit cycle or the homoclinic loop is stable.*

Rewriting $H(x, y)$ as

$$H(x, y) = \frac{1}{2}(y^2 + (x - 1)^2) + \frac{1}{3}(x - 1)^3,$$

then from (2.5) we get

$$M(h, \lambda) = 2\pi(\lambda + 1) \left(h + \frac{1}{6} \right) - \pi \left(h + \frac{1}{6} \right)^2 + \dots.$$

Therefore, from Theorem 2.2 and Corollary 2.1 we obtain:

Lemma 3.2 *There exist a constant $\epsilon_1 > 0$ and a C^1 function $\lambda_1^*(\epsilon) = -1 + O(\epsilon)$, such that for $0 < \epsilon \leq \epsilon_1$ and $|\lambda + 1| \leq \epsilon_1$ system (3.6) has a unique limit cycle near the center $(1, 0)$ if and only if $\lambda > \lambda_1^*(\epsilon)$. Moreover, the limit cycle is stable when it exists.*

Now our next step is to prove that the limit cycle of the above two lemmas exists uniquely for all $\lambda_1^*(\epsilon) < \lambda < \lambda^*(\epsilon)$. For this purpose we rewrite the function $M(h, \lambda)$ as

$$M(h, \lambda) = I_0(h)(\lambda - P(h)), \quad P(h) = -I_1(h)/I_0(h).$$

The following lemma provides the properties of the function $P(h)$ due to Bogdanov [1].

Lemma 3.3 *The function $P(h)$ is continuous on $[-\frac{1}{6}, 0]$, differentiable on $[-\frac{1}{6}, 0)$ with $P(-\frac{1}{6}) = -1$, $P(0) = -\frac{6}{7}$, $P'(-\frac{1}{6}) = \frac{1}{2}$ and $P'(h) > 0$ for $(-\frac{1}{6}, 0)$.*

Proof From the proofs of Lemmas 3.1 and 3.2 we have $P(0) = -\frac{6}{7}$ and $P(-\frac{1}{6}) = -1$. Note that along L_h we get

$$y dy = x dx - x^2 dx, \quad \frac{\partial y}{\partial h} = \frac{1}{y}.$$

By writing $I_1(h)$ into the form of definite integral in x we find

$$I_1'(h) = \oint_{L_h} \frac{x dx}{y} = \oint_{L_h} \frac{x dx - y dy}{y} = \oint_{L_h} \frac{x^2 dx}{y}.$$

Then by integration by parts we have

$$\begin{aligned}
I_0(h) &= - \oint_{L_h} x dy = - \oint_{L_h} \frac{x^2 dx - x^3 dx}{y} \\
&= \oint_{L_h} \frac{1}{y} (3h + \frac{3}{2}x^2 - \frac{3}{2}y^2 - x^2) dx \\
&= 3hI'_0(h) + \frac{1}{2}I'_1(h) - \frac{3}{2}I_0(h).
\end{aligned}$$

Hence

$$5I_0 = 6hI'_0 + I'_1. \quad (3.7)$$

Similarly, integrating by parts the function I_1 we get

$$35I_1 = 6hI'_0 + 6(1 + 5h)I'_1. \quad (3.8)$$

Solving I'_0 and I'_1 from (3.7) and (3.8) yields to the so-called Picard-Fuchs equations as follows

$$I'_1 = \frac{1}{1 + 6h}(7I_1 - I_0), \quad I'_0 = \frac{1}{6h(1 + 6h)}[6(1 + 5h)I_0 - 7I_1].$$

Therefore we obtain

$$\frac{dP}{dh} = \frac{-1}{I_0^2}[I_0I'_1 - I_1I'_0] = \frac{-7P^2 + 6(2h - 1)P + 6h}{6h(1 + 6h)}.$$

This means that the function $P = P(h)$ defines an orbit connecting the saddle $(-\frac{1}{6}, -1)$ and the node $(0, -\frac{6}{7})$ in the plane of the quadratic system

$$\begin{aligned}
\dot{h} &= 6h(1 + 6h), \\
\dot{P} &= -7P^2 + 6(2h - 1)P + 6h.
\end{aligned} \quad (3.9)$$

It follows that the function $P(h)$ is continuously differentiable on the interval $[-\frac{1}{6}, 0)$.

Consider the isocline equation

$$-7P^2 + 6(2h - 1)P + 6h = 0.$$

It is a hyperbola with two branches L_1 and L_2 with L_1 lying above L_2 . It is easy to see that L_1 passes through the points $(0, 0)$ and $(-\frac{1}{6}, -\frac{1}{7})$, while L_2 passes through the points $(0, -\frac{6}{7})$ and $(-\frac{1}{6}, -1)$. Note that for $-\frac{1}{6} < h < 0$ we have

$$\begin{aligned}
\dot{h} &= 6h(1 + 6h) < 0, \\
\dot{P} &= -7P^2 + 6(2h - 1)P + 6h \begin{cases} > 0 & \text{between } L_1 \text{ and } L_2, \\ = 0 & \text{on } L_1 \text{ or } L_2, \\ < 0 & \text{above } L_1 \text{ or below } L_2. \end{cases}
\end{aligned}$$

It implies that the curve $P = P(h)$ is below L_2 for $-\frac{1}{6} < h < 0$. Thus $P'(h) = \frac{dP}{dh} = \frac{\dot{P}}{h} > 0$. Furthermore the linear variational equation of system (3.9) at the saddle $(-\frac{1}{6}, -1)$ is

$$\dot{x} = -6x, \quad \dot{y} = -6x + 6y,$$

which has the separatrix $y = \frac{1}{2}x$. Hence $P'(-\frac{1}{6}) = \frac{1}{2}$. This completes the proof.

On the other hand, the variational equation of system (3.9) at the degenerate node $(0, -\frac{6}{7})$ has the form

$$\dot{x} = 6x, \quad \dot{y} = -\frac{30}{7}x + 6y,$$

from which we get $P'(0 - 0) = \infty$.

Noting that $0 \leq x \leq 3/2$ and $|y| \leq 1/\sqrt{3}$ along the homoclinic loop L_0 we can suppose in system (3.6) that $-\varepsilon^* \leq x \leq 3/2 + \varepsilon^*$, $|y| \leq 1/\sqrt{3} + \varepsilon^*$ and $0 < \varepsilon < \varepsilon^*$ for a small constant $\varepsilon^* > 0$. In this case there exists a constant $N > 0$ such that the divergence of (3.6) is positive or negative (and hence there is no limit cycle) if $|\lambda| > N$ and $0 < \varepsilon < \varepsilon^*$. For example, we can take $N = 3$ for a suitable ε^* since

$$|\operatorname{div}(3.6)| = |\varepsilon(\lambda + x + O(\varepsilon))| > \varepsilon(|\lambda| - 2 + O(\varepsilon)).$$

That is, in system (3.6) we can assume $|\lambda| \leq N$ to discuss the existence of limit cycles. With this assumption and applying Lemmas 3.1-3.3 we can prove the following theorem.

Theorem 3.1 *There exist a constant $\varepsilon^* > 0$ and two differentiable functions $\lambda^*(\varepsilon) = -\frac{6}{7} + O(\varepsilon)$ and $\lambda_1^*(\varepsilon) = -1 + O(\varepsilon)$ such that for all $0 < \varepsilon < \varepsilon^*$ and $|\lambda| \leq N$, system (3.6) has*

- (1) *no limit cycles for $\lambda > \lambda^*(\varepsilon)$ or $\lambda \leq \lambda_1^*(\varepsilon)$;*
- (2) *a stable homoclinic loop near L_0 for $\lambda = \lambda^*(\varepsilon)$;*
- (3) *a unique (stable) limit cycle for $\lambda_1^*(\varepsilon) < \lambda < \lambda^*(\varepsilon)$.*

Proof. Correspondingly we need only to prove the following three statements:

(i) There exists a constant $\varepsilon_1^* > 0$, such that for $0 < \varepsilon < \varepsilon_1^*$ system (3.6) has no limit cycles when either $\lambda^*(\varepsilon) < \lambda \leq N$ or $-N \leq \lambda \leq \lambda_1^*(\varepsilon)$.

(ii) There exists a constant $\varepsilon_2^* > 0$ such that for $0 < \varepsilon < \varepsilon_2^*$ and $\lambda = \lambda^*(\varepsilon)$ system (3.6) has a stable homoclinic loop near L_0 .

(iii) There exists a constant $\varepsilon_3^* > 0$ such that for $0 < \varepsilon < \varepsilon_3^*$ and $\lambda_1^*(\varepsilon) < \lambda < \lambda^*(\varepsilon)$ system (3.6) has a unique limit cycle, and it is stable.

When the three statements are proved, we can take $\varepsilon^* = \min\{\varepsilon_1^*, \varepsilon_2^*, \varepsilon_3^*\}$ to complete the proof.

We shall prove the statements by contradiction. If statement (i) is not true, then there exist sequences ε_n and λ_n satisfying $\varepsilon_n > 0$ and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and $\lambda^*(\varepsilon_n) < \lambda_n \leq N$, or $-N \leq \lambda_n \leq \lambda_1^*(\varepsilon_n)$ such that for $(\varepsilon, \lambda) = (\varepsilon_n, \lambda_n)$ system (3.6) has a limit cycle L_n^* . Note that $\{\lambda_n\}$ and $\{L_n^*\}$ are all bounded for all n . We can suppose that $\lambda_n \rightarrow \lambda^*$ and $L_n^* \rightarrow \Gamma^*$ (as $n \rightarrow +\infty$).

Clearly $-\frac{6}{7} \leq \lambda^* \leq N$, or $-N \leq \lambda^* \leq -1$, and Γ^* should be a closed invariant curve of system $(3.6)_{\varepsilon=0}$. This means that Γ^* is defined by an equation $H(x, y) = h^*$ for some $h^* \in [-\frac{1}{6}, 0]$. By Theorems 3.1-3.3 we have necessarily $M(h^*, \lambda^*) = 0$, or $\lambda^* = P(h^*)$. By Lemma 3.3 one obtains $-1 \leq \lambda^* \leq -\frac{6}{7}$. Then by Lemmas 3.1 and 3.2 we have either $\lambda^* = -\frac{6}{7}$ and Γ^* should be the loop L_0 , or $\lambda^* = -1$ and Γ^* should be the center $(1, 0)$. In the first case we have $-\frac{6}{7} - \epsilon_0 < \lambda_n < -\frac{6}{7} + \epsilon_0$ and $0 < \varepsilon_n < \epsilon_0$ for n sufficiently large, which, together with Lemma 3.1, implies that for $(\varepsilon, \lambda) = (\varepsilon_n, \lambda_n)$ system (3.6) has no limit cycles near L_0 because $\lambda_n > \lambda^*(\varepsilon_n)$, a contradiction. In the second case a contradiction occurs similarly by Lemma 3.2. Then statement (i) follows.

Statement (ii) is direct from Lemma 3.1.

For statement (iii), first by Poincaré-Bendixson theorem one can see that for $\lambda_1^*(\varepsilon) < \lambda < \lambda^*(\varepsilon)$ with $\varepsilon > 0$ small system (3.6) has a limit cycle. We shall prove the uniqueness of the limit cycle by reduction to a contradiction. That is, we suppose that the uniqueness conclusion is not true. Then there exists sequences $\varepsilon_n > 0$ and $\lambda_n > 0$ satisfying $\varepsilon_n \rightarrow 0$ and $\lambda_1^*(\varepsilon_n) < \lambda_n < \lambda^*(\varepsilon_n)$ such that for $(\varepsilon, \lambda) = (\varepsilon_n, \lambda_n)$ system (3.6) has two limit cycles $L_n^{(1)}$ and $L_n^{(2)}$. As before we can assume that as $n \rightarrow \infty$ we have $\lambda_n \rightarrow \lambda^*$ with $-1 \leq \lambda^* \leq -\frac{6}{7}$, and $L_n^{(j)} \rightarrow \Gamma_j$ for $j = 1, 2$. Then Γ_j satisfies $H(x, y) = h_j$ for some $h_j \in [-\frac{1}{6}, 0]$, $j = 1, 2$. Hence, $M(h_j, \lambda^*) = 0$, and then $\lambda^* = P(h_1) = P(h_2)$. Thus $h_1 = h_2$ by Lemma 3.3. Therefore, a contradiction must occur in the same way than in Lemma 3.2 for $h_1 = -1$, Lemma 3.1 for $h_1 = -6/7$ or Lemma 3.2 and Theorem 1.1 for $h_1 \in (-1, -6/7)$. This ends the proof of Theorem 3.1.

From the above proof, one can see that each one of Lemmas 3.1-3.3 is indispensable and plays its own role in the proof. In other words any one of them is not sufficient to deduce the statements of Theorem 3.1.

Note that $\mu_2 = -\delta|\mu_1|^{\frac{1}{2}} = -(2\lambda + 1)|\mu_1|^{\frac{1}{2}}$, $\varepsilon = \sqrt{2}|\mu_1|^{\frac{1}{4}}$. Back to system (3.5) and we have from Theorem 3.1 the following result.

Theorem 3.2 *There exist a constant $\epsilon_0 > 0$, a neighborhood U of the origin and two continuous functions $\psi_1(\mu_1) = (-\mu_1)^{\frac{1}{2}} + O(|\mu_1|^{\frac{3}{4}})$ (corresponding to Hopf bifurcation) and $\psi_2(\mu_1) = \frac{5}{7}(-\mu_1)^{\frac{1}{2}} + O(|\mu_1|^{\frac{3}{4}})$ (corresponding to homoclinic bifurcation), such that for*

$0 \leq -\mu_1 \leq \epsilon_0$ and $|\mu_2| \leq \epsilon_0$, system (3.5) has a unique limit cycle in U if and only if $\psi_2(\mu_1) < \mu_2 < \psi_1(\mu_1)$.

4 Some examples with two limit cycles

In this section we give two examples which have two limit cycles. However the first order Melnikov function of them has only one nontrivial zero. The first example is a system of Liénard type

$$\dot{x} = y - \varepsilon(\varepsilon x - x^3 + x^5), \quad \dot{y} = -x, \quad (4.1)$$

where $\varepsilon > 0$ is small. The unperturbed system of it is a linear center with the first integral $H(x, y) = x^2 + y^2$. Let L_h denote the circle defined by $x^2 + y^2 = h$ with $h > 0$. Then by (2.2) the first order Melnikov function is

$$M(h) = \oint_{L_h} (-x^3 + x^5) dy.$$

Note that L_h can be represented as $x = \sqrt{h} \sin t$, $y = \sqrt{h} \cos t$. A direct computation shows that

$$\begin{aligned} M(h) &= \int_0^{2\pi} [h^2 \sin^4 t - h^3 \sin^6 t] dt \\ &= h^2 [\pi/2 + \frac{1}{4} \int_0^{2\pi} \cos^2(2t) dt] - h^3 [\pi/4 + \frac{3}{8} \int_0^{2\pi} \cos^2(2t) dt] \\ &= \frac{3\pi}{4} h^2 (1 - 5h/6), \end{aligned}$$

which has a unique simple positive zero $h = 6/5$. It follows from Theorem 2.1 that for $\varepsilon > 0$ small system (4.1) has a unique limit cycle $\Gamma_{1\varepsilon}$ near the circle $x^2 + y^2 = 6/5$.

Noting that $\text{div}(4.1) = \varepsilon(3x^2 - 5x^4 - \varepsilon)$, we have

$$\oint_{\Gamma_{1\varepsilon}} \text{div}(4.1) dt = \varepsilon[-9\pi/5 + O(\varepsilon)] < 0,$$

for $0 < \varepsilon \ll 1$. This implies that the limit cycle $\Gamma_{1\varepsilon}$ is stable.

On the other hand, the origin is a stable focus for $0 < \varepsilon \ll 1$ since

$$\text{div}(4.1)|_{(x,y)=(0,0)} = -\varepsilon^2.$$

Therefore there must have an unstable limit cycle, denoted by $\Gamma_{2\varepsilon}$, inside $\Gamma_{1\varepsilon}$. By Theorems 2.1 and 2.2 we obtain that $\Gamma_{1\varepsilon}$ and $\Gamma_{2\varepsilon}$ are the only limit cycles of system (4.1).

From the above discussion one can see that the following system

$$\dot{x} = y - \varepsilon(\varepsilon x - x^3), \quad \dot{y} = -x,$$

has a unique limit cycle Γ_ε for $0 < \varepsilon \ll 1$. However its first order Melnikov function is $M(h) = \frac{3\pi}{4}h^2$, which has no positive zero. This fact implies that for any $h_0 > 0$ there exists $\varepsilon_0 > 0$ such that for $0 < \varepsilon < \varepsilon_0$ all orbits starting from a point in the region $x^2 + y^2 > h_0$ are unbounded positively. And, by Theorems 2.1 and 2.2 the limit cycle Γ_ε is unique and approaches the origin as $\varepsilon \rightarrow 0$.

Our second example is

$$\begin{aligned} \dot{x} &= y + \varepsilon x[\varepsilon + x(a - H(x, y))], \\ \dot{y} &= x - x^2, \end{aligned} \tag{4.2}$$

where

$$H(x, y) = \frac{1}{2}(y^2 - x^2) + \frac{1}{3}x^3.$$

The unperturbed system of system (4.2) has a saddle at the origin, a center at $(1, 0)$, and a family of closed orbits given by

$$L_h : H(x, y) = h, \quad h \in \left(-\frac{1}{6}, 0\right).$$

The limit of L_h as $h \rightarrow 0$, denoted by L_0 , is a homoclinic loop satisfying $H(x, y) = 0$. By (2.2) the first order Melnikov function has the form

$$M(h, a) = I(h)[a - h], \tag{4.3}$$

where

$$I(h) = \oint_{L_h} (-x^2)dy = 2 \oint_{L_h} xydx.$$

Clearly $I(-\frac{1}{6}) = 0$. Noting that $yy_h = 1$ along L_h , one has

$$I'(h) = \oint_{L_h} \frac{2x}{y}dx = \oint_{L_h} 2xdt > 0, \tag{4.4}$$

which yields $I(h) > 0$ for all $h \in (-\frac{1}{6}, 0)$. Hence $M(h, a)$ has at most one zero on the interval $(-\frac{1}{6}, 0)$. From (4.4) one also has $I'(-\frac{1}{6}) = 4\pi$, from which it follows

$$I(h) = 4\pi\left(h + \frac{1}{6}\right) + O\left(\left|h + \frac{1}{6}\right|^2\right),$$

for $h + \frac{1}{6} > 0$ small. Hence by (4.3) we obtain

$$M(h, a) = 4\pi\left(h + \frac{1}{6}\right)\left[a + \frac{1}{6} - \left(1 + K\left(a + \frac{1}{6}\right)\right)\left(h + \frac{1}{6}\right)\right] + O\left(\left|h + \frac{1}{6}\right|^3\right),$$

for $h + \frac{1}{6} > 0$ small, where K is a constant. By Corollary 2.1 there exists a function $\varphi_c(\varepsilon) = -\frac{1}{6} + O(\varepsilon)$ such that for $\varepsilon + |a + \frac{1}{6}|$ small system (4.2) has a unique limit cycle, say $\Gamma_{\varepsilon,a}$, in a neighborhood of the center $(1, 0)$ if and only if $a > \varphi_c(\varepsilon)$. On the (ε, a) -plane the curve $a = \varphi_c(\varepsilon)$ defines a Hopf bifurcation curve. Note that by (2.5)

$$\begin{aligned} \operatorname{div}(4.2)|_C &= \varepsilon[c_{00} + O(\varepsilon)] \\ &= \varepsilon\left[\frac{b_0}{2\pi} + O(\varepsilon)\right] \\ &= \varepsilon[2(a + 1/6) + O(\varepsilon)], \end{aligned}$$

where C denotes the focus of system (4.2) near the center $(1,0)$. Hence

$$\operatorname{div}(4.2)|_C > 0 \Leftrightarrow a > \varphi_c(\varepsilon). \quad (4.5)$$

Thus the limit cycle $\Gamma_{\varepsilon,a}$ is stable.

Now we consider the problem: How does the stable limit cycle $\Gamma_{\varepsilon,a}$ vary when the parameter a increases from $\varphi_c(\varepsilon)$?

To understand the behavior of $\Gamma_{\varepsilon,a}$ as a increases, let $a \in (-\frac{1}{6}, 0)$. In this case the function $M(h, a)$ has a unique simple zero $h = a$ for $h \in (-\frac{1}{6}, 0)$. This implies that for $\varepsilon > 0$ small system (4.2) has a unique limit cycle near L_a . Along the limit cycle we have

$$\begin{aligned} \oint \operatorname{div}(4.2)dt &= \varepsilon\left[\oint_{L_a} x^2(x - x^2)dt + O(\varepsilon)\right] \\ &= -\varepsilon[I(a) + O(\varepsilon)] \\ &< 0 \end{aligned}$$

for $\varepsilon > 0$ small. Hence the limit cycle is stable. Then it is clear that when a increases from $\varphi_c(\varepsilon)$ the stable limit cycle $\Gamma_{\varepsilon,a}$ enlarges constantly, and it approaches to L_a as $\varepsilon \rightarrow 0$.

We consider an additional problem: what happens for a near zero?

First noting that $M(0, a) = I(0)a = \frac{72}{35}a$, by Theorem 2.4(1) and its proof there exists a function $\varphi_h(\varepsilon) = O(\varepsilon)$ such that (4.2) has a homoclinic loop Γ_ε^* if and only if $a = \varphi_h(\varepsilon)$. Moreover for a near $\varphi_h(\varepsilon)$ the orbit behavior of system (4.2) near L_0 is as shown in Fig.4.1.

For $\operatorname{div}(4.2)|_{x=y=0, a=\varphi_h(\varepsilon)} = \varepsilon^2$ the homoclinic loop Γ_ε^* is unstable. On the other hand, by (4.5) the focus C is unstable for $\varphi_c(\varepsilon) < a \leq \varphi_h(\varepsilon)$. Then it follows that system (4.2) has a stable limit cycle, denoted by $\bar{\Gamma}_\varepsilon$, inside the loop Γ_ε^* for $a = \varphi_h(\varepsilon)$. In the same way, from Fig. 4.1, the stable limit cycle $\Gamma_{\varepsilon,a}$ obtained before always exists for




Fig. 4.1. Phase portraits of system (4.2) near L_0 .




Fig. 4.2. Global behavior of system (4.2).

$\varphi_c(\varepsilon) < a < \varphi_h(\varepsilon)$. Furthermore, for $0 < a - \varphi_h(\varepsilon) \ll 1$ the homoclinic loop Γ_ε^* has broken and generated an unstable limit cycle $\Gamma_{\varepsilon,a}^*$, see Fig. 4.2.

By Theorem 2.3, $\Gamma_{\varepsilon,a}$ and $\Gamma_{\varepsilon,a}^*$ are the only limit cycles of system (4.2).

Finally, for any given constant $a_0 > 0$, we have $M(h, a) > 0$ for all $h \in [-\frac{1}{6}, 0]$ and $a \geq a_0$. Hence there exists $\varepsilon_0 = \varepsilon_0(a_0) > 0$ such that for $0 < \varepsilon < \varepsilon_0$ and $a \geq a_0$ system (4.2) has no limit cycles. Thus there exists a function $\varphi_{sn}(\varepsilon) = O(\varepsilon) > \varphi_h(\varepsilon)$ such that for $a = \varphi_{sn}(\varepsilon)$ system (4.2) has a semi-stable limit cycle which is the limit of $\Gamma_{\varepsilon,a}$ and $\Gamma_{\varepsilon,a}^*$ when $a \rightarrow \varphi_{sn}(\varepsilon)$ from left. The function $\varphi_{sn}(\varepsilon)$ corresponds to the saddle-node bifurcation of limit cycles in system (4.2). Therefore we have three bifurcation curves on (ε, a) -plane: Hopf bifurcation $a = \varphi_c(\varepsilon)$, homoclinic bifurcation $a = \varphi_h(\varepsilon)$ and saddle-node bifurcation of limit cycles $a = \varphi_{sn}(\varepsilon)$. The bifurcation diagram is shown in Fig. 4.3.

We have seen from examples (4.1) and (4.2) that the study of the uniqueness of limit




Fig. 4.3. Bifurcation diagram of limit cycles of system (4.2).

cycles near the boundary of the period annulus in the Bogdanov-Takens bifurcation is necessary and non-trivial.

Acknowledgments

The first and the third authors are partially supported by National Natural Sciences Foundation 11431008, 11771296 and 11571090. The second author is partially supported by the MINECO grants MTM2013-40998-P and MTM2016-77278-P (FEDER) and the AGAUR grant 2014-SGR568.

References

- [1] R.I. Bogdanov, The versal deformation of a singular point of a vector field on the plane in the case of zero eigenvalues, in "Proceedings of Petrovskii Seminar, Vol. 2, Moscow State University, Moscow, 37-65". In Russian (English translation: Selecta Math. Soviet., 1(1981), 389-421).
- [2] J. Carr, Applications of Center Manifold Theory, Springer-Verlag, New York, 1982.
- [3] S.N. Chow, J.K. Hale, Methods of Bifurcation Theory, Springer-Verlag, New York, 1996.
- [4] S-N. Chow, C. Li, D. Wang, Normal Forms and Bifurcations of Planar Vector Fields (Cambridge University Press), 1994.

- [5] J. Guckenheimer, P. Holmes, *Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields*, Springer-Verlag, New York, 1983.
- [6] M. Han, Bifurcation of limit cycles and the cusp of order n , *Acta Mathematica Sinica, New Series* 13(1997), 64-75.
- [7] M. Han, On Hopf cyclicity of planar systems, *Journal of Mathematical Analysis and Applications* 245(2000), 404-422.
- [8] M. Han, *Bifurcation Theory of Limit Cycles*, Science Press, Beijing, 2012.
- [9] M. Han, S. Gu, *Theory and Methods of Nonlinear Systems*, Science Press, Beijing, 2001 (In Chinese).
- [10] M. Han, D. Zhu, *Bifurcation Theory of Ordinary Differential Equations*, Coal Mine Industry Publishing House, Beijing, 1994 (In Chinese).
- [11] M. Han, J. Yang, A. Tarta, Y. Gao, Limit cycles near homoclinic and heteroclinic loops, *J. Dynamics and Differential Equations* 20(2008), 923-944.
- [12] J. Li, Hilbert's 16th problem and bifurcations of planar polynomial vector fields. *Int. J. Bifurcation Chaos* 13(2003), 47-106.
- [13] Yuri A. Kuznetsov, *Elements of Applied Bifurcation Theory*, Springer-Verlag New York, 1995, 1998, 2004.
- [14] D. Luo, M. Han, D. Zhu, Uniqueness of limit cycles bifurcated from singular closed orbits (I), *Acta Math. Sinica* 3(1992), 404-414.
- [15] V. K. Melnikov, On the stability of the center for time periodic perturbations, *Trans. Moscow Math. Soc.*, 12(1963), 1-57.
- [16] R. Roussarie, On the number of limit cycles which appear by perturbation of separatrix loop of planar vector fields, *Bol. Soc. Bras. Mat.* 17(1986), 67-101.
- [17] Y. Tian, M. Han, Hopf and homoclinic bifurcations for near-Hamiltonian systems, *J. Differential Equations* 262(2017), 3214-3234.
- [18] Y. Ye, et al., *Theory of limit cycles*, Shanghai Science and Technology Publishing House, Shanghai 1984(In Chinese), *Transl. Math. Monographs*, Vol. 66, Amer. Math. Soc., Providence RI, 1986.

- [19] Z. Zhang, B. Li, High order Melnikov functions and the problem of uniformity in global bifurcation, *Annali di Mat. Pura ed Appl. (IV)*, vol. CLX(1992), 181-212.
- [20] Z. Zhang, C. Li, Z. Zheng, W. Li, *Elementary Theory of Bifurcations of Vector Fields* (Higher Education Press, Beijing), 1997 (in Chinese).
- [21] A. Zhao, M. Li, M. Han *Fundamental Theory of Differential Equations* Science Press, Beijing 2011 (In Chinese).