



## **An in Vivo Mouse Model to Investigate the Effect of Local Anesthetic Nanomedicines on Axonal Conduction and Excitability**

**Moldovan, Mihai; Alvarez, Susana; Rothe, Christian; Andresen, Thomas Lars; Urquhart, Andrew; Lange, Kai H. W.; Krarup, Christian**

*Published in:*  
Frontiers in Neuroscience

*Link to article, DOI:*  
[10.3389/fnins.2018.00494](https://doi.org/10.3389/fnins.2018.00494)

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Moldovan, M., Alvarez, S., Rothe, C., Andresen, T. L., Urquhart, A., Lange, K. H. W., & Krarup, C. (2018). An in Vivo Mouse Model to Investigate the Effect of Local Anesthetic Nanomedicines on Axonal Conduction and Excitability. *Frontiers in Neuroscience*, 12, [494]. DOI: 10.3389/fnins.2018.00494

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# An *in Vivo* Mouse Model to Investigate the Effect of Local Anesthetic Nanomedicines on Axonal Conduction and Excitability

Mihai Moldovan<sup>1,2\*</sup>, Susana Alvarez<sup>1</sup>, Christian Rothe<sup>3</sup>, Thomas L. Andresen<sup>4</sup>, Andrew Urquhart<sup>4</sup>, Kai H. W. Lange<sup>3</sup> and Christian Krarup<sup>1,2</sup>

<sup>1</sup> Department of Neuroscience, University of Copenhagen, Copenhagen, Denmark, <sup>2</sup> Department of Clinical Neurophysiology, Rigshospitalet, Copenhagen, Denmark, <sup>3</sup> Department of Anesthesia, Nordsjællands Hospital, Hillerød, Denmark, <sup>4</sup> Department for Micro- and Nanotechnology, Technical University of Denmark, Lyngby, Denmark

## OPEN ACCESS

### Edited by:

Ioan Opris,  
University of Miami, United States

### Reviewed by:

Angelika Lampert,  
Uniklinik RWTH Aachen, Germany  
Mario Valentino,  
University of Malta, Malta

### \*Correspondence:

Mihai Moldovan  
moldovan@sund.ku.dk

### Specialty section:

This article was submitted to  
Neural Technology,  
a section of the journal  
Frontiers in Neuroscience

**Received:** 20 April 2018

**Accepted:** 02 July 2018

**Published:** 26 July 2018

### Citation:

Moldovan M, Alvarez S, Rothe C, Andresen TL, Urquhart A, Lange KHW and Krarup C (2018) An *in Vivo* Mouse Model to Investigate the Effect of Local Anesthetic Nanomedicines on Axonal Conduction and Excitability. *Front. Neurosci.* 12:494. doi: 10.3389/fnins.2018.00494

Peripheral nerve blocks (PNBs) using local anesthetic (LA) are superior to systemic analgesia for management of post-operative pain. An insufficiently short PNB duration following single-shot LA can be optimized by development of extended release formulations among which liposomes have been shown to be the least toxic. *In vivo* rodent models for PNB have focused primarily on assessing behavioral responses following LA. In a previous study in human volunteers, we found that it is feasible to monitor the effect of LA *in vivo* by combining conventional conduction studies with nerve excitability studies. Here, we aimed to develop a mouse model where the same neurophysiological techniques can be used to investigate liposomal formulations of LA *in vivo*. To challenge the validity of the model, we tested the motor PNB following an unilamellar liposomal formulation, filled with the intermediate-duration LA lidocaine. Experiments were carried out in adult transgenic mice with fluorescent axons and with fluorescent tagged liposomes to allow *in vivo* imaging by probe-based confocal laser endomicroscopy. Recovery of conduction following LA injection at the ankle was monitored by stimulation of the tibial nerve fibers at the sciatic notch and recording of the plantar compound motor action potential (CMAP). We detected a delayed recovery in CMAP amplitude following liposomal lidocaine, without detrimental systemic effects. Furthermore, CMAP threshold-tracking studies of the distal tibial nerve showed that the increased rheobase was associated with a sequence of excitability changes similar to those found following non-encapsulated lidocaine PNB in humans, further supporting the translational value of the model.

**Keywords:** liposomes, peripheral nerve block, threshold-tracking, *in vivo* imaging, lidocaine

## INTRODUCTION

Adequate pain management has been shown to improve the rate and quality of patient recovery following surgery (Zaslansky et al., 2015). Peripheral nerve blocks (PNBs) using local anesthetic (LA) are superior to opioid analgesia which have a high rate of undesirable systemic effects (Wu and Raja, 2011; Aguirre et al., 2012). Voltage-gated Na<sup>+</sup> channel (VGSC) blockers acting as LA (e.g.,

bupivacaine, lidocaine, etc.) are the most widely used drugs in PNBs. Single PNB only lasts several hours (Wu and Raja, 2011). Catheter-based PNBs using LA infusions can extend analgesia (Madsen et al., 2018); however, their efficiency is limited by mechanical factors and can expose patients to large quantities of LA (Ilfeld, 2017). Alternatively, several LA carriers with extended release have been developed, among which liposomes have been shown to be the least toxic (Weiniger et al., 2012). A multilamellar liposome formulation of bupivacaine (Exparel) has so far failed to show PNB of major nerves (Hadzic et al., 2016) and remains of limited indication (Vyas et al., 2016). Further experimental work is required to understand the *in vivo* effects of liposomal LA nanomedicines.

*In vivo* rodent models for PNB have focused primarily on assessing behavioral responses following LA such as the motor performance on an inverted mesh (Leszczynska and Kau, 1992), the tail flick latency to a thermal stimulus (Grant et al., 1993), or the vocalization responses to an electrical shock (Grant et al., 2000). Such sensory-motor behavioral techniques were successfully used to investigate the protracted effect of liposomal LA (Epstein-Barash et al., 2009), and nevertheless, the effect on axonal function remains poorly investigated. In a previous study in human volunteers, we found that it is feasible to monitor the effect of LA *in vivo* by combining conventional conduction studies with nerve excitability studies by threshold-tracking (Moldovan et al., 2014). Here, we aimed to develop a translational mouse model where the same neurophysiological techniques can be used to investigate liposomal formulations of LA *in vivo*. To challenge the validity of the model, we tested an unilamellar liposomal formulation, consisting of a single lipid bilayer (Betageri and Parsons, 1992) which has faster release profile than multilamellar liposomes (Silva et al., 2016). The liposomes were filled with the LA lidocaine which has a shorter duration of action than bupivacaine in extended-release formulations (Huynh et al., 2015).

## MATERIALS AND METHODS

### Mice, Anesthesia, and Experimental Design

Investigations *in vivo* were carried out in adult mice of C57Bl background (3- to 5-month-old) with axonal expression of the Yellow Fluorescent Protein (YFP) reporter (homozygote Thy1-YFP) obtained from the Jackson Laboratory, Bar Harbor, ME, United States. General anesthesia was ensured using a 1:1 mixture of Hypnorm/Midazolam (5 mg/ml). A volume of 0.1 ml/10 g from the mixture was injected subcutaneously for induction, and then maintained with 50% hourly for up to 4 h as needed.

Using *in vivo* neurophysiological recordings (Figure 1), we monitored the recovery of tibial nerve conduction (Figure 2) and excitability (Figure 3) following injection of the LA. Slow subcutaneous injection of a volume of 50  $\mu$ l at the ankle engulfed the tibial nerve in an easily observable bubble (Figure 1C) ensuring the reproducibility of the LA exposure without the need of ultrasound visualization as in the case of human PNBs (Moldovan et al., 2014). The

monitoring of the systemic toxicity was carried out using a 1-channel electrocardiogram (ECG) recorded between the forepaws (Figure 1B) (MP150 with an ECG100C module, Biopac Systems Inc., United States).

At the completion of the electrophysiological experiments, the lateral aspect of the leg was exposed for imaging studies (Figure 4) after which the mice were killed by cervical dislocation.

This study was carried out in accordance with the recommendations of directive 2010/63/EU of the European parliament and of the council on the protection of animals used for scientific purposes. The protocol was approved by the Danish Animal Experiments Inspectorate.

### Electrophysiological Setup


For *in vivo* electrophysiological investigations, the mice were placed on a temperature-controlled pad (HB 101/2, LSI Letica) set to 37°C (Moldovan and Krarup, 2006). Stimulation and recording were carried out using percutaneously inserted Pt-needle electrodes (Moldovan and Krarup, 2006; Wild et al., 2018). Electrical stimuli generated from a constant current stimulator (DS4, Digitimer Ltd.) were delivered proximally to the sciatic nerve at the sciatic notch (Figures 1A,B) and distal to the tibial nerve at the ankle (Figures 1A,C). The evoked compound muscle potential (CMAP) of the plantar muscle was recorded using a bandpass filter of 10 Hz–6 kHz (Neurolog NL820A amplifier with NL844 Pre-Amplifier, Digitimer Ltd., United Kingdom). A ground reference electrode was inserted into the back of the mouse. The near-nerve electrode placement was carried out ensuring the lowest threshold current.

The signals were digitized into a PC-based recording and control system via a multifunction I/O device (PCI-6221, National Instruments Corporation Ltd., United Kingdom). The amplitudes of the CMAP evoked from the proximal (Figure 1D) and distal (Figure 1E) stimulation sites were measured peak-to-peak (ppA). The CMAP latency was measured to the first take-off from baseline (Figures 1D,E).

It should be noted that both from the proximal stimulation site (Figure 1D) as well as from the distal stimulation site (Figure 1E), the evoked CMAP was followed by a secondary motor response (“F-wave”) resulting from motoneuronal backfiring following antidromic activation in humans (Magladery and McDougal, 1950) and rodents (Meinck, 1976; Robertson et al., 1993; Moldovan et al., 2012). Although the F-waves were not specifically quantified, their presence was considered as an indication of the functional integrity of the motoneurons and the alpha-motor fibers.


### Nerve Conduction Studies

We have previously established a setup for the continuous monitoring of tibial nerve conduction by stimulation at the ankle for up to 4 h (Moldovan and Krarup, 2006; Alvarez et al., 2008). Nevertheless, overcoming the partial voltage-gate Na<sup>+</sup> channel block by lidocaine could require very large stimulation currents (Moldovan et al., 2014). Given the small distances, using such high currents at ankle could cause the stimulus to “jump” and activate the unanesthetized part of the



tibial nerve. To avoid this potential confounder, we tested the changes in conduction across the PNB by proximal stimulation (**Figures 1A,B**). Stimulation and recording were controlled using a custom control software developed in MATLAB (version R2013b, MathWorks Inc., United States). We delivered negative

rectangular stimuli of 1 ms duration at 0.5 Hz and recorded 1000 ms sweeps with a sampling frequency of 125 KHz allowing synchronized CMAP (**Figure 2A**) and ECG recordings (**Figure 2C**). The amplitudes at proximal stimulation were averaged over 1-min and expressed, relative to the 1st minute




prior to injection (**Figure 2B**). The stimulation current was set at twofold the current required to obtain a 100% ppA CMAP and kept constant for the entire duration of the recording.

## Nerve Excitability Studies

The changes in nerve excitability at the PNB site were monitored by distal stimulation at the ankle (**Figures 1A,C**) and tracking changes in the threshold current (**Figures 3A,B**) required to evoke a 40% ppA CMAP (Bostock et al., 1998) using QtracS

stimulation software for recording and QtracP software for analysis (©Institute of Neurology, London, United Kingdom). We used that same TRONDH multiple excitability protocol (Kiernan and Bostock, 2000), that we used in our human PNB study (Moldovan et al., 2014). We have previously adapted this protocol for use in mice (Moldovan et al., 2009, 2011). This allowed recording of the full sequence of excitability measures (**Figure 3C**) as previously described in detail (Bostock et al., 1998; Kiernan and Bostock, 2000; Kiernan et al., 2000): charge–duration relationship to measure the rheobase, threshold



electrotonus to measure accommodation to shifts in depolarizing and hyperpolarizing membrane potential, current–threshold relationship to measure input conductance, and the recovery of excitability following conduction of the action potential to measure refractoriness.


## Imaging Studies

*In vivo* imaging (**Figure 4**) was carried out using probe-based confocal laser endomicroscopy (pCLE) using a Cellvizio single band, with a 488-nm laser (Mauna Kea Technologies, France) which allows for the investigation of fluorescent blood vessels and

axons (Wong et al., 2009). The frames were captured via a S-1500 objective (**Figure 4A**) and exported using the Cellvizio Viewer for PC (version 1.4.2).

## Liposomal LA Formulation

We formed an unilamellar liposomes using the established methods of freeze drying lipids followed by rehydration in lidocaine hydrochloride (2% w/v) phosphate-buffered saline. This solution was pressure extruded through 100 nm filters to reduce liposome size polydispersity. The liposomes comprised of hydrogenated soy L- $\alpha$ -phosphatidylcholine (HSPC), cholesterol



**FIGURE 4 |** Imaging of the tibial nerve block area by probe-based confocal laser endomicroscopy (pCLE) at 90 min after liposomal lidocaine (L. Lip). **(A)** The ankle was first dissected to allow access of the imaging probe (Probe) to the subcutaneous tissue. Note the scattering of the blue laser light used for imaging; **(B)** Superficial veins were fluorescent consistent with the absorption of the fluorescent liposomes. The arrow identifies the same venous confluence in light imaging and pCLE imaging; **(C)** With further advancement of the imaging probe between the muscle planes (blunt dissection), the tibial nerve could be readily identified due to its fluorescent fibers (axonal YFP reporter expression). Occasionally, liposomal accumulations (star) could be identified near the fibers.

(Chol), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-methoxy (polyethylene glycol)-2000 (DSPE-PEG2000), and 1-palmitoyl-2-(dipyrrometheneboron difluoride)undecanoyl-sn-glycero-3-phosphocholine (TopFluor PC) in a mol% ratio of 54.8:40:5:0.2, respectively. Polyethylene glycol conjugated lipid was introduced into the liposome bilayer to increase residency time (Eriksen et al., 2017) and TopFluor PC was incorporated

to allow visualization. The resulting liposome size (Mean  $\pm$  SD) was found to be  $196.2 \pm 2.2$  nm with a size polydispersity index of  $0.134 \pm 0.03$  as determined by dynamic light scattering. A non-encapsulated lidocaine salt solution (20 mg/ml, L. HCl) served as control. All chemical components were purchased from commercial suppliers (Sigma-Aldrich, DE, and Avanti Polar Lipids, United States).

## RESULTS

### Recovery of Conduction

Proximal stimulation of the tibial nerve fibers at the sciatic notch (**Figures 1A,B**) evoked a CMAP (Mean  $\pm$  SD) with a ppA of  $12 \pm 4$  mV and a latency of  $1.7 \pm 0.1$  ms (**Figure 1D**). Distal stimulation of the tibial nerve fibers at the ankle (**Figures 1A,C**) evoked a CMAP (Mean  $\pm$  SD) with a ppA of  $16 \pm 5$  mV and a latency of  $1.2 \pm 0.1$  ms (**Figure 1E**). Note that the proximal CMAP was not larger than the distal CMAP indicating that the CMAP measurements reflected tibial-nerve innervated muscles with minimal contamination (if any) from common peroneal nerve innervated muscles.

An experiment with non-encapsulated lidocaine is detailed in **Figure 2A**. Within 3 min following LA injection, there was a complete abolishment of the CMAP and F-waves evoked by proximal stimulation which recovered completely within 30 min (**Figure 2A**). The time-course of CMAP amplitude (ppA) recovery, measured relative to the minute before LA, is presented in **Figure 2B**. In contrast, our model was able to detect a slower recovery after liposomal lidocaine, with good reproducibility in repeated studies (**Figure 2B**). Furthermore, we found that following liposomal lidocaine the CMAP evoked by proximal stimulation recovered rapidly to above 50% ppA whereas the subsequent recovery was much slower (**Figure 1B**), so that a full recovery was observed after 3 h (data not shown). Although liposomes were absorbed, as indicated by pCLE (**Figure 4B**), there were no ECG signs of bradycardia (**Figure 2C**) that could indicate a confounding effect of lidocaine toxicity.

### Recovery of Excitability

During the slow CMAP recovery phase after liposomal lidocaine (**Figure 2B**), liposomal accumulations could be demonstrated by pCLE near the tibial nerve fibers at the ankle (**Figure 4C**) suggesting a persistent release. An experiment with uncaged lidocaine is detailed in **Figure 3**. We found that during the partial CMAP recovery phase, evoking the maximal CMAP by distal stimulation in the nerve exposed to LA required a much larger stimulation current (**Figure 3A**) with a right-shift in the stimulus-response curve (**Figure 3B**). At that time, the current required to evoke the CMAP from the proximal stimulation site remained undistinguishable from the value prior to LA (data not shown).

To explore the mechanisms of impaired tibial nerve excitability at the LA site, we carried out multiple measures of excitability by “threshold-tracking” (**Figure 3C**). Consistent with the larger stimulus current required to evoke the CMAP, we found a marked increase in rheobase (Weiss, 1901; Bostock et al., 1998), as indicated by the slope of the charge–duration relationship (**Figure 3C**). Moreover, the observed increase in rheobase after liposomal lidocaine was associated with (**Figure 3C**) a reduction in threshold changes during both depolarizing and hyperpolarizing threshold electrotonus, a marked increase in the current–threshold slope (input conductance) on depolarization, and an increased in refractoriness of the recovery cycle (**Figure 3C**).

## DISCUSSION

We developed an *in vivo* mouse model to explore the effect of local anesthetic nanomedicines on axonal function. We tested the validity of the model by comparing the effect of a liposomal lidocaine formulation versus non-encapsulated lidocaine on motor axon function and found a slower conduction recovery along the tibial nerve following liposomal lidocaine. Furthermore, the model opened the possibility to combine conduction studies with “threshold-tracking” excitability studies in the same nerve, allowing a translational mechanistic insight into the liposomal release *in vivo*.

Assessment of the effect of liposomal local anesthetic nanomedicines on sciatic nerve block in rodents has been previously carried out previously by behavioral measures (Yin et al., 2016). The sensory (thermal nociceptive) blockade was investigated in a hot-plate paradigm (Sagie and Kohane, 2010), whereas the motor blockade was investigated using the extensor postural thrust test (Sagie and Kohane, 2010). Although these behavioral studies provided a valuable approach to distinguish the motor versus sensory selectivity of the nanomedicine, they offered little information on the actual changes in axonal function. It is long known that the susceptibility to an anesthetic conduction block is different between neuronal populations, reflecting differences in membrane properties (Gasser and Erlanger, 1929). Nevertheless, motor population was found to be at least as good an indicator for changes in the duration of PNB as the sensory fibers (Dietz and Jaffe, 1997; Gokin et al., 2001) even though they do not directly reflect the analgesic effect. Furthermore, the fact that the myelinated motor axon population is more functionally homogenous than the sensory fiber population (Gasser and Erlanger, 1929), can convey an advantage in characterizing the excitability changes.

The LA lidocaine is thought to impair the function of the voltage-gated Na<sup>+</sup> channels (VGSC) (Sheets and Hanck, 2007) reducing the axonal “safety factor” for conduction (Tasaki, 1953) in excitable tissues of the nervous system (Nathan and Sears, 1961) as well as the heart (Austen and Moran, 1965). Although the pCLE imaging studies indicated liposome accumulation within veins (**Figure 4B**), the ECG (**Figure 2C**) remained within normal limits (Ho et al., 2011). We therefore do not think that the observed effects on axonal function were confounded by lidocaine cardiotoxicity (Fox and Kenmore, 1967).

We confirmed an increase in the threshold for electrical stimulation during recovery after LA (Gasser and Erlanger, 1929; Gasser and Grundfest, 1939). Similar deviations in excitability were found in our previous study using non-encapsulated lidocaine in humans (Moldovan et al., 2014) which supports the translational value of our mouse model. These excitability deviations differed from those observed with selective VGSC block following accidental tetrodotoxin poisoning in humans (Kiernan et al., 2005). It is likely that the protracted period of reduced nerve excitability following LA (Tabatabai and Booth, 1990) reflected both the partial recovery of the Na<sup>+</sup> conductance as well as an impairment of the axolemmal passive electrical properties (Moldovan et al., 2014) due to a physical effect of lidocaine dissolved within the membrane (Kassahun et al., 2010).



In further studies, excitability changes could allow the detection of early signs of toxicity (Moldovan et al., 2009) in combination with imaging of morphological changes (Beirowski et al., 2005; Alvarez et al., 2008, 2013).

## AUTHOR CONTRIBUTIONS

CK and KL initiated the study. MM made the neurophysiological control software. AU made the liposomal lidocaine formulation.

## REFERENCES

- Aguirre, J., Del Moral, A., Cobo, I., Borgeat, A., and Blumenthal, S. (2012). The role of continuous peripheral nerve blocks. *Anesthesiol. Res. Pract.* 2012:560879. doi: 10.1155/2012/560879
- Alvarez, S., Moldovan, M., and Krarup, C. (2008). Acute energy restriction triggers Wallerian degeneration in mouse. *Exp. Neurol.* 212, 166–178. doi: 10.1016/j.expneurol.2008.03.022
- Alvarez, S., Moldovan, M., and Krarup, C. (2013). Prolonged high frequency electrical stimulation is lethal to motor axons of mice heterozygously deficient for the myelin protein P(0) gene. *Exp. Neurol.* 247, 552–561. doi: 10.1016/j.expneurol.2013.02.006
- Austen, W. G., and Moran, J. M. (1965). Cardiac and peripheral vascular effects of lidocaine and procainamide. *Am. J. Cardiol.* 16, 701–707. doi: 10.1016/0002-9149(65)90054-8
- Beirowski, B., Adalbert, R., Wagner, D., Grumme, D. S., Addicks, K., Ribchester, R. R., et al. (2005). The progressive nature of Wallerian degeneration in wild-type and slow Wallerian degeneration (Wlds) nerves. *BMC Neurosci.* 6:6. doi: 10.1186/1471-2202-6-6
- Betageri, G. V., and Parsons, D. L. (1992). Drug encapsulation and release from multilamellar and unilamellar liposomes. *Int. J. Pharm.* 81, 235–241. doi: 10.1016/0378-5173(92)90015-T
- Bostock, H., Cikurel, K., and Burke, D. (1998). Threshold tracking techniques in the study of human peripheral nerve. *Muscle Nerve* 21, 137–158. doi: 10.1002/(SICI)1097-4598(199802)21:2<137::AID-MUS1>3.0.CO;2-C
- Dietz, F. B., and Jaffe, R. A. (1997). Bupivacaine preferentially blocks ventral root axons in rats. *Anesthesiology* 86, 172–180. doi: 10.1097/00000542-199701000-00021
- Epstein-Barash, H., Shichor, I., Kwon, A. H., Hall, S., Lawlor, M. W., Langer, R., et al. (2009). Prolonged duration local anesthesia with minimal toxicity. *Proc. Natl. Acad. Sci. U.S.A.* 106, 7125–7130. doi: 10.1073/pnas.0900598106
- Eriksen, A. Z., Brewer, J., Andresen, T. L., and Urquhart, A. J. (2017). The diffusion dynamics of PEGylated liposomes in the intact vitreous of the ex vivo porcine eye: a fluorescence correlation spectroscopy and biodistribution study. *Int. J. Pharm.* 522, 90–97. doi: 10.1016/j.ijpharm.2017.03.003
- Fox, J. L., and Kenmore, P. I. (1967). The effect of ischemia on nerve conduction. *Exp. Neurol.* 17, 403–419. doi: 10.1016/0014-4886(67)90127-6
- Gasser, H. S., and Erlanger, J. (1929). The role of fiber size in the establishment of a nerve block by pressure or cocaine. *Am. J. Physiol.* 88, 581–591. doi: 10.1152/ajplegacy.1929.88.4.581
- Gasser, H. S., and Grundfest, H. (1939). Axon diameters in relation to the spike dimensions and the conduction velocity in mammalian A fibers. *Am. J. Physiol.* 127, 393–414. doi: 10.1152/ajplegacy.1939.127.2.393
- Gokin, A. P., Philip, B., and Strichartz, G. R. (2001). Preferential block of small myelinated sensory and motor fibers by lidocaine: in vivo electrophysiology in the rat sciatic nerve. *Anesthesiology* 95, 1441–1454. doi: 10.1097/00000542-200112000-00025
- Grant, G. J., Piskoun, B., Lin, A., and Bansinath, M. (2000). An in vivo method for the quantitative evaluation of local anesthetics. *J. Pharmacol. Toxicol. Methods* 43, 69–72. doi: 10.1016/S1056-8719(00)00079-4
- Grant, G. J., Zakowski, M. I., Vermeulen, K., Langerman, L., Ramanathan, S., and Turndorf, H. (1993). Assessing local anesthetic effect using the mouse tail flick test. *J. Pharmacol. Toxicol. Methods* 29, 223–226. doi: 10.1016/1056-8719(93)90029-E

All the authors contributed to the experiments, analyzed the data, discussed the results, and wrote the manuscript.

## FUNDING

The project was supported by the Danish Medical Research Council grants (4092-00100/2014 and 7016-00165/2017), the Lundbeck Foundation, and the Foundation for Research in Neurology and Jytte and Kaj Dahlboms Foundation.

- Hadzic, A., Minkowitz, H. S., Melson, T. I., Berkowitz, R., Uskova, A., Ringold, F., et al. (2016). Liposome bupivacaine femoral nerve block for postsurgical analgesia after total knee arthroplasty. *Anesthesiology* 124, 1372–1383. doi: 10.1097/ALN.0000000000001117
- Ho, D., Zhao, X., Gao, S., Hong, C., Vatner, D. E., and Vatner, S. F. (2011). Heart rate and electrocardiography monitoring in mice. *Curr. Protoc. Mouse Biol.* 1, 123–139. doi: 10.1002/9780470942390.mo100159
- Huynh, T. M., Marret, E., and Bonnet, F. (2015). Combination of dexamethasone and local anaesthetic solution in peripheral nerve blocks: a meta-analysis of randomised controlled trials. *Eur. J. Anaesthesiol.* 32, 751–758. doi: 10.1097/EJA.0000000000000248
- Ilfeld, B. M. (2017). Continuous peripheral nerve blocks: an update of the published evidence and comparison with novel, alternative analgesic modalities. *Anesth. Analg.* 124, 308–335. doi: 10.1213/ANE.0000000000001581
- Kassahun, B. T., Murashov, A. K., and Bier, M. (2010). A thermodynamic mechanism behind an action potential and behind anesthesia. *Biophys. Rev. Lett.* 5, 35–41. doi: 10.1142/S1793048010001123
- Kiernan, M. C., and Bostock, H. (2000). Effects of membrane polarization and ischaemia on the excitability properties of human motor axons. *Brain* 123, 2542–2551. doi: 10.1093/brain/123.12.2542
- Kiernan, M. C., Burke, D., Andersen, K. V., and Bostock, H. (2000). Multiple measures of axonal excitability: a new approach in clinical testing. *Muscle Nerve* 23, 399–409. doi: 10.1002/(SICI)1097-4598(200003)23:3<399::AID-MUS12>3.0.CO;2-G
- Kiernan, M. C., Isbister, G. K., Lin, C. S., Burke, D., and Bostock, H. (2005). Acute tetrodotoxin-induced neurotoxicity after ingestion of puffer fish. *Ann. Neurol.* 57, 339–348. doi: 10.1002/ana.20395
- Leszczynska, K., and Kau, S. T. (1992). A sciatic nerve blockade method to differentiate drug-induced local anesthesia from neuromuscular blockade in mice. *J. Pharmacol. Toxicol. Methods* 27, 85–93. doi: 10.1016/1056-8719(92)90026-W
- Madsen, M. H., Christiansen, C. B., Rothe, C., Andreasen, A. M., Lundstrom, L. H., and Lange, K. H. W. (2018). Local anesthetic injection speed and common peroneal nerve block duration: a randomized controlled trial in healthy volunteers. *Reg. Anesth. Pain Med.* 43, 467–473. doi: 10.1097/AAP.0000000000000759
- Magladery, J. W., and McDougal, D. B. (1950). Electrophysiological studies of nerve and reflex activity in normal man. I. Identification of certain reflexes in the electromyogram and the conduction velocity of peripheral nerve fibres. *Bull. Johns Hopkins Hosp.* 86, 265–290.
- Meinck, H. M. (1976). Occurrence of the H reflex and the F wave in the rat. *Electroencephalogr. Clin. Neurophysiol.* 41, 530–533. doi: 10.1016/0013-4694(76)90064-X
- Moldovan, M., Alvarez, S., and Krarup, C. (2009). Motor axon excitability during Wallerian degeneration. *Brain* 132(Pt 2), 511–523. doi: 10.1093/brain/awn332
- Moldovan, M., Alvarez, S., Pinchenko, V., Klein, D., Nielsen, F. C., Wood, J. N., et al. (2011). Nav1.8 channelopathy in mutant mice deficient for myelin protein zero is detrimental to motor axons. *Brain* 134(Pt 2), 585–601. doi: 10.1093/brain/awq336
- Moldovan, M., Alvarez, S., Pinchenko, V., Marklund, S., Graffmo, K. S., and Krarup, C. (2012). Nerve excitability changes related to axonal degeneration in amyotrophic lateral sclerosis: insights from the transgenic SOD1(G127X) mouse model. *Exp. Neurol.* 233, 408–420. doi: 10.1016/j.expneurol.2011.11.008

- Moldovan, M., and Krarup, C. (2006). Evaluation of Na<sup>+</sup>/K<sup>+</sup> pump function following repetitive activity in mouse peripheral nerve. *J. Neurosci. Methods* 155, 161–171. doi: 10.1016/j.jneumeth.2005.12.015
- Moldovan, M., Lange, K. H., Aachmann-Andersen, N. J., Kjaer, T. W., Olsen, N. V., and Krarup, C. (2014). Transient impairment of the axolemma following regional anaesthesia by lidocaine in humans. *J. Physiol.* 592(Pt 13), 2735–2750. doi: 10.1113/jphysiol.2014.270827
- Nathan, P. W., and Sears, T. A. (1961). Some factors concerned in differential nerve block by local anaesthetics. *J. Physiol.* 157, 565–580. doi: 10.1113/jphysiol.1961.sp006743
- Robertson, A., Day, B., Pollock, M., and Collier, P. (1993). The neuropathy of elderly mice. *Acta Neuropathol.* 86, 163–171. doi: 10.1007/BF00334883
- Sagie, I., and Kohane, D. S. (2010). Prolonged sensory-selective nerve blockade. *Proc. Natl. Acad. Sci. U.S.A.* 107, 3740–3745. doi: 10.1073/pnas.0911542107
- Sheets, M. F., and Hanck, D. A. (2007). Outward stabilization of the S4 segments in domains III and IV enhances lidocaine block of sodium channels. *J. Physiol.* 582(Pt 1), 317–334. doi: 10.1113/jphysiol.2007.134262
- Silva, C. B., Groppo, F. C., Santos, C. P., Serpe, L., Franz-Montan, M., Paula, E., et al. (2016). Anaesthetic efficacy of unilamellar and multilamellar liposomal formulations of articaine in inflamed and uninflamed tissue. *Br. J. Oral Maxillofac. Surg.* 54, 295–300. doi: 10.1016/j.bjoms.2016.01.005
- Tabatabai, M., and Booth, A. M. (1990). Effects of lidocaine on the excitability and membrane properties of the nerve cell soma. *Clin. Physiol. Biochem.* 8, 289–296.
- Tasaki, I. (1953). *Nervous Transmission*. Springfield, IL: Charles C. Thomas.
- Vyas, K. S., Rajendran, S., Morrison, S. D., Shakir, A., Mardini, S., Lemaine, V., et al. (2016). Systematic review of liposomal bupivacaine (Exparel) for postoperative analgesia. *Plast. Reconstr. Surg.* 138, 748e–756e. doi: 10.1097/PRS.0000000000002547
- Weiniger, C. F., Golovanevski, L., Domb, A. J., and Ickowicz, D. (2012). Extended release formulations for local anaesthetic agents. *Anaesthesia* 67, 906–916. doi: 10.1111/j.1365-2044.2012.07168.x
- Weiss, G. (1901). Sur la possibilité de rendre comparables entre eux les appareils servant à l'excitation électrique. *Arch. Ital. Biol.* 35, 413–446.
- Wild, B. M., Morris, R., Moldovan, M., Krarup, C., Krishnan, A. V., and Arnold, R. (2018). *In vivo* electrophysiological measurement of the rat ulnar nerve with axonal excitability testing. *J. Vis. Exp.* 6:132. doi: 10.3791/56102
- Wong, F., Fan, L., Wells, S., Hartley, R., Mackenzie, F. E., Oyeboode, O., et al. (2009). Axonal and neuromuscular synaptic phenotypes in Wld(S), SOD1(G93A) and ostes mutant mice identified by fiber-optic confocal microendoscopy. *Mol. Cell. Neurosci.* 42, 296–307. doi: 10.1016/j.mcn.2009.08.002
- Wu, C. L., and Raja, S. N. (2011). Treatment of acute postoperative pain. *Lancet* 377, 2215–2225. doi: 10.1016/S0140-6736(11)60245-6
- Yin, Q., Ke, B., Chen, X., Guan, Y., Feng, P., Chen, G., et al. (2016). Effects of liposomes charge on extending sciatic nerve blockade of N-ethyl bromide of lidocaine in rats. *Sci. Rep.* 6:38582. doi: 10.1038/srep38582
- Zaslansky, R., Rothaug, J., Chapman, C. R., Backstrom, R., Brill, S., Fletcher, D., et al. (2015). PAIN OUT: the making of an international acute pain registry. *Eur. J. Pain* 19, 490–502. doi: 10.1002/ejp.571

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Moldovan, Alvarez, Rothe, Andresen, Urquhart, Lange and Krarup. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.