



Preservation methods throughout the value chain: Fish oil.

Jacobsen, Charlotte

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Jacobsen, C. (Author). (2018). Preservation methods throughout the value chain: Fish oil.. Sound/Visual production (digital), Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Preservation methods for fish oil

Charlotte Jacobsen

Professor and group leader

Research group for Bioactives – Analysis and Application

chja@food.dtu.dk




Agenda

- Intro to antioxidants
- Different examples on effects of antioxidants on fish oil oxidation
 - Tocopherols
 - Citric acid vs tocopherols vs ascorbyl palmitate (AP)
 - Sesamol vs oryzanol vs rosemary extract vs BHT
 - Propyl gallate and citric acid vs rosemary extract and AP
 - Tocopherol and rosemary vs tocopherol
- Research needs

Antioxidants - Mechanisms




Lipidoxidation (Autoxidation)



Primary antioxidants

Radical scavengers

Prevention of initiation and propagation




Termination of lipid oxidation reactions



Figure: With courtesy of Ann-Dorit M Sørensen, DTU Food

Antioxidants - Mechanisms




Secondary antioxidants

- Metal chelation
 - Scavenging oxygen
- Not illustrated in the figure...***
- Singlet oxygen quencher
 - Synergism between antioxidants

Figure: With courtesy of Ann-Dorit M Sørensen, DTU Food

Antioxidants - Mechanisms

Examples on AO which can work as primary or secondary AO




Ascorbic acid

Radical scavenger

Oxygen scavenger


Synergy with tocopherol

Reducing $Fe^{3+} \rightarrow Fe^{2+}$



EDTA

Metal chelator




BHT

Radical scavenger

Lycopene

Singlet oxygen quencher




Natural vs synthetic antioxidants

- Synthetic (examples)
 - Propyl gallate
 - BHT
 - Ascorbyl palmitate


- Natural (examples)
 - Tocopherol
 - Ascorbic acid
 - Rosemary extract (rosmarinic acid, carnosol and carnosic acid)
 - Sesamol
 - Oryzanol
 - (Citric acid)

Effect of different tocopherol isomers on and different concentrations on fish oil oxidation (PV and CD) (stripped oils)



Delta- and gamma-toc more efficient than alpha-toc

Kulås and Ackman, *J. Agric. Food Chem* (2001) 49, 1724-1729



Lower concentrations of alpha-tocopherol more efficient than higher ones

Figure 2. Effect of α -tocopherol (α TOH) on the formation of primary oxidation products in anchovy oil during storage at 30 °C: (A) peroxide value measurements; (B) conjugated diene measurements. Data points are means \pm standard deviation, $n = 3$.

Effect of tocopherol homologues on different volatile oxidation products in stripped fish oil

Tab. 2. Values for variables found to be important in the principal component analysis of high-level tocopherol samples*.

PCA variable	α TOH (1000 ppm, day 5)	δ TOH (1000 ppm, day 5)
Total volatiles	11.0 \pm 0.9 ppm	11.2 \pm 0.4 ppm
Hydrocarbons	1.46 \pm 0.08 ppm	0.71 \pm <0.01 ppm
Propanal	0.37 \pm 0.07 ppm	0.70 \pm 0.02 ppm
<i>t,c/t,t-2,4-Heptadienal</i>	7.3 \pm 1.5**	5.0 \pm 0.5***
Diunsaturated/saturated aldehydes	0.61 \pm 0.07	0.29 \pm 0.04

* The values are mean of two samples \pm the difference between the mean and each sample value. Concentrations are relative to the ethyl heptanoate internal standard.

** Day 8: 7.5 \pm 0.5 (mean \pm standard deviation, n = 3).

*** Day 8: 5.1 \pm 0.3 (mean \pm standard deviation, n = 3).


Different tocopherol homologues will change the ratio between different volatiles that are formed

Comparison of the efficacy of citric acid, ascorbyl palmitate and alpha-tocopherol in fish oil by Oxidograph

Oil without antioxidant (NA)

Antioxidants in 3 different concentrations



Citric acid (CA), Ascorbyl palmitate (AP) and alpha-tocopherol (AT)



Ascorbyl palmitate and citric acid were equally efficient (200 mg/kg)

Data from DTU Food

Effect of sesamol, oryzanol, rosemary extract or BHT on volatiles in stripped fish oil




8.4 mM Sesamol SM, Rosemary Extract (RE) or Oryzanol (OZ) vs 0.84 mM BHT

- Rosemary extract and sesamol equally efficient and both better than BHT at 50 °C
- At 30 °C, rosemary extract was slightly better than sesamol (data not shown)

Faner et al, EJLST 118, 2016, 885-897

Rosemary and AP vs propyl gallate and citric acid

Increasing shelf life of fish oil




Rosemary plus ascorbyl palmitate almost as efficient as propyl gallate and citric acid

From Dupont presentation Marine Lipider meeting Århus 2014

Rosemary and AP vs propyl gallate and citric acid


Increasing shelf life of salmon oil



CONFIDENTIAL 13

From Dupont presentation Marine Lipider meeting Århus 2014

Effect of high concentrations of rosemary and tocopherol on lipid oxidation in fish oil



High doses of rosemary extract and tocopherol almost completely inhibited formation of volatiles

Thomsen et al., *Eur. J Lipid Sci. Technol.*, (2017), 119.

Research needs

- Most studies reported in the literature are carried out with stripped or refined oils

More research should be carried out with unrefined fish oil with different fatty acid compositions and levels of natural antioxidants. Research should address the following questions:

- How can BHT (or ethoxyquin if still used) be replaced by natural antioxidants in unrefined oil?
- Are there differences in antioxidant efficacy in refined vs unrefined oils?
- How is antioxidant efficacy influenced by fatty acid composition and presence of endogenous antioxidants or prooxidants in unrefined fish oils?