



Itaconimides as Novel Quorum Sensing Inhibitors of *Pseudomonas aeruginosa*

Fong, July; Mortensen, Kim Thollund; Nørskov, Amalie; Qvortrup, Katrine; Yang, Liang; Tan, Choon Hong; Nielsen, Thomas E.; Givskov, Michael

Published in:

Frontiers in Cellular and Infection Microbiology

Link to article, DOI:

[10.3389/fcimb.2018.00443](https://doi.org/10.3389/fcimb.2018.00443)

Publication date:

2019

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link back to DTU Orbit](#)

Citation (APA):

Fong, J., Mortensen, K. T., Nørskov, A., Qvortrup, K., Yang, L., Tan, C. H., ... Givskov, M. (2019). Itaconimides as Novel Quorum Sensing Inhibitors of *Pseudomonas aeruginosa*. *Frontiers in Cellular and Infection Microbiology*, 8, [443]. DOI: 10.3389/fcimb.2018.00443

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Itaconimides as Novel Quorum Sensing Inhibitors of *Pseudomonas aeruginosa*

July Fong^{1†}, Kim T. Mortensen^{2†}, Amalie Nørskov², Katrine Qvortrup², Liang Yang^{1,3*}, Choon Hong Tan^{1,4}, Thomas E. Nielsen^{1,5} and Michael Givskov^{1,5*}

¹ Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, Singapore, Singapore, ² Department of Chemistry, Technical University of Denmark, Lyngby, Denmark, ³ Southern University of Science and Technology, Guangdong, China, ⁴ Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore, Singapore, ⁵ Department of Immunology and Microbiology, Costerton Biofilm Center, University of Copenhagen, Copenhagen, Denmark

OPEN ACCESS

Edited by:

You-Hee Cho,
CHA University, South Korea

Reviewed by:

Eric Déziel,
Institut National de la Recherche
Scientifique (INRS), Canada
Daniel Angel Ortiz,
Laboratory Corporation of America
Holdings (LabCorp), United States

*Correspondence:

Liang Yang
yangliang@ntu.edu.sg
Michael Givskov
mgivskov@sund.ku.dk

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Clinical Microbiology,
a section of the journal
Frontiers in Cellular and Infection
Microbiology

Received: 06 October 2018

Accepted: 11 December 2018

Published: xx December 2018

Citation:

Fong J, Mortensen KT, Nørskov A,
Qvortrup K, Yang L, Tan CH,
Nielsen TE and Givskov M (2018)
Itaconimides as Novel Quorum
Sensing Inhibitors of *Pseudomonas*
aeruginosa.
Front. Cell. Infect. Microbiol. 8:443.
doi: 10.3389/fcimb.2018.00443

Pseudomonas aeruginosa is known as an opportunistic pathogen that often causes persistent infections associated with highly antibiotic-resistance and biofilms formation. Chemical interference with bacterial cell-cell communication, termed quorum sensing (QS), has been recognized as an attractive approach to control infections and address the drug resistance problems currently observed worldwide. Instead of imposing direct selective pressures on bacterial growth, the right bioactive compounds can preferentially block QS-based communication and attenuate cascades of bacterial gene expression and production of virulence factors, thus leading to reduced pathogenicity. Herein, we report on the potential of itaconimides as quorum sensing inhibitors (QSI) of *P. aeruginosa*. An initial hit was discovered in a screening program of an in-house compound collection, and subsequent structure-activity relationship (SAR) studies provided analogs that could reduce expression of central QS-regulated virulence factors (elastase, rhamnolipid, and pyocyanin), and also successfully lead to the eradication of *P. aeruginosa* biofilms in combination with tobramycin. Further studies on the cytotoxicity of compounds using murine macrophages indicated no toxicity at common working concentrations, thereby pointing to the potential of these small molecules as promising entities for antimicrobial drug development.

Keywords: quorum sensing, biofilm, itaconimides, antivirulence, chemical biology

INTRODUCTION

Pseudomonas aeruginosa is an opportunistic pathogen and a major cause of nosocomial infections in patients with pneumonia, chronic wounds, urinary tract infections, and intensive care units (ICUs) (Vincent et al., 1995). As an opportunistic pathogen, this organism is highly adaptive, versatile and exhibits remarkable resistance toward many antimicrobial agents. Resistance is a distinctive characteristic of *P. aeruginosa*, due to its ability to express multiple resistance mechanisms, including enzymes and efflux pumps (Poole, 2001; Lister et al., 2009). (U.S. Centers for Disease Control and Prevention (CDC), 2013) estimated that more than 20,000 deaths per year are attributed to antibiotic resistance cases. It has become a global issue, and we are threatened with the slow progress of new antibiotics development and lack of preventive measure for the spread of resistance.

In addition, *P. aeruginosa* is well-known to form biofilms, which have been identified as a major underlying cause of persistent infections in immunocompromised patients, chronic wounds as well as on medical devices like implants, catheters, tubes, artificial hip, and many more (Costerton et al., 1999). Biofilms infections are often characterized by their broad range resistance toward host defense mechanisms and antibiotic therapy. This results in prolonged treatment, complications in clinical outcomes, and additional socio-economic burdens. In cystic fibrosis patients, chronic biofilm infections of *P. aeruginosa* can cause premature death despite intensive antibiotic therapy care (Bjarnsholt et al., 2009).

The bacteria in biofilms often exhibit different phenotypic and genetic variants as compared to their planktonic counterparts. In the biofilm mode of life, bacterial cells are enclosed within a matrix of extracellular polymeric substances (EPS) comprises of exopolysaccharides, proteins, deoxyribonucleic acid (DNA), lipids or surfactants, and macromolecules that are self-produced by the cells (Flemming and Wingender, 2010). All of these could render antibiotics impenetrable, chelated or sequestered, and diminish the efficacy of the treatment. The presence of persister cells in biofilms also contributes to multidrug resistance property of biofilms (Lewis, 2007). Overall, the complex biology of biofilms represents a tremendous challenge to develop therapeutic agents that could successfully prevent or eradicate biofilms-associated infections.

The cell-to-cell communication system called quorum sensing (QS) has been reported to play major roles for establishing persistent, biofilm based infections (Hentzer et al., 2003; Alhede et al., 2009; Van Gennip et al., 2009; Chiang et al., 2013). The QS system in *P. aeruginosa* utilizes acyl homoserine lactones (AHLs) as signal molecules and comprises the Lux homologs LasRI and RhlRI. LasI synthase is responsible for the synthesis of *N*-(3-oxododecanoyl) homoserine lactone (3-oxo-C12-HSL), which will bind to its receptor LasR and activate transcription of genes responsible for virulence such as *lasB*, *apr*, and *toxA* (Gambello et al., 1993; Passador et al., 1993). The *las* system also positively regulates *rhl* system, where RhlI directs the synthesis of *N*-butanoylhomoserine lactone (C4-HSL) that would bind to its receptor RhlR and subsequently activate gene expression of QS target genes (Brint and Ohman, 1995; Pearson et al., 1995; Pesci et al., 1997). In addition, there is also a third signaling molecule “pseudomonas quinolone signal” (PQS) which is intertwined between the *las* and *rhl* systems (Schertzer et al., 2009). Recently, a fourth signal molecules called Integrative Quorum Sensing Signal (IQS) has been reported, which could overtake the central *las* system under phosphate depletion condition (Lee et al., 2013). QS defective *P. aeruginosa* mutants are attenuated as compared to the wild-type strain, and their biofilms are more susceptible toward antibiotics treatment and host immune system as compared to the wild-type (Pearson et al., 2000; Hentzer et al., 2002, 2003).

As QS governs various patterns of genes expression to control virulence and biofilm formation, it has been proposed that interfering with the communication system could be a promising strategy for the control and prevention of bacterial infections (Hentzer et al., 2003). Quorum sensing inhibitors (QSI) are

compounds that interfere with QS pathways, reduce expression of QS-controlled genes and attenuate infecting bacteria. As such compounds do not affect the growth of bacteria, these molecular entities pose lower selective pressure on bacteria and lower the risk of resistance development. Recent exploration of new classes of QSI comprises natural products, synthetic molecules, and enzymes that may quench or inactivate QS signals (Dong et al., 2001; Hentzer et al., 2002; Jakobsen et al., 2012; Fong et al., 2017). Unfortunately, no clinical candidates have yet been developed for therapy.

In the present study, we report a new class of small molecules that disrupt QS pathways in *P. aeruginosa*. The structurally related itaconimides and citraconimides have previously been reported to inhibit growth of mycobacteria (Balganesh et al., 1999). We synthesized a range of small molecules by an iterative structure-activity relationship (SAR) study and found two promising candidates for further biological investigation. The efficacies of these compounds were tested on *P. aeruginosa* QS bioreporter strains (*lasB-gfp*; *rhlA-gfp*; *pqsA-gfp*) and also on QS-controlled virulence phenotypes, such as elastase, pyocyanin, and rhamnolipid production.

MATERIALS AND METHODS

General Information

All chemicals were purchased from Sigma Aldrich and used without further purification. For biological studies, synthesized compounds were prepared in DMSO as 10 mM stock solution and stored at -20°C until further usage. Overnight culture of bacteria was grown in Lysogeny broth (LB) which consisted of 1% tryptone, 0.5% yeast extract, 0.5% NaCl. For bioreporter assay, strains were grown in ABTGC (AB minimal medium supplemented with 0.2% glucose and 0.2% casamino acids) (Clark and Maaloe, 1967) to minimize fluorescence interference. ABTG (with no casamino acid) medium was used for biofilms study in flow chambers. Strains used in this study can be found in **Table 1**. Summary of chemical synthesis and spectroscopy data can be found in **Supplementary Material**.

Bioreporter QS Assay

From its frozen stock, compounds were diluted appropriately to their working concentration in ABTGC medium. Experiments were done as previously reported (Fong et al., 2017). Briefly, 200 μL of compounds was pipetted into the first rows of 96-well plates (Nunc, Denmark), followed by two-fold serial dilution to the rest of the rows. The last two rows were allocated to solvent control (DMSO 0.1%) and blank (media control). Overnight cultures of *P. aeruginosa* bioreporter strains were diluted to optical density at 600 nm (OD_{600}) of 0.02 ($\sim 2.5 \times 10^8$ CFU/mL). Next, 100 μL of the bacteria culture was added into each well to make final OD_{600} of 0.01. The plate was incubated at 37°C for 16 h, with time-point measurement of GFP fluorescence (excitation 485 nm, emission 535 nm) and OD_{600} recorded at every 15 min using Tecan Infinite 200 Pro plate reader (Tecan Group Ltd, Männedorf, Switzerland). The data were exported into excel files, and IC_{50} value calculation was determined using GraphPad Prism 6 software. For IC_{50} values determination, the

Q5 229 **TABLE 1** | Bacterial strains used in this study.

230	231	232	233
Strains or plasmids	Relevant genotype and/or characteristics ^a		
234	PAO1	ATCC <i>Pseudomonas aeruginosa</i> Hentzer et al., 2002	
235	PAO1- <i>gfp</i>	GFP-tagged wild-type <i>Pseudomonas aeruginosa</i> Yang et al., 2007	
236	PAO1- <i>lasB-gfp</i>	PAO1 containing <i>lasB-gfp</i> (ASV) reporter fusion Hentzer et al., 2002	
237	PAO1- <i>rhlA-gfp</i>	PAO1 containing <i>rhlA-gfp</i> (ASV) reporter fusion Yang et al., 2007	
238	PAO1- <i>pqsA-gfp</i>	PAO1 containing <i>pqsA-gfp</i> (ASV) reporter fusion Yang et al., 2009	
239	PAO1 Δ <i>lasI</i> Δ <i>rhlI</i>	Gm ^a ; PAO1 <i>lasI</i> and <i>rhlI</i> mutant Hentzer et al., 2003	
240	PAO1 Δ <i>lasR</i>	PAO1 <i>lasR</i> mutant Hentzer et al., 2003	
241	Δ <i>lasR-rhlA-gfp</i>	PAO1 <i>lasR</i> mutant containing <i>rhlA-gfp</i> (ASV) reporter fusion Tan et al., 2013	
242	Δ <i>lasR-pqsA-gfp</i>	PAO1 <i>lasR</i> mutant containing <i>pqsA-gfp</i> (ASV) reporter fusion Tan et al., 2013	

243 ^aDescription of the strains' antibiotic resistance. Gm, gentamicin; Carb, carbenicillin resistance.

244 GFP/OD₆₀₀ values were taken at the time point between 4 and 245 6 h, where inhibition started to occur. All experiments were done 246 in triplicate manner and repeated at least twice to confirm the 247 results.

248 QS-Regulated Virulence Factor Assays

249 Elastase activity was measured using EnzChekElastase kit 250 (Invitrogen, USA), following the manufacturer's instruction. 251 Rhamnolipid was extracted and quantified using method 252 reported by Koch et al. with modifications (Koch et al., 253 1991). Briefly, overnight cultures of *P. aeruginosa* were diluted 254 into ABTGC medium (OD₆₀₀ = 0.01), with and without the 255 presence of compounds (DMSO as control). Cultures were grown 256 overnight at 37°C, 200 rpm. Rhamnolipid was extracted from 257 the supernatant with diethyl ether (twice), and organic fractions 258 were concentrated to yield yellowish-white solids. The solids were 259 re-suspended in deionized water and added with 0.19% (w/v) 260 orcinol in 50% H₂SO₄. It was then heated at 80°C for 20–30 min 261 to give dark orange color. Absorbance was measured at 421 nm 262 and the values were normalized with cell density at OD₆₀₀. 263 Pyocyanin was extracted from overnight culture of *P. aeruginosa* 264 grown in Kings Medium A Base [MilliQ water supplemented 265 with proteose peptone (20 g/L), potassium sulfate (10 g/L), 266 magnesium chloride, anhydrous (1.640 g/L), and glycerol (10% 267 v/v)]. Supernatants were collected and extracted with chloroform 268 and 0.2 M HCl. The presence of pyocyanin would turn the HCl 269 solution into pinkish color. Absorbance was measured at 520 nm 270 and normalized with cell density OD₆₀₀ values. Experiments were 271 done in triplicate manner and repeated at least twice to confirm 272 the results.

273 Biofilm Experiments

274 GFP-tagged *P. aeruginosa* were grown in ABTG medium 275 and flowed through flow chambers as previously described 276

(Sternberg and Tolker-Nielsen, 2006). Each flow chamber is 286 consisted of three-channel flow cells that were supplied with 287 a flow of medium and oxygen, while waste medium would be 288 directed into a waste flask. Briefly, overnight cultures were diluted 289 1,000 times in ABTG medium and injected into each channel for 290 1 h incubation time without flow. Next, the medium was allowed 291 to flow into the flow cells and the velocity was maintained at 0.2 292 mm/s using Cole-Palmer peristaltic pump. Biofilms were grown 293 for 72 h before treatment with compounds for further 48 h. To 294 visualize dead cells, 300 μ L of propidium iodide (PI) stain was 295 injected into each flow cells. Biofilm images were taken with LSM 296 confocal laser scanning microscope (Carl Zeiss, Germany) at 20x 297 objective lens. Microscopy images were processed with IMARIS 298 software (Bitplane AG, Zurich, Switzerland). Experiments were 299 done in triplicate manner and repeated at least twice to confirm 300 the results.

301 Cytotoxicity Assay



















302 Toxicity assay was done as previously reported (Fong et al., 303 2017). Murine macrophage RAW264.7 cell lines were 304 grown in Dulbecco's Modified Eagle's Medium (DMEM, 305 Life Technologies), supplemented with 10% fetal bovine 306 serum (Gibco) at 37°C and 5% CO₂. The cells were passaged 307 into 96-well microplates, with each well containing 1 \times 308 10⁴ macrophages. After 16 h, the cells were washed with 309 phosphate-buffered saline (PBS) and treated with compounds 310 at varying concentration in 100 μ L DMEM. The plate was 311 incubated at 37°C and 5% CO₂ for further 16 h. Resazurin 312 was then added into each well to reach final concentration 313 of 10 μ M for cell viability measurement. The live cells would 314 be able to convert the dye to red color, whereby dead cells 315 would remain as blue color. Absorbance was taken at 595 nm 316 using Tecan Infinite 200 Pro plate reader (Tecan Group 317 Ltd, Männedorf, Switzerland). Experiments were done in 318 triplicate manner and repeated at least twice to confirm the 319 results.

320 RESULTS

321 Synthesis and SAR Study

322 Recently, we discovered that 3-methylene-1- 323 tetradecylpyrrolidine-2,5-dione (Table 2, 1a) displayed QSI 324 activity against *P. aeruginosa* QS reporter strain (carrying a 325 *lasB-gfp* fusion). Different variation on the left-hand side of 326 1a proved that the exo-cyclic double bond was essential for 327 its biological activity. This led us to focus on the synthesis 328 and biological evaluation of *N*-substituted itaconimide analogs 329 against *P. aeruginosa* (Scheme 1). The procedure for the 330 synthesis of the itaconimides have been described by Cava 331 et al. (1961) and Leow et al. (2008). The commercially available 332 itaconic anhydride was treated with anilines (1.0 equiv.) 333 in CHCl₃ to afford the corresponding α -itaconamic acids 334 (Kyung et al., 1974). The resulting acids were subsequently 335 treated with Ac₂O (3.5 equiv.) and NaOAc (0.5 equiv.) at 336 elevated temperature to afford a mixture of itaconimide and 337 the isomerized product citraconimide. Generally, the yield for 338 the aniline derivatives was higher than that of the aliphatic 339

TABLE 2 | Summary of the synthesized compounds and the corresponding yields.

Compound	R	a (%) ^a	b (%) ^a	Compound	R	a (%) ^a	b (%) ^a
1		50	14	10		35	15
2		37	– ^b	11		61	9
3		17	– ^b	12		37	20
4		10	– ^b	13		40	19
5		47	– ^b	14		8	14
6		49	29	15		16	– ^b
7		– ^c		16		13	10
8		31	40	17		4	12
9		24	14	18		2	11

^aYields based on starting itaconic anhydride and purified after cyclization by flash column chromatography.

^bNot isolated.


^cCommercial sample.

amines. An overview of the synthesized compounds is provided below (Table 2).

QSI Activities of Itaconimides

The synthesized compounds were tested for their QS inhibitory activity against the *P. aeruginosa lasB-gfp* reporter strain. None

of our synthesized compounds showed antibiotic properties (Supplementary Figures 1, 2). We found that both the *p*-bromophenyl and tetradecyl-substituted itaconimide (12a and 18a) showed strong QS inhibition activity against the *lasB-gfp* reporter strain (Figure 1A). Both compounds inhibit the expression of *lasB-gfp* in a dose-dependent manner (Figure 1B).



SCHEME 1 | Synthesis of itaconimides and citraconimides.

18a was proved to be the most active compound synthesized in this series, with almost 25-fold higher activity (IC₅₀) compared to our hit compound (**Figures 1C,D**).

To address the specificity of our compounds, we also tested our compounds against PAO1 wild-type (WT) and $\Delta lasR$ harboring either *rhlA-gfp* or *pqsA-gfp* fusions. In this case, we would be able to determine if the compounds could affect other QS pathways in *las*-dependent or independent manner. Both *rhlA* and *pqsA* are the first genes of *rhl* operon and *pqs* operon that code for the production of the rhamnolipid and PQS precursor molecules (Ochsner et al., 1994; Gallagher et al., 2002). To eliminate false positive, we also tested both compounds with a *gfp*-tagged *P. aeruginosa* (expresses GFP constitutively) as control and did not observe any reduction in the fluorescence signals (**Supplementary Figure 3**).

Compounds **12a** and **18a** were found to inhibit expression of *rhlA-gfp* and *pqsA-gfp* in PAO1 WT. IC₅₀ values calculated for **12a** are $6.67 \pm 0.27 \mu\text{M}$ for *rhlA-gfp* and $2.51 \pm 0.19 \mu\text{M}$ for *pqsA-gfp*, whereby compound **18a** provides IC₅₀ values of $0.61 \pm 0.04 \mu\text{M}$ for *rhlA-gfp* and 0.143 ± 0.13 for *pqsA-gfp* (**Figure 2**). At $10 \mu\text{M}$, compound **12a** was observed to inhibit *rhlA-gfp* more strongly in $\Delta lasR$ (78% inhibition) as compared to the wild-type (58% inhibition). In $\Delta lasR$ *pqsA-gfp* reporter strain, the inhibition only happened in the later stage. Meanwhile, compound **18a** was found to inhibit both *rhlA-gfp* and *pqsA-gfp* efficiently in $\Delta lasR$ (**Figure 3**). Therefore, it is likely that two compounds have different mechanisms, and compound **18a** affect multiple QS pathways in *las*-independent pathway.

Effects of Itaconimides on Virulence Production

Next, we investigated the effects of synthesized compounds on virulence factors produced by *P. aeruginosa*, notably elastase, rhamnolipid, and pyocyanin. The three virulence factors are under QS-regulation, therefore they could be a good indicator for evaluating antivirulence activities of our compounds. We also included QS mutants as control strains, where they are defective in producing quorum sensing signals hence lower level of virulence production. At $10 \mu\text{M}$, both compounds were able to reduce all three virulence factors production. Elastase level was reduced almost half, and rhamnolipid and pyocyanin productions were abolished almost to the same level as mutant strains (**Figure 4**). The results showed that both compounds could indeed lower the production of virulence factors.

Effects of Itaconimides on *P. aeruginosa* Biofilms

QS has been shown to play important roles in biofilms matrix formation (Davey et al., 2003; Sakuragi and Kolter, 2007). Mutants lacking QS often form flat and undifferentiated biofilms that could be easily cleared by antimicrobial agents and host immune system (Wu et al., 2001; Christensen et al., 2007). Here, we investigated if the itaconimides could eradicate biofilms formation when used in combination with tobramycin antibiotic.

From **Figure 5**, it can be seen that treatment with tobramycin alone ($10 \mu\text{g/ml}$) couldn't clear the whole population of *P. aeruginosa* biofilms. The dead cells only appeared on the surface of the biofilms, which indicated that the antibiotic could not penetrate into the biofilms. In addition, biofilms often contain extracellular DNA (eDNA) which could chelate aminoglycoside antibiotics and render it inactive (Chiang et al., 2013). At $10 \mu\text{M}$, itaconimide **12a** could remarkably kill the base population of the biofilms, but not the surface. When added together with tobramycin, the combination treatment successfully eradicated the whole population of biofilms, which appeared red. Compound **18a** did not show any synergistic effect when used together with tobramycin, which could be due to poor solubility in the aqueous media.

Cytotoxic Effects of Itaconimides on Macrophages

We next evaluated the potential application of compounds **12a** and **18a** for therapeutic application. To test the cytotoxic effects, both compounds were added into murine macrophage RAW2647 cell lines, and the cell viability was measured after 16h incubation time. Both compounds were observed to be toxic at $40 \mu\text{M}$, but not cytotoxic at their working concentration ($10 \mu\text{M}$ and lower, **Figure 6**). This indicates that they could be further used for subsequent *in vivo* experiments at their relevant concentration.

DISCUSSION

In this study, we present the investigation of itaconimides as a novel QSI against *P. aeruginosa*. The approach of using QSI molecules to attenuate the virulence of pathogenic bacteria has advantages over conventional antibiotics therapy. Instead of targeting the growth of bacteria (bactericidal or bacteriostatic strategies), QSI compounds pose lesser selective




FIGURE 1 | (A) Effects of compounds **12a** and **18a** on *PAO1-lasB-gfp* reporter strain. DMSO 0.1% was used as solvent control. **(B)** Dose-response effects of compounds on *P. aeruginosa lasB-gfp*. **(C)** IC₅₀ values of **12a** and **18a** **(D)** on *lasB-gfp*. Calculation was done using GraphPad Prism 6 software, taken at time point between 4 and 6 h when QS inhibition started to occur. All experiments were done in triplicate manner (technical and biological replicates), only representative data are shown.




FIGURE 2 | QSI inhibition, dose-response and IC₅₀ values of compounds **12a** and **18a** on *P. aeruginosa rhlA-gfp* **(A)** and *pqSA-gfp* **(B)**. DMSO 0.1% was used as solvent control. For the IC₅₀ values, calculation was done using GraphPad Prism 6 software, taken at time point between 4 and 6 h when QS inhibition started to occur. All experiments were done in triplicate manner (technical and biological replicates), only representative data are shown.

pressure for the development of resistant mutants. In several studies, it has been shown that mice treated with QSI afford better survival profile and lower bacteria count loads when compared to the control group (Hentzer et al., 2002, 2003; Jakobsen et al., 2013; Fong et al., 2017). With limited options of treatment available, antivirulence therapy could be a viable approach to mitigate the future crisis of antibiotic resistance.

In this study, a structure-activity relationship (SAR) study was performed to investigate the QSI activities of itaconimides and citraconimide analogs. We screened the compounds against QS reporter strain *lasB-gfp*, where gene expression is under control of the QS *las* system. We first investigated the inhibitory properties of the two isomers with respect to *lasB-gfp* expression. Interestingly, the itaconimide scaffold was found to be more active than the citraconimide. Next, we investigated the influence




FIGURE 3 | QSI inhibition of compounds **12a** and **18a** on PAO1 *ΔlasR rhlA-gfp* (A) and *pqsA-gfp* (B). DMSO 0.1% was used as solvent control. All experiments were done in triplicate manner (technical and biological replicates), only representative data are shown.




FIGURE 4 | Effects of compounds **12a** and **18a** on elastase (A), rhamnolipid (B), and pyocyanin production (C). Compounds were tested at final concentration of 10 μM. All experiments were done in triplicate manner (technical and biological replicates). Error bars are means ± SDs. **** $p < 0.0001$, Student's t -test.





FIGURE 5 | *P. aeruginosa* biofilms formed on flow cells for 72 h, followed by treatment with medium containing antibiotic (tobramycin 10 μg/ml) and compounds (10 μM) for further 48 h. Live cells are *P. aeruginosa* tagged with GFP which appeared as green, and dead cells appeared as red. Scale bars, 50 μm. Experiments were done in triplicate manner (technical and biological replicates), only representative images were shown.

of electron donating groups (EDGs) and electron withdrawing groups (EWGs) on the phenyl group. Whereas, *p*-anisidine moiety failed to improve activity, the more electron-withdrawing 4-chloro derivative induced a 1.4-fold inhibition of PAO1-*lasB-gfp*. This guided us to explore anilines with different EWGs. The most potent compound in this series was **12a** from 4-bromoaniline that results in a 4.4-fold increase over the parent aniline analog. To our surprise, the 4-fluoroaniline analog **13a** did not afford a more potent inhibitor. Alongside the findings that 4-bromoaniline provided a more potent analog, we also found that the octylamine analog **14a** was more active than our hit compound **1a**. Lastly, we also included aliphatic chains to

mimic the long alkyl chain of C4-HSL and 3-oxo-C12-HSL. On this note, a second series was synthesized from commercially available aliphatic amines with varying carbon lengths ($n = 4, 6, 8, 10, 12, 14$). A clear trend was observed, where longer alkyl chain resulted in lower IC₅₀ values. The outcome was quite interesting, as IC₅₀ values of the citraconimide substituents of these analogs (**16b**, **17b**, and **18b**) were significantly lowered. The most potent compound was found to be the tetradecylamine analog, **18a**, being the most active of both series. By substituting the naphthalene group with either 4-bromoaniline or long chain alkyl amines, IC₅₀ values against PAO1-*lasB-gfp* were significantly reduced to the low micromolar range ($2.41 \pm$



0.99 μM for **12a** and $0.17 \pm 0.04 \mu\text{M}$ for **18a**). In summary, our SAR studies revealed two structurally important variations for the itaconimides that are highly important for QSI activity (Table 3).

Previous studies have emphasized the advantage of therapeutics that suppress multiple QS pathways (Fong et al., 2018). Indeed, *lasR* mutants are commonly found in patients suffering from cystic fibrosis and other clinical setting (Hamood et al., 1996; Cabrol et al., 2003; Marvig et al., 2014). Nevertheless, the loss-of-function *lasR* mutants continuously express virulence traits so *P. aeruginosa* could use other pathways to bypass LasR in controlling pathogenicity (Dekimpe and Déziel, 2009; Lee et al., 2013). Using different QS reporter strains in both PAO1 WT and ΔlasR , it can be deduced that our compounds do not specifically inhibit one QS pathway. Both compounds could inhibit expression of *rhlA-gfp* and *pqsA-gfp* in *lasR* mutant. The inhibition of *pqsA-gfp* in ΔlasR was less apparent for compound **12a**, which could indicate that the compound inhibits PQS system through *las*-dependent manner. The itaconimide analogs presented here hold unique potential as broad target QSIs to control infections via an anti-virulence strategy. Future work will aim to elucidate the mechanism on how both compounds inhibit QS in *P. aeruginosa*.

Next, we also tested the effects of our compounds on the production of various virulence factors, such as elastase, rhamnolipid, and pyocyanin production. Elastase is one of the major proteases produced by *P. aeruginosa*, involved in host tissue damage and host immune responses (Kamath et al., 2002). Rhamnolipid is also an essential virulence factor and plays several key roles in biofilms formation, swarming, and in particular host immune evasion. It promotes rapid necrotic killing of polymorphonuclear (PMNs) leukocytes and also infiltration of respiratory epithelial cells (Zulianello et al., 2006; Jensen et al., 2007). *P. aeruginosa* also secretes pyocyanin, blue redox-active secondary metabolite that has several deleterious effects on mammalian cells (Lau et al., 2004), which is also regulated by QS under *pqs* system. When tested at 10 μM , the compounds could

TABLE 3 | Summary of IC_{50} values of itaconimide (a) and citraconimide (b) analogs against PAO1-*lasB-gfp*.

Compound	a	b
1	7.37 \pm 0.71	–
2	23.52 \pm 0.81	–
3	10.27 \pm 0.05	–
4	–	–
5	17.95 \pm 0.49	–
6	10.67 \pm 0.49	–
7	–	NA
8	7.45 \pm 0.65	–
9	7.65 \pm 0.37	–
10	6.53 \pm 0.61	–
11	17.68 \pm 0.16	–
12	2.41 \pm 0.99	–
13	11.89 \pm 0.34	–
14	26.66 \pm 0.98	–
15	1.71 \pm 0.34	–
16	0.30 \pm 0.05	2.01 \pm 0.39
17	0.30 \pm 0.17	0.91 \pm 0.20
18	0.17 \pm 0.04	0.53 \pm 0.12

All IC_{50} values were reported as μM . Experiments were done in triplicate manner and repeated at least twice to confirm the results. NA, Not available.

reduce production of various virulence factors controlled by QS. This shows potential of our compounds as antivirulence agent of *P. aeruginosa*.

P. aeruginosa biofilms are highly resistant to most antibiotics, including the last-resort polymyxin antibiotic available, colistin (Chua et al., 2016). As QS has definite role in biofilms development, it has been proposed that QSI compounds could be used as prophylactic treatment for biofilms infections. One such case is using azithromycin, which inhibits QS and biofilm formation at sub-MIC concentration. Promising results were observed in pulmonary infections and CF patients in many clinical trials data upon treatment of low-dose of AZM (2 $\mu\text{g}/\text{ml}$) (Hansen et al., 2005; Fleet et al., 2013). In this study, we utilized a combination of our compounds with the aminoglycoside tobramycin to treat *P. aeruginosa* biofilms grown on flow chambers for 3 days. We chose to study tobramycin because of its clinical relevance to cystic fibrosis (CF) patients. The clinical isolates from lung patients often confer resistance to aminoglycoside antibiotics (Hurley et al., 1995; Saiman et al., 1996). Our data also shows that treatment with tobramycin alone only kills the upper layer of biofilms. Combined with **12a**, we observed that the whole population of biofilms was eradicated. The results offer promising application of QSI in combination with antibiotics as a control for biofilm-associated infections.

It has been reported that stringent response, which provides rapid adaptation to environmental stresses, regulates QS network and also bacteria's survival in biofilms of *P. aeruginosa* (van Delden et al., 2001; Nguyen et al., 2011; Schafhauser et al., 2014). QS-deficient mutants also have lower catalase and superoxide dismutase activities, and therefore more sensitive to oxidative stress (Hassett et al., 1999). Through our biofilms experiment, we

could observe some cells death upon treatment with compound **12a** in the base of the biofilms. This raises the possibility that our compound could also target stringent response and other stress-response genes, which resulted in the observed killing effect. For compound **18a**, it contains long alkyl chain, which may render its solubility and penetration into the biofilms matrix. Future study will investigate how our compounds could synergize with other antibiotics to treat biofilms from other clinical isolates.

Lastly, we investigated the cytotoxic profile of our compounds on macrophages. The cytotoxicity of the compounds is important if they are to be used in animal studies for subsequent drug development. Results indicated that compound **12a** and **18a** are not toxic up to 40 μ M concentration. Still, further studies are needed to qualify the efficacy of the compounds in mice models for their potential as anti-biofilm agents, as well as their pharmacodynamic and pharmacokinetic profiles.

In conclusion, we report the novel use of itaconimides as antivirulence compounds for *P. aeruginosa*. These compounds suppress the *las*, *rhl*, and *pqs* QS systems of *P. aeruginosa*, and effectively abolish virulence expression activities. Compounds **12a** and **18a** showed low micromolar IC₅₀ values against all three QS reporter strains with only little toxicity against macrophages at the administrated concentration. Moreover, a synergistic effect with tobramycin was observed for the killing of *P. aeruginosa*

biofilms, including otherwise the tolerant and hard to target sub-population cells. Overall, our findings point to a new class of hit compounds of relevance to the development of new drugs against the superbug *P. aeruginosa*.

AUTHOR CONTRIBUTIONS

LY, TN, and MG designed methods and experiments. JF and KM performed the experiments and analyzed the data. AN, KQ, CT, and TN were in charge of the chemical synthesis. JF and KM wrote the paper and carefully revised by LY, TN, and MG. All authors have contributed to, read, and approved the manuscript.

FUNDING

This research was supported by the National Research Foundation and Ministry of Education Singapore under its Research Centre of Excellence Program and AcRF Tier 2 (MOE2016-T2-1-010) from Ministry of Education, Singapore.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcimb.2018.00443/full#supplementary-material>

REFERENCES

- Alhede, M., Bjarnsholt, T., Jensen, P. Ø., Phipps, R. K., Moser, C., Christophersen, L., et al. (2009). *Pseudomonas aeruginosa* recognizes and responds aggressively to the presence of polymorphonuclear leukocytes. *Microbiology* 155, 3500–3508. doi: 10.1099/mic.0.031443-0
- Balganesh, M., Ethirajulu, K., Ganguly, B. S., Janakiraman, R., Kaur, P., Kajipalya, R., et al. (1999). Mycobacterial inhibitors. WO1999065483A1.
- Bjarnsholt, T., Jensen, P. Ø., Fiandaca, M. J., Pedersen, J., Hansen, C. R., Andersen, C. B., et al. (2009). *Pseudomonas aeruginosa* biofilms in the respiratory tract of cystic fibrosis patients. *Pediatr. Pulmonol.* 44, 547–558. doi: 10.1002/ppul.21011
- Brint, J. M., and Ohman, D. E. (1995). Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhII, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *J. Bacteriol.* 177. doi: 10.1128/jb.177.24.7155-7163.1995
- Cabrol, S., Olliver, A., Pier, G. B., Andreumont, A., and Ruimy, R. (2003). Transcription of quorum-sensing system genes in clinical and environmental isolates of *Pseudomonas aeruginosa*. *J. Bacteriol.* 185, 7222–7230. doi: 10.1128/JB.185.24.7222-7230.2003
- Cava, M. P., Deana, A. A., Muth, K., and Mitchell, M. J. (1961). N-Phenylmaleimide. *Org. Synth.* 41:93. doi: 10.15227/orgsyn.041.0093
- Chiang, W.-C., Nilsson, M., Jensen, P. Ø., Høiby, N., Nielsen, T. E., Givskov, M., et al. (2013). Extracellular DNA shields against aminoglycosides in *Pseudomonas aeruginosa* biofilms. *Antimicrob. Agents Chemother.* 57, 2352–2361. doi: 10.1128/AAC.00001-13
- Christensen, L. D., Moser, C., Jensen, P. Ø., Rasmussen, T. B., Christophersen, L., Kjelleberg, S., et al. (2007). Impact of *Pseudomonas aeruginosa* quorum sensing on biofilm persistence in an *in vivo* intraperitoneal foreign-body infection model. *Microbiology* 153, 2312–2320. doi: 10.1099/mic.0.2007/006122-0
- Chua, S. L., Yam, J. K. H., Hao, P., Adav, S. S., Salido, M. M., Liu, Y., et al. (2016). Selective labelling and eradication of antibiotic-tolerant bacterial populations in *Pseudomonas aeruginosa* biofilms. *Nat. Commun.* 7:10750. doi: 10.1038/ncomms10750

- Clark, D. J., and Maaloe, O. (1967). DNA replication and the division cycle in *Escherichia coli*. *J. Mol. Biol.* 23, 99–112. doi: 10.1016/S0022-2836(67)80070-6
- Costerton, J. W., Stewart, P. S., and Greenberg, E. P. (1999). Bacterial biofilms: a common cause of persistent infections. *Science* 284, 1318–1322. doi: 10.1126/science.284.5418.1318
- Davey, M. E., Caiazza, N. C., and O'toole, G. A. (2003). Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* 185, 1027–1036. doi: 10.1128/JB.185.3.1027-1036.2003
- Dekimpe, V., and Déziel, E. (2009). Revisiting the quorum-sensing hierarchy in *Pseudomonas aeruginosa*: the transcriptional regulator RhlR regulates LasR-specific factors. *Microbiology* 155, 712–723. doi: 10.1099/mic.0.022764-0
- Dong, Y.-H., Wang, L.-H., Xu, J.-L., Zhang, H.-B., Zhang, X.-F., and Zhang, L.-H. (2001). Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature* 144, 813–817. doi: 10.1038/35081101
- Fleet, J. E., Guha, K., Piper, S., Banya, W., Bilton, D., and Hodson, M. E. (2013). A retrospective analysis of the impact of azithromycin maintenance therapy on adults attending a UK cystic fibrosis clinic. *J. Cyst. Fibros.* 2013, 49–53. doi: 10.1016/j.jcf.2012.05.010
- Flemming, H.-C., and Wingender, J. (2010). The biofilm matrix. *Nat. Rev. Microbiol.* 8, 623–633. doi: 10.1038/nrmicro2415
- Fong, J., Yuan, M., Jakobsen, T. H., Mortensen, K. T., Santos, M. M. S. D., Chua, S. L., et al. (2017). Disulfide bond-containing ajoene analogues as novel quorum sensing inhibitors of *Pseudomonas aeruginosa*. *J. Med. Chem.* 60, 215–227. doi: 10.1021/acs.jmedchem.6b01025
- Fong, J., Zhang, C., Yang, R., Boo, Z. Z., Tan, S. K., Nielsen, T. E., et al. (2018). Combination therapy strategy of quorum quenching enzyme and quorum sensing inhibitor in suppressing multiple quorum sensing pathways of *P. aeruginosa*. *Sci. Rep.* 8:1155. doi: 10.1038/s41598-018-19504-w
- Gallagher, L. A., Mcknight, S. L., Kuznetsova, M. S., Pesci, E. C., and Manoil, C. (2002). Functions required for extracellular quinolone signaling by *Pseudomonas aeruginosa*. *J. Bacteriol.* 184, 6472–6480. doi: 10.1128/JB.184.23.6472-6480.2002

- Gambello, M. J., Kaye, S., and Iglewski, B. H. (1993). LasR of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (*apr*) and an enhancer of exotoxin A expression. *Infect. Immun.* 61, 1180–1184. doi: 10.1128/IAI.61.7.1180-1184.1993
- Hamood, A. N., Griswold, J., and Colmer, J. (1996). Characterization of elastase-deficient clinical isolates of *Pseudomonas aeruginosa*. *Infect. Immun.* 64, 3154–3160. doi: 10.1128/IAI.64.12.3154-3160.1996
- Hansen, C. R., Pressler, T., Koch, C., and Høiby, N. (2005). Long-term azitromycin treatment of cystic fibrosis patients with chronic *Pseudomonas aeruginosa* infection; an observational cohort study. *J. Cyst. Fibros* 4, 35–40. doi: 10.1016/j.jcf.2004.09.001
- Hassett, D. J., Ma, J. F., Elkins, J. G., McDermott, T. R., Ochsner, U. A., West, S. E. H., et al. (1999). Quorum sensing in *Pseudomonas aeruginosa* controls expression of catalase and superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. *Mol. Microbiol.* 34, 1082–1093. doi: 10.1046/j.1365-2958.1999.01672.x
- Hentzer, M., Riedel, K., Rasmussen, T. B., Heydorn, A., Andersen, J. B., Parsek, M. R., et al. (2002). Inhibition of quorum sensing in *Pseudomonas aeruginosa* biofilm bacteria by a halogenated furanone compound. *Microbiology* 148, 87–102. doi: 10.1099/00221287-148-1-87
- Hentzer, M., Wu, H., Andersen, J. B., Riedel, K., Rasmussen, T. B., Bagge, N., et al. (2003). Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* 22, 3803–3815. doi: 10.1093/emboj/cdg366
- Hurley, J. C., Miller, G. H., and Smith, A. L. (1995). Mechanism of amikacin resistance in *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Diagn. Microbiol. Infect. Dis.* 22, 331–336. doi: 10.1016/0732-8893(95)00138-6
- Jakobsen, T. H., Bjarnsholt, T., Jensen, P. Ø., Givskov, M., and Høiby, N. (2013). Targeting quorum sensing in *Pseudomonas aeruginosa* biofilms: current and emerging inhibitors. *Future Microbiol.* 8, 901–921. doi: 10.2217/fmb.13.57
- Jakobsen, T. H., Gennip, M. V., Phipps, R. K., Shanmugham, M. S., Christensen, L. D., Alhede, M., et al. (2012). Ajoene, a sulfur-rich molecule from garlic, inhibits genes controlled by quorum sensing. *Antimicrob. Agents Chemother.* 56:2314. doi: 10.1128/AAC.05919-11
- Jensen, P. Ø., Bjarnsholt, T., Phipps, R., Rasmussen, T. B., Calum, H., Christoffersen, L., et al. (2007). Rapid necrotic killing of polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of rhamnolipid by *Pseudomonas aeruginosa*. *Microbiology* 153, 1329–1338. doi: 10.1099/mic.0.2006/003863-0
- Kamath, S., Kapatral, V., and Chakrabarty, A. M. (2002). Cellular function of elastase in *Pseudomonas aeruginosa*: role in the cleavage of nucleoside diphosphate kinase and in alginate synthesis. *Mol. Microbiol.* 30, 933–941. doi: 10.1046/j.1365-2958.1998.01121.x
- Koch, A. K., Kappeli, O., Fiechter, A., and Reiser, J. (1991). Hydrocarbon assimilation and biosurfactant production in *Pseudomonas aeruginosa* mutants. *J. Bacteriol.* 173, 4212–4219. doi: 10.1128/jb.173.13.4212-4219.1991
- Kyung, J. H., Cha, S., and Clapp, L. B. (1974). Identification of isomeric amides of itaconic acid by proton magnetic resonance spectroscopy. *Organ. Magn. Res.* 6, 466–468. doi: 10.1002/mrc.1270060816
- Lau, G. W., Hassett, D. J., Ran, H., and Kong, F. (2004). The role of pyocyanin in *Pseudomonas aeruginosa* infection. *Trends Mol. Med.* 10, 599–606. doi: 10.1016/j.molmed.2004.10.002
- Lee, J., Wu, J., Deng, Y., Wang, J., Wang, C., Wang, J., et al. (2013). A cell-cell communication signal integrates quorum sensing and stress response. *Nat. Chem. Biol.* 9, 339–343. doi: 10.1038/nchembio.1225
- Leow, D., Lin, S., Chittimalla, S. K., Fu, X., and Tan, C. H. (2008). Enantioselective protonation catalyzed by a chiral bicyclic guanidine derivative. *Angew. Chem. Int. Ed.* 47, 5641–5645. doi: 10.1002/anie.200801378
- Lewis, K. (2007). Persister cells, dormancy and infectious disease. *Nat. Rev. Microbiol.* 5, 48–56. doi: 10.1038/nrmicro1557
- Lister, P. D., Wolter, D. J., and Hanson, N. D. (2009). Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin. Microbiol. Rev.* 22, 582–610. doi: 10.1128/CMR.00040-09
- Marvig, R. L., Sommer, L. M., Molin, S., and Johansen, H. K. (2014). Convergent evolution and adaptation of *Pseudomonas aeruginosa* within patients with cystic fibrosis. *Nat. Genet.* 47, 1–9. doi: 10.1038/ng.3148
- Nguyen, D., Joshi-Datar, A., Lepine, F., Bauerle, E., Olakanmi, O., Beer, K., et al. (2011). Active starvation responses mediate antibiotic tolerance in biofilms and nutrient-limited bacteria. *Science* 334, 982–986. doi: 10.1126/science.1211037
- Ochsner, U. A., Fiechter, A., and Reiser, J. (1994). Isolation, characterization, and expression in *Escherichia coli* of the *Pseudomonas aeruginosa* *rhlAB* genes encoding a rhamnolipid biosurfactant synthesis. *J. Biol. Chem.* 269, 19787–19795.
- Passadori, L., Cook, J. M., Gambello, M. J., Rust, L., and Iglewski, B. H. (1993). Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science* 260, 1127–1130. doi: 10.1126/science.8493556
- Pearson, J. P., Feldman, M., Iglewski, B. H., and Prince, A. (2000). *Pseudomonas aeruginosa* cell-to-cell signaling is required for virulence in a model of acute pulmonary infection. *Infect. Immun.* 68, 4331–4334. doi: 10.1128/IAI.68.7.4331-4334.2000
- Pearson, J. P., Passadori, L., Iglewski, B. H., and Greenberg, E. P. (1995). A second N-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U.S.A.* 92, 1490–1494. doi: 10.1073/pnas.92.5.1490
- Pesci, E. C., Pearson, J. P., Seed, P. C., and Iglewski, B. H. (1997). Regulation of *las* and *rhl* quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* 179, 3127–3132. doi: 10.1128/jb.179.10.3127-3132.1997
- Poole, K. (2001). Multidrug efflux pumps and antimicrobial resistance in *Pseudomonas aeruginosa* and related organisms. *J. Mol. Microbiol. Biotechnol.* 3, 255–264.
- Saiman, L., Mehar, F., Niu, W. W., Neu, H. C., Shaw, K. J., Miller, G., et al. (1996). Antibiotic susceptibility of multiply resistant *Pseudomonas aeruginosa* isolated from patients with cystic fibrosis, including candidates for transplantation. *Clin. Infect. Dis.* 23, 532–537. doi: 10.1093/clinids/23.3.532
- Sakuragi, Y., and Kolter, R. (2007). Quorum-sensing regulation of the biofilm matrix genes (*pel*) of *Pseudomonas aeruginosa*. *J. Bacteriol.* 189, 5383–5386. doi: 10.1128/JB.00137-07
- Schafhauser, J., Lepine, F., McKay, G., Ahlgren, H. G., Khakimova, M., and Nguyen, D. (2014). The stringent response modulates 4-hydroxy-2-alkylquinoline biosynthesis and quorum-sensing hierarchy in *Pseudomonas aeruginosa*. *J. Bacteriol.* 196, 1641–1650. doi: 10.1128/JB.01086-13
- Schertzer, J. W., Boulette, M. L., and Whiteley, M. (2009). More than a signal: non-signaling properties of quorum sensing molecules. *Trends Microbiol.* 17, 189–195. doi: 10.1016/j.tim.2009.02.001
- Sternberg, C., and Tolker-Nielsen, T. (2006). Growing and analyzing biofilms in flow cells. *Curr. Protoc. Microbiol.* Chapter 1, Unit 1B 2. doi: 10.1002/9780471729259.mc01b02s00
- Tan, S. Y., Chua, S.-L., Chen, Y., Rice, S. A., Kjelleberg, S., Nielsen, T. E., et al. (2013). Identification of five structurally unrelated quorum-sensing inhibitors of *Pseudomonas aeruginosa* from a natural-derivative database. *Antimicrob. Agents Chemother.* 57, 5629–5641. doi: 10.1128/AAC.00955-13
- U.S. Centers for Disease Control and Prevention (CDC) (2013). *Antibiotic Resistance Threats in the United States*.
- van Delden, C., Comte, R., and Bally, M. (2001). Stringent response activates quorum sensing and modulates cell density-dependent gene expression in *Pseudomonas aeruginosa*. *J. Bacteriol.* 183, 5376–5384. doi: 10.1128/JB.183.18.5376-5384.2001
- Van Gennip, M., Christensen, L. D., Alhede, M., Phipps, R., Jensen, P. Ø., Christophersen, L., et al. (2009). Inactivation of the *rhlA* gene in *Pseudomonas aeruginosa* prevents rhamnolipid production, disabling the protection against polymorphonuclear leukocytes. *APMIS* 117, 537–546. doi: 10.1111/j.1600-0463.2009.02466.x
- Vincent, J.-L., Bihari, D. J., Suter, P. M., Bruining, H. A., White, J., Nicolas-Chanoine, M. H., et al. (1995). The prevalence of nosocomial infection in intensive care units in Europe. results of the European prevalence of infection in intensive care (epic) study. epic international advisory committee. *J. Am. Med. Assoc.* 274, 639–644. doi: 10.1001/jama.1995.03530080055041
- Wu, H., Song, Z., Givskov, M., Doring, G., Worlitzsch, D., Mathee, K., et al. (2001). *Pseudomonas aeruginosa* mutations in *lasI* and *rhlII* quorum sensing systems result in milder chronic lung infection. *Microbiology* 147, 1105–1113. doi: 10.1099/00221287-147-5-1105
- Yang, L., Barken, K. B., Skindersoe, M. E., Christensen, A. B., Givskov, M., and Tolker-Nielsen, T. (2007). Effects of iron on DNA release and biofilm

1141 development by *Pseudomonas aeruginosa*. *Microbiology* 153, 1318–1328.
 1142 doi: 10.1099/mic.0.2006/004911-0

1143 Yang, L., Rybtke, M. T., Jakobsen, T. H., Hentzer, M., Bjarnsholt, T., Givskov,
 1144 M., et al. (2009). Computer-aided identification of recognized drugs as
 1145 *Pseudomonas aeruginosa* quorum-sensing inhibitors. *Antimicrob. Agents
 Chemother.* 53, 2432–2443. doi: 10.1128/AAC.01283-08

1146 Zulianello, L., Canard, C., Kohler, T., Caille, D., Lacroix, J.-S., and
 1147 Meda, P. (2006). Rhamnolipids are virulence factors that promote
 1148 early infiltration of primary human airway epithelia by *Pseudomonas
 aeruginosa*. *Infect. Immun.* 74, 3134–3147. doi: 10.1128/IAI.
 1149 01772-05

1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197

Conflict of Interest Statement: The authors declare that the research was
 conducted in the absence of any commercial or financial relationships that could
 be construed as a potential conflict of interest.

Copyright © 2018 Fong, Mortensen, Nørskov, Qvortrup, Yang, Tan, Nielsen and
 Givskov. This is an open-access article distributed under the terms of the Creative
 Commons Attribution License (CC BY). The use, distribution or reproduction in
 other forums is permitted, provided the original author(s) and the copyright owner(s)
 are credited and that the original publication in this journal is cited, in accordance
 with accepted academic practice. No use, distribution or reproduction is permitted
 which does not comply with these terms.

1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254