

#### The high energy particle flux distribution of Jupiter measured by the ASCs of the Magnetometer Investigation

Jørgensen, John Leif; Jørgensen, Peter Siegbjørn; Shushkova, J.; Merayo, José M.G.; Denver, Troelz; Herceg, Matija; Jørgensen, Finn E.; Connerney, E. P.; Bolton, S. J.; Levin, S. M.

Publication date: 2018

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Jørgensen, J. L., Jørgensen, P. S., Shushkova, J., Merayo, J. M. G., Denver, T., Herceg, M., ... Levin, S. M. (2018). The high energy particle flux distribution of Jupiter measured by the ASCs of the Magnetometer Investigation. Poster session presented at Asia Oceania Geosciences Society 15th Annual Meeting (AOGS 2018), Honolulu, United States.

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Technical University of Denmark



# The high energy particle flux distribution of Jupiter measured by the ASCs of the Magnetometer Investigation



Credit: NASA/GSFC/JPL-Caltech/DTU Space

#### Introduction

The Juno mission offers an excellent opportunity to study the flux and distribution of very high energy particles moving the Jovian magnetosphere. The Magnetometer Investigation is equipped with two star trackers collocated with each of the two vector-magnetometers at the end of one of the solar panels. The primary objective for the star cameras is to provide accurate attitude recovery data for the magnetic mapping mission. The star trackers were provided with limited shielding, which in turn result in sensitivity to >15MeV electrons and 85MeV protons.

The star trackers, reporting the number of particles penetrating to the CCD at 4Hz, thus realize a very high energy particle detector.

### micro Advanced Stellar Compass µASC

- Designed and produced by the Measurement and Instrumentation Systems (DTU)
- one of the most successful star tracker worldwide
- autonomously calculates attitude based on all bright stars in the CHUs Field of View
- provides 22 true solutions per second with one single CHU
- absolute accuracy of < 1 arc second
- operating on many satellite missions without a single hardware or functional failure

J. L. Jørgensen<sup>1</sup>, P. S. Jørgensen<sup>1</sup>, J. Shushkova<sup>1</sup>, J.M.G. Merayo<sup>1</sup>, T. Denver<sup>1</sup>, M. Herceg<sup>1</sup>, F. Jørgensen<sup>1</sup>, E.P. Connerney<sup>2,3</sup>, S.J. Bolton<sup>4</sup>, S.M. Levin<sup>5</sup> <sup>1</sup> Technical University of Denmark (Denmark), <sup>2</sup> NASA Goddard Space Flight Center (USA), <sup>4</sup> Southwest Research Institute (USA), <sup>5</sup> Jet Propulsion Laboratory (USA)

## Calibration

The µASC response to the high energy irradiation is fully calibrated using 1-2 MeV electron and 30-200 MeV proton irradiation with varied incidence angle, flux, dose rate, temperature, TID, and DDD doses.



Calibration test setup inside mounted the thermal chamber consist of a pre-irradiated rotatable CCD under test, the star stimulator optics and grating, and a movable Sr90 e<sup>-</sup> source.

The shield-depth of the CCD onboard sensor Juno is analyzed using a Monte Carlo simulator. Here red rays show where materials are traversed.

Calibrating images acquired during irradiation with high flux of electrons (left) and heavies (right).



### Image analysis. Deposited energy

A track left by an energetic particle that hit the CCD is ~50 pixels long (~0.7 mm or 160 mg/cm<sup>2</sup>) and has 4 or 5 splits



due to recoil nuclei produced by the incident particle in silicon. Estimation of the primary energy of a particle that could traverse at least 0.7 mm in Si behind the heavy shielding will involve the transport flux modelling for the µASC, the calibration curves of the CCD and inflight image analysis.

Heavy particle with incoming E>> 85MeV

#### **Moons and Radiation**

The flybys of Juno orbiting Jupiter in highly elliptical have a period of about 53 days (left, in J2000). The Juno orbit in SIII coordinates (right) shows the effect of Jupiter's rotation period (~10h).





J2000 coordinates  $[R_{J}]$ 

SIII coordinates [R]]

The wiggle plot below shows Juno's position in magnetic coordinates (Rho, Z) in [R<sub>1</sub>]. In the trace of the orbit, the flux



measured by the <sup>v</sup>µASC is displayed with a color scale. The major Jovian radial moons distance have been <sup>a</sup> also added in the plot addition to the model the Of magnetodisc.

Juno's crossing of the magnetic field lines which intersect the Jovian moon orbits is shown. These crossings are correlated with observations of the µASC particle counter.



JPL (SwR

### Jovian magnetic field

Jupiter's magnetic field model (JRM09) by Connerney (2018) describes the field with spherical harmonics up to degree 10. It is based on measurements performed near Jupiter by the Juno mission. The JRM09 model shows a more complex Jovian magnetic field in comparison to earlier models (eg. VIP4), and a clear asymmetry within the northern and southern hemispheres. For larger distances to the planet (>10R<sub>1</sub>), the magnetic field can be represented well by a 10.5° tilted magnetic dipole and there is no apparent differences wrt VIP4 model.



We have calibrated and demonstrated the use of the µASC as a high energy particle detector in the Jovian magnetosphere. As the forthcoming Juno flybys will cover more spatial regions near Jupiter, we expect to build a model of the high energy electron radiation environment near the planet solely based on the data from the star trackers.

Ref.: Jørgensen, P. S., Jørgensen, J. L., & Denver, T, 2004. µAsc: a miniature sta tracker, Proceedings of The 4S Symposium: Small Satellites, Systems and Services (ESA SP-571). 20 - 24 September 2004, La Rochelle, France. Editor: E Warmbein. Published on CDROM., id.55.1, 2004ESASP.571E..55J