

## Research Article

# Nicotine-Cadmium Interaction Alters Exploratory Motor Function and Increased Anxiety in Adult Male Mice

Duyilemi Chris Ajonijebu,<sup>1</sup> Philip Adeyemi Adeniyi,<sup>2</sup> Adeshina Oloruntoba Adekeye,<sup>2</sup> Babawale Peter Olatunji,<sup>3</sup> Azeez Olakunle Ishola,<sup>4</sup> and Olalekan Michael Ogundele<sup>2</sup>

<sup>1</sup> Department of Physiology, College of Medicine and Health Sciences, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria

<sup>2</sup> Department of Anatomy, Cell Biology and Neuroscience Unit, College of Medicine and Health Sciences, Afe Babalola University, College Building II, Room G14, Km 8.5 Afe Babalola Way, PMB 5454, Ado-Ekiti, Ekiti State, Nigeria

<sup>3</sup> Department of Biological Sciences, College of Sciences, Afe Babalola University, Ado-Ekiti, Nigeria

<sup>4</sup> Department of Anatomy, College of Health Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria

Correspondence should be addressed to Olalekan Michael Ogundele; [mikealslaw@hotmail.com](mailto:mikealslaw@hotmail.com)

Received 17 May 2014; Accepted 27 August 2014; Published 13 November 2014

Academic Editor: Yasuji Matsuoka

Copyright © 2014 Duyilemi Chris Ajonijebu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this study we evaluated the time dependence in cadmium-nicotine interaction and its effect on motor function, anxiety linked behavioural changes, serum electrolytes, and weight after acute and chronic treatment in adult male mice. Animals were separated randomly into four groups of  $n = 6$  animals each. Treatment was done with nicotine, cadmium, or nicotine-cadmium for 21 days. A fourth group received normal saline for the same duration (control). Average weight was determined at 7-day interval for the acute (D1-D7) and chronic (D7-D21) treatment phases. Similarly, the behavioural tests for exploratory motor function (open field test) and anxiety were evaluated. Serum electrolytes were measured after the chronic phase. Nicotine, cadmium, and nicotine-cadmium treatments caused no significant change in body weight after the acute phase while cadmium-nicotine and cadmium caused a decline in weight after the chronic phase. This suggests the role of cadmium in the weight loss observed in tobacco smoke users. Both nicotine and cadmium raised serum  $\text{Ca}^{2+}$  concentration and had no significant effect on  $\text{K}^{+}$  ion when compared with the control. In addition, nicotine-cadmium treatment increased bioaccumulation of  $\text{Cd}^{2+}$  in the serum which corresponded to a decrease in body weight, motor function, and an increase in anxiety.

## 1. Introduction

Wide arrays of behavioural and physiological responses have reportedly been observed following nicotine use either in tobacco smoke or as a pure drug [1]. In human abusers and rodent models, weight loss, anxiety, depression, and motor dysfunctions are among the most frequently reported behavioural changes [2–4]. Long and short term use of nicotine in tobacco smoke alter synaptic function, neurotransmitter level, and the anatomical structure of the brain to varying extents. Similarly, the addictive effect of nicotine in tobacco smoke is known to potently alter the expression of cholinergic receptors, serum level of ions, and neurotransmission (central and peripheral) [5]. Primarily, nicotine,

being a structural analogue of acetylcholine, potentiates the receptors and facilitates the increase in intracellular  $\text{Ca}^{2+}$  concentration, vesicle formation, and neurotransmission at synapses and neuromuscular junctions. However, prolonged excitation of cholinergic synapses by nicotine leads to excitotoxicity and synaptic dysfunction, observed in prolonged use and abuse of nicotine [6–11].

Although nicotine gets into the body through various means—such as chewing of tobacco leaves and absorption of nicotine through the skin [12, 13]—the most predominant route of nicotine use is through tobacco smoke in cigarettes and pipes (respiratory) [14, 15]. Interestingly, several studies have reported the presence of cadmium, carbon monoxide, lead, copper, silicon, and alkaloids in the tobacco smoke

[14, 16]. Despite the striking evidence of the bioaccumulation of cadmium in tobacco smoke, coupled with its ability to inhibit cellular respiration in neurons, the metabolic and behavioural changes associated tobacco smoke has often been restricted to nicotine and its associated functions. Although cadmium is a trace element in the blood, previous studies have shown that the level of cadmium in the serum of smokers is usually 30–50% higher than that of nonsmokers [17–19]. Furthermore, overproduction of carbon dioxide, due to cadmium-mediated mitochondria stress, has been reported to induce synaptic inhibition at cholinergic endings in the nervous system [18, 19]. Consequently, the reactive oxygen species (ROS), generated from cadmium blockade of *cytochrome a3* and *cytochrome C*, participates in the elevation of cellular calcium concentration, peroxidation of neuronal membrane, and degeneration of neurons [20–23].

The bioavailability and effect of nicotine and (or) cadmium in the blood are known to be dependent on the duration of exposure to these substances either solely or combined in tobacco smoke [24–26]. Similarly, differences in the magnitude of addiction or degenerative changes seen in the brain, after a time-dependent tobacco smoke abuse, vary with the sex, age, and the developmental stage of the individual [27, 28]. Females, generally, are more sensitive to nicotine addiction than males, which also show less toxicity compared with the females [27]. Consequently, the long term effect of tobacco abuse in males has been found to be linked with irrational behaviour, violence, anxiety, mood disorders, movement disorders, and weight loss [27, 29–31]. Furthermore, movement and anxiety are affected due the effect of nicotine on reward brain regions and brain stress systems involving dopaminergic regulation, epinephrine, neuropeptide Y, and neuromuscular nicotinic stimulation [31–34].

Although the effect of nicotine (tobacco) on metabolism, anxiety, motor function, and brain reward has been studied extensively, the specific role of cadmium-nicotine treatment on these parameters (metabolism, anxiety and motor function) are yet to be elucidated, comparatively, the effect of cadmium, nicotine, and a combination of nicotine-cadmium treatment in male adult mice. We investigated pharmacologically the effect of nicotine and (or) cadmium treatment on the body weight, motor function, and anxiety linked behavioural changes during acute and chronic treatment phases. Furthermore, we sought to identify the synergistic effect of nicotine and cadmium on metabolism, anxiety, and exploratory motor dysfunction seen in tobacco abuse.

## 2. Materials and Methods

Cadmium chloride and nicotine were a generous gift from Dr. Maria Deluca of the University of Cagliari, while the normal saline used for the preparation was procured from Kanada Pharmaceuticals, Nigeria. The test chemicals were prepared weekly in normal saline and applied within the same period.

**2.1. Treatment.** Male adult albino Swiss mice ( $N = 24$ ), each weighing between 19 and 22 gms, were used for this study. The animals were obtained from the animal holding

facility of Obafemi Awolowo University, Nigeria, and allowed to acclimatize for 7 days before the commencement of treatment. Subsequently, four (4) separate groups of  $n = 6$  animals each were created at random and maintained in a standard laboratory environment with controlled humidity, pressure, and temperature. Treatment protocols were in accordance with the ethical requirements of the Animal Use and Care Committee of the Afe Babalola University, Nigeria. Animals (all groups) were treated once daily for a duration of 21 days (D1–D21); days 1–7 represent the acute treatment while days 7–21 are the chronic treatment phase. A group of  $n = 6$  mice received 1.4 mg/Kg BW of cadmium chloride prepared in normal saline (intramuscular). Another group received 0.4 mg/Kg BW of nicotine (subcutaneous) [35, 36] while a combined treatment group received nicotine (subcutaneous; 0.4 mg/Kg) and cadmium (intramuscular 1.4 mg/Kg) [37, 38]. The control was treated with normal saline (subcutaneous) for the duration of the experiment (Table S1 in the Supplementary Material available online at <http://dx.doi.org/10.1155/2014/359436>).

**2.2. Weight Measurement.** Average weight per group (gms) was measured at days 1, 7, 14, and 21 using a sensitive weighing balance (Jenway) and plotted (ANOVA) to compare weight changes for the treatment groups versus the control.

**2.3. Open Field Test.** This was done to determine exploratory motor function relative to anxiety linked behavior in a single test [39]. The animals were trained in the open field area (OFA) continually for 4 days before the actual test to facilitate acclimatization to the test equipment and environment. In addition, the animals were assisted to move or navigate the open field area when necessary during the training phase. OFA was made of white wood with dimensions, 100 cm wide, 100 cm long, and 45 cm high, and was marked with dark lines dividing the floor into 16 uniform squares and a center square. A high definition video recording system (SONY) was placed in position to capture all corners and sides of the OFA in order to record animal movement and behavior. For the actual test, animals were placed in the OFA and allowed to explore for 5 minutes following which analysis was done for motor and anxiety related functions.

**2.4. Evaluation of Motor Function.** This was done by estimating the average number of lines crossed per animal/per group for the duration of the test (5 minutes) and was expressed as the frequency of lines crossed after the acute and chronic treatment phases.

**2.5. Anxiety Linked Behaviors.** This was evaluated in relation to the observed motor functions in the OFA. Center square duration, frequency of grooming, and rearing were estimated and analyzed after the acute and chronic treatment phases to determine the level of anxiety linked with exploratory motor behavior.

**2.6. Blood Electrolyte Concentration.** At the end of the chronic phase, the animals were euthanized humanely to

obtain blood through cardiac puncture. Subsequent processing involved the use of anticoagulant (heparin), followed by centrifugation at 4°C to obtain the cold serum. The concentrations of  $\text{Ca}^{2+}$ ,  $\text{K}^+$ ,  $\text{Na}^+$ , and  $\text{Cd}^{2+}$  were determined in a mass atomic spectrometry as previously described [40]. Serum was diluted and spiked with enriched radioisotopes following which the primary standards were measured in analogue mode for  $n = 6$  samples per group.

**2.7. Statistical Analysis.** Data was plotted in ANOVA for all behavioral tests, weight, and electrolyte concentrations from mass spectrometry. Statistical significance was determined using GraphPad Prism (Version 6) ( $*P < 0.05$ ) and expressed as mean  $\pm$  SEM (standard error of mean).

### 3. Results


**3.1. Change in Average Weight.** The comparative weight analysis for nicotine, cadmium, or a combined treatments showed that nicotine treatment caused a significant increase in body weight ( $P < 0.05$ ) while cadmium ( $P < 0.001$ ) treatment induced a decline in body weight when compared with the control. Similarly, a combined nicotine-cadmium treatment decreased the body weight of the animals versus the control ( $P < 0.01$ ) during the acute phase (D1–D7). However, following chronic administration (D7–D21) further decline in weight was observed in the cadmium and cadmium-nicotine treatment groups when compared with the control ( $P < 0.01$ ). Nicotine treatment caused a significant increase in average body weight during the chronic treatment phase recording a significance of  $P < 0.05$  versus the control (Figure 1). These findings suggest that the weight loss associated with tobacco smoke is a result of cadmium or tobacco cadmium synergy rather than nicotine only; nicotine treatment increased the body weight both in the chronic and acute phases.

#### 3.2. Metal Ion Analysis in Serum

**3.2.1. Sodium ( $\text{Na}^+$ ).** Spectrum analysis showed that cadmium treatment increased serum  $\text{Na}^+$  ion level significantly above the values observed for the control. No significant change was observed in the nicotine and nicotine-cadmium treatment group when compared with the control (Figure 2(a)).

**3.2.2. Calcium ( $\text{Ca}^{2+}$ ).** The concentration of calcium in the serum of the treatment groups showed no significant change among the treatment groups for the duration of treatment (21 days). The control group recorded the lowest calcium concentration in the blood when compared with the treatments. This suggests the role of calcium in the activity of nicotine and cadmium and the significance of raised calcium levels in both cadmium and nicotine toxicity (Figure 2(b)).

**3.2.3. Potassium ( $\text{K}^+$ ).** Although alterations in potassium levels were statistically insignificant, empirically, cadmium treatment and nicotine-cadmium treatment groups showed an increase in  $\text{K}^+$  ion level when compared with the nicotine



**FIGURE 1:** Comparative body weight analysis for Cd, Nc, Cd-Nc, and control group after the acute and chronic treatment phases. As a result of acute nicotine treatment, an increase in body weight was observed in the acute (D1–D7) and chronic treatment phases (D7–D21). However, comparing the nicotine treated animals and the control, only a slight change in body weight was seen ( $P < 0.05$ ). Administration of cadmium caused an increase in body weight after the acute phase; subsequently, a decline in body weight was seen after chronic treatment with cadmium ( $P < 0.001$ ) versus the control. Similarly, combined treatment with nicotine and cadmium induced an increase in body weight after the acute treatment phase, followed by a decline after chronic treatment. In addition, the decrease in body weight observed in the nicotine-cadmium treatment group was lower than the average body weight for nicotine treatment but higher than the weight recorded in cadmium treatment group. The average weight/group was expressed as mean  $\pm$  standard error of mean (SEM) ( $*P < 0.001$ ,  $\text{Cd}^{2+}$  versus Nic;  $^{\wedge}P > 0.05$ ,  $\text{Cd}^{2+}$  versus  $\text{Cd}^{2+} + \text{Nic}$ ;  $^{\times}P > 0.05$ , Nic versus  $\text{Cd}^{2+} + \text{Nic}$ ;  $^{\#}P < 0.001$ ,  $\text{Cd}^{2+}$  versus control;  $^{\S}P > 0.05$ , Nic versus control; and  $^{\alpha}P > 0.05$ ,  $\text{Cd}^{2+} + \text{Nic}$  versus control).

treatment and the control. The potassium ion concentration change also correlated with  $\text{Ca}^{2+}$  ion level change in the treatment and control groups (Figure 2(c)).

**3.2.4. Cadmium ( $\text{Cd}^{2+}$ ).** The cadmium concentration in the cadmium and nicotine treatments reduced significantly when compared with the control and the nicotine-cadmium treatment groups. In the cadmium-nicotine treatment, blood  $\text{Cd}^{2+}$  level increased significantly when compared with the control, cadmium, and nicotine treatment groups (Figure 2(d)). This suggests the role of nicotine in increased bioaccumulation of  $\text{Cd}^{2+}$  in the blood of tobacco smokes. Also, we observed that cadmium treatment was associated with reduced serum  $\text{Cd}^{2+}$ ; although the mechanism involved is unclear, it is suspected to have resulted from the rapid clearance from the blood in urine and other physiological means (positive feedback).

**3.3. Motor Function.** The open field test (OFT) was employed to evaluate exploratory motor functions after the acute phase (D1–D7) and the chronic phase (D7–D21). Subsequent analysis involved determination of the frequency of lines




FIGURE 2: Concentration (mg/dL) of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$  and cadmium in the blood. (a) Bar chart showing the level for  $\text{Na}^+$  in blood (mg/mL). An increase in  $\text{Na}^+$  was associated with cadmium treatment when compared with the control ( $P < 0.05$ ). (b) An empirical increase in  $\text{Ca}^{2+}$  concentration was observed in the blood of the treatment groups when compared with the control ( $P < 0.05$ ). (c) Concentration of  $\text{K}^+$  in blood; no significant change was seen in the  $\text{K}^+$  ion concentration after nicotine and (or) cadmium treatment. (d) Bar chart demonstrating the blood level of cadmium in the treatment groups and control. An increase in  $\text{Cd}^{2+}$  was observed in the nicotine-cadmium treatment versus the control ( $P < 0.05$ ).

crossed in the open field area (OFA) for the nicotine, cadmium, nicotine-cadmium, and control groups. After the acute treatment phase, nicotine treatment caused an increase in exploratory motor activity when compared with the cadmium and nicotine-cadmium treatment groups ( $P < 0.05$ ). Similarly, the cadmium and nicotine-cadmium treatment recorded a significant decline in motor function versus the control ( $P < 0.01$ ). After chronic treatment (D21), the nicotine group recorded an increase in motor function when compared with the control ( $P < 0.05$ ). Also, the nicotine-cadmium treatment showed an increase in motor function versus the control ( $P < 0.05$ ) while cadmium treatment recorded a significant decline in exploratory motor function versus the control ( $P < 0.001$ ) (Figure 3).

**3.4. Centre Square Duration (CSD).** No significant change was observed in the CSD after the acute treatment phase (Figure 4(a)) when the treatment groups were compared with the control. However, after the chronic treatment phase, nicotine, cadmium, and nicotine-cadmium treatment showed a significant decrease in CSD ( $P < 0.001$ ) when compared with the control. This was observed as an increase

in time spent exploring the corners and walls of the open field area ( $\uparrow$ anxiety) (Figure 4(a)).

**3.5. Frequency of Rearing.** No significant change was seen after the chronic phase when the treatment groups were compared with the control. Similarly, the cadmium treatment group showed a decrease in the frequency of rearing versus the control ( $P < 0.001$ ). This was similar to our observations in frequency of lines crossed (Figure 3) and the CSD (Figure 4(a)). Generally, the cadmium treatment group showed a decline in motor function and an increase in anxiety related behaviour when compared with the nicotine or nicotine-cadmium treatment (Figure 4(b)).

**3.6. Frequency of Grooming.** No significant change was observed in the treatment groups after the chronic and acute phase versus the control (Figure 4(c)).

**3.7. Analysis of Anxiety in OFT.** From these findings, we deduced that a time-dependent relationship exists between the use of nicotine and (or) cadmium and the observed increase in anxiety or decreased motor function. The






FIGURE 3: Motor function analysis in open field test (OFT) for  $\text{Cd}^{2+}$  and nicotine toxicity. Bar chart (acute phase) representing the frequency of lines crossed in OFT for the treatment groups and control after the acute (D1–D7) and chronic treatment phases (D7–D21). No significant change in motor function was observed between the treatment groups and control after the acute phase. However, after the chronic phase, the cadmium ( $P < 0.01$ ) treated group showed a decline in motor function while the nicotine treatment showed an increase in exploratory motor function when compared with the control ( $P < 0.05$ ). Similarly, the nicotine-cadmium treatment recorded a significant increase in motor function when compared with the cadmium treatment group ( $P < 0.05$ ) and control.

most significant decline in exploratory motor function and increase in anxiety were observed after chronic treatment of cadmium while no significance was recorded after acute treatment. In addition, nicotine increased motor function and anxiety after the acute and chronic phases when compared with the control. A combination of nicotine and cadmium reduced the exploratory motor activity and increased anxiety in the animals after the chronic phase, thus suggesting the complimentary role of cadmium in observed anxiety related behavioural changes seen in prolonged tobacco use (Figures 4(a)–4(c)).

#### 4. Discussion

Taken together, the outcomes of this study prove that cadmium is involved in the weight, anxiety, and motor changes observed in tobacco (nicotine) use. We observed that nicotine treatment increased the body weight after the acute and chronic treatment phases while cadmium caused a decrease in weight versus the control. Interestingly, nicotine-cadmium treatment, similar to tobacco use, caused a decline in body weight after the chronic treatment phase. Subsequent analysis reveals that an increase in exploratory motor activity was associated with nicotine use after the acute and chronic treatment phases; however, an increase in anxiety was also observed in this group. Consequently, cadmium and nicotine-cadmium treatment induced a decline in exploratory motor activity and increased anxiety after the chronic phase, further suggesting the complimentary effect of cadmium on nicotine induced anxiety and decreased motor coordination observed in chronic tobacco smoke exposure.

Furthermore, the effect of cadmium and nicotine was also examined on electrolyte balance and bioaccumulation of cadmium in the serum after the chronic phase (D21). We observed that chronic cadmium treatment increased serum  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  ion concentration when compared with the control (Figures 2(a) and 2(b)). Similarly, nicotine increased serum  $\text{Ca}^{2+}$  ion level, while no changes in  $\text{K}^{+}$  and  $\text{Cd}^{2+}$  ion concentration were observed in nicotine and cadmium treatment groups (Figures 2(c) and 2(d)). Combined nicotine-cadmium treatment did not cause any significant change in serum  $\text{Ca}^{2+}$ ,  $\text{Na}^{+}$ , and  $\text{K}^{+}$  ion concentrations when compared with the control (Figures 2(a)–2(c)). Interestingly, the nicotine-cadmium treatment increased the serum level of  $\text{Cd}^{2+}$  ion significantly versus the control, nicotine, and cadmium treatment groups (Figure 2(d)); this suggests the role of nicotine in bioaccumulation of cadmium observed in chronic exposure to tobacco smoke. The observed effects of nicotine, cadmium, and nicotine-cadmium combination suggest a rapid change in metabolism which is dependent on the dose and duration of exposure [41, 42]. Furthermore, nicotine and cadmium as separate entities have been implicated in neurotoxicity and cell death by altering energy metabolism, signal transduction, reward, and induction of brain stress [43–46]. Previous studies have shown that the anxiety and exploratory motor changes associated with acute and chronic tobacco use involved alteration in neurotransmitters such as GABA [47, 48], dopamine [49, 50], and norepinephrine in brain [51]. Similarly, the nicotine in tobacco smoke affects the level of acetylcholine and epinephrine in the neuromuscular junction, arteries, glands, and other viscera [51, 52].

Dopaminergic  $\text{D}_1$  and  $\text{D}_2$  stimulation in the hippocampus has been implicated in nicotine-mediated anxiety and reward system [46–48, 53]. Nicotinic modulation of dopaminergic system is believed to be involved in increased motor function, brain reward, addiction, anxiety, and late onset Parkinson's disease observed in smokers [54, 55]. However, nicotinic stimulation of acetylcholine receptors induces the release of acetylcholine and norepinephrine in the central and peripheral nervous system and has been found to involve transient calcium release and release of high frequency pulses associated with anxiety and motor dysfunctions [56]. Interestingly, cadmium and the associated carbon dioxide produced in oxidative stress have been linked with the inhibition of synaptic activity at cholinergic endings [57, 58]. Although, its actual role in the modulation of neurotransmitter and reward system is elusive, our findings suggest nicotine increases the serum concentration of cadmium when both molecules are coadministered. As a result, both nicotinic stimulation of cholinergic receptors and cadmium induced oxidative stress are capable of inducing elevated calcium and sodium ions concentration in the serum (Figure 2), an effect that is linked to the observed anxiety and altered motor function in these groups.

Nicotine-mediated cholinergic stimulation also involves the activation of neuropeptide Y and cAMP, which represents calcium dependent mechanisms involved in cholinergic hippocampal-mediated anxiety, epinephrine-mediated anxiety, and cholinergic neuromuscular stimulation [59–62].




FIGURE 4: (a) Anxiety related behavioral analysis (I) in open field test (OFT) for  $Cd^{2+}$  and nicotine toxicity. Bar chart representing the center square duration (CSD) in OFT for the treated groups and control after the acute (D1–D7) and chronic (D7–D21) treatment phases. The duration spent in the center square of the open field area was estimated to determine the mean duration for each group at the end of the acute and chronic phases. After the acute treatment phase, no significant change in anxiety linked behavior was seen in the nicotine and (or) cadmium treatment groups. After chronic treatment, cadmium and nicotine treatment increased anxiety linked behavior in the animals, as shown by a significant decrease in CSD (D7–D21). Similarly, a combined nicotine-cadmium treatment also caused an increase in anxiety linked behavior (reduced CSD) when compared with the control ( $^{*}P < 0.001$ ). (b) Anxiety related behavioral analysis (II) in open field test (OFT) for the treatment and control groups. The mean frequency of rearing was determined for the nicotine and (or) treatment groups versus the control after the acute (D1–D7) and chronic (D7–D21) treatment phases. No significant change in the frequency of rearing for the treatment groups after the acute treatment phase. After the chronic phase, subsequent tests show a significant reduction in rearing time ( $\uparrow$  anxiety) for the cadmium treatment group when compared with the control, nicotine, and nicotine-cadmium treatment groups ( $^{***}P < 0.001$ ). (c) Anxiety related behavioral analysis (III) in open field test (OFT) for  $Cd^{2+}$  and (or) nicotine toxicity. Bar chart representing the frequency of grooming in OFT for the treatment groups and control after the acute (D1–D7) and chronic (D7–D21) treatment phases. No significant change in the frequency of grooming was observed after the acute treatment phase. Similarly, no significant change was seen in the frequency of grooming after the chronic treatment phase for all treatment groups versus the control.

Similarly, cadmium inhibits *cytochrome C* and *a3* in the mitochondria leading to ROS-dependent calcium release, anxiety, and synaptic excitotoxicity [63, 64]. Other studies have shown that cadmium affects the level of monoamine in the hypothalamus and hippocampus without a corresponding change in norepinephrine [63]. Interestingly, an increase in anxiety index was observed in the cadmium treated

animals, similar to our findings. However, the mechanism of nicotine-induced bioaccumulation of cadmium and its relationship with the anxiety linked behavioural changes remains elusive. Based on the outcome of this study and previous studies, we hypothesise that cadmium-mediated synaptic inhibition, monoaminergic alterations, and induced calcium-shift are important in anxiety linked behaviour and

decline in motor functions. As a result it complements the neural and behavioural changes attributed to nicotine in long term tobacco use.

## 5. Conclusion

Based on the outcome of this study, we infer that the motor and anxiety related behavioural changes observed in prolonged tobacco use in adult male mice are a result of the synergistic effect of nicotine-cadmium on synaptic plasticity, neurotransmitter function, and brain reward system. Also, nicotine and cadmium increased serum  $\text{Ca}^{2+}$  ions and caused no significant change in  $\text{K}^+$  ion concentration. However, combined nicotine-cadmium showed that nicotine facilitates bioaccumulation of cadmium in serum, resulting in a decrease in motor function and increased anxiety when compared with the nicotine and cadmium treatment groups. In addition, nicotine induced increase in weight during the acute and chronic phases while cadmium and cadmium-nicotine treatment caused a decrease in weight after the chronic phase. Ultimately, we conclude that nicotine and cadmium interaction in tobacco use is time-dependent and affects metabolism, electrolytes, and neurotransmitter functions that translate into the observed increased anxiety, decreased weight and altered motor function after chronic treatment.

## Conflict of Interests

The authors hereby declare there is no conflict of interests associated with this study or any of the procedures and materials used for the purpose of the study.

## Acknowledgments

The authors acknowledge Dr. Maria A. Deluca of Department of Biological Science, University of Cagliari, for her generous gift of the nicotine salt and also Mr. Oso of the Department of Chemical Sciences, Afe Babalola University, who worked with them in the preparation of salts and reagents. They will also like to appreciate Professor Micaela Morelli of the University of Cagliari for hosting Philip Adeniyi during his visit for the CEAN-ISON fellowship. This work was funded by the ABUAD-DOR for Neurotoxicology Research Grant issued to the College of Medicine and Health Sciences, Afe Babalola University.

## References

- [1] L. Robertson, R. McGee, and R. J. Hancox, "Smoking cessation and subsequent weight change," *Nicotine & Tobacco Research*, 2014.
- [2] A. Stanton and G. Grimshaw, "Tobacco cessation interventions for young people," *The Cochrane Database of Systematic Reviews*, vol. 8, Article ID CD003289, 2013.
- [3] S. E. Jones and S. Hamilton, "Introducing a new stop smoking service in an acute UK hospital: a qualitative study to evaluate service user experience," *European Journal of Oncology Nursing*, vol. 17, no. 5, pp. 563–569, 2013.
- [4] T. Raupach, P. H. J. Hoogsteder, and C. P. O. van Schayck, "Nicotine vaccines to assist with smoking cessation: current status of research," *Drugs*, vol. 72, no. 4, pp. e1–e16, 2012.
- [5] H. S. Menossi, A. E. Goudriaan, C. D. A.-M. Périco et al., "Neural bases of pharmacological treatment of nicotine dependence—insights from functional brain imaging: a systematic review," *CNS Drugs*, vol. 27, no. 11, pp. 921–941, 2013.
- [6] S. Zhang, K. Hyrc, S. Wang, and B. M. Wice, "Xenin-25 increases cytosolic free calcium levels and acetylcholine release from a subset of myenteric neurons," *The American Journal of Physiology: Gastrointestinal and Liver Physiology*, vol. 303, no. 12, pp. G1347–G1355, 2012.
- [7] N. da Silva, C. E. Herron, K. Stevens, C. A. B. Jollimore, S. Barnes, and M. E. Kelly, "Metabotropic receptor-activated calcium increases and store-operated calcium influx in mouse Müller cells," *Investigative Ophthalmology and Visual Science*, vol. 49, no. 7, pp. 3065–3073, 2008.
- [8] H. Wang, W. Sun, J. Ma, Y. Pan, L. Wang, and W. Zhang, "Polycystin-1 mediates mechanical strain-induced osteoblastic mechanoresponses via potentiation of intracellular calcium and Akt/ $\beta$ -catenin pathway," *PLoS ONE*, vol. 9, no. 3, Article ID e91730, 2014.
- [9] S. N. Grishin, R. R. Kamaliev, A. Y. Teplov, and A. U. Ziganshin, "Opposite effect of ATP on contraction force of tonic and phasic skeletal muscles in frogs," *Bulletin of Experimental Biology and Medicine*, vol. 151, no. 3, pp. 280–283, 2011.
- [10] Y.-M. Yang and G.-T. Liu, "Damaging effect of cigarette smoke extract on primary cultured human umbilical vein endothelial cells and its mechanism," *Biomedical and Environmental Sciences*, vol. 17, no. 2, pp. 121–134, 2004.
- [11] D. Pogocki, T. Ruman, M. Danilczuk, M. Celuch, and E. Wałajtyś-Rode, "Application of nicotine enantiomers, derivatives and analogues in therapy of neurodegenerative disorders," *European Journal of Pharmacology*, vol. 563, no. 1–3, pp. 18–39, 2007.
- [12] N. Miura, D. Yuki, N. Minami, A. Takehi, and M. Onozawa, "Pharmacokinetic analysis of nicotine when using non-combustion inhaler type of tobacco product in Japanese adult male smokers," *Regulatory Toxicology and Pharmacology*, vol. 67, no. 2, pp. 198–205, 2013.
- [13] X. M. Shao, B. Xu, J. Liang, X. Xie, Y. Zhu, and J. L. Feldman, "Nicotine delivery to rats via lung alveolar region-targeted aerosol technology produces blood pharmacokinetics resembling human smoking," *Nicotine and Tobacco Research*, vol. 15, no. 7, pp. 1248–1258, 2013.
- [14] M. A. Babizhayev, "The detox strategy in smoking comprising nutraceutical formulas of non-hydrolyzed carnosine or carnitine used to protect human health," *Human and Experimental Toxicology*, vol. 33, no. 3, pp. 284–316, 2014.
- [15] S. Storage, M. A. Mandelkern, J. Phuong, M. Kozman, M. K. Neary, and A. L. Brody, "A positive relationship between harm avoidance and brain nicotinic acetylcholine receptor availability," *Psychiatry Research*, vol. 214, no. 3, pp. 415–421, 2013.
- [16] E. Kleszczewska, J. Hejza, and T. Kleszczewski, "Determination of cadmium level in passive tobacco smokers schizophrenia patients," *Przegląd Lekarski*, vol. 63, no. 10, pp. 998–1001, 2006.
- [17] P. Checconi, R. Sgarbanti, I. Celestino et al., "The environmental pollutant cadmium promotes influenza virus replication in MDCK cells by altering their redox state," *International Journal of Molecular Sciences*, vol. 14, no. 2, pp. 4148–4162, 2013.

- [18] M. González-Estechea, E. Trasobares, M. Fuentes et al., "Blood lead and cadmium levels in a six hospital employee population. PESA study, 2009," *Journal of Trace Elements in Medicine and Biology*, vol. 25, supplement 1, pp. S22–S29, 2011.
- [19] M. E. Culbreth, J. A. Harrill, T. M. Freudenrich, W. R. Mundy, and T. J. Shafer, "Comparison of chemical-induced changes in proliferation and apoptosis in human and mouse neuroprogenitor cells," *NeuroToxicology*, vol. 33, no. 6, pp. 1499–1510, 2012.
- [20] Y. Kurata, O. Katsuta, T. Doi et al., "Chronic cadmium treatment induces tubular nephropathy and osteomalacic osteopenia in ovariectomized cynomolgus monkeys," *Veterinary Pathology*, vol. 51, no. 5, pp. 919–931, 2013.
- [21] Y.-F. Song, Z. Luo, Y.-X. Pan, X. Liu, C. Huang, and Q.-L. Chen, "Effects of copper and cadmium on lipogenic metabolism and metal element composition in the javelin goby (*Synechogobius hasta*) after single and combined exposure," *Archives of Environmental Contamination and Toxicology*, vol. 67, no. 2, pp. 167–180, 2014.
- [22] S. Haeuem and A. El Hani, "Effect of cadmium on lipid peroxidation and on some antioxidants in the liver, kidneys and testes of rats given diet containing cadmium-polluted radish bulbs," *Journal of Toxicologic Pathology*, vol. 26, no. 4, pp. 359–364, 2013.
- [23] B. Wang and Y. Du, "Cadmium and its neurotoxic effects," *Oxidative Medicine and Cellular Longevity*, vol. 2013, Article ID 898034, 12 pages, 2013.
- [24] K. P. Timms, D. E. Rivera, L. M. Collins, and M. E. Piper, "A dynamical systems approach to understanding self-regulation in smoking cessation behavior change," *Nicotine and Tobacco Research*, vol. 16, no. 2, pp. S159–S168, 2014.
- [25] E. A. Kiyatkin, "Critical role of peripheral sensory systems in mediating the neural effects of nicotine following its acute and repeated exposure," *Review of Neuroscience*, vol. 25, no. 2, pp. 207–221, 2014.
- [26] S. P. Barrett and C. Darredeau, "The acute effects of nicotine on the subjective and behavioural responses to denicotinized tobacco in dependent smokers," *Behavioural Pharmacology*, vol. 23, no. 3, pp. 221–227, 2012.
- [27] S. E. Santiago and K. J. Huffman, "Prenatal nicotine exposure increases anxiety and modifies sensorimotor integration behaviors in adult female mice," *Neuroscience Research*, vol. 79, no. 1, pp. 41–51, 2014.
- [28] H. Yamada, M. Bishnoi, K. F. M. Keijzers et al., "Preadolescent tobacco smoke exposure leads to acute nicotine dependence but does not affect the rewarding effects of nicotine or nicotine withdrawal in adulthood in rats," *Pharmacology Biochemistry and Behavior*, vol. 95, no. 4, pp. 401–409, 2010.
- [29] J. Wu, M. Gao, J. X. Shen, W. X. Shi, A. M. Oster, and B. S. Gutkin, "Cortical control of VTA function and influence on nicotine reward," *Biochemical Pharmacology*, vol. 86, no. 8, pp. 1173–1180, 2013.
- [30] J. Ren, J. Sun, Y. Zhang et al., "Down-regulation of Decapping Protein 2 mediates chronic nicotine exposure-induced locomotor hyperactivity in *Drosophila*," *PLoS ONE*, vol. 7, no. 12, Article ID e52521, 2012.
- [31] B. N. Cohen, E. D. W. Mackey, S. R. Grady et al., "Nicotinic cholinergic mechanisms causing elevated dopamine release and abnormal locomotor behavior," *Neuroscience*, vol. 200, pp. 31–41, 2012.
- [32] C. V. Abobo, J. Ma, and D. Liang, "Effect of menthol on nicotine pharmacokinetics in rats after cigarette smoke inhalation," *Nicotine and Tobacco Research*, vol. 14, no. 7, pp. 801–808, 2012.
- [33] M. Sofuoğlu, A. I. Herman, H. Nadim, and P. Jatlow, "Rapid nicotine clearance is associated with greater reward and heart rate increases from intravenous nicotine," *Neuropsychopharmacology*, vol. 37, no. 6, pp. 1509–1516, 2012.
- [34] Z. X. Xi, K. Spiller, and E. L. Gardner, "Mechanism-based medication development for the treatment of nicotine dependence," *Acta Pharmacologica Sinica*, vol. 30, no. 6, pp. 723–739, 2009.
- [35] A. C. Harris, C. Mattson, D. Shelley, and M. G. LeSage, "Restraint stress attenuates nicotine's locomotor stimulant but not discriminative stimulus effects in rats," *Pharmacology Biochemistry and Behavior*, vol. 124, pp. 92–100, 2014.
- [36] F. Panin, A. Lintas, and M. Diana, "Nicotine-induced increase of dopaminergic mesoaccumbal neuron activity is prevented by acute restraint stress. In vivo electrophysiology in rats," *European Neuropsychopharmacology*, vol. 24, no. 7, pp. 1175–1180, 2014.
- [37] K. Łukawski, B. Nieradko, and M. Sieklucka-Dziuba, "Effects of cadmium on memory processes in mice exposed to transient cerebral oligemia," *Neurotoxicology and Teratology*, vol. 27, no. 4, pp. 575–584, 2005.
- [38] United States Environmental Protection Agency, "Acute exposure guide line level for cadmium," 2014, <http://www.epa.gov/oppt/aegl/pubs/rest303.html>.
- [39] M. Mesfin, K. Asres, and W. Shibeshi, "Evaluation of anxiolytic activity of the essential oil of the aerial part of *Foeniculum vulgare* Miller in mice," *BMC Complementary and Alternative Medicine*, vol. 14, no. 1, p. 310, 2014.
- [40] S. Döker, M. Hazar, M. Uslu, İ. Okan, E. Kafkas, and I. I. Boşgelmez, "Influence of training frequency on serum concentrations of some essential trace elements and electrolytes in male swimmers," *Biological Trace Element Research*, vol. 158, no. 1, pp. 15–21, 2014.
- [41] A. V. Ivanina, E. P. Sokolov, and I. M. Sokolova, "Effects of cadmium on anaerobic energy metabolism and mRNA expression during air exposure and recovery of an intertidal mollusk *Crassostrea virginica*," *Aquatic Toxicology*, vol. 99, no. 3, pp. 330–342, 2010.
- [42] H. Lu, G. Yuan, Z. Yin et al., "Effects of subchronic exposure to lead acetate and cadmium chloride on rat's bone: Ca and Pi contents, bone density, and histopathological evaluation," *International Journal of Clinical and Experimental Pathology*, vol. 7, no. 2, pp. 640–647, 2014.
- [43] V. Prabhakar, G. Jayakrishnan, S. V. Nair, and B. Ranganathan, "Determination of trace metals, moisture, pH and assessment of potential toxicity of selected smokeless tobacco products," *Indian Journal of Pharmaceutical Sciences*, vol. 75, no. 3, pp. 262–269, 2013.
- [44] B. Bodereau-Dubois, O. List, D. Calas-List et al., "Transmembrane potential polarization, calcium influx, and receptor conformational state modulate the sensitivity of the imidacloprid-insensitive neuronal insect nicotinic acetylcholine receptor to neonicotinoid insecticides," *Journal of Pharmacology and Experimental Therapeutics*, vol. 341, no. 2, pp. 326–339, 2012.
- [45] M. F. Borgerding, J. A. Bodnar, G. M. Curtin, and J. E. Swauger, "The chemical composition of smokeless tobacco: a survey of products sold in the United States in 2006 and 2007," *Regulatory Toxicology and Pharmacology*, vol. 64, no. 3, pp. 367–387, 2012.
- [46] N. B. Kroemer, A. Guevara, S. Vollstädt-Klein, and M. N. Smolka, "Nicotine alters food-cue reactivity via networks extending from the hypothalamus," *Neuropsychopharmacology*, vol. 38, no. 11, pp. 2307–2314, 2013.



- [47] A. P. Varani and G. N. Balerio, "GABAB receptors involvement in the effects induced by nicotine on anxiety-related behaviour in mice," *Pharmacological Research*, vol. 65, no. 5, pp. 507–513, 2012.
- [48] S. D. Iñiguez, B. L. Warren, E. M. Parise et al., "Nicotine exposure during adolescence induces a depression-like state in adulthood," *Neuropsychopharmacology*, vol. 34, no. 6, pp. 1609–1624, 2009.
- [49] M. R. Zarrindast, N. Naghdi-Sedeh, M. Nasehi, H. Sahraei, F. Bahrami, and F. Asadi, "The effects of dopaminergic drugs in the ventral hippocampus of rats in the nicotine-induced anxiogenic-like response," *Neuroscience Letters*, vol. 475, no. 3, pp. 156–160, 2010.
- [50] S. Weiss, M. Nosten-Bertrand, J. M. McIntosh, B. Giros, and M.-P. Martres, "Nicotine improves cognitive deficits of dopamine transporter knockout mice without long-term tolerance," *Neuropsychopharmacology*, vol. 32, no. 12, pp. 2465–2478, 2007.
- [51] F. Beiranvand, C. Zlabinger, A. Orr-Urtreger, R. Ristl, S. Huck, and P. Scholze, "Nicotinic acetylcholine receptors control acetylcholine and noradrenaline release in the rodent habenulo-interpeduncular complex," *British Journal of Pharmacology*, 2014.
- [52] S. Zevin and N. L. Benowitz, "Drug interactions with tobacco smoking. An update," *Clinical Pharmacokinetics*, vol. 36, no. 6, pp. 425–438, 1999.
- [53] M. Haass, G. Richardt, T. Brenn, E. Schomig, and A. Schomig, "Nicotine-induced release of noradrenaline and neuropeptide Y in guinea-pig heart: role of calcium channels and protein kinase C," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 344, no. 5, pp. 527–531, 1991.
- [54] A. Kucinski, S. Wersinger, E. K. Stachowiak et al., "Neuronal nicotinic receptor agonists ameliorate spontaneous motor asymmetries and motor discoordination in a unilateral mouse model of Parkinson's disease," *Pharmacology Biochemistry and Behavior*, vol. 111, pp. 1–10, 2013.
- [55] M. Ohashi, A. Saitoh, M. Yamada, J. I. Oka, and M. Yamada, "Riluzole in the prelimbic medial prefrontal cortex attenuates veratrine-induced anxiety-like behaviors in mice," *Psychopharmacology*, 2014.
- [56] C. Le Magueresse and E. Cherubini, "Presynaptic calcium stores contribute to nicotine-elicited potentiation of evoked synaptic transmission at CA3-CA1 connections in the neonatal rat hippocampus," *Hippocampus*, vol. 17, no. 4, pp. 316–325, 2007.
- [57] N. Djulančić, V. Radojičić, and M. Srbinovska, "The influence of tobacco blend composition on carbon monoxide formation in mainstream cigarette smoke," *Arhiv za Higijenu Rada i Toksikologiju*, vol. 64, no. 1, pp. 107–113, 2013.
- [58] S. M. Bierbower and R. L. Cooper, "The mechanistic action of carbon dioxide on a neural circuit and NMJ communication," *Journal of Experimental Zoology A: Ecological Genetics and Physiology*, vol. 319, no. 6, pp. 340–354, 2013.
- [59] S. Nuutinen, E. Ekokoski, E. Lahdensuo, and R. K. Tuominen, "Nicotine-induced upregulation of human neuronal nicotinic  $\alpha 7$ -receptors is potentiated by modulation of cAMP and PKC in SH-EPI-h $\alpha 7$  cells," *European Journal of Pharmacology*, vol. 544, no. 1–3, pp. 21–30, 2006.
- [60] M. F. Ferrari, E. F. Coelho, K. L. Farizatto, G. Chadi, and D. R. Fior-Chadi, "Modulation of tyrosine hydroxylase, neuropeptide Y, glutamate, and substance P in ganglia and brain areas involved in cardiovascular control after chronic exposure to nicotine," *International Journal of Hypertension*, vol. 2011, Article ID 216464, 9 pages, 2011.
- [61] C. Aydin, O. Oztan, and C. Isgor, "Effects of a selective Y2R antagonist, JNJ-31020028, on nicotine abstinence-related social anxiety-like behavior, neuropeptide Y and corticotropin releasing factor mRNA levels in the novelty-seeking phenotype," *Behavioural Brain Research*, vol. 222, no. 2, pp. 332–341, 2011.
- [62] C. Aydin, O. Oztan, and C. Isgor, "Vulnerability to nicotine abstinence-related social anxiety-like behavior: molecular correlates in neuropeptide Y, Y2 receptor and corticotropin releasing factor," *Neuroscience Letters*, vol. 490, no. 3, pp. 220–225, 2011.
- [63] M. L. Leret, J. A. San Millán, and M. T. Antonio, "Perinatal exposure to lead and cadmium affects anxiety-like behaviour," *Toxicology*, vol. 186, no. 1–2, pp. 125–130, 2003.
- [64] E. G. Moreira, I. Vassilieff, and V. S. Vassilieff, "Developmental lead exposure: behavioral alterations in the short and long term," *Neurotoxicology and Teratology*, vol. 23, no. 5, pp. 489–495, 2001.



**Hindawi**  
Submit your manuscripts at  
<http://www.hindawi.com>

