

Article

Changes in testing for and incidence of celiac disease in the UK: a population-based cohort study

West, Joe, Otete, Harmony, Sultan, Alyshah A and Crooks, Colin J

Available at <http://clock.uclan.ac.uk/27718/>

West, Joe, Otete, Harmony ORCID: 0000-0003-2467-2605, Sultan, Alyshah A and Crooks, Colin J (2019) Changes in testing for and incidence of celiac disease in the UK: a population-based cohort study. Epidemiology . ISSN 1044-3983

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1097/EDE.0000000000001006>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Epidemiology Publish Ahead of Print

DOI:10.1097/EDE.0000000000001006

Changes in testing for and incidence of celiac disease in the UK: a population-based cohort study

Abbreviated title: Incidence of Celiac disease

Professor Joe West ^{1,4*}, Dr Harmony Otete ^{2,6}, Dr Alyshah A Sultan ³, Dr Colin J Crooks ^{4,5}

*Corresponding author:

1 Division of Epidemiology and Public Health, Clinical Sciences Building 2, City Hospital Campus, University of Nottingham, NG5 1PB, UK. E-mail: joe.west@nottingham.ac.uk, Telephone: +44 (0)1158231345*

2 School of Pharmacy, University of Nottingham, UK

3 Research Institute for Primary Care & Health Sciences, Primary Care Sciences, Keele University, UK

4 NIHR Nottingham Biomedical Research Centre at the Nottingham University Hospitals NHS Trust and University of Nottingham, Nottingham, UK.

5 Nottingham Digestive Diseases Centre, School of Medicine, University of Nottingham

6 School of Medicine & Dentistry, Harrington building, University of Central Lancashire, Preston PR17BH

Key words: incidence: celiac disease; prevalence; serologic tests

Source of funding: Coeliac UK

Conflict of interests: None declared

Author contributions:

All authors meet the following criteria in that they carried out substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND drafting the work or revising it critically for important intellectual content; AND final approval of the version to be published; AND agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Keywords: celiac disease; incidence; prevalence; trends

Data and code: The data used in the article were acquired from the Clinical Practice Research Datalink under a license to the University Of Nottingham. These data are not permitted to be distributed to third parties, but are available from CPRD following acquisition of the appropriate approvals and payment.

Word count: 600

To the Editor:

The diagnosis rates of celiac disease differ substantially between countries¹. Intriguingly, there has been recent evidence from Olmstead County, USA, and Finland that in the last 5-10 years incidence has leveled off or even declined^{2, 3}. In most populations the prevalence also varies widely, with serologic prevalence from 0% to 1.87% and clinical prevalence from 0.9 to 12.9 per 100000¹. Understanding of why this variation exists is minimal, yet one of the key aspects governing incidence rates of any disease are factors related to the health system, such as the availability and use of diagnostic tests. We previously reported rising incidence rates of celiac disease⁴ from 1990 to 2011 with differences related to socioeconomic deprivation in the UK. Although national guidance on recognition and diagnosis of celiac disease published in 2009⁵ suggested widening the patient groups that should be tested for celiac disease, National Health System (NHS) financial constraints could have hindered implementation of these guidelines. Indeed, in the USA researchers have observed that over the period 2000-2010 there was a marked decrease in treated prevalence of many diseases alongside a sustained period of reduced spending on health care⁶.

We used the UK Clinical Practice Research Datalink GOLD (Independent Scientific Advisory Committee approval 16_130) to estimate the European (EUROSTAT EU-27 plus EFTA 2013 population⁷) age-standardized incidence rates of celiac disease⁸ 2005-2015 and the corresponding rates of serologic testing (Anti-Tissue Transglutaminase antibody (TTG) and anti-Endomysial antibody (EMA)) for the disease. We used Joinpoint analysis⁹ to examine statistical evidence of changes in the rates of diagnosis and testing during this period. We estimated celiac disease point prevalence based on all contributing patients at 30 June 2015 and estimated incidence rate ratios (IRR) using Poisson regression for testing and incidence rates.

There were 8177 incident cases of celiac diseases diagnosed among 45,539,211 million person-years. The overall incidence rate between 2005 and 2015 was 18 per 100,000 person-years, serological testing rate was 118 per 100,000 person-years, and point prevalence on the 30th June 2015 was 0.30% (95% CI 0.30-0.31). Incidence rates of celiac disease were highest in people aged between 60 and 69 years (23 per 100,000 person-years) whereas the rate of serologic testing was highest in those aged 20-29 (233 per 100,000 person-years). For the calendar period 2005-2015 there was an increase in European age-standardized incidence rates from 2005 until 2012 and then a plateau effect (Figure 1). Serologic testing increased and then decreased during the same period (Figure 2). Joinpoint analysis identified that there were changes in the rates of both diagnosis and testing at 2012 (95% CI 2007-2013) and 2011 (95% CI 2010-2013) respectively. The Joinpoint analysis is presented in the table.

In this study we found that European age-standardized rates of diagnosis of celiac disease and serological testing have, since 2011, respectively leveled off and declined, while prevalence increased from 0.24%⁴ to 0.3%.

This could be because, since 2010, the UK NHS has been operating under a period of financial austerity. While health funding has been forecast to grow 1.2 per cent in real terms between 2009/10 and 2020/21 this is below the long-term average increase in health spending of approximately 4 per cent a year since the NHS was established in 1948¹⁰. Alternatively, clinicians based in primary care could be carrying out more targeted use of testing in certain age or at-risk groups, leading to an overall reduction in testing. We may have missed some tests carried out in secondary care as we did not have access to these. We found some evidence that testing did vary by age, disproportionately to disease incidence, in that the highest testing rate was in those aged 20-29 yet the highest incidence rate was in the 60-69 year old group. Finally, it

is possible that following several years of increasing diagnosis rates prior to 2011⁴ that the threshold of clinically identifiable celiac disease in the UK has been reached and a steady-state incidence rate obtained.

ACCEPTED

References

1. Kang JY, Kang AH, Green A, et al. Systematic review: worldwide variation in the frequency of celiac disease and changes over time. *Aliment Pharmacol Ther* 2013.
2. Virta LJ, Saarinen MM, Kolho KL. Declining trend in the incidence of biopsy-verified celiac disease in the adult population of Finland, 2005-2014. *Aliment Pharmacol Ther* 2017;46:1085-1093.
3. Ludvigsson JF, Rubio-Tapia A, van Dyke CT, et al. Increasing incidence of celiac disease in a North American population. *Am J Gastroenterol* 2013;108:818-24.
4. West J, Fleming KM, Tata LJ, et al. Incidence and prevalence of celiac disease and dermatitis herpetiformis in the UK over two decades: population-based study. *Am J Gastroenterol* 2014;109:757-68.
5. Celiac disease. Recognition and assessment of celiac disease. NICE clinical guideline 86. London: National Institute for Health and Clinical Excellence, 2009:1-86.
6. Cox C, Dunn A, Rittmueller R, et al. A new way of measuring health costs sheds light on recent health spending trends. Volume 2018: Peterson-Kaiser, 2016.
7. Report of Eurostat's task force : 2013 edition. Luxembourg: Publications Office of the European Union, 2013:121.
8. West J. Celiac disease: studies of its frequency and consequence. *Epidemiology and Public Health*. Nottingham: Nottingham, 2005:162.
9. Kim HJ, Fay MP, Feuer EJ, et al. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med* 2000;19:335-51 (correction: 2001;20:655).
10. King's Fund. The NHS budget and how it has changed. Volume 2019, 2018.

Table. Joinpoint analysis for Celiac disease and Serological testing rates.

	Segment	Lower Endpoint	Upper Endpoint	APC	Lower CI	Upper CI	Test Statistic (t)	Prob > t
Celiac disease	1	2005	2012	5.0 [^]	3.1	7.1	6.4	0.0
	2	2012	2015	-0.3	-7.2	7.2	-0.1	0.9
Serology	1	2005	2011	15.4 [^]	12.4	18.5	13.4	0.0
	2	2011	2015	-13.9 [^]	-18.3	-9.2	-7.0	0.0

APC – Annual Percentage Change

[^] - indicates that the APC is significantly different from zero at the alpha = 0.05

Figure Legends.

Figure 1. European age-standardized incidence rates of celiac disease per 100,000 person-years




Figure 2. European age-standardized serological testing rates per 100,000 person-years (TTG and EMA) for celiac disease

