

Research Article

Solutions and Improved Perturbation Analysis for the Matrix Equation $X - A^* X^{-p} A = Q$ ($p > 0$)

Jing Li

School of Mathematics and Statistics, Shandong University, Weihai, Weihai 264209, China

Correspondence should be addressed to Jing Li; xlijing@sdu.edu.cn

Received 16 November 2012; Revised 2 May 2013; Accepted 7 May 2013

Academic Editor: Carlos Vazquez

Copyright © 2013 Jing Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The nonlinear matrix equation $X - A^* X^{-p} A = Q$ with $p > 0$ is investigated. We consider two cases of this equation: the case $p \geq 1$ and the case $0 < p < 1$. In the case $p \geq 1$, a new sufficient condition for the existence of a unique positive definite solution for the matrix equation is obtained. A perturbation estimate for the positive definite solution is derived. Explicit expressions of the condition number for the positive definite solution are given. In the case $0 < p < 1$, a new sharper perturbation bound for the unique positive definite solution is derived. A new backward error of an approximate solution to the unique positive definite solution is obtained. The theoretical results are illustrated by numerical examples.

1. Introduction

In this paper, we consider the Hermitian positive definite solution of the nonlinear matrix equation

$$X - A^* X^{-p} A = Q, \quad (1)$$

where A , Q , and X are $n \times n$ complex matrices, Q is a positive definite matrix, and $p > 0$. This type of nonlinear matrix equations arises in the analysis of ladder networks, the dynamic programming, control theory, stochastic filtering, statistics, and many applications [1–7].

In the last few years, (1) was investigated in some special cases. For the nonlinear matrix equations $X - A^* X^{-1} A = Q$ [8–12], $X - A^* X^{-2} A = Q$ [13, 14], $X - A^* X^{-n} A = Q$ [15, 16], and $X^s - A^* X^{-t} A = Q$ [17], there were many contributions in the literature to the solvability, numerical solutions, and perturbation analysis. In addition, the related equations $X + A^* X^{-1} A = Q$ [9–11, 18–23], $X + A^* X^{-2} A = Q$ [13, 24, 25], $X + A^* X^{-n} A = Q$ [16, 26], $X^s + A^* X^{-t} A = Q$ [17, 27–30], $X + A^* X^{-q} A = Q$ [31–33], and $X \pm \sum_{i=1}^m A_i^* X^{-1} A_i = Q$ [34–36] were studied by many scholars.

In [31], a sufficient condition for the equation $X - A^* X^{-p} A = Q$ ($0 < p \leq 1$) to have a unique positive definite solution was provided. When the coefficient matrix A is nonsingular, several sufficient conditions for the equation $X -$

$A^* X^{-q} A = Q$ ($q \geq 1$) to have a unique positive definite solution were given in [37]. When the coefficient matrix A is an arbitrary complex matrix, necessary conditions and sufficient conditions for the existence of positive definite solutions for the equation $X - A^* X^{-q} A = Q$ ($q \geq 1$) were derived in [38]. Li and Zhang in [39] proved that there always exists a unique positive definite solution to the equation $X - A^* X^{-p} A = Q$ ($0 < p < 1$). They also obtained a perturbation bound and a backward error of an approximate solution for the unique solution of the equation $X - A^* X^{-p} A = Q$ ($0 < p < 1$).

As a continuation of the previous results, the rest of the paper is organized as follows. Section 2 gives some preliminary lemmas that will be needed to develop this work. In Section 3, a new sufficient condition for (1) with $p \geq 1$ having a unique positive definite solution is derived. In Section 4, a perturbation bound for the positive definite solution to (1) with $p \geq 1$ is given. In Section 5, applying the integral representation of matrix function, we also discuss the explicit expressions of condition number for the positive definite solution to (1) with $p \geq 1$. Furthermore, in Section 6, a new sharper perturbation bound for the unique positive definite solution to (1) with $0 < p < 1$ is given. In Section 7, a new backward error of an approximate solution to (1) with $0 < p < 1$ is obtained. Finally, several numerical examples are presented in Section 8.

We denote by $\mathcal{C}^{n \times n}$ the set of $n \times n$ complex matrices, by $\mathcal{H}^{n \times n}$ the set of $n \times n$ Hermitian matrices, by I the identity matrix, by $\|\cdot\|$ the spectral norm, by $\|\cdot\|_F$ the Frobenius norm and by $\lambda_{\max}(M)$ and $\lambda_{\min}(M)$ the maximal and minimal eigenvalues of M , respectively. For $A = (a_{ij}) \in \mathcal{C}^{n \times n}$ with columns a_i and a matrix B , $A \otimes B = (a_{ij}B)$ is a Kronecker product, and $\text{vec } A$ is a vector defined by $\text{vec } A = (a_1^T, \dots, a_n^T)^T$. For $X, Y \in \mathcal{H}^{n \times n}$, we write $X \geq Y$ (resp., $X > Y$) if $X - Y$ is Hermitian positive semidefinite (resp., definite). Let $\bar{\kappa} = \lambda_{\max}(A^*A)$, $\underline{\kappa} = \lambda_{\min}(A^*A)$.

2. Preliminaries

In this section we quote some preliminary lemmas that we use later.

Lemma 1 (see [39, Lemma 3.2]). *For every positive definite matrix $X \in \mathcal{H}^{n \times n}$ and $0 < p < 1$, then*

- (i) $X^{-p} = (\sin p\pi/\pi) \int_0^\infty (\lambda I + X)^{-1} \lambda^{-p} d\lambda$.
- (ii) $X^{-p} = (\sin p\pi/p\pi) \int_0^\infty (\lambda I + X)^{-1} X (\lambda I + X)^{-1} \lambda^{-p} d\lambda$.

Lemma 2 (see [39, Theorem 2.5]). *There exists a unique positive definite solution X of $X - A^*X^{-p}A = Q$ ($0 < p < 1$) and the iteration*

$$X_0 > 0, \quad X_n = Q + A^*X_{n-1}^{-p}A, \quad n = 1, 2, \dots \quad (2)$$

converges to X .

Lemma 3 (see [32, Lemma 2]). (i) *If $X \in \mathcal{H}^{n \times n}$, then $\|e^{-X}\| = e^{-\lambda_{\min}(X)}$.*

(ii) *If $X \in \mathcal{H}^{n \times n}$ and $r > 0$, then $X^{-r} = (1/\Gamma(r)) \int_0^\infty e^{-sX} s^{r-1} ds$.*

(iii) *If $A, B \in \mathcal{C}^{n \times n}$, then $e^{A+B} - e^A = \int_0^1 e^{(1-t)A} B e^{t(A+B)} dt$.*

3. A Sufficient Condition for the Existence of a Unique Solution of $X - A^*X^{-p}A = Q$ ($p \geq 1$)

In this section, we derive a new sufficient condition for the existence of a unique solution of $X - A^*X^{-p}A = Q$ ($p \geq 1$) beginning with the lemma.

Lemma 4 (see [38, Theorem 5, Remark 4]). *If*

$$\beta > (p\bar{\kappa})^{1/(p+1)}, \quad (3)$$

then (1) has a unique positive definite solution $X \in [\beta I, \alpha I]$, where α and β are, respectively, positive solutions of the following equations:

$$(x - \lambda_{\max}(Q)) \left(\lambda_{\min}(Q) + \frac{\underline{\kappa}}{x^p} \right)^p = \bar{\kappa}, \quad (4)$$

$$(x - \lambda_{\min}(Q)) \left(\lambda_{\max}(Q) + \frac{\bar{\kappa}}{x^p} \right)^p = \underline{\kappa}. \quad (5)$$

Furthermore,

$$\lambda_{\min}(Q) \leq \beta \leq \alpha. \quad (6)$$

Theorem 5. *If*

$$\left((p\bar{\kappa})^{1/(p+1)} - \lambda_{\min}(Q) \right) \left(\lambda_{\max}(Q) + \frac{\bar{\kappa}}{(p\bar{\kappa})^{p/(p+1)}} \right)^p < \underline{\kappa}, \quad (7)$$

$$\frac{(\lambda_{\min}(Q)p)^{p/(p+1)} \lambda_{\max}^{1/(p+1)}(Q)}{(\bar{\kappa})^{1/(p+1)}} + 1 - p > 0, \quad (8)$$

then (1) has a unique positive definite solution.

Proof. We first prove $\beta > (p\bar{\kappa})^{1/(p+1)}$, where β is the positive solution to (5). Let

$$f(x) = (x - \lambda_{\min}(Q)) \left(\lambda_{\max}(Q) + \frac{\bar{\kappa}}{x^p} \right)^p - \underline{\kappa}. \quad (9)$$

By computation, we obtain

$$f'(x) = \frac{\bar{\kappa}}{x^p} \left(\lambda_{\max}(Q) + \frac{\bar{\kappa}}{x^p} \right)^{p-1} \times \left(\frac{\lambda_{\max}(Q)}{\bar{\kappa}} x^p + p^2 \lambda_{\min}(Q) x^{-1} + 1 - p^2 \right). \quad (10)$$

Define

$$g(x) = \frac{\lambda_{\max}(Q)}{\bar{\kappa}} x^p + p^2 \lambda_{\min}(Q) x^{-1} + 1 - p^2. \quad (11)$$

Then $g(x)$ is decreasing on $[0, \lambda_{\min}(Q)p\bar{\kappa}/(\lambda_{\max}(Q))^{1/(p+1)}]$ and increasing on $[\lambda_{\min}(Q)p\bar{\kappa}/(\lambda_{\max}(Q))^{1/(p+1)}, +\infty)$, which implies that

$$g_{\min} = g \left(\left(\frac{\lambda_{\min}(Q)p\bar{\kappa}}{\lambda_{\max}(Q)} \right)^{1/(p+1)} \right) = (1+p) \left(\frac{(\lambda_{\min}(Q)p)^{p/(p+1)} \lambda_{\max}^{1/(p+1)}(Q)}{(\bar{\kappa})^{1/(p+1)}} + 1 - p \right). \quad (12)$$

According to the condition (8), it follows that $g_{\min} > 0$. Note that

$$f'(x) = \frac{\bar{\kappa}}{x^p} \left(\lambda_{\max}(Q) + \frac{\bar{\kappa}}{x^p} \right)^{p-1} g(x), \quad (13)$$

which implies that $f(x)$ is increasing on $(0, +\infty)$. Considering the condition (7), one sees that $f((p\bar{\kappa})^{1/(p+1)}) < 0$. Combining that and the definition of β in Lemma 4, we obtain $\beta > (p\bar{\kappa})^{1/(p+1)}$. By Lemma 4, (1) has a unique positive definite solution. \square

4. Perturbation Bound for

$$X - A^*X^{-p}A = Q \quad (p \geq 1)$$

Li and Zhang in [39] proved that there always exists a unique positive definite solution to the equation

$X - A^* X^{-p} A = Q$ ($0 < p < 1$). They also obtained a perturbation bound for the unique solution. But their approach becomes invalid for the case of $p \geq 1$. Since the equation $X - A^* X^{-p} A = Q$ ($p \geq 1$) does not always have a unique positive definite solution, there are two difficulties for a perturbation analysis of the equation $X - A^* X^{-p} A = Q$ ($p \geq 1$). One difficulty is how to find some reasonable restrictions on the coefficient matrices of perturbed equation ensuring that this equation has a unique positive definite solution. The other difficulty is how to find an expression of ΔX which is easy to handle.

Assume that the coefficient matrix A is perturbed to $\tilde{A} = \Delta A + A$. Let $\tilde{X} = \Delta X + X$ with $\Delta X \in \mathcal{H}^{n \times n}$ satisfying the perturbed equation

$$\tilde{X} - \tilde{A}^* \tilde{X}^{-p} \tilde{A} = Q, \quad p \geq 1. \tag{14}$$

In the following, we derive a perturbation estimate for the positive definite solution to the matrix equation $X - A^* X^{-p} A = Q$ ($p \geq 1$) beginning with the lemma.

Lemma 6 (see [38, Corollary 1. Remark 4]). *If*

$$p\|A\|^2 < \lambda_{\min}^{p+1}(Q), \tag{15}$$

then (1) has a unique positive definite solution X . Moreover, $X \geq \lambda_{\min}(Q)I$.

Theorem 7. *If*

$$\|A\| < \sqrt{\frac{\lambda_{\min}^{p+1}(Q)}{p}}, \quad \|\Delta A\| < \sqrt{\frac{\lambda_{\min}^{p+1}(Q)}{p}} - \|A\|, \tag{16}$$

then

$$X - A^* X^{-p} A = Q, \quad \tilde{X} - \tilde{A}^* \tilde{X}^{-p} \tilde{A} = Q \tag{17}$$

have unique positive definite solutions X and \tilde{X} , respectively. Furthermore,

$$\frac{\|\tilde{X} - X\|}{\|X\|} \leq \frac{(2\|A\| + \|\Delta A\|)}{\lambda_{\min}^{p+1}(Q) - p\|A\|^2} \|\Delta A\| \equiv \varrho. \tag{18}$$

Proof. By (16), it follows that $\|\tilde{A}\| \leq \|A\| + \|\Delta A\| \leq \sqrt{\lambda_{\min}^{p+1}(Q)}/p$. According to Lemma 6, the condition (16) ensures that (1) and (14) have unique positive definite solutions X and \tilde{X} , respectively. Furthermore, we obtain that

$$X \geq \lambda_{\min}(Q)I, \quad \tilde{X} \geq \lambda_{\min}(Q)I. \tag{19}$$

Subtracting (14) from (1) gives

$$\begin{aligned} \Delta X &= \tilde{A}^* \tilde{X}^{-p} \tilde{A} - A^* X^{-p} A \\ &= A^* (\tilde{X}^{-p} - X^{-p}) A + \Delta A^* \tilde{X}^{-p} A + \tilde{A}^* \tilde{X}^{-p} \Delta A. \end{aligned} \tag{20}$$

By Lemma 3 and inequalities in (19), we have

$$\begin{aligned} &\|\Delta X + A^* X^{-p} A - A^* \tilde{X}^{-p} A\| \\ &= \left\| \Delta X + A^* \frac{1}{\Gamma(p)} \int_0^\infty (e^{-sX} - e^{-s\tilde{X}}) s^{p-1} ds A \right\| \\ &= \left\| \Delta X + A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\tilde{X}} \Delta X e^{-tsX} dt s^p ds A \right\| \\ &\geq \|\Delta X\| - \frac{\|A\|^2 \|\Delta X\|}{\Gamma(p)} \int_0^\infty \int_0^1 \|e^{-(1-t)s\tilde{X}}\| \|e^{-tsX}\| dt s^p ds \\ &\geq \|\Delta X\| - \frac{\|A\|^2 \|\Delta X\|}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\lambda_{\min}(\tilde{X})} e^{-ts\lambda_{\min}(X)} dt s^p ds \\ &\geq \|\Delta X\| - \frac{\|A\|^2 \|\Delta X\|}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\lambda_{\min}(Q)} e^{-ts\lambda_{\min}(Q)} dt s^p ds \\ &= \|\Delta X\| - \frac{\|A\|^2 \|\Delta X\|}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-s\lambda_{\min}(Q)} dt s^p ds \\ &= \|\Delta X\| - \frac{\Gamma(p+1)}{\Gamma(p)} \cdot \frac{\|A\|^2 \|\Delta X\|}{\lambda_{\min}^{p+1}(Q)} \\ &= \frac{\lambda_{\min}^{p+1}(Q) - p\|A\|^2}{\lambda_{\min}^{p+1}(Q)} \|\Delta X\|. \end{aligned} \tag{21}$$

Noting (16), we have

$$\lambda_{\min}^{p+1}(Q) - p\|A\|^2 > 0. \tag{22}$$

Combining (20) and (21), one sees that

$$\begin{aligned} \frac{\lambda_{\min}^{p+1}(Q) - p\|A\|^2}{\lambda_{\min}^{p+1}(Q)} \|\Delta X\| &\leq \|\Delta A^* \tilde{X}^{-p} A + \tilde{A}^* \tilde{X}^{-p} \Delta A\| \\ &\leq (\|\Delta A\| + 2\|A\|) \|\Delta A\| \|\tilde{X}^{-p}\| \\ &\leq (\|\Delta A\| + 2\|A\|) \|\Delta A\| \lambda_{\min}^{-p}(Q), \end{aligned} \tag{23}$$

which implies that

$$\frac{\|\Delta X\|}{\|X\|} \leq \frac{(\|\Delta A\| + 2\|A\|)}{\lambda_{\min}^{p+1}(Q) - p\|A\|^2} \|\Delta A\|. \tag{24}$$

□

5. Condition Number for

$$X - A^* X^{-p} A = Q \quad (p \geq 1)$$

A condition number is a measurement of the sensitivity of the positive definite stabilizing solutions to small changes in the coefficient matrices. In this section, we apply the theory

of condition number developed by Rice [40] to derive explicit expressions of the condition number for the matrix equation $X - A^* X^{-p} A = Q$ ($p \geq 1$).

Here we consider the perturbed equation

$$\tilde{X} - \tilde{A}^* \tilde{X}^{-p} \tilde{A} = \tilde{Q}, \quad p \geq 1, \tag{25}$$

where \tilde{A} and \tilde{Q} are small perturbations of A and Q in (1), respectively.

Suppose that $p\|A\|^2 < \lambda_{\min}^{p+1}(Q)$ and $p\|\tilde{A}\|^2 < \lambda_{\min}^{p+1}(\tilde{Q})$. According to Lemma 6, (1) and (25) have unique positive definite solutions X and \tilde{X} , respectively. Let $\Delta X = \tilde{X} - X$, $\Delta Q = \tilde{Q} - Q$, and $\Delta A = \tilde{A} - A$.

Subtracting (25) from (1) gives

$$\begin{aligned} \Delta X &= \tilde{A}^* \tilde{X}^{-p} \tilde{A} - A^* X^{-p} A + \Delta Q \\ &= A^* (\tilde{X}^{-p} - X^{-p}) A + \Delta A^* \tilde{X}^{-p} A + \tilde{A}^* \tilde{X}^{-p} \Delta A + \Delta Q \\ &= -A^* \frac{1}{\Gamma(p)} \int_0^\infty (e^{-sX} - e^{-s\tilde{X}}) s^{p-1} ds A \\ &\quad + \Delta A^* \tilde{X}^{-p} A + \tilde{A}^* \tilde{X}^{-p} \Delta A + \Delta Q \\ &= -A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\tilde{X}} (\tilde{X} - X) e^{-tsX} dt s^p ds A \\ &\quad + \Delta A^* \tilde{X}^{-p} A + \tilde{A}^* \tilde{X}^{-p} \Delta A + \Delta Q \\ &= -A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 (e^{-(1-t)s\tilde{X}} - e^{-(1-t)sX}) \\ &\quad \times \Delta X e^{-tsX} dt s^p ds A + \Delta Q \\ &\quad - A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)sX} \Delta X e^{-tsX} dt s^p ds A \\ &\quad - (\tilde{A}^* X^{-p} \Delta A - \tilde{A}^* (X + \Delta X)^{-p} \Delta A) \\ &\quad + \tilde{A}^* X^{-p} \Delta A - (\Delta A^* X^{-p} A - \Delta A^* (X + \Delta X)^{-p} A) \\ &\quad + \Delta A^* X^{-p} A \\ &= A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 \int_0^1 e^{-(1-m)(1-t)sX} \Delta X e^{-m(1-t)s\tilde{X}} \\ &\quad \times \Delta X e^{-tsX} dm(1-t) dt s^{p+1} ds A \\ &\quad + \Delta Q \\ &\quad - A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)sX} \Delta X e^{-tsX} dt s^p ds A \\ &\quad + \Delta A^* X^{-p} \Delta A + A^* X^{-p} \Delta A + \Delta A^* X^{-p} A \\ &\quad - \tilde{A}^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s(X+\Delta X)} \Delta X e^{-tsX} dt s^p ds \Delta A \\ &\quad - \Delta A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s(X+\Delta X)} \Delta X e^{-tsX} dt s^p ds A. \end{aligned} \tag{26}$$

Therefore,

$$\begin{aligned} \Delta X + A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)sX} \Delta X e^{-tsX} dt s^p ds A \\ = E + h(\Delta X), \end{aligned} \tag{27}$$

where

$$\begin{aligned} B &= X^{-p} A, \\ E &= \Delta Q + (B^* \Delta A + \Delta A^* B) + \Delta A^* X^{-p} \Delta A, \\ h(\Delta X) &= A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 \int_0^1 e^{-(1-m)(1-t)sX} \\ &\quad \times \Delta X e^{-m(1-t)s\tilde{X}} \Delta X e^{-tsX} \\ &\quad \times dm(1-t) dt s^{p+1} ds A \\ &\quad - \tilde{A}^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s(X+\Delta X)} \Delta X e^{-tsX} dt s^p ds \Delta A \\ &\quad - \Delta A^* \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s(X+\Delta X)} \Delta X e^{-tsX} dt s^p ds A. \end{aligned} \tag{28}$$

Lemma 8. *If*

$$p\|A\|^2 < \lambda_{\min}^{p+1}(Q), \tag{29}$$

then the linear operator $\mathbf{V} : \mathcal{H}^{n \times n} \rightarrow \mathcal{H}^{n \times n}$ defined by

$$\begin{aligned} \mathbf{V}W &= W + \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 A^* e^{-(1-t)sX} W e^{-tsX} A dt s^p ds, \\ &W \in \mathcal{H}^{n \times n} \end{aligned} \tag{30}$$

is invertible.

Proof. Define the operator $\mathbf{R} : \mathcal{H}^{n \times n} \rightarrow \mathcal{H}^{n \times n}$ by

$$\begin{aligned} \mathbf{R}Z &= \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 A^* e^{-(1-t)sX} Z e^{-tsX} A dt s^p ds, \\ &Z \in \mathcal{H}^{n \times n}, \end{aligned} \tag{31}$$

it follows that

$$\mathbf{V}W = W + \mathbf{R}W. \tag{32}$$

Then, \mathbf{V} is invertible if and only if $I + \mathbf{R}$ is invertible.

According to Lemma 3 and the condition (29), we have

$$\begin{aligned}
 & \| \mathbf{R}W \| \\
 & \leq \| A \|^2 \| W \| \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 \| e^{-(1-t)sX} \| \| e^{-tsX} \| dt s^p ds \\
 & = \| A \|^2 \| W \| \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\lambda_{\min}(X)} e^{-ts\lambda_{\min}(X)} dt s^p ds \\
 & \leq \| A \|^2 \| W \| \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 e^{-(1-t)s\lambda_{\min}(Q)} e^{-ts\lambda_{\min}(Q)} dt s^p ds \\
 & = \| A \|^2 \| W \| \frac{1}{\Gamma(p)} \int_0^\infty e^{-s\lambda_{\min}(Q)} s^p ds \\
 & = \frac{\rho \| A \|^2}{\lambda_{\min}^{p+1}(Q)} \| W \| < \| W \|,
 \end{aligned} \tag{33}$$

which implies that $\| \mathbf{R} \| < 1$ and $I + \mathbf{R}$ is invertible. Therefore, the operator \mathbf{V} is invertible. \square

Thus, we can rewrite (27) as

$$\begin{aligned}
 \Delta X & = \mathbf{V}^{-1} \Delta Q + \mathbf{V}^{-1} (B^* \Delta A + \Delta A^* B) \\
 & \quad + \mathbf{V}^{-1} (\Delta A^* X^{-p} \Delta A) + \mathbf{V}^{-1} (h(\Delta X)),
 \end{aligned} \tag{34}$$

$$\begin{aligned}
 \Delta X & = \mathbf{V}^{-1} \Delta Q + \mathbf{V}^{-1} (B^* \Delta A + \Delta A^* B) \\
 & \quad + O(\|(\Delta A, \Delta Q)\|_F^2), \quad (\Delta A, \Delta Q) \rightarrow 0.
 \end{aligned} \tag{35}$$

By the theory of condition number developed by Rice [40], we define the condition number of the Hermitian positive definite solution X to the matrix equation $X - A^* X^{-p} A = Q$ ($p \geq 1$) by

$$c(X) = \lim_{\delta \rightarrow 0} \sup_{\|(\Delta A/\eta, \Delta Q/\rho)\|_F \leq \delta} \frac{\| \Delta X \|_F}{\xi \delta}, \tag{36}$$

where ξ , η , and ρ are positive parameters. Taking $\xi = \eta = \rho = 1$ in (36) gives the absolute condition number $c_{\text{abs}}(X)$, and taking $\xi = \| X \|_F$, $\eta = \| A \|_F$, and $\rho = \| Q \|_F$ in (36) gives the relative condition number $c_{\text{rel}}(X)$.

Substituting (35) into (36), we get

$$\begin{aligned}
 c(X) & = \frac{1}{\xi} \max_{\substack{(\Delta A/\eta, \Delta Q/\rho) \neq 0 \\ \Delta A \in \mathcal{C}^{n \times n}, \Delta Q \in \mathcal{H}^{n \times n}}} \frac{\| \mathbf{V}^{-1} (\Delta Q + B^* \Delta A + \Delta A^* B) \|_F}{\|(\Delta A/\eta, \Delta Q/\rho)\|_F} \\
 & = \frac{1}{\xi} \max_{\substack{(E, H) \neq 0 \\ E \in \mathcal{C}^{n \times n}, H \in \mathcal{H}^{n \times n}}} \frac{\| \mathbf{V}^{-1} (\rho H + \eta (B^* E + E^* B)) \|_F}{\| (E, H) \|_F}.
 \end{aligned} \tag{37}$$

Let V be the matrix representation of the linear operator \mathbf{V} . It follows from Lemma 4.3.2. in [41] that

$$\text{vec}(\mathbf{V}W) = V \cdot \text{vec} W. \tag{38}$$

By Lemma 4.3.1. in [41], we have

$$\begin{aligned}
 & \text{vec}(\mathbf{V}W) \\
 & = \left(I \otimes I + \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 (e^{-tsX} A)^T \otimes (A^* e^{-(1-t)sX}) dt s^p ds \right) \\
 & \quad \cdot \text{vec} W.
 \end{aligned} \tag{39}$$

Then,

$$\begin{aligned}
 V & = I \otimes I + \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 (e^{-tsX} A)^T \\
 & \quad \otimes (A^* e^{-(1-t)sX}) dt s^p ds.
 \end{aligned} \tag{40}$$

Let

$$\begin{aligned}
 V^{-1} & = S + i\Sigma, \\
 V^{-1} (I \otimes B^*) & = V^{-1} (I \otimes (X^{-p} A)^*) = U_1 + i\Omega_1, \\
 V^{-1} (B^T \otimes I) \Pi & = V^{-1} ((X^{-p} A)^T \otimes I) \Pi = U_2 + i\Omega_2,
 \end{aligned} \tag{41}$$

$$\begin{aligned}
 S_c & = \begin{bmatrix} S & -\Sigma \\ \Sigma & S \end{bmatrix}, \quad U_c = \begin{bmatrix} U_1 + U_2 & \Omega_2 - \Omega_1 \\ \Omega_1 + \Omega_2 & U_1 - U_2 \end{bmatrix}, \\
 \text{vec} H & = x + iy, \quad \text{vec} E = a + ib,
 \end{aligned} \tag{42}$$

$$g = (x^T, y^T, a^T, b^T)^T,$$

where $x, y, a, b \in \mathcal{R}^M$, $S, \Sigma, U_1, U_2, \Omega_1, \Omega_2 \in \mathcal{R}^{n^2 \times n^2}$, $M = (E, H)$, $i = \sqrt{-1}$, Π is the vec-permutation matrix, that is,

$$\text{vec} E^T = \Pi \text{vec} E. \tag{43}$$

Furthermore, we obtain that

$$\begin{aligned}
 c(X) & = \frac{1}{\xi} \max_{M \neq 0} \frac{\| \mathbf{V}^{-1} (\rho H + \eta (B^* E + E^* B)) \|_F}{\| (E, H) \|_F} \\
 & = \frac{1}{\xi} \max_{M \neq 0} \left\| \rho V^{-1} \text{vec} H + \eta V^{-1} \right. \\
 & \quad \times \left((I \otimes B^*) \text{vec} E + (B^T \otimes I) \text{vec} E^* \right) \| \\
 & \quad \times (\| \text{vec} (E, H) \|)^{-1} \\
 & = \frac{1}{\xi} \max_{M \neq 0} \left\| \rho (S + i\Sigma) (x + iy) \right. \\
 & \quad \left. + \eta [(U_1 + i\Omega_1) (a + ib) \right. \\
 & \quad \left. + (U_2 + i\Omega_2) (a - ib)] \right\| \\
 & \quad \times (\| \text{vec} (E, H) \|)^{-1} \\
 & = \frac{1}{\xi} \max_{g \neq 0} \frac{\| (\rho S_c, \eta U_c) g \|}{\| g \|} \\
 & = \frac{1}{\xi} \| (\rho S_c, \eta U_c) \|, \quad E \in \mathcal{C}^{n \times n}, H \in \mathcal{H}^{n \times n}.
 \end{aligned} \tag{44}$$

Then, we have the following theorem.

Theorem 9. *If $p\|A\|^2 < \lambda_{\min}^{p+1}(Q)$, then the condition number $c(X)$ defined by (36) has the explicit expression*

$$c(X) = \frac{1}{\xi} \|(\rho S_c, \eta U_c)\|, \tag{46}$$

where the matrices S_c and U_c are defined by (40)-(41).

Remark 10. From (46) we have the relative condition number

$$c_{\text{rel}}(X) = \frac{\|(\|Q\|_F S_c, \|A\|_F U_c)\|}{\|X\|_F}. \tag{47}$$

5.1. The Real Case. In this subsection, we consider the real case, that is, where all the coefficients matrices A, Q of the matrix equation $X - A^* X^{-p} A = Q$ ($p \geq 1$) are real. In such a case the corresponding solution X is also real. Similar arguments as in Theorem 9 give the following theorem.

Theorem 11. *Let A, Q be real, $c(X)$ the condition number defined by (36). If $p\|A\|^2 < \lambda_{\min}^{p+1}(Q)$, then $c(X)$ has the explicit expression*

$$c(X) = \frac{1}{\xi} \|(\rho S_r, \eta U_r)\|, \tag{48}$$

where

$$S_r = \left(I \otimes I + \frac{1}{\Gamma(p)} \int_0^\infty \int_0^1 (e^{-tsX} A)^T \otimes (A^T e^{-(1-t)sX}) dt s^p ds \right)^{-1}, \tag{49}$$

$$U_r = S_r [I \otimes (A^T X^{-p}) + ((A^T X^{-p}) \otimes I) \Pi].$$

Proof. Let

$$\begin{aligned} S_r &= V^{-1}, \\ U_r &= V^{-1} \left((I \otimes B^T) + (B^T \otimes I) \Pi \right) \\ &= S_r \left((I \otimes (X^{-p} A)^T) + ((X^{-p} A)^T \otimes I) \Pi \right) \\ \text{vec } H &= x, \quad \text{vec } E = a, \quad g = (x^T, a^T)^T, \end{aligned} \tag{50}$$

where $x, a \in \mathcal{R}^n$, $M = (E, H)$, $\mathbf{i} = \sqrt{-1}$, Π is the vec-permutation matrix, that is,

$$\text{vec } E^T = \Pi \text{vec } E. \tag{51}$$

It follows from (44) that

$$\begin{aligned} c(X) &= \frac{1}{\xi} \max_{M \neq 0} \frac{\|V^{-1} (\rho H + \eta (B^T E + E^T B))\|_F}{\|(E, H)\|_F} \\ &= \frac{1}{\xi} \max_{M \neq 0} \left\| \rho V^{-1} \text{vec } H \right. \\ &\quad \left. + \eta V^{-1} \left((I \otimes B^T) \text{vec } E + (B^T \otimes I) \text{vec } E^T \right) \right\| \\ &\quad \times (\|\text{vec } (E, H)\|)^{-1} \\ &= \frac{1}{\xi} \max_{g \neq 0} \frac{\|(\rho S_r, \eta U_r) g\|}{\|g\|} \\ &= \frac{1}{\xi} \|(\rho S_r, \eta U_r)\|. \end{aligned} \tag{52}$$

□

Remark 12. In the real case the relative condition number is given by

$$c_{\text{rel}}(X) = \frac{\|(\|Q\|_F S_r, \|A\|_F U_r)\|}{\|X\|_F}. \tag{53}$$

6. New Perturbation Bound for

$$X - A^* X^{-p} A = Q \quad (0 < p < 1)$$

Here, we consider the perturbed equation

$$\tilde{X} - \tilde{A}^* \tilde{X}^{-p} \tilde{A} = \tilde{Q}, \quad 0 < p < 1, \tag{54}$$

where \tilde{A} and \tilde{Q} are small perturbations of A and Q in (1), respectively. We assume that X and \tilde{X} are the solutions of (1) and (54), respectively. Let $\Delta X = \tilde{X} - X$, $\Delta Q = \tilde{Q} - Q$, and $\Delta A = \tilde{A} - A$.

In this section, we develop a new perturbation bound for the solution of (1) which is sharper than that in [39, Theorem 3.1].

Subtracting (1) from (54), using Lemma 1, we have

$$\begin{aligned} \Delta X + \frac{\sin p\pi}{\pi} \int_0^\infty [(\lambda I + X)^{-1} A]^* \\ \times \Delta X [(\lambda I + X)^{-1} A] \lambda^{-p} d\lambda = E + h(\Delta X), \end{aligned} \tag{55}$$

where

$$B = X^{-p} A,$$

$$E = \Delta Q + (B^* \Delta A + \Delta A^* B) + \Delta A^* X^{-p} \Delta A,$$

$$\begin{aligned} h(\Delta X) &= \frac{\sin p\pi}{\pi} A^* \int_0^\infty (\lambda I + X)^{-1} \\ &\quad \times \Delta X (\lambda I + X + \Delta X)^{-1} \\ &\quad \times \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda \end{aligned}$$

$$\begin{aligned}
 & - \frac{\sin p\pi}{\pi} \tilde{A}^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X + \Delta X)^{-1} \lambda^{-p} d\lambda \Delta A \\
 & - \frac{\sin p\pi}{\pi} \Delta A^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X + \Delta X)^{-1} \lambda^{-p} d\lambda A.
 \end{aligned} \tag{56}$$

By Lemma 5.1 in [39], the linear operator $\mathbf{L} : \mathcal{H}^{n \times n} \rightarrow \mathcal{H}^{n \times n}$ defined by

$$\begin{aligned}
 \mathbf{L}W &= W + \frac{\sin p\pi}{\pi} \int_0^\infty [(\lambda I + X)^{-1} A]^* \\
 & \quad \times W [(\lambda I + X)^{-1} A] \lambda^{-p} d\lambda, \tag{57} \\
 & \quad W \in \mathcal{H}^{n \times n}
 \end{aligned}$$

is invertible.

We also define operator $\mathbf{P} : \mathcal{C}^{n \times n} \rightarrow \mathcal{H}^{n \times n}$ by

$$\begin{aligned}
 \mathbf{P}Z &= \mathbf{L}^{-1} (B^* Z + Z^* B), \\
 Z &\in \mathcal{C}^{n \times n}, \quad i = 1, 2, \dots, m.
 \end{aligned} \tag{58}$$

Thus, we can rewrite (55) as

$$\Delta X = \mathbf{L}^{-1} \Delta Q + \mathbf{P} \Delta A + \mathbf{L}^{-1} (\Delta A^* X^{-p} \Delta A) + \mathbf{L}^{-1} (h(\Delta X)). \tag{59}$$

Define

$$\|\mathbf{L}^{-1}\| = \max_{\substack{W \in \mathcal{H}^{n \times n} \\ \|W\|=1}} \|\mathbf{L}^{-1}W\|, \quad \|\mathbf{P}\| = \max_{\substack{Z \in \mathcal{C}^{n \times n} \\ \|Z\|=1}} \|\mathbf{P}Z\|. \tag{60}$$

Now we denote

$$\begin{aligned}
 l &= \|\mathbf{L}^{-1}\|^{-1}, \quad \zeta = \|X^{-1}\|, \quad \xi = \|X^{-p}\|, \\
 n &= \|\mathbf{P}\|, \quad \eta = p\xi \|A\|^2, \\
 \epsilon &= \frac{1}{l} \|\Delta Q\| + n \|\Delta A\| + \frac{\xi}{l} \|\Delta A\|^2, \\
 \sigma &= \frac{p}{l} \zeta \xi (2 \|A\| + \|\Delta A\|) \|\Delta A\|.
 \end{aligned} \tag{61}$$

Theorem 13. *If*

$$\sigma < 1, \quad \epsilon < \frac{l(1-\sigma)^2}{\zeta \left(l + l\sigma + 2\eta + 2\sqrt{(l\sigma + \eta)(\eta + l)} \right)}, \tag{62}$$

then

$$\begin{aligned}
 & \|\bar{X} - X\| \\
 & \leq \frac{2l\epsilon}{l(1 + \zeta\epsilon - \sigma) + \sqrt{l^2(1 + \zeta\epsilon - \sigma)^2 - 4l\zeta\epsilon(l + \eta)}} \equiv \mu_*.
 \end{aligned} \tag{63}$$

Proof. Let

$$\begin{aligned}
 f(\Delta X) &= \mathbf{L}^{-1} \Delta Q + \mathbf{P} \Delta A + \mathbf{L}^{-1} (\Delta A^* X^{-p} \Delta A) \\
 & \quad + \mathbf{L}^{-1} (h(\Delta X)).
 \end{aligned} \tag{64}$$

Obviously, $f : \mathcal{H}^{n \times n} \rightarrow \mathcal{H}^{n \times n}$ is continuous. The condition (62) ensures that the quadratic equation

$$\zeta(l + \eta)x^2 - l(1 + \zeta\epsilon - \sigma)x + l\epsilon = 0 \tag{65}$$

in x has two positive real roots. The smaller one is

$$\mu_* = \frac{2l\epsilon}{l(1 + \zeta\epsilon - \sigma) + \sqrt{l^2(1 + \zeta\epsilon - \sigma)^2 - 4l\zeta\epsilon(l + \eta)}}. \tag{66}$$

Define $\Omega = \{\Delta X \in \mathcal{H}^{n \times n} : \|\Delta X\| \leq \mu_*\}$. Then for any $\Delta X \in \Omega$, by (62), we have

$$\begin{aligned}
 & \|X^{-1} \Delta X\| \\
 & \leq \|X^{-1}\| \|\Delta X\| \\
 & \leq \zeta \mu_* \leq \zeta \cdot \frac{2\epsilon}{1 + \zeta\epsilon - \sigma} = 1 + \frac{\zeta\epsilon + \sigma - 1}{1 + \zeta\epsilon - \sigma} \\
 & \leq 1 + \left(\zeta \cdot l(1 - \sigma)^2 / \zeta \left(l + l\sigma + 2\eta + 2\sqrt{(l\sigma + \eta)(\eta + l)} \right) \right. \\
 & \quad \left. + \sigma - 1 \right) \times (1 + \zeta\epsilon - \sigma)^{-1} \\
 & \leq 1 + \frac{l(1 - \sigma)^2 + (\sigma - 1)(l + l\sigma + 2\eta)}{(l + l\sigma + 2\eta)(1 + \zeta\epsilon - \sigma)} \\
 & = 1 + \frac{-2(1 - \sigma)(l\sigma + \eta)}{(l\sigma + l + 2\eta)(1 + \zeta\epsilon - \sigma)} < 1.
 \end{aligned} \tag{67}$$

It follows that $I - X^{-1} \Delta X$ is nonsingular and

$$\|I - X^{-1} \Delta X\| \leq \frac{1}{1 - \|X^{-1} \Delta X\|} \leq \frac{1}{1 - \zeta \|\Delta X\|}. \tag{68}$$

Therefore

$$\begin{aligned}
 & \|f(\Delta X)\| \\
 & \leq \frac{1}{l} \|\Delta Q\| + n \|\Delta A\| \\
 & \quad + \frac{\xi}{l} \|\Delta A_i\|^2 + \frac{p}{l} \zeta \xi \|A\|^2 \frac{\|\Delta X\|^2}{1 - \zeta \|\Delta X\|} \\
 & \quad + \frac{p}{l} \zeta \xi (2 \|A\| + \|\Delta A\|) \|\Delta A\| \cdot \frac{\|\Delta X\|}{1 - \zeta \|\Delta X\|} \\
 & \leq \epsilon + \frac{\sigma \|\Delta X\|}{1 - \zeta \|\Delta X\|} + \frac{\eta \zeta \|\Delta X\|^2}{l(1 - \zeta \|\Delta X\|)} \\
 & \leq \epsilon + \frac{\sigma \mu_*}{1 - \zeta \mu_*} + \frac{\eta \zeta \mu_*^2}{l(1 - \zeta \mu_*)} = \mu_*,
 \end{aligned} \tag{69}$$

for $\Delta X \in \Omega$, in which the last equality is due to the fact that μ_* is a solution to (65). That is $f(\Omega) \subseteq \Omega$. According to Schauder fixed point theorem, there exists $\Delta X_* \in \Omega$ such that $f(\Delta X_*) = \Delta X_*$. It follows that $X + \Delta X_*$ is a Hermitian solution of (54). By Lemma 2, we know that the solution of (54) is unique. Then $\Delta X_* = \tilde{X} - X$ and $\|\tilde{X} - X\| \leq \mu_*$. \square

7. New Backward Error for

$$X - A^* X^{-p} A = Q \quad (0 < p < 1)$$

In this section, we evaluate a new backward error estimate for an approximate solution to the unique solution, which is sharper than that in [39, Theorem 4.1], .

Theorem 14. *Let $\tilde{X} > 0$ be an approximation to the solution X of (1). If $\|\tilde{X}^{-p/2} A\|^2 \|\tilde{X}^{-1}\| < 1$ and the residual $R(\tilde{X}) \equiv Q + A^* \tilde{X}^{-p} A - \tilde{X}$ satisfies*

$$\begin{aligned} \|R(\tilde{X})\| &\leq \frac{\theta_1}{2} \min \left\{ 1, \frac{\theta_1}{2\lambda_{\min}(\tilde{X})} \right\}, \quad \text{where} \\ \theta_1 &= \left(1 - \|\tilde{X}^{-p/2} A\|^2 \|\tilde{X}^{-1}\| \right) \lambda_{\min}(\tilde{X}) \\ &\quad + \|R(\tilde{X})\| > 0, \end{aligned} \tag{70}$$

then

$$\begin{aligned} \|\tilde{X} - X\| &\leq \theta \|R(\tilde{X})\|, \quad \text{where} \\ \theta &= \frac{2\lambda_{\min}(\tilde{X})}{\theta_1 + \sqrt{\theta_1^2 - 4\lambda_{\min}(\tilde{X}) \|R(\tilde{X})\|}}. \end{aligned} \tag{71}$$

To prove the above theorem, we first verify the following lemma.

Lemma 15. *For every positive definite matrix $X \in \mathcal{H}^{n \times n}$, $0 < p < 1$, if $X + \Delta X \geq (1/\nu)I > 0$, then*

$$\begin{aligned} \|A^* ((X + \Delta X)^{-p} - X^{-p}) A\| \\ \leq p (\|\Delta X\| + \nu \|\Delta X\|^2) \|X^{-p/2} A\|^2 \|X^{-1}\|. \end{aligned} \tag{72}$$

Proof. It follows from Lemma 1 that

$$\begin{aligned} \|A^* ((X + \Delta X)^{-p} - X^{-p}) A\| \\ = \|A^* \left(\frac{\sin p \pi}{\pi} \int_0^\infty ((\lambda I + X + \Delta X)^{-1} (\lambda I + X)^{-1}) \right. \\ \left. \times \lambda^{-p} d\lambda \right) A\| \\ \leq \frac{\sin p \pi}{\pi} \left(\|A^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \right) \\ + \frac{\sin p \pi}{\pi} \left(\|A^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X + \Delta X)^{-1} \right. \\ \left. \times \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \right). \end{aligned} \tag{73}$$

Note that $\Delta X \leq \|\Delta X\|I$, $X + \Delta X \geq (1/\nu)I > 0$, and $X^{-p} = (\sin p\pi/p\pi) \int_0^\infty (\lambda I + X)^{-1} X (\lambda I + X)^{-1} \lambda^{-p} d\lambda$, we have

$$\begin{aligned} &\|A^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \\ &= \|A^* \int_0^\infty (\lambda I + X)^{-1} X X^{-1} \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \\ &\leq \|A^* \int_0^\infty (\lambda I + X)^{-1} X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \|X^{-1}\| \|\Delta X\| \\ &= \frac{p\pi}{\sin(p\pi)} \cdot \|A^* X^{-p} A\| \|X^{-1}\| \|\Delta X\|, \end{aligned} \tag{74}$$

$$\begin{aligned} &\|A^* \int_0^\infty (\lambda I + X)^{-1} \Delta X (\lambda I + X + \Delta X)^{-1} \\ &\quad \times \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \\ &= \|A^* \int_0^\infty (\lambda I + X)^{-1} X X^{-1} \\ &\quad \times \Delta X (\lambda I + X + \Delta X)^{-1} \Delta X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A\| \\ &\leq \|A^* \left(\int_0^\infty (\lambda I + X)^{-1} X (\lambda I + X)^{-1} \lambda^{-p} d\lambda A \right)\| \\ &\quad \cdot \nu \|\Delta X\|^2 \|X^{-1}\| \\ &= \frac{p\pi}{\sin(p\pi)} \cdot \|A^* X^{-p} A\| \cdot \nu \|\Delta X\|^2 \|X^{-1}\|. \end{aligned} \tag{75}$$

A combination of (73)–(75) gives

$$\begin{aligned} &\|A^* ((X + \Delta X)^{-p} - X^{-p}) A\| \\ &\leq p \|A^* X^{-p} A\| \|X^{-1}\| \|\Delta X\| \\ &\quad + p \|A^* X^{-p} A\| \nu \|\Delta X\|^2 \|X^{-1}\| \\ &= p (\|\Delta X\| + \nu \|\Delta X\|^2) \|X^{-p/2} A\|^2 \|X^{-1}\|. \end{aligned} \tag{76}$$

Here, we have used the result $\|A^* X^{-p} A\| = \|(X^{-p/2} A)^* (X^{-p/2} A)\| = \|X^{-p/2} A\|^2$ to derive the last equality (refer to [42, Problem 11. Page 312]). \square

Proof of Theorem 14. Let

$$\Psi = \{ \Delta X \in \mathcal{H}^{n \times n} : \|\Delta X\| \leq \theta \|R(\tilde{X})\| \}. \tag{77}$$

TABLE 1: Assumptions check for Example 1 with different values of j .

j	4	5	6	7
ass ₁	0.0455	0.0456	0.0456	0.0456
ass ₂	0.9999	1.0000	1.0000	1.0000
ass ₃	0.3957	0.3959	0.3959	0.3959

Obviously, Ψ is a nonempty bounded convex closed set. Let

$$g(\Delta X) = A^* \left((\bar{X} + \Delta X)^{-p} - \bar{X}^{-p} \right) A + R(\bar{X}). \quad (78)$$

Evidently $g : \Psi \mapsto \mathcal{H}^{n \times n}$ is continuous. The condition (70) ensures that the equation

$$y^2 - \left[(1 - \|\bar{X}^{-p/2} A\|^2 \|\bar{X}^{-1}\|) \lambda_{\min}(\bar{X}) + \|R(\bar{X})\| \right] y + \lambda_{\min}(\bar{X}) \|R(\bar{X})\| = 0 \quad (79)$$

in y has two positive real roots. The smaller one is $y_* = 2\lambda_{\min}(\bar{X})\|R(\bar{X})\|/\theta_1 + \sqrt{\theta_1^2 - 4\lambda_{\min}(\bar{X})\|R(\bar{X})\|}$, where $\theta_1 = (1 - \|\bar{X}^{-p/2} A\|^2 \|\bar{X}^{-1}\|)\lambda_{\min}(\bar{X}) + \|R(\bar{X})\|$.

We will prove that $g(\Psi) \subseteq \Psi$. For every $\Delta X \in \Psi$, we have

$$\Delta X \geq -\theta \|R(\bar{X})\| I. \quad (80)$$

Hence,

$$\bar{X} + \Delta X \geq \bar{X} - \theta \|R(\bar{X})\| I \geq (\lambda_{\min}(\bar{X}) - \theta \|R(\bar{X})\|) I. \quad (81)$$

Using (70) and (71), one sees that

$$\begin{aligned} \theta \|R(\bar{X})\| &= \frac{2\lambda_{\min}(\bar{X}) \|R(\bar{X})\|}{\theta_1 + \sqrt{\theta_1^2 - 4\lambda_{\min}(\bar{X}) \|R(\bar{X})\|}} \\ &< \frac{2\lambda_{\min}(\bar{X}) \|R(\bar{X})\|}{\theta_1} < \lambda_{\min}(\bar{X}). \end{aligned} \quad (82)$$

Therefore, $(\lambda_{\min}(\bar{X}) - \theta \|R(\bar{X})\|) I > 0$.

According to (72), we obtain

$$\begin{aligned} &\|g(\Delta X)\| \\ &\leq p \left(\|\Delta X\| + \frac{\|\Delta X\|^2}{\lambda_{\min}(\bar{X}) - \theta \|R(\bar{X})\|} \right) \\ &\quad \times \|\bar{X}^{-p/2} A\|^2 \|\bar{X}^{-1}\| + \|R(\bar{X})\| \\ &\leq \left(\theta \|R(\bar{X})\| + \frac{(\theta \|R(\bar{X})\|)^2}{\lambda_{\min}(\bar{X}) - \theta \|R(\bar{X})\|} \right) \\ &\quad \times \left(p \|\bar{X}^{-p/2} A\|^2 \|\bar{X}^{-1}\| + \|R(\bar{X})\| \right) \\ &= \theta \|R(\bar{X})\|, \end{aligned} \quad (83)$$

for $\Delta X \in \Psi$, in which the last equality is due to the fact that $\theta \|R(\bar{X})\|$ is a solution to (79). That is $g(\Psi) \subseteq \Psi$. By Brouwer fixed point theorem, there exists a $\Delta X \in \Psi$ such that $g(\Delta X) = \Delta X$. Hence $\bar{X} + \Delta X$ is a solution of (1). Moreover, by Lemma 2, we know that the solution X of (1) is unique. Then

$$\|\bar{X} - X\| = \|\Delta X\| \leq \theta \|R(\bar{X})\|. \quad (84)$$

□

8. Numerical Examples

To illustrate the theoretical results of the previous sections, in this section four simple examples are given, which were carried out using MATLAB 7.1. For the stopping criterion we take $\varepsilon_{k+1}(X) = \|X_k - A^* X_k^{-p} A - Q\| < 1.0e - 10$.

Example 1. We consider the matrix equation

$$X - A^* X^{-1/3} A = I, \quad (85)$$

where

$$A = \frac{A_0}{\|A_0\|}, \quad A_0 = \begin{pmatrix} 2 & 0.95 \\ 0 & 1 \end{pmatrix}. \quad (86)$$

Suppose that the coefficient matrix A is perturbed to $\tilde{A} = A + \Delta A$, where

$$\Delta A = \frac{10^{-j}}{\|C^T + C\|} (C^T + C) \quad (87)$$

and C is a random matrix generated by MATLAB function `randn`.

We compare our own result $\mu_*/\|X\| \triangleq \text{err}_2$ in Theorem 13 with the perturbation bound $\xi_* \triangleq \text{err}_1$ proposed in [39, Theorem 3.1].

The assumption in [39, Theorem 3.1] is

$$\text{ass}_1 = \sqrt{\|A\|^2 + \zeta} - \|A\| - \|\Delta A\| > 0. \quad (88)$$

The assumptions in Theorem 13 are

$$\begin{aligned} \text{ass}_2 &= 1 - \sigma > 0, \\ \text{ass}_3 &= \frac{l(1 - \sigma)^2}{\zeta \left(l + \sigma l + 2\eta + 2\sqrt{(l\sigma + \eta)(\eta + l)} \right)} - \epsilon > 0. \end{aligned} \quad (89)$$

By computation, we list them in Table 1.

The results listed in Table 1 show that the assumptions in Theorem 3.1 [39] and Theorem 13 are satisfied.

By Theorem 3.1 in [39] and Theorem 13, we can compute the relative perturbation bounds $\text{err}_1, \text{err}_2$, respectively. These results averaged as the geometric mean of 10 randomly perturbed runs. Some results are listed in Table 2.

The results listed in Table 2 show that the perturbation bound err_2 given by Theorem 13 is fairly sharp, while the bound err_1 given by Theorem 3.1 in [39] is conservative.

TABLE 2: Perturbation bounds for Example 1 with different values of j .

j	4	5	6	7
$\frac{\ \bar{X} - X\ }{\ X\ }$	6.8119×10^{-5}	4.2332×10^{-6}	4.3287×10^{-7}	5.5767×10^{-8}
err ₁	2.6003×10^{-4}	2.1375×10^{-5}	1.9229×10^{-6}	2.7300×10^{-7}
err ₂	8.8966×10^{-5}	6.5825×10^{-6}	7.2867×10^{-7}	9.3455×10^{-8}

TABLE 3: Results for Example 2 with different values of k .

k	4	5	6	7
$\ \bar{X}_k - X\ $	6.2131×10^{-6}	1.5830×10^{-7}	8.2486×10^{-9}	6.0132×10^{-10}
$\nu_* \ R(\bar{X}_k)\ $	2.5930×10^{-5}	6.6257×10^{-7}	3.5697×10^{-8}	2.4646×10^{-9}
κ_1	4.1734	4.1856	4.3277	4.0986
$\theta \ R(\bar{X}_k)\ $	7.0053×10^{-6}	1.7900×10^{-7}	9.6440×10^{-9}	6.6583×10^{-10}
κ_2	1.1275	1.1308	1.1692	1.1073

Example 2. Consider the equation

$$X - A^* X^{-3/4} A = Q, \tag{90}$$

for

$$A = \begin{pmatrix} 0.2 & -0.2 \\ 0.1 & 0.1 \end{pmatrix}, \quad Q = \begin{pmatrix} 0.8939 & 0.2987 \\ 0.1991 & 0.6614 \end{pmatrix}. \tag{91}$$

Choose $\bar{X}_0 = 3Q$. Let the approximate solution \bar{X}_k be given with the iterative method (2), where k is the iteration number. Assume that the solution X of (1) is unknown.

We compare our own result with the backward error proposed in Theorem 4.1 [39].

The residual $R(\bar{X}_k) \equiv Q + A^* \bar{X}_k^{-P} A - \bar{X}_k$ satisfies the conditions in Theorem 4.1 [39] and in Theorem 14.

By Theorem 4.1 in [39], we can compute the backward error bound

$$\begin{aligned} \|\bar{X}_k - X\| &\leq \nu_* \|R(\bar{X}_k)\|, \quad \text{where} \\ \nu_* &= \frac{2 \|\bar{X}_k\| \|\bar{X}_k^{-1}\|}{1 - (3/4) \|\bar{X}_k^{-3/8} A \bar{X}_k^{-1/2}\|^2}. \end{aligned} \tag{92}$$

By Theorem 14, we can compute the new backward error bound

$$\begin{aligned} \|\bar{X}_k - X\| &\leq \theta \|R(\bar{X}_k)\|, \quad \text{where} \\ \theta &= \frac{2\lambda_{\min}(\bar{X}_k)}{\theta_1 + \sqrt{\theta_1^2 - 4\lambda_{\min}(\bar{X}_k) \|R(\bar{X}_k)\|}}, \\ \theta_1 &= \left(1 - \|\bar{X}_k^{-3/8} A\|^2 \|\bar{X}_k^{-1}\|\right) \lambda_{\min}(\bar{X}_k) + \|R(\bar{X}_k)\|. \end{aligned} \tag{93}$$

Let

$$\kappa_1 = \frac{\nu_* \|R(\bar{X}_k)\|}{\|\bar{X}_k - X\|}, \quad \kappa_2 = \frac{\theta \|R(\bar{X}_k)\|}{\|\bar{X}_k - X\|}. \tag{94}$$

Some results are shown in Table 3.

From the results listed in Table 3 we see that the new backward error bound $\theta \|R(\bar{X}_k)\|$ is sharper and closer to the actual error than the backward error bound $\nu_* \|R(\bar{X}_k)\|$ in [39]. Moreover, we see that the backward error $\theta \|R(\bar{X})\|$ for an approximate solution \bar{X} seems to be independent of the conditioning of the solution X .

Example 3. We consider the matrix equation

$$X - A^* X^{-3} A = 5I, \tag{95}$$

where

$$A = \frac{A_0}{\|A_0\|}, \quad A_0 = \begin{pmatrix} 2 & 0.95 \\ 0 & 1 \end{pmatrix}. \tag{96}$$

We now consider the perturbation bounds for the solution X when the coefficient matrix A is perturbed to $\bar{A} = A + \Delta A$, where

$$\Delta A = \frac{10^{-j}}{\|C^T + C\|} (C^T + C) \tag{97}$$

and C is a random matrix generated by MATLAB function randn.

The conditions in Theorem 7 are satisfied.

By Theorem 7, we can compute the relative perturbation bound ϱ with different values of j . These results averaged as the geometric mean of 10 randomly perturbed runs. Some results are listed in Table 4.

The results listed in Table 4 show that the perturbation bound ϱ given by Theorem 7 is fairly sharp.

Example 4. Consider the matrix equation $X - A^* X^{-3} A = Q$, where

$$A = \begin{pmatrix} 0.5 & 0.55 - 10^{-k} \\ 1 & 1 \end{pmatrix}, \quad Q = \begin{pmatrix} 5 & 1 \\ 1 & 5 \end{pmatrix}. \tag{98}$$

By Remark 12, we can compute the relative condition number $c_{\text{rel}}(X)$. Some results are listed in Table 5.

Table 5 shows that the unique positive definite solution X is well conditioned.

TABLE 4: Results for Example 3 with different values of j .

j	4	5	6	7
$\frac{\ \bar{X} - X\ }{\ X\ }$	1.1892×10^{-7}	2.1101×10^{-8}	2.4085×10^{-9}	1.6847×10^{-10}
ϱ	2.0791×10^{-7}	3.5353×10^{-8}	3.9573×10^{-9}	3.2580×10^{-10}

TABLE 5: Results for Example 4 with different values of k .

k	1	3	5	7	9
$c_{rel}(X)$	1.2510	1.0991	1.0009	1.0009	1.0009

Acknowledgments

The author would like to express her gratitude to the referees for their fruitful comments. This research project was funded by the National Nature Science Foundation of China (11201263), the Nature Science Foundation of Shandong Province (ZR2012AQ004), and Independent Innovation Foundation of Shandong University (IIFSDU), China.

References

[1] W. N. Anderson, Jr., G. B. Kleindorfer, P. R. Kleindorfer, and M. B. Woodrooffe, "Consistent estimates of the parameters of a linear system," *Annals of Mathematical Statistics*, vol. 40, pp. 2064–2075, 1969.

[2] W. N. Anderson, Jr., T. D. Morley, and G. E. Trapp, "The cascade limit, the shorted operator and quadratic optimal control," in *Linear Circuits, Systems and Signal Processing: Theory and Application*, C. I. Byrnes, C. F. Martin, and R. E. Saeks, Eds., pp. 3–7, North-Holland, New York, NY, USA, 1988.

[3] R. S. Bucy, "A priori bounds for the Riccati equation," in *Proceedings of the Berkley Symposium on Mathematical Statistics and Probability, Vol. III: Probability Theory*, pp. 645–656, University of California Press, Berkeley, Calif, USA, 1972.

[4] B. L. Buzbee, G. H. Golub, and C. W. Nielson, "On direct methods for solving Poisson's equations," *SIAM Journal on Numerical Analysis*, vol. 7, pp. 627–656, 1970.

[5] D. V. Ouellette, "Schur complements and statistics," *Linear Algebra and its Applications*, vol. 36, pp. 187–295, 1981.

[6] W. Pusz and S. L. Woronowicz, "Functional calculus for sesquilinear forms and the purification map," *Reports on Mathematical Physics*, vol. 8, no. 2, pp. 159–170, 1975.

[7] J. Zabczyk, "Remarks on the control of discrete-time distributed parameter systems," *SIAM Journal on Control and Optimization*, vol. 12, pp. 721–735, 1974.

[8] A. Ferrante and B. C. Levy, "Hermitian solutions of the equation $X = Q + NX^{-1}N^*$," *Linear Algebra and its Applications*, vol. 247, pp. 359–373, 1996.

[9] V. I. Hasanov, "Notes on two perturbation estimates of the extreme solutions to the equations $X \pm A^*X^{-1}A = Q$," *Applied Mathematics and Computation*, vol. 216, no. 5, pp. 1355–1362, 2010.

[10] V. I. Hasanov and I. G. Ivanov, "On two perturbation estimates of the extreme solutions to the equations $X \pm A^*X^{-1}A = Q$," *Linear Algebra and its Applications*, vol. 413, no. 1, pp. 81–92, 2006.

[11] V. I. Hasanov, I. G. Ivanov, and F. Uhlig, "Improved perturbation estimates for the matrix equations $X \pm A^*X^{-1}A = Q$," *Linear Algebra and its Applications*, vol. 379, pp. 113–135, 2004.

[12] J. Li and Y. H. Zhang, "The Hermitian positive definite solution and its perturbation analysis for the matrix equation $X - A^*X^{-1}A = Q$," *Mathematica Numerica Sinica*, vol. 30, no. 2, pp. 129–142, 2008 (Chinese).

[13] I. G. Ivanov, V. I. Hasanov, and B. V. Minchev, "On matrix equations $X \pm A^*X^{-2}A = I$," *Linear Algebra and its Applications*, vol. 326, no. 1–3, pp. 27–44, 2001.

[14] Y. Zhang, "On Hermitian positive definite solutions of matrix equation $X - A^*X^{-2}A = I$," *Journal of Computational Mathematics*, vol. 23, no. 4, pp. 408–418, 2005.

[15] V. I. Hasanov and I. G. Ivanov, "On the matrix equation $X - A^*X^{-n}A = I$," *Applied Mathematics and Computation*, vol. 168, no. 2, pp. 1340–1356, 2005.

[16] V. I. Hasanov and I. G. Ivanov, "Solutions and perturbation estimates for the matrix equations $X \pm A^*X^{-n}A = Q$," *Applied Mathematics and Computation*, vol. 156, no. 2, pp. 513–525, 2004.

[17] X. Liu and H. Gao, "On the positive definite solutions of the matrix equations $X^s \pm A^T X^{-t} A = I_n$," *Linear Algebra and its Applications*, vol. 368, pp. 83–97, 2003.

[18] J. C. Engwerda, "On the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = I$," *Linear Algebra and its Applications*, vol. 194, pp. 91–108, 1993.

[19] J. C. Engwerda, A. C. M. Ran, and A. L. Rijkeboer, "Necessary and sufficient conditions for the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = Q$," *Linear Algebra and its Applications*, vol. 186, pp. 255–275, 1993.

[20] C.-H. Guo and P. Lancaster, "Iterative solution of two matrix equations," *Mathematics of Computation*, vol. 68, no. 228, pp. 1589–1603, 1999.

[21] J.-g. Sun and S.-F. Xu, "Perturbation analysis of maximal solution of the matrix equation $X + A^*X^{-1}A = P$, II," *Linear Algebra and its Applications*, vol. 362, pp. 211–228, 2003.

[22] X. Zhan, "Computing the extremal positive definite solutions of a matrix equation," *SIAM Journal on Scientific Computing*, vol. 17, no. 5, pp. 1167–1174, 1996.

[23] X. Zhan and J. Xie, "On the matrix equation $X + A^*X^{-1}A = I$," *Linear Algebra and its Applications*, vol. 247, pp. 337–345, 1996.

[24] I. G. Ivanov and S. M. El-sayed, "Properties of positive definite solutions of the equation $X + A^*X^{-2}A = I$," *Linear Algebra and its Applications*, vol. 279, no. 1-3, pp. 303–316, 1998.

[25] Y. Zhang, "On Hermitian positive definite solutions of matrix equation $X + A^*X^{-2}A = I$," *Linear Algebra and its Applications*, vol. 372, pp. 295–304, 2003.

[26] V. I. Hasanov, "On positive definite solutions of the family of matrix equations $X + A^*X^{-n}A = Q$," *Journal of Computational and Applied Mathematics*, vol. 193, no. 1, pp. 277–301, 2006.

[27] J. Cai and G. L. Chen, "On the Hermitian positive definite solution of nonlinear matrix equation $X^s - A^*X^{-t}A = Q$,"

- Applied Mathematics and Computation*, vol. 217, pp. 2448–2456, 2010.
- [28] X. Duan and A. Liao, “On the existence of Hermitian positive definite solutions of the matrix equation $X^s - A^* X^{-t} A = Q$,” *Linear Algebra and its Applications*, vol. 429, no. 4, pp. 673–687, 2008.
- [29] X. Yin, S. Liu, and L. Fang, “Solutions and perturbation estimates for the matrix equation $X^s - A^* X^{-t} A = Q$,” *Linear Algebra and its Applications*, vol. 431, no. 9, pp. 1409–1421, 2009.
- [30] D. Zhou, G. Chen, G. Wu, and X. Zhang, “Some properties of the nonlinear matrix equation $X^s - A^* X^{-t} A = Q$,” *Journal of Mathematical Analysis and Applications*, vol. 392, no. 1, pp. 75–82, 2012.
- [31] V. I. Hasanov, “Positive definite solutions of the matrix equations $X \pm A^* X^{-q} A = Q$,” *Linear Algebra and its Applications*, vol. 404, pp. 166–182, 2005.
- [32] J. F. Wang, Y. H. Zhang, and B. R. Zhu, “The Hermitian positive definite solutions of the matrix equation $X + A^* X^{-q} A = I (q > 0)$,” *Mathematica Numerica Sinica*, vol. 26, no. 1, pp. 61–72, 2004 (Chinese).
- [33] X. Yin, S. Liu, and T. Li, “On positive definite solutions of the matrix equation $X + A^* X^{-q} A = Q (0 < p \leq 1)$,” *Taiwanese Journal of Mathematics*, vol. 16, no. 4, pp. 1391–1407, 2012.
- [34] X. Duan, A. Liao, and B. Tang, “On the nonlinear matrix equation $X - \sum_{i=1}^m A_i^* X^{-1} A_i = I$,” *Linear Algebra and its Applications*, vol. 429, no. 1, pp. 110–121, 2008.
- [35] X. Duan, C. Li, and A. Liao, “Solutions and perturbation analysis for the nonlinear matrix equation $X + \sum_{i=1}^m A_i^* X^{-1} A_i = I$,” *Applied Mathematics and Computation*, vol. 218, no. 8, pp. 4458–4466, 2011.
- [36] Y.-m. He and J.-h. Long, “On the Hermitian positive definite solution of the nonlinear matrix equation $X + \sum_{i=1}^m A_i^* X^{-1} A_i = I$,” *Applied Mathematics and Computation*, vol. 216, no. 12, pp. 3480–3485, 2010.
- [37] X. Yin and S. Liu, “Positive definite solutions of the matrix equations $X \pm A^* X^{-q} A = Q (q \geq 1)$,” *Computers & Mathematics with Applications*, vol. 59, no. 12, pp. 3727–3739, 2010.
- [38] H. Xiao and J. T. Wang, “On the matrix equation $X - A^* X^{-p} A = Q (p > 1)$,” *Chinese Journal of Engineering Mathematics*, vol. 26, no. 2, pp. 305–309, 2009.
- [39] J. Li and Y. Zhang, “Perturbation analysis of the matrix equation $X - A^* X^{-q} A = Q$,” *Linear Algebra and its Applications*, vol. 431, no. 9, pp. 1489–1501, 2009.
- [40] J. R. Rice, “A theory of condition,” *SIAM Journal on Numerical Analysis*, vol. 3, pp. 287–310, 1966.
- [41] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Posts and Telecom press, Beijing, China, 2005.
- [42] R. A. Horn and C. R. Johnson, *Matrix Analysis*, Posts and Telecom press, Beijing, China, 2005.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

