

Review Article

The Cosmic History of Black Hole Growth from Deep Multiwavelength Surveys

Ezequiel Treister^{1,2} and C. Megan Urry^{3,4,5}

¹Departamento de Astronomía, Universidad de Concepción, Casilla 160-C, Concepción 4070386, Chile

²Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

³Yale Center for Astronomy and Astrophysics (YCAA), P.O. Box 208121, New Haven, CT 06520, USA

⁴Department of Physics, Yale University, P.O. Box 208121, New Haven, CT 06520, USA

⁵Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520, USA

Correspondence should be addressed to Ezequiel Treister, etreiste@astroudec.cl

Received 21 September 2011; Accepted 30 November 2011

Academic Editor: Francesca Civano

Copyright © 2012 E. Treister and C. M. Urry. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Significant progress has been made in the last few years on understanding how supermassive black holes form and grow. In this paper, we begin by reviewing the spectral signatures of active galactic nuclei (AGN) ranging from radio to hard X-ray wavelengths. We then describe the most commonly used methods to find these sources, including optical/UV, radio, infrared, and X-ray emission, and optical emission lines. We then describe the main observational properties of the obscured and unobscured AGN population. Finally, we summarize the cosmic history of black hole accretion, that is, when in the history of the universe supermassive black holes were getting most of their mass. We finish with a summary of open questions and a description of planned and future observatories that are going to help answer them.

1. Introduction

Astrophysical black holes come in a wide range of masses, from $\gtrsim 3M_{\odot}$ for stellar mass black holes [1] to $\sim 10^{10}M_{\odot}$ for so-called supermassive black holes [2, 3]. The best evidence for the existence of a supermassive black hole can be found in the center of the Milky Way galaxy, where from dynamical studies, the mass of the Sgr A* source was established to be $\sim 4.4 \times 10^6 M_{\odot}$ [4, 5].

Evidence for the existence of supermassive black holes has also been found in other massive nearby galaxies [6], mostly from resolved stellar and gas kinematics. For active galaxies, it has been possible to use the technique known as reverberation mapping [7–9]. From these observations, a clear correlation has been established between the mass of the central black hole and properties of the host galaxy such as stellar mass in the spheroidal component [10], luminosity [11], velocity dispersion [12, 13] and mass of the dark matter halo [14]. The fact that such correlations exist, even though these components have very different spatial

scales, suggests a fundamental relationship between black hole formation and galaxy evolution. Furthermore, it is now well established by simulations [15] that the energy output from the growing central black hole can play a significant role in the star formation history of the host galaxy. In particular, theory suggests that nuclear activity regulates star formation either by removing all the gas [16, 17] or by heating it [18]. It is therefore obvious that a complete study of galaxy evolution requires a comprehensive understanding of black hole growth.

Most current black hole formation models tell us that the first black hole seeds formed at $z \gtrsim 15$. While the exact mechanism for the formation of the first black holes is not currently known, there are several prevailing theories (see the comprehensive reviews by Rees [19] and Volonteri [20] for more details). One of the most popular possibilities is that the first black hole seeds are the remnants of the first generation of stars, the so-called population III stars, formed out of primordial ultralow metallicity gas. These black holes formed at $z \sim 20$ and have typical masses $\sim 100\text{--}1,000M_{\odot}$.

This scenario has problems explaining the very high masses, of $\sim 10^9 M_\odot$, estimated for supermassive black holes in $z \sim 6$ optically selected quasars [21]. Alternatively, the first black holes could have formed directly as the result of gas-dynamical processes. It is possible for metal-free gas clouds with $T_{\text{vir}} \gtrsim 10^4$ K and suppressed H_2 formation to collapse very efficiently [22], possibly forming massive black hole seeds with $M \sim 10^4$ – $10^5 M_\odot$ as early as $z \sim 10$ – 15 . If instead the UV background is not enough to suppress the formation of H_2 , the gas will fragment and form “normal” stars in a very compact star cluster. In that case, star collisions can lead to the formation of a very massive star, that will then collapse and form a massive black hole seed with mass $\sim 10^2$ – $10^4 M_\odot$ [23].

Given the current masses of $10^{6-9} M_\odot$, most black hole growth happens in the active galactic nuclei (AGN) phase [2, 24]. With typical bolometric luminosities $\sim 10^{45-48} \text{ erg s}^{-1}$, AGN are amongst the most luminous emitters in the universe, particularly at high energies and radio wavelengths. These luminosities are a significant fraction of the Eddington luminosity—the maximum luminosity for spherical accretion beyond which radiation pressure prevents further growth—for a $10^{8-9} M_\odot$ central black hole. A significant fraction of the total black hole growth, $\sim 60\%$ [25], happens in the most luminous AGN, quasars, which are likely triggered by the major merger of two massive galaxies [26]. In an AGN phase, which lasts $\sim 10^8$ years, the central supermassive black hole can gain up to $\sim 10^7$ – $10^8 M_\odot$, so even the most massive galaxies will have only a few of these events over their lifetime. Further black hole growth, mostly in low-luminosity (low Eddington rate) AGN, is likely due to stochastic accretion of cold gas, mostly in spiral galaxies [27].

According to the AGN unification paradigm [28, 29], a large fraction of these sources, $\sim 75\%$ locally, are heavily obscured by optically and geometrically thick axisymmetric material, which explains many of the observed differences among different types of active galaxies. In addition, luminosity [30] and cosmic epoch [31] play a significant role. One constraint on the fraction of obscured AGN and its evolution comes from the spectral shape of the extragalactic X-ray “background” (XRB). Thanks to deep X-ray observations at $E \lesssim 10$ keV performed by *Chandra* and *XMM-Newton*, a very large fraction of the X-ray background, $\sim 80\%$, has been resolved into point sources [32], the vast majority of them AGN [33]. Several studies, the first of them ~ 20 years ago [34], have used a combination of obscured and unobscured AGN to explain the spectral shape and normalization of the X-ray background with overall good results. The latest AGN population synthesis models [35, 36] assume an average ratio of obscured to unobscured AGN of $\sim 3:1$ locally, increasing towards lower luminosities and higher redshifts, as well as a fraction of Compton-thick sources (CT; $N_H > 10^{24} \text{ cm}^{-2}$) of ~ 5 – 10% , consistent with the value observed at higher energies, $E = 10$ – 100 keV, of $\sim 5\%$ by INTEGRAL in the local universe [35, 37], lower by factors of ~ 3 than expectations of previous population synthesis models [38, 39].

In this paper, we review multiwavelength methods used to trace the growth of SMBHs (Section 2), the known properties of unobscured and obscured AGN (Sections 3 and 4,




FIGURE 1: Average unobscured radio-quiet AGN spectrum from far-infrared to hard X-rays, as compiled by Manners (2002; [41]). The contributions from each component are shown separately together with the total AGN emission (*solid line*). Most of the AGN radiation appears in three regions: infrared (reemission from dust; *dashed red line*), UV and optical (accretion disk; *dot-dot-dot-dashed blue line*), and X-rays (hot corona and reflection from the accretion disk; *dashed green and dot-dashed cyan lines*). At longer wavelengths, $>100 \mu\text{m}$, a starburst component associated with the host galaxy dominates (*dotted magenta line*).

resp.), the cosmic history of black hole accretion (Section 5), and prospects for future observations (Section 6). Throughout this paper, we assume a Λ CDM cosmology with $h_0 = 0.7$, $\Omega_m = 0.27$, and $\Omega_\Lambda = 0.73$, in agreement with the most recent cosmological observations [40].

2. How to Trace SMBH Growth?

One of the main features of the AGN emission is that it covers a very wide range of wavelengths, from radio to Gamma-rays (Figure 1). While unobscured sources are easily detectable and identified by their UV and soft X-ray continuum and their broad optical emission lines, obscured AGN can only be found at longer, mid-IR, wavelengths or in hard X-rays. Of course, selections at different wavelengths have different biases. For example, while radio surveys are not particularly affected by obscuration, they are most likely to detect radio-loud sources, which are only $\sim 10\%$ of the total AGN population at bright fluxes [42]. However, combining different multiwavelength techniques gets us closer to a complete picture. Below, we briefly describe the most popular AGN selection methods, their advantages, and drawbacks.

2.1. Optical/UV Continuum. Rest-frame optical/UV selection of AGN, in particular for high-luminosity unobscured quasars, is particularly efficient because the spectral shapes of




FIGURE 2: Composite rest-frame optical/UV spectrum for the optically selected quasars in the Sloan Digital Sky Survey, from the work of Vanden Berk et al. [49]. The *dashed* and *dotted* lines show power-law fits to the continuum emission.

stars and quasars at these wavelengths produce very different broadband colors due to the presence of the “big blue bump” [43] in quasar spectra from $\sim 100 \text{ \AA}$ to $\sim 1 \mu\text{m}$ (Figure 2). This emission is often attributed to the thermal radiation with temperatures $\sim 30,000 \text{ K}$ originating in the accretion disk [44]. This unique spectral shape has been used in the past to identify quasars with great success by optical surveys such as the Palomar-Green (PG) survey [45], the 2 degree field QSO redshift survey [46], or more recently the Sloan Digital Sky Survey (SDSS; [47]), which has found now more than 1 million quasars [48].

However, samples selected in the optical are far from complete, as emission at these wavelengths is strongly affected by reddening or extinction, and most AGN are obscured along our line of sight. Furthermore, for lower-luminosity sources, the optical light from the host galaxy can easily outshine the nuclear emission. This is particularly important for ground-based observations and high-redshift sources, for which it is very hard to separate the nonthermal and stellar components spatially.

2.2. Radio. Historically, identification of AGN based on their radio emission has been very important. In fact, the first discovered quasar, 3C 273, was originally classified as a radio source [71]. In spite of this, radio selection can be very problematic. In radio-loud sources (defined as $f_{5 \text{ GHz}}/f_B > 10$ [72]) radio emission is associated with a strong, nonthermal, component, probably originating in a beamed collimated relativistic jet [73]. In radio-quiet sources, which are typically ~ 3 orders of magnitude fainter at these wavelengths [74], the radio emission corresponds to the long-wavelength tail of the far-infrared dust emission. As a consequence, radio-selected

samples are necessarily biased towards radio-loud sources, which represent only $\sim 10\%$ of the overall AGN population.

2.3. Optical Emission Lines. As first reported by Baldwin et al. [81], the photoionizing spectrum of a power law continuum source, such as an AGN, produces very different emission line intensity ratios when compared with that of typical star-forming regions (mostly due to O and B stars). Hence, emission lines can be used to identify the presence of AGN even in galaxies in which the optical/continuum does not show any direct AGN signature, due to obscuration and/or low luminosity. Because the AGN ionizing emission reaches material even a few kiloparsecs away from the nuclear region, this selection technique is less sensitive to circumnuclear obscuration and thus provides a more complete AGN view when compared with, for example, optical/UV continuum selection. This technique was used successfully in the SDSS [50, 51] to extend the low-redshift AGN sample to lower luminosities. Emission line ratios and diagnostic regions can be seen in Figure 3. Emission-line selection can also be used at higher redshifts, as shown by the DEEP2 galaxy redshift survey, which selected a sample of 247 AGN at $z \sim 1$ from optical spectroscopy using the DEIMOS spectrograph at the Keck observatory [82].

While this is an efficient AGN selection technique, optical spectroscopy is very expensive in telescope time and is only feasible for relatively bright emission line regions. This selection may be incomplete at the low luminosity end, if the host galaxy can outshine the high-ionization emission lines. It is currently very difficult to extend this selection beyond $z \sim 1$, as the relevant emission lines move to observed-frame near-IR wavelengths, where current-generation spectrographs are significantly affected by atmospheric emission, do not cover wide field of views, and have limited multiobject capabilities.

2.4. X-Rays. As was found more than 30 years ago, AGN are ubiquitous X-ray emitters [89]. Their X-ray emission extends from $\sim 0.1 \text{ keV}$ to $\sim 300 \text{ keV}$ and is attributed to inverse-Compton scattering due to high-energy electrons in a hot corona, surrounding the accretion disk. The high-energy cutoff at $\sim 100\text{--}300 \text{ keV}$ is presumably due to a cutoff in the energy distribution of the electrons in the hot corona. AGN are typically $\sim 1\text{--}5$ orders of magnitude more luminous in X-rays than normal galaxies, which makes them the dominant extragalactic population at these wavelengths. Most AGN are obscured by photoelectric absorption by gas along the line of sight, which preferentially affects emission at the lower energies. This is often parametrized by the amount of neutral hydrogen column density in the line of sight. Figure 4 shows typical AGN X-ray spectra including the power law component and high energy cutoff, for different levels of photoelectric absorption.

Deep X-ray surveys with *Chandra* and *XMM-Newton* have found the largest AGN densities, $\sim 7,000 \text{ deg}^{-2}$ [90], $\sim 10\text{--}20$ times higher than even the deepest optical surveys. Still, X-ray selected AGN samples are still biased against the most obscured sources. In fact, even the deepest *Chandra* surveys can miss more than half of the total AGN due to a




FIGURE 3: AGN selection diagrams based on optical emission line ratios. Figure obtained from Kewley et al. [50]. Divisions between star-forming galaxies and AGN are shown by the *dashed* [51] and *solid* [52] lines; bona-fide AGN sit at the upper right in these distributions.




FIGURE 4: Typical AGN X-ray spectrum from 0.1 to 100 keV. Each spectrum includes a power law with photon index $\Gamma = 1.9$, a Compton reflection hump, peaking at ~ 30 keV, a high-energy cutoff at 300 keV, and photoelectric absorption. Numbers for each curve indicate the amount of absorption, given as $\log(N_H)$, with N_H in units of atoms cm^{-2} .

combination of obscuration and/or low luminosity [91]. Observing at higher energies helps to alleviate the effects of obscuration. Wide-area surveys performed by the *Swift* [66] and *INTEGRAL* [92] satellites have detected a large number of AGN in the local universe at $E = 10\text{--}100$ keV, where all but the most obscured, Compton-thick, sources emit strongly. Unfortunately, due to their relatively high flux limits ($\sim 2\text{--}3$ order of magnitudes shallower than the deeper *Chandra* observations), surveys at these energies are so far limited to low redshifts only.

2.5. Infrared. Intrinsic AGN emission is not particularly strong at near/mid-IR wavelengths. Radiation coming from the accretion disk, often characterized as a power-law, while very strong at UV and optical wavelengths decreases rapidly beyond $\sim 1\ \mu\text{m}$ [93]. However, as was originally found by

IRAS [74] and latter confirmed by *ISO* [94] and *Spitzer* [95], AGN are luminous IR sources. This is commonly attributed to reemission of absorbed energy by dust. This radiation can be found starting at $\sim 2\text{--}3\ \mu\text{m}$, which corresponds to the dust sublimation temperature, about $\sim 1,000\text{--}2,000$ K [96], and extends to $\sim 100\ \mu\text{m}$, at the tail of the black body spectrum for a $\sim 100\text{--}1,000$ K distribution [97], where the emission from the host galaxy typically starts to dominate. Typical AGN infrared luminosities are $10^{44\text{--}46}$ erg s^{-1} and thus they represent a significant fraction, $\sim 30\%$ on average [98], of the bolometric luminosity.

One clear advantage of infrared AGN selection is that this reemission is mostly isotropic, and hence both obscured and unobscured sources have similar detection probabilities. Therefore, it provides a complementary approach to the most common selection techniques described above, which are less sensitive to obscured sources. However, star forming galaxies are very luminous at these wavelengths as well, so that the host galaxy can easily outshine the central emission, in particular for low-luminosity sources, hence yielding a low efficiency in detecting AGN [99]. Furthermore, the selection function in infrared studies is more complicated, as the probability of detecting the AGN depends on the properties of the host galaxy, such as the amount of dusty star-formation, which is in principle independent of the nuclear luminosity.

3. Unobscured AGN

Because quasars are the most luminous, and thus easily detectable, members of the AGN family, the luminosity function of optical quasars has been well determined for years. In particular, it was found that the number of quasars evolve strongly [100] and peak at $z \approx 2$ [101]. This evolution has been modeled either as a pure luminosity evolution (PLE), in which the characteristic luminosity changes with redshift while the shape of the luminosity function remains the same [102], or a pure density evolution (PDE), so that only the normalization of the luminosity function depends on redshift [100]; however, it was quickly discovered that




FIGURE 5: Binned optical quasar luminosity function from the work of Croom et al. ([53]; their Figure 11) in six redshift intervals, from $z = 0.4$ to $z = 2.1$. Measurements were obtained from the 2dF QSO Redshift Survey (2QZ; [54] *blue*), the 2dF-SDSS LRG and QSO survey (2SLAQ; [55]; *green*), and combining the latter with the general SDSS QSO sample ([56]; *red*). The dotted line shows the measured luminosity function in the $1.53 < z < 1.81$ range for reference. The strong evolution of the QSO luminosity function is clearly seen in this figure. This evolution is best fitted by a LEDE model, as described in the text.

at least in the PG quasar survey, the shape of luminosity function also evolves with redshift and thus neither a PLE nor a PDE provides a good description [45].

In Figure 5 we show one of the latest measurements of the quasar luminosity function at $0.4 < z < 2.6$, from Croom et al. [53]. They conclude that a luminosity-dependent density evolution provides a better fit to the optical quasar luminosity function, than either a PLE or PDE. Similar conclusions were reached by studying X-ray selected sources using soft X-ray (0.5–2 keV) observations [103] or hard X-ray (2–10 keV) data [79, 104] which includes both obscured and unobscured AGN. However, taking advantage of the large number of sources in their sample, $\sim 10,000$, Croom et al. [53] concluded that the best fit to the observed luminosity function is obtained by using a model based on a luminosity

evolution + density evolution (LEDE). The most important difference between a LEDE and a PLE fit is a change in amplitude and bright-end slope at high redshift.

An important conclusion obtained from observations of the quasar luminosity function is the evidence for “cosmic downsizing” [105], that is, that the most massive black holes get most of their mass at high redshift, while at low redshift only low mass black holes are still growing. This is observed both in optical [53] and hard X-ray luminosity functions [104, 105], thus indicating that this result is independent of obscuration. Recent deep optical surveys such as the Great Observatories Origins Deep Survey (GOODS; [106]), the Cosmic Evolution Survey (COSMOS; [59]), the NOAO Deep Wide Field Survey (NDWFS), and the Deep Lens Survey (DLS; [58]) have produced significant advances in extending




FIGURE 6: (a) $z \sim 4$ quasar luminosity function, from the work of Glikman et al. [57]. The *filled red circles* show the measurements presented on that work, while the *open red circles* were reported previously by the same group [58] and show the change with increased spectroscopic completeness. The *triangles* were obtained from the SDSS quasar sample at $z = 4.25$ [56]. The *blue squares* are the space densities of $z \sim 4$ QSOs from Ikeda et al. [59] and the *dot-dashed line* is their best-fit double power law. The lower right-hand legend lists the best-fit parameters to a double power-law (*solid line*) shaded region represents the 1σ uncertainties. Dashed and dotted lines show the $z \sim 3$ QLF [60], representing different fits to the observed QLF. (b) Quasar space density as a function of redshift, from the work of Ikeda et al. [59]. *Dotted lines* used the combined 2SLAQ, SDSS, SWIRE, NDFWS and DLS samples, while the *dashed lines* combine the COMOS and 2SLAQ sources. While AGN downsizing is clearly visible at $z < 2.5$, at higher redshifts the situation is more uncertain.

the quasar luminosity function to higher redshifts, $z > 3$. Figure 6 shows the quasar luminosity function at $z \sim 4$ and the redshift dependence of the quasar spatial density [58, 106]. While the presence of downsizing is clear up to $z \sim 2.5$, at higher redshifts it is less convincing, most likely due to poor statistics and incompleteness. As argued by Glikman et al. [57], the slope of the faint end of the quasar luminosity function is critical in determining the contribution of these sources to the ionization of the intergalactic medium. Based on current measurements, quasars contribute $\sim 60\%$ of the ionizing photons at $z \sim 4$ and thus are the dominant source at this redshift.

At even higher redshifts, $z \sim 5-6$, current deep surveys do not cover enough area to detect a significant number of sources. However, wide-area survey such as SDSS [107] and the Canada-France High- z Quasar Survey [108] have been able to find a sizable sample, ~ 40 high-luminosity quasars, at these high redshifts. According to these samples, there is a large decrease in the number density of high-redshift quasars, when compared to $z \sim 2$, suggesting that the peak of the quasar activity is at $z \sim 2.5$ [56]. The decline towards higher redshifts is given by $10^{-0.47z}$ [107] from $z = 3$ to $z = 6$. A similar trend is also observed for high-luminosity sources in

X-ray surveys [109], which are also dominated by unobscured sources. This indicates that due to their very low spatial density, unobscured quasars do not contribute significantly to the early hydrogen reionization of the intergalactic medium at $z \sim 6$ [107, 108], in contrast to the situation at $z \sim 4$.

4. Obscured Accretion

The space density and evolution of the unobscured AGN population has been well studied, mostly from optical and soft X-ray surveys. However, we know that a large fraction of the SMBH growth happens in heavily obscured systems. Observations of the nearest AGN suggest that the local ratio of obscured to unobscured sources is $\sim 4:1$ [110]. A similarly high fraction of obscured AGN has been used to explain the spectrum and normalization of the extragalactic XRB, as shown by the latest AGN population synthesis models [35, 38, 39, 111]. The XRB gives an integral constraint to the AGN population and its evolution; the most recent deep surveys show that $\sim 90\%$ of the observed 2–8 keV XRB radiation can be attributed to resolved AGN [32]. In Figure 7, we show the latest AGN population synthesis models for




FIGURE 7: Observed spectrum of the extragalactic XRB from HEAO-1 [61], Chandra [32], XMM [62], INTEGRAL [63], and Swift [64] data. The *thick black solid line* shows the population synthesis model for the XRB spectrum of Treister et al. [35]. *Red, blue, and thin black solid lines* show the contribution to this model from unobscured, obscured Compton thin, and CT AGN, respectively.

the XRB which uses a local ratio of obscured to unobscured AGN of $\sim 3:1$, plus a luminosity and redshift dependence, as described below [35]. The largest uncertainty in such models stems from the normalization mismatch of the data in the 1–10 keV energy range.

A possible dependence of the fraction of obscured AGN on luminosity was first suggested nearly 20 years ago [30] and confirmed since then by hard X-ray surveys [104, 105, 112]. A possible physical explanation is the so-called “receding torus,” in which the size of the inner opening angle depends on luminosity [30, 113]. More recent observations found a luminosity dependence of the ratio of mid-IR to bolometric flux for unobscured AGN, consistent with this idea [98]. Alternatively, it has been proposed that the observed dependence of the obscured fraction on luminosity could be explained either by the effects of photoionization on the X-ray obscuring matter [114] or by the Eddington limit on a clumpy torus [65]. In Figure 8 we show the observed fraction of obscured AGN as a function of luminosity obtained by combining data from deep *Chandra* X-ray surveys [115]. A consistent result is observed for AGN selected in a hard X-ray ($E = 14\text{--}195\text{ keV}$) survey, as shown in Figure 8(b), indicating that this trend is not due to selection biases.

In the Figure 8(a) we compare the observed dependence of the fraction of obscured sources on luminosity with the expectations for different geometrical parameters of the

obscuring material. If the height of the torus is roughly independent of luminosity, the change in covering fraction is due to a change in inner radius (the original “receding torus” model), hence a rough $L^{-1/2}$ dependence for the contrast should be expected [30, 116]. If the effects of radiation pressure are incorporated, in the case of a clumpy torus, a $L^{-1/4}$ dependence is expected. As can be seen in Figure 8, a $L^{-1/2}$ dependence is too steep compared to observed data. This implies that the height of the obscuring material cannot be independent of the source luminosity and provides evidence for a radiation-limited structure.

The dependence of the fraction of obscured AGN on redshift is more controversial. While some studies [31, 80, 111, 117, 118] found a small increase in the fraction of obscured AGN at higher redshifts, other results suggest that this fraction is constant [104, 119]. These discrepancies can be understood due to a combination of small samples and the use of N_H to classify AGN, which produces a well-known redshift bias [119]. The fraction of obscured AGN as a function of redshift for a large, $\sim 2,000$ sources, X-ray selected sample [115], using optical emission lines to separate obscured and unobscured AGN, is shown in Figure 9. It increases significantly with redshift, roughly as $(1+z)^\alpha$, with $\alpha = 0.3\text{--}0.5$ (thin dashed lines, bottom panel, Figure 9; best fit, $\alpha \simeq 0.4$, thick dashed line). This value of α does not change significantly if a different host galaxy evolution is assumed, and it is consistent with the value of 0.3 reported by other studies [111, 117, 118].

Since star-forming galaxies may be expected to have more dust, the increase in the relative fraction of obscured AGN at high redshift may be due to an increase in the contribution to obscuration by galactic dust. By combining hard X-ray and mid-infrared observations, a similar ratio of hard X-ray to mid-infrared flux for obscured and unobscured AGN has been found [120], contrary to the predictions of the simplest AGN unification paradigm, in which the obscuration comes from the dusty torus and therefore the mid-infrared emission is reduced due to self-absorption. This result can be explained if the obscuration comes from a much more extended region, that is, kiloparsec, galactic scales rather than a compact parsec-scale torus. Furthermore, signatures for extended absorbing regions have been detected in nearby galaxies like NGC 1068 [121] and NGC 4151 [122]. Heavy absorption at kiloparsec scales has routinely been found in ultraluminous infrared galaxies (ULIRGs), which suffer a very strong evolution [123]. Hence, it seems likely that the change in the relative fraction of obscured AGN could be related to galactic-scale absorption, in particular since some ULIRGs also contain an obscured AGN (e.g., Arp 220; [124]).

Below, we review in more detail our current knowledge of the obscured AGN population at three different cosmic epochs: $z \simeq 0$, $z = 1\text{--}3$ and $z > 6$.

4.1. Obscured AGN in the Local Universe. Nearby AGN are found in the so-called Seyfert galaxies [125], which are known to host low luminosity and/or obscured nuclear activity [110, 126]. These growing supermassive black holes have been identified because of their high-ionization optical emission lines and in some cases their blue UV/optical




FIGURE 8: Ratio of obscured to total AGN as a function of hard X-ray luminosity. The black circles with *dashed error bars* show the obscured AGN fraction from the extended *Chandra* deep field south (ECDF-S) alone, while the *gray circles* show the results obtained using the meta-analysis by Treister and Urry [31]. Black circles with *solid error bars* show the fraction obtained combining these two samples. The dependence in the AGN population synthesis model of Gilli et al. [39] for the intrinsic and observed fractions of obscured AGN is shown by the *dashed* and *dotted red lines*; the dependence used by Treister and Urry [31] is shown by the *solid blue line*. The *dashed magenta line* shows the expected dependence for a radiation-limited torus [65], while the *dotted green line* shows the expectation for the original “receding torus” [30], both normalized to the observed value in the 10^{42-43} erg s^{-1} bin. (b) Same as (a) but using an AGN sample at $z \sim 0$ selected in hard X-rays from *Swift*/BAT observations [37]. The fact that the same luminosity dependence is observed in both samples indicates that it is not due to selection effects.

continuum. The first nearby AGN catalogues, produced ~ 40 years ago [127], contained ~ 200 quasars. Roughly speaking, $\sim 5\text{--}15\%$ of the galaxies near the Milky Way contain an active nucleus [110], and $\sim 75\%$ of these active galaxies are obscured. In fact, two of the three nearest AGN are Compton-thick (NGC 4945 and the Circinus galaxy [128]). Hence, optical surveys are not particularly efficient in unveiling this accretion, while observations at other wavelengths, in particular in the infrared [129], and hard X-rays, are more complete.

Surveys at hard X-ray energies, $E > 15$ keV, have been very successful in providing the most complete AGN samples in the local Universe. As long as the neutral hydrogen column density is lower than $\sim 10^{24}$ cm^{-2} , the direct AGN emission is mostly unaffected at these energies. Current observations at $E > 10$ keV with the International Gamma-Ray Astrophysics Laboratory (INTEGRAL; [130]) and Swift [131] satellites are available only at relatively high fluxes and hence low redshifts, $z < 0.05$.

Using the IBIS coded-mask telescope [132], INTEGRAL surveyed $\sim 80\%$ of the sky down to a flux of 5 mCrab in the 17–60 keV. Krivonos et al. [67] report the properties of 130 AGN detected in these all-sky observations. A large number of unidentified sources remain in this full INTEGRAL catalog (48) but only seven are found at high galactic latitude ($|b| > 5^\circ$) and thus are likely of extragalactic origin. Five of the 130 known AGN are Compton thick. Using similar observations

from the all-sky Swift/BAT survey, a catalog of 103 AGN [66] contains five AGN with estimated N_H greater than 10^{24} cm^{-2} . However, we caution that some of these N_H measurements were obtained by fitting a single absorbed power law to the X-ray spectrum, while heavily absorbed AGN have more complex spectra [133, 134], so the N_H estimates are likely to be lower limits.

Figure 10 shows the cumulative number counts of AGN, with CT sources shown separately, as a function of hard X-ray flux. In order to avoid the necessity of specifying a standard spectrum to convert fluxes to different energy bands, we show the INTEGRAL and Swift sources separately, but note that there is good agreement (within $\sim 40\%$) in the normalization between the two distributions if a standard band conversion is assumed. At these high fluxes the slope of the $\log N\text{--}\log S$ is Euclidean, implying a uniform spatial distribution, as expected given the low redshifts of these sources. The number of CT AGN found by these surveys is surprisingly low, compared to the sample of known CT AGN in the local universe, most likely due to the effects of obscuration even at these high energies [35, 37, 68]. A study of optically selected local Seyfert 2 galaxies with hard X-ray information [126] found 12 CT AGN in a total of 45 Seyfert galaxies. Three were detected by *Chandra* and/or *XMM*, while the rest are mostly reflection-dominated sources, too faint to be detected by either INTEGRAL or Swift even though they are nearby, moderate-luminosity AGN. This suggests




FIGURE 9: Fraction of obscured AGN as a function of redshift. *Upper panel:* direct measurements using the sources on the ECDF-S field only (black circles with dashed error bars), from the sample of Treister and Urry ([31]; gray circles) and combining both samples (black circles with solid error bars). The expected observed fraction for an intrinsic fraction of 3 : 1 obscured to unobscured AGN, accounting for optical and X-ray selection effects, is shown by the black solid line. As can be seen, while the observed fraction of obscured AGN declines toward higher redshifts, if the X-ray and optical selection effects and the luminosity dependence of the obscured AGN fraction are taken into account, this decline should be even stronger. *Bottom panel:* Inferred fraction of obscured AGN relative to an intrinsically constant fraction after correcting for selection effect and including the luminosity dependence of the obscured AGN fraction. Symbols are the same as for the upper panel. The corrected fraction of obscured AGN increases with redshift following a power law of the form $(1+z)^\alpha$ with $\alpha = 0.4 \pm 0.1$.

that even hard X-ray surveys miss quite a few Compton, thick AGN.

The observed fraction of CT AGN in the INTEGRAL and Swift/BAT hard X-ray selected samples is low, $\sim 5\%$. A very similar and consistent value, 4.6%, was recently obtained from a sample of 307 objects detected in the three-year all-sky Swift/BAT survey [37]. This was initially surprising, since previous AGN population synthesis models that can explain the XRB used much higher CT fractions of $\sim 15\text{--}20\%$ [38, 39], that is, factors of 3-4 higher. We now know that even observations at these high energies can be affected by obscuration, if the column density is high enough. For example, $\sim 50\%$ of the source flux in the 15–55 keV range can be lost if $\log N_H > 24.5$ [135]. As pointed out by Malizia et al. [68], and as can be clearly seen in Figure 11, the INTEGRAL all-sky observations, which have a similar flux limit as the Swift/BAT images, show a steep decline in the number of obscured sources, from $\sim 80\%$ at $z < 0.015$ to $\sim 20\text{--}30\%$ at higher redshifts. Also, all the CT AGN in this sample were found at $z < 0.015$. Hence, these authors concluded that the INTEGRAL

observations are affected by obscuration at larger distances and that the intrinsic fraction of CT sources is $\sim 25\%$, as observed at $z < 0.015$. However, it is worth mentioning that these additional sources, because of their very high column densities, do not contribute significantly to the XRB, although they certainly contribute to black hole growth.

The cumulative contribution of CT AGN to the XRB, as a function of redshift, determined from population synthesis models, is shown in Figure 12. As can be seen, the total contribution of CT AGN to the XRB is $\sim 9\%$, and about 50% of it comes from sources at $z < 0.7$. Similarly, only $\sim 2\%$ of the XRB is provided by CT AGN at $z > 1.4$, while CT AGN at $z > 2$ contribute only $\lesssim 1\%$ to the XRB. Conversely, the 5% uncertainty in the absolute measurement of the XRB intensity translates into an uncertainty of a factor of ~ 5 in the number of CT AGN at $z > 2$. Hence, the number of CT AGN at high redshift is largely unconstrained by the XRB.

4.2. Obscured AGN at Intermediate Redshifts ($1 \lesssim z \lesssim 3$).

As shown in the previous section, the number of heavily obscured AGN at intermediate redshifts, $z \gtrsim 1$, is largely unconstrained by the XRB due to model degeneracies, or by current X-ray surveys at $E > 10$ keV, which do not have the required sensitivity. NuSTAR will change this situation dramatically. However, for now we are forced to use alternative methods to determine the amount of black hole growth occurring in these sources. We explore here two of these techniques, which have been particularly successful: X-ray stacking and mid-IR AGN selection.

Using the deepest available X-ray observations obtained with Chandra, as in the example shown in Figure 13, a number of moderately obscured AGN have been identified at $z > 1$. The observed-frame hard X-ray band (2–8 keV band) at these redshifts covers higher rest-frame energies, thus making them less affected by obscuration. For example, the vast majority of the sources the GOODS fields with high X-ray to optical flux ratios are obscured at $z \sim 2$ [136–138]. Furthermore, from X-ray spectral analysis, ~ 30 CT AGN candidates have been identified in the CDF-S [70] and CDF-N [139]. However, it is clear that X-ray selection remains highly incomplete for obscured sources at these redshifts [91].

Because much of the energy absorbed at optical to X-ray wavelengths is later reemitted in the mid-to far-IR, it is expected that AGN, in particular the most obscured ones, should be very bright mid-IR sources [95]. Sources having mid-IR excesses, relative to their rest-frame optical and UV emission, have been identified as potential CT AGN candidates at $z \sim 2$ [75, 77, 140–142]. However, because of the strong connection between vigorous star formation and AGN activity in the most luminous infrared sources [26], the relative contribution of these two processes remains uncertain and controversial [143–145]. Significant progress has been made by virtue of deep Spitzer observations, in particular using the $24 \mu\text{m}$ band. At $z \sim 1\text{--}2$, this emission corresponds to rest-frame wavelengths of $\sim 10 \mu\text{m}$, where the contrast between AGN and star formation is largest. In order to look for high-luminosity obscured AGN missed by X-ray observations, Fiore et al. [78, 141] defined the “mid-IR excess” region as $f_{24}/f_R > 1000$ and $R-K > 4.5$ (Vega), as shown




FIGURE 10: LogN-logS distribution for AGN detected at high energies ($E > 10$ keV). The *gray line* in the (a) shows the AGN in the well-defined Swift/BAT samples in the 14–195 keV band [66], while the (b) shows the INTEGRAL sources [67] in the 17–60 keV band. *Solid squares* show the 15 sources detected in the ultradeep 3 Msec INTEGRAL observations of the XMM-LSS field. *Solid circles* mark the CT AGN detected with Swift (a) and INTEGRAL (b); the fraction is $\sim 5\%$. The *black solid lines* show the expected AGN LogN-logS from the most complete AGN population synthesis model [38], which at these high fluxes has a Euclidean slope. The *dashed lines* mark the Euclidean slope normalized to the number of Swift and INTEGRAL CT AGN (5% of the total). The *gray lower limits* show the previously known transmission-dominated AGN with hard X-ray observations, not detected in the INTEGRAL or Swift surveys. These are lower limits since they were selected from pointed observations and are thus highly incomplete.




FIGURE 11: Fraction of obscured AGN ($N_H > 10^{22} \text{ cm}^{-2}$) as a function of redshift for a sample of AGN selected at high energies from INTEGRAL observations, as shown by Malizia et al. [68]. All of these sources have X-ray luminosities lower than $10^{46} \text{ erg s}^{-1}$. The remarkably strong decline in this fraction at $z \sim 0.015$ is likely due to selection effects and not intrinsic to the AGN population. That the fraction of CT AGN in the first bin is $\sim 20\%$, in contrast to the $\sim 5\%$ found overall suggests that current hard X-ray surveys are not sensitive enough to observe CT AGN beyond $z \sim 0.01$. This will improve dramatically with NuSTAR [69].

in Figure 14. As argued by several authors, the fraction of CT AGN in these infrared-excess samples is very high, $>70\%$ [75, 78, 141].

Because sources in the mid-IR excess region are, by definition, very faint at optical wavelengths, it has been very difficult to use optical spectroscopy to measure accurate redshifts. Instead, most surveys have had to rely on (hopefully) accurate photometric redshifts [146, 147]. The distribution of photometric redshifts for the sources in the mid-IR excess region in the ECDF-S is shown in Figure 15; most mid-IR excess sources have $1 < z < 3$. While the majority of these sources are not detected in X-rays, a significant signal is found in X-ray stacks [75, 78, 141]. As shown in Figure 16, the strong stacked detection at $E > 5$ keV clearly indicates the presence of a large number of heavily obscured AGN in this infrared-excess subsample. Specifically, Treister et al. [75] reported that heavily obscured AGN were $\sim 80\text{--}90\%$ of the mid-IR-excess sources in the ECDF-S. A similarly high fraction of $\sim 80\%$ was found in the CDF-S [141] and other fields [78]. Optical spectral fitting of these sources indicates evidence for substantial young stellar populations, younger than 100 Myrs [75]. This suggests that these sources are simultaneously experiencing significant star formation and heavily obscured AGN activity. The best-fit stellar masses for ECDF-S infrared-excess sources range between 10^9 and $10^{12} M_\odot$ with a median stellar mass of $\sim 10^{11} M_\odot$ [75]. Hence, in general these are very massive galaxies.

In order to study in more detail the evolution of the CT AGN space density, in Figure 17 we present the existing measurements of the CT AGN space density as a function




FIGURE 12: Cumulative fractional contribution of CT AGN to the XRB in the 14–195 keV Swift/BAT band as a function of redshift from population synthesis models [35]. As shown by the vertical dashed lines, 50%, 80%, and 90% of the total CT AGN contribution come from sources at $z < 0.7$, 1.4, and 2.0, respectively. Only $\sim 1\%$ of the total XRB intensity comes from CT AGN at $z > 2$. The data point at $z \sim 0$ corresponds to the contribution to the XRB by the CT AGN detected by Swift/BAT, while the data points at high redshift were obtained from the CT AGN in the CDF-S [70]. Solid error bars correspond to transmission-dominated sources only, while the data points with dashed error bars include all the sources in the sample.




FIGURE 13: 4Msec *Chandra* observations of the CDF-S. This is currently the deepest view of the X-ray sky. There are ~ 760 individual sources in ~ 450 arcmin². This is a smooth enhanced image corresponding to the full (0.5–8 keV) X-ray band. Image and data obtained from <http://cxc.harvard.edu/cdo/cdfs.html>.

of redshift. Reasonable agreement, in particular at $z < 1$, is found between both observed values and existing hard X-ray luminosity functions and evolution [79]. However, at $z > 1.7$ and high luminosity, a clear discrepancy is found. Treister et al. [35] concluded that this difference of a factor of 2–3 could be due either to incompleteness in the Swift/BAT and INTEGRAL CT AGN samples at $z = 0$ used to fix the luminosity function normalization (because




FIGURE 14: The ratio of 24-micron to R -band optical flux as a function of $R-K$ color for the sources in the E-CDFS field [75]. Contours show the location of all the Spitzer 24 μm sources. *Red points* show the location of X-ray sources with hard X-ray spectra (hardness ratio $\text{HR} > -0.3$, defined as $\text{HR} = (H - S)/(H + S)$ where S and H are the background-subtracted counts in the soft and hard bands, resp.), while *blue points* have $\text{HR} < -0.3$. *Small black points* show the 193 sources in the IR-red excess region that were not detected individually in X-rays.

reflection-dominated AGN are missed) or to contamination by other types of sources in the observed values at high redshifts. However, after adding the measurements obtained using the infrared-excess sources in the ECDF-S, it appears not only that the systematic difference is still present but perhaps more importantly that there is a strong increase in the number of CT AGN from $z \approx 1.7$ to 2.4. This is not described by any existing luminosity function. It is unlikely that this evolution is due to selection effects, as results from different fully independent surveys and selection techniques are combined in Figure 17, namely, X-ray-selected sources [70], 24 μm -selected sources [75, 78], and a sample of CT AGN found using mid-IR spectroscopy [77]. This result can be interpreted in the context of galaxy evolution models [148], where quasar activity is driven by galaxy mergers and the supermassive black hole is initially completely surrounded by dust, before radiation pressure removes it and a “classical” unobscured quasar is visible [25].

Several groups [149] have found that the fraction of galaxies containing an AGN is a strong function of their IR luminosity. In Figure 18 we present the stacked spectra for the sources in the CDF-S, grouped in bins of IR luminosity [150]. We can see by comparing these spectra that the relative emission at $E \gtrsim 5$ keV, where we expect the AGN emission to dominate even for heavily obscured sources, changes with IR luminosity. In other words, there is a clear trend, with stronger high energy X-ray emission at increasing IR luminosity. The spectra shown in Figure 18 cannot be directly interpreted, as the detector-plus-telescope response information is lost after the conversion to rest-frame energy and stacking. Hence, simulations assuming different intrinsic X-ray spectra have to be used in order to constrain the nature of the sources dominating the coadded signal.




FIGURE 15: Photometric redshift distribution for 90 sources in the ECDF-S in the mid-IR excess region. The *solid histogram* shows the distribution for the sources not detected in X-rays, while the *dashed hatched histogram* considers only the X-ray-detected sources. A KS test shows that it is perfectly likely ($\sim 16\%$) that these two distributions were drawn from the same parent distribution. The *dotted histogram* shows the slightly lower redshift distribution (divided by 1,000) for all the sources with a $24\ \mu\text{m}$ detection and a measured photometric redshift in the ECDF-S.

The observed stacked spectral shape cannot be explained by any plausible starburst spectrum. An AGN component dominating at $E > 5\ \text{keV}$, is required. The average intrinsic rest-frame 2–10 keV AGN luminosity needed to explain the observed spectrum, assuming that every source in the sample contains an AGN of the same luminosity, is $6 \times 10^{42}\ \text{erg s}^{-1}$ for sources with $L_{\text{IR}} > 10^{11} L_{\odot}$, $3 \times 10^{42}\ \text{erg s}^{-1}$ for sources with $L_{\text{IR}} > 5 \times 10^{10} L_{\odot}$, $5 \times 10^{41}\ \text{erg s}^{-1}$ for $5 \times 10^{10} L_{\odot} > L_{\text{IR}} > 10^{10} L_{\odot}$, and $7 \times 10^{41}\ \text{erg s}^{-1}$ for $L_{\text{IR}} > 10^{10} L_{\odot}$. All of these are (intrinsically) very low-luminosity AGN; even if there is a range, it is extremely unlikely to include high-luminosity quasars like those discussed in previous stacking papers. This is not too surprising, actually, because the surveyed volume (even to high redshift) is small, so rare objects like high-luminosity quasars do not appear. If the heavily obscured AGN in these stacked samples have the same median intrinsic luminosity as the X-ray-detected sources with similar IR luminosities, this would indicate that 15% of the galaxies with $L_{\text{IR}} > 10^{11} L_{\odot}$ contain a heavily obscured AGN. This fraction is $\sim 10\%$ in the $L_{\text{IR}} > 5 \times 10^{10} L_{\odot}$ and $5 \times 10^{10} L_{\odot} > L_{\text{IR}} > 10^{10} L_{\odot}$ samples. For sources with $L_{\text{IR}} > 10^{10} L_{\odot}$ this fraction is $< 5\%$. This extra AGN activity (in addition to the X-ray detected sources) can account for $\sim 22\%$ of the total black hole accretion. Adding this to the obscured black hole growth in X-ray-detected AGN [151], we confirm that most of this growth, $\sim 70\%$, is significantly obscured and missed by even the deepest X-ray surveys [25, 91].




FIGURE 16: Stacked background-subtracted Chandra counts as a function of rest-frame energy, for sources with $f_{24}/f_R > 1000$ and $R-K > 4.5$ in the 4 Msec CDF-S field (*filled circles*; [75]). The *cyan dashed lines (stars)* show the simulated spectra for the high-mass X-ray binary (HMXB) population normalized using the relation between star-formation rate and X-ray luminosity [76]. The *blue dashed lines (open squares)* show simulated thermal spectra corresponding to a black body with $kT = 0.7\ \text{keV}$. An absorbed AGN spectrum, given by a power law with $\Gamma = 1.9$ and a fixed $N_H = 10^{24}\ \text{cm}^{-2}$, is shown by the *red dashed lines (open circles)*. In addition, a scattered AGN component, characterized by a 1% reflection of the underlying unobscured power law, is shown by the *green dashed lines (open triangles)*. The resulting summed spectrum (*black solid lines*) is in very good agreement with the observed counts. The strong detection in the stacked spectrum at $E > 5\ \text{keV}$ confirms the presence of a significant number of heavily obscured AGN in these IR-excess objects [75].

4.3. *Obscured AGN at High Redshifts, $z > 3$.* As mentioned in Section 3, most measurements of black hole accretion at high redshift, $z > 3$, come from optical observations of unobscured sources. This is not only because obscured sources are obviously fainter at most wavelengths, but also because large areas have to be covered in order to survey a significant volume at high redshifts. As can be seen in Figure 19, there is a clear decline in the number of luminous quasars at $z > 2$, although the decline is shallower for X-rays compared to optical surveys, since X-ray selection is less biased [79, 85]. However, it is important to point out that: (i) these results are limited to the highest luminosity sources, $\log l_x > 44.5\ \text{erg s}^{-1}$, which do not represent the average AGN and do not contribute much to the extragalactic XRB [35] and (ii) only relatively unobscured sources are included. In particular, heavily obscured, Compton-thick, AGN are systematically underrepresented in these surveys. As we will describe, these missing populations can have a significant impact in our understanding of cosmic supermassive black hole growth.

In order to search for the presence of growing supermassive black holes in young galaxies, Treister et al. [152] stacked X-ray images of $z > 6$ galaxy candidates selected based on the optical and near-IR dropout techniques, selected from a




FIGURE 17: Space density of Compton thick AGN as a function of redshift, as published by Treister et al. [75]. *Filled triangles* show the ECDF-S results [75]. *Squares*: X-ray-selected sources in the CDF-S [70]. *Star*: Measurement obtained using mid-IR spectroscopy [77]. *Pentagons*: Values obtained using mid-IR excess sources in COSMOS [78]. *Solid lines* show the expected space density of Compton thick AGN from the luminosity function of Yencho et al. [79], with the overall normalization fixed to the results of the INTEGRAL and Swift/BAT surveys [35], while the *dashed lines* show the expectations based on the luminosity function of Della Ceca et al. [80]. *Red symbols* show measurements and expectations for $L_X > 10^{43}$ erg s $^{-1}$ sources, while the *blue symbols* are for $L_X > 10^{44}$ erg s $^{-1}$. While for the lower luminosity sources a good agreement is found between observations and expectations, higher luminosity sources at $z > 1.8$ lie well above the luminosity function.

sample of 197 galaxies, 151 in the CDF-S and 46 in the CDF-N from Bouwens et al. [153]. Using the 4 Msec Chandra observations of the CDF-S, and the 2 Msec data available on the CDF-N, this corresponds to a total exposure time of $\sim 7 \times 10^8$ seconds (~ 23 years). Significant detections in both the soft and hard X-ray bands were obtained. However, these detections have recently been questioned (after the submission of this paper) by several authors [154–156] due to a possible bias in the background subtraction technique used by Treister et al. While a full discussion is clearly beyond the scope of this paper, we note that a full analysis, using an optimal weighting scheme (as in [152]), as well as considering the effects of faint, undetected, sources in the background, remains to be done. At a minimum, the results presented below can be considered as upper limits.

The corresponding average rest-frame 2–10 keV luminosity, derived from the observed-frame hard band, is 6.8×10^{42} erg s $^{-1}$. Since none of these sources were individually detected in X-rays, at least 30% of the galaxies in this sample likely contain an AGN [152]. Furthermore, there is a factor ~ 9 difference between the fluxes measured in the observed-frame soft and hard bands. The only explanation for this relatively large flux ratio in the hard-to-soft bands is

very high levels of obscuration. As can be seen in Figure 20, at $z \sim 6$, a minimum column density of $N_H \simeq 10^{24}$ cm $^{-2}$, that is, Compton thick obscuration, is required. Given that this ratio is observed in a stacked X-ray spectrum, this implies that there are very few sources with significantly lower levels of obscuration, which in turn means that these sources must be nearly Compton-thick along most directions ($\sim 4\pi$ obscuration). Similar sources have also been observed in the local universe [157] but appear to be rare. Furthermore, for $z \lesssim 3$ we know that the fraction of obscured AGN increases with decreasing luminosity [92, 104, 117] and increasing redshift [31, 111]. Hence, it is not entirely surprising that the sources studied here, given their low luminosities and high redshifts, are heavily obscured. In fact, the discovery of a Compton-thick AGN at $z \sim 5$ selected using the dropout technique has been recently reported [158].

This relatively large number of growing supermassive black holes at $z \gtrsim 6$ is contrary to the picture obtained from optical observations of high-luminosity quasars at similar distances, which are much rarer [21, 88]. This is particularly important in our understanding of the early hydrogen reionization, which can be either due to young stars and/or growing supermassive black holes [159]. The results presented by Treister et al. [152] show that while growing supermassive black holes do not contribute much to hydrogen reionization, this is not because their numbers drop steeply at $z > 4$ as previously suggested [160], but because large amounts of obscuration found in these sources imply that UV and soft X-rays do not escape.

5. The Cosmic History of Black Hole Accretion

Direct black hole mass measurements, either through stellar or gas dynamics, are available for only a few nearby galaxies. However, thanks to the tight correlation between mass of the supermassive black hole and other properties such as velocity dispersion and others, it has been possible to estimate the black hole mass function at $z \simeq 0$ [161–164]. This is commonly done starting from the observed galaxy luminosity or velocity function and assuming either a constant black hole to stellar mass ratio [161] or the M - σ relation [163]. Both the overall shape of the black hole mass function and the integrated black hole mass density, which can only be computed at $z \simeq 0$, can be used to infer properties of the AGN population. This was first used in the so-called “Soltan’s argument” [24], which says that the intrinsic bolometric AGN luminosity, L , is directly linked to the amount of mass accreted by the black hole, \dot{M}_{acc} :

$$L = \epsilon \dot{M}_{\text{acc}} c^2, \quad (1)$$

where ϵ is the accretion efficiency and c is the speed of light. A typical value assumed for the efficiency is $\sim 10\%$ [24, 163].

Recent comparisons of the black hole mass function to the distribution inferred from the observed AGN luminosity indicate that the average efficiency is 8%, the Eddington ratio is $\sim 50\%$, and the average lifetime of the visible AGN phase is $\sim 10^8$ years [163, 164]. By studying the black hole mass distribution at the high mass end, $M > 10^9 M_\odot$, Natarajan




FIGURE 18: Stacked background-subtracted Chandra counts as a function of rest-frame energy for sources with low (a) and high (b) IR luminosity (black filled circles). The cyan dashed lines (stars), blue dashed lines (open squares), red dashed lines (open circles) and green dashed lines (open triangles) show the same model components as in Figure 16. The resulting summed spectra (black solid lines) are in very good agreement with the observed counts. The strong detection in the stacked spectrum at $E > 5$ keV, in particular at the higher IR luminosities, confirms the presence of a significant number of heavily obscured AGN in these samples.

and Treister [165] found that the observed number of ultra-massive black holes is significantly lower than the number density inferred from the AGN hard X-ray luminosity function. They concluded that this is evidence for an upper limit to the black hole mass, which can be explained by the presence of a self-regulation mechanism.

The observed black hole mass density at $z \approx 0$, obtained by integrating the black hole mass function, ranges from 2.9×10^5 [162] to $4.6^{+1.9}_{-1.4} \times 10^5 M_{\odot} \text{Mpc}^{-3}$ [163]; most recently, Shankar et al. [166] found $3.2\text{--}5.4 \times 10^5 M_{\odot} \text{Mpc}^{-3}$. For comparison, integrating the AGN hard X-ray LF, including the number of Compton-thick AGN constrained by INTEGRAL and Swift/BAT observations, Treister et al. [35] obtained a value of $4.5 \times 10^5 M_{\odot} \text{Mpc}^{-3}$, perfectly consistent with the observed value, indicating that at least locally X-ray-detected AGN can account for most or all of the black hole growth.

The black hole mass function can be measured observationally for unobscured, high-luminosity AGN at higher redshift, taking advantage of the known correlation between black hole mass and observational quantities such as luminosity and emission line width [167]. These correlations are calibrated using more direct black hole measurements, available for a few, mostly local, sources [8, 9]. The large number of unobscured quasars with optical spectroscopy provided by the SDSS and other optical surveys has been very

useful in determining the black hole mass function up to high redshifts, [168–170], and for lower luminosity sources using deeper surveys such as the AGN and Galaxy Evolution Survey (AGES; [171]) and COSMOS [172]. Using these black hole mass functions as a function of redshift as constraints, recently Natarajan [173] concluded that the observational data are inconsistent with the hypothesis that these black holes are created as the remnants of population III stars. Instead, they argue that massive seeding models are required [174].

While a clear picture of the history of black hole growth is emerging, significant uncertainties still remain. In particular, while the spectral shape and intensity of the extragalactic X-ray background have been used to constrain the AGN population, the number of heavily obscured accreting super-massive black holes beyond $z \sim 1$ is not properly bounded. Infrared and deep X-ray selection methods have been useful in that sense, but have not provided a final answer, due to confusion with star-forming galaxies in the infrared and the effects of obscuration in X-rays. At higher redshifts, the situation is even more unclear, and only a few, very rare, high luminosity quasars are known. Unless high-redshift AGN luminosity functions are pathological, these extreme sources do not represent the typical growing black holes in the early universe. As a consequence, and in spite of recent advances




FIGURE 19: Number density of X-ray-selected AGN as a function of redshift for sources with $\log l_{2-10} > 44.5$ erg s $^{-1}$, as published by Brusa et al. [83]. Lines show AGN luminosity functions [39, 79, 84–86] while individual measurements are shown by green circles [87] and black squares [86]. The red star at $z = 6$ was obtained from the optical luminosity function assuming no evolution in α_{ox} . The black-shaded area shows the observations of optically bright SDSS QSOs [88].

[152, 173], the formation mechanism for the first black holes in the universe is still unknown.

6. Prospects

Scheduled for launch in February 2012, NuSTAR will be the first focusing high-energy ($E = 5\text{--}80$ keV) X-ray mission, reaching flux limits ~ 100 times fainter than *INTEGRAL* or *Swift*/BAT observations and comparable to *Chandra* and *XMM-Newton* at lower energies. During the first two years of operations, NuSTAR will likely observe, as part of the guaranteed time program, two extragalactic fields: the ECDF-S and the central 1 deg 2 part of COSMOS, for a total of 3.1 Msec each. These deep high-energy observations will enable to obtain a nearly complete AGN survey, including heavily obscured Compton-thick sources, up to $z \sim 1.5$ [175]. A similar mission, ASTRO-H [176], will be launched by Japan in 2014. Both missions will provide angular resolutions. $\lesssim 1'$, in combination with observations at longer wavelengths will allow for the detection and identification of most growing supermassive black holes at $z \sim 1$.

There is little doubt that the Atacama Large Millimeter Array (ALMA) will revolutionize our understanding of galaxy evolution. Sources of mm and submm emission traced by ALMA include thermal emission of the warm/cold dust, which traces star formation, synchrotron radiation associated with relativistic particles, and free-free radiation from HII regions. In particular, CO rotational transition lines have been used to trace the spatial distribution, kinematics, temperature, and mass of the molecular gas [177]. The sensitivity




FIGURE 20: Expected ratio of the observed-frame hard-to-soft flux as a function of obscuring neutral Hydrogen column density (N_H). The black solid line was derived assuming an intrinsic power law spectrum with slope $\Gamma = 1.9$ and photoelectric absorption. The gray zone shows the measured ratio for the stack of galaxies at $z \approx 6$ and the ± 1 standard deviation limits. A column density of $N_H \approx 10^{24}$ cm $^{-2}$, that is, Compton-thick obscuration, is required to explain the observed hard-to-soft X-ray flux ratio [152].

of ALMA will allow for the detection of luminous IR galaxies ($L_{\text{IR}} > 10^{11} L_{\odot}$), which have been found to often host a heavily obscured AGN [150], up to $z \sim 10$. Furthermore, with ALMA it will be possible to study separately the molecular dust surrounding the central black hole and those in star-forming regions in the host galaxy. Due to their limited sensitivity and relatively bad angular resolution, currently available mm/sub-mm telescopes are not ideal to study star-forming regions even in nearby galaxies. This will dramatically change thanks to ALMA, which will have orders of magnitude better sensitivity and HST-like angular resolution. The first call for ALMA observations was released on March 31, 2011 for observations starting on September 30, 2011. It is expected that the complete array will be in full operation in 2013. The superb spatial resolution and sensitivity of ALMA will allow uniquely identify of the optical/near-IR counterpart of the mm-submm sources. Furthermore, ALMA will directly provide the redshift of the mm-submm sources through the detection of CO rotational transition lines, up to very high redshifts. Combining these new data with existing multiwavelength information will finally allow us to complete the census of supermassive black hole growth since the epoch of cosmic reionization.

Acknowledgments

Support for the work of E. Treiser was provided by the National Aeronautics and Space Administration through

Chandra/Einstein Postdoctoral Fellowship Award number PF8-90055, issued by the Chandra X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory (SAO) for and on behalf of the National Aeronautics Space Administration under Contract no. NAS8-03060. Additional support for this work was provided by NASA through Chandra Award SP1-12005X. E. Treister received partial support from Center of Excellence in Astrophysics and Associated Technologies (PFB 06). C. M. Urry acknowledges support from NSF Grants AST-0407295, AST-0449678, AST-0807570, and Yale University.

References

- [1] J. A. Orosz, "Inventory of black hole binaries," in *A Massive Star Odyssey: From Main Sequence to Supernova*, K. van der Hucht, A. Herrero, and C. Esteban, Eds., vol. 212 of *IAU Symposium*, p. 365, 2003.
- [2] D. Lynden-Bell, "Galactic nuclei as collapsed old quasars," *Nature*, vol. 223, no. 5207, pp. 690–694, 1969.
- [3] T. R. Lauer, S. M. Faber, D. Richstone et al., "The masses of nuclear black holes in luminous elliptical galaxies and implications for the space density of the most massive black holes," *Astrophysical Journal*, vol. 662, no. 2, pp. 808–834, 2007.
- [4] R. Genzel, A. Eckart, T. Ott, and F. Eisenhauer, "On the nature of the dark mass in the centre of the Milky Way," *Monthly Notices of the Royal Astronomical Society*, vol. 291, no. 1, pp. 219–234, 1997.
- [5] A. M. Ghez, B. L. Klein, M. Morris, and E. E. Becklin, "High proper-motion stars in the vicinity of sagittarius A*: evidence for a supermassive black hole at the center of our galaxy," *Astrophysical Journal*, vol. 509, no. 2, pp. 678–686, 1998.
- [6] J. Kormendy and D. Richstone, "Inward bound—the search for supermassive black holes in galactic nuclei," *Annual Review of Astronomy and Astrophysics*, vol. 33, no. 1, pp. 581–624, 1995.
- [7] J. N. Bahcall, B.-Z. Kozlovsky, and E. E. Salpeter, "On the time dependence of emission-line strengths from a photoionized nebula," *Astrophysical Journal*, vol. 171, p. 467, 1972.
- [8] R. D. Blandford and C. F. McKee, "Reverberation mapping of the emission line regions of Seyfert galaxies and quasars," *Astrophysical Journal*, vol. 255, pp. 419–439, 1982.
- [9] B. M. Peterson, "Reverberation mapping of active galactic nuclei," *Astronomical Society of the Pacific*, vol. 105, pp. 247–268, 1993.
- [10] A. Marconi and L. K. Hunt, "The relation between black hole mass, bulge mass, and near-Infrared luminosity," *Astrophysical Journal*, vol. 589, no. 1, pp. L21–L24, 2003.
- [11] J. Magorrian, S. Tremaine, D. Richstone et al., "The demography of massive dark objects in galaxy centers," *Astronomical Journal*, vol. 115, no. 6, pp. 2285–2305, 1998.
- [12] L. Ferrarese and D. Merritt, "A fundamental relation between supermassive black holes and their host galaxies," *Astrophysical Journal*, vol. 539, no. 1, pp. L9–L12, 2000.
- [13] K. Gebhardt, R. Bender, G. Bower et al., "A relationship between nuclear black hole mass and galaxy velocity dispersion," *Astrophysical Journal*, vol. 539, no. 1, pp. L13–L16, 2000.
- [14] L. Ferrarese, "Beyond the bulge: a fundamental relation between supermassive black holes and dark matter halos," *Astrophysical Journal*, vol. 578, no. 1, pp. 90–97, 2002.
- [15] V. Springel, T. Di Matteo, and L. Hernquist, "Modelling feedback from stars and black holes in galaxy mergers," *Monthly Notices of the Royal Astronomical Society*, vol. 361, no. 3, pp. 776–794, 2005.
- [16] P. F. Hopkins, L. Hernquist, T. J. Cox, T. Di Matteo, B. Robertson, and V. Springel, "A unified, merger-driven model of the origin of starbursts, quasars, the cosmic X-ray background, supermassive black holes, and galaxy spheroids," *Astrophysical Journal Supplement Series*, vol. 163, no. 1, pp. 1–49, 2006.
- [17] N. Menci, A. Fontana, E. Giallongo, A. Grazian, and S. Salimbeni, "The abundance of distant and extremely red galaxies: the role of AGN feedback in hierarchical models," *Astrophysical Journal*, vol. 647, no. 2, pp. 753–762, 2006.
- [18] D. J. Croton, V. Springel, S. D. M. White et al., "The many lives of active galactic nuclei: cooling flows, black holes and the luminosities and colours of galaxies," *Monthly Notices of the Royal Astronomical Society*, vol. 365, no. 1, pp. 11–28, 2006.
- [19] M. J. Rees, "Structure and properties of nearby galaxies," in *Proceedings of the Symposium*, vol. 1, pp. 237–244, Bad Muenstereifel, Germany, 1978.
- [20] M. Volonteri, "Formation of supermassive black holes," *Astronomy and Astrophysics Review*, vol. 18, no. 3, pp. 279–315, 2010.
- [21] C. J. Willott, L. Albert, D. Arzoumanian et al., "Eddington-limited accretion and the black hole mass function at redshift 6," *Astronomical Journal*, vol. 140, no. 2, pp. 546–560, 2010.
- [22] V. Bromm and A. Loeb, "Formation of the first supermassive black holes," *Astrophysical Journal*, vol. 596, no. 1, pp. 34–46, 2003.
- [23] B. Devecchi and M. Volonteri, "Formation of the first nuclear clusters and massive black holes at high redshift," *Astrophysical Journal*, vol. 694, no. 1, pp. 302–313, 2009.
- [24] A. Soltan, "Masses of quasars," *Monthly Notices of the Royal Astronomical Society*, vol. 200, pp. 115–122, 1982.
- [25] E. Treister, P. Natarajan, D. B. Sanders, C. Megan Urry, K. Schawinski, and J. Kartaltepe, "Major galaxy mergers and the growth of supermassive black holes in quasars," *Science*, vol. 328, no. 5978, pp. 600–602, 2010.
- [26] D. B. Sanders, B. T. Soifer, J. H. Elias et al., "Ultraluminous infrared galaxies and the origin of quasars," *Astrophysical Journal*, vol. 325, pp. 74–91, 1988.
- [27] P. F. Hopkins and L. Hernquist, "Fueling low-level AGN activity through stochastic accretion of cold gas," *Astrophysical Journal*, vol. 166, no. 1, pp. 1–36, 2006.
- [28] R. Antonucci, "Unified models for active galactic nuclei and quasars," *Annual Review of Astronomy and Astrophysics*, vol. 31, no. 1, pp. 473–521, 1993.
- [29] C. M. Urry and P. Padovani, "Unified schemes for radio-loud active galactic nuclei," *Astronomical Society of the Pacific*, vol. 107, p. 803, 1995.
- [30] A. Lawrence, "The relative frequency of broad-lined and narrow-lined active galactic nuclei—implications for unified schemes," *Monthly Notices of the Royal Astronomical Society*, vol. 252, pp. 586–592, 1991.
- [31] E. Treister and C. M. Urry, "The evolution of obscuration in active galactic nuclei," *Astrophysical Journal*, vol. 652, no. 2, pp. L79–L82, 2006.
- [32] R. C. Hickox and M. Markevitch, "Absolute measurement of the unresolved cosmic X-ray background in the 0.5–8 keV band with Chandra," *Astrophysical Journal*, vol. 645, no. 1, pp. 95–114, 2006.

- [33] R. F. Mushotzky, L. L. Cowie, A. J. Barger, and K. A. Arnaud, "Resolving the extragalactic hard X-ray background," *Nature*, vol. 404, no. 6777, pp. 459–464, 2000.
- [34] G. Setti and L. Woltjer, "Active galactic nuclei and the spectrum of the X-ray background," *Astronomy and Astrophysics*, vol. 224, pp. L21–L23, 1989.
- [35] E. Treister, C. M. Urry, and S. Virani, "The space density of compton thick AGN and the X-ray background," *Astrophysical Journal*, vol. 696, pp. 110–120, 2009.
- [36] A. R. Draper and D. R. Ballantyne, "Balancing the cosmic energy budget: the cosmic X-ray background, blazars, and the compton thick active galactic nucleus fraction," *Astrophysical Journal*, vol. 707, no. 1, pp. 778–786, 2009.
- [37] D. Burlon, M. Ajello, J. Greiner, A. Comastri, A. Merloni, and N. Gehrels, "Three-year swift-BAT survey of active galactic nuclei: reconciling theory and observations?" *Astrophysical Journal*, vol. 728, p. 58, 2011.
- [38] E. Treister and C. Megan Urry, "Active galactic nuclei unification and the X-ray background," *Astrophysical Journal*, vol. 630, no. 1, pp. 115–121, 2005.
- [39] R. Gilli, A. Comastri, and G. Hasinger, "The synthesis of the cosmic X-ray background in the Chandra and XMM-Newton era," *Astronomy and Astrophysics*, vol. 463, no. 1, pp. 79–96, 2007.
- [40] G. Hinshaw, J. L. Weiland, R. S. Hill et al., "Five-year wilkinson microwave anisotropy probe observations: data processing, sky maps and basic results," *Astrophysical Journal Supplement Series*, vol. 180, no. 2, pp. 225–245, 2009.
- [41] J. C. Manners, *Obscuration and X-ray variability of active galactic nuclei*, Ph.D. thesis, University of Edinburgh, 2002.
- [42] J. J. Condon, Q. F. Yin, T. X. Thuan, and T. Boller, "The ROSAT/IRAS galaxy sample revisited," *Astronomical Journal*, vol. 116, no. 6, pp. 2682–2716, 1998.
- [43] G. A. Shields, "Thermal continuum from accretion disks in quasars," *Nature*, vol. 272, no. 5655, pp. 706–708, 1978.
- [44] M. A. Malkan and W. L. W. Sargent, "The ultraviolet excess of Seyfert 1 galaxies and quasars," *Astrophysical Journal*, vol. 254, pp. 22–37, 1982.
- [45] M. Schmidt and R. F. Green, "Quasar evolution derived from the Palomar bright quasar survey and other complete quasar surveys," *Astrophysical Journal*, vol. 269, pp. 352–374, 1983.
- [46] B. J. Boyle, T. Shanks, S. M. Croom et al., "The 2dF QSO Redshift Survey-I. The optical luminosity function of quasar-stellar objects," *Monthly Notices of the Royal Astronomical Society*, vol. 317, no. 4, pp. 1014–1022, 2000.
- [47] G. T. Richards, X. Fan, D. P. Schneider et al., "Colors of 2625 quasars at $0 < z < 5$ measured in the sloan digital sky survey photometric system," *Astronomical Journal*, vol. 121, no. 5, pp. 2308–2330, 2001.
- [48] G. T. Richards, A. D. Myers, A. G. Gray et al., "Efficient photometric selection of quasars from the sloan digital sky survey. II. $\sim 1,000,000$ quasars from data release 6," *Astrophysical Journal Supplement Series*, vol. 180, no. 1, pp. 67–83, 2009.
- [49] D. E. Vanden Berk, G. T. Richards, A. Bauer et al., "Composite quasar spectra from the sloan digital sky survey," *Astronomical Journal*, vol. 122, no. 2, pp. 549–564, 2001.
- [50] L. J. Kewley, B. Groves, G. Kauffmann, and T. Heckman, "The host galaxies and classification of active galactic nuclei," *Monthly Notices of the Royal Astronomical Society*, vol. 372, no. 3, pp. 961–976, 2006.
- [51] G. Kauffmann, T. M. Heckman, C. Tremonti et al., "The host galaxies of active galactic nuclei," *Monthly Notices of the Royal Astronomical Society*, vol. 346, no. 4, pp. 1055–1077, 2003.
- [52] L. J. Kewley, M. A. Dopita, R. S. Sutherland, C. A. Heisler, and J. Trevena, "Theoretical modeling of starburst galaxies," *Astrophysical Journal*, vol. 556, no. 1, pp. 121–140, 2001.
- [53] S. M. Croom, G. T. Richards, T. Shanks et al., "The 2dF-SDSS LRG and QSO survey: the QSO luminosity function at $0.4 < z < 2.6$," *Monthly Notices of the Royal Astronomical Society*, vol. 399, no. 4, pp. 1755–1772, 2009.
- [54] S. M. Croom, R. J. Smith, B. J. Boyle et al., "The 2dF QSO Redshift Survey-XII. The spectroscopic catalogue and luminosity function," *Monthly Notices of the Royal Astronomical Society*, vol. 349, no. 4, pp. 1397–1418, 2004.
- [55] G. T. Richards, S. M. Croom, S. F. Anderson et al., "The 2dF-SDSS LRG and QSO (2SLAQ) Survey: the $z < 2.1$ quasar luminosity function from 5645 quasars to $g = 21.85$," *Monthly Notices of the Royal Astronomical Society*, vol. 360, no. 3, pp. 839–852, 2005.
- [56] G. T. Richards, M. A. Strauss, X. Fan et al., "The sloan digital sky survey quasar survey: Quasar luminosity function from data release 3," *Astronomical Journal*, vol. 131, no. 6, pp. 2766–2787, 2006.
- [57] E. Glikman, S. G. Djorgovski, D. Stern, A. Dey, B. T. Jannuzi, and K.-S. Lee, "The faint end of the Quasar luminosity function at $z \sim 4$: implications for ionization of the intergalactic medium and cosmic downsizing," *Astrophysical Journal*, vol. 728, p. L26, 2011.
- [58] E. Glikman, M. Bogosavljević, S. G. Djorgovski et al., "The faint end of the quasar luminosity function at $z \sim 4$," *Astrophysical Journal*, vol. 710, no. 2, pp. 1498–1514, 2010.
- [59] H. Ikeda, T. Nagao, K. Matsuoka et al., "Probing the faint end of the Quasar luminosity function at $z \sim 4$ in the COSMOS field," *Astrophysical Journal*, vol. 728, p. L25, 2011.
- [60] B. Siana, M. Del Carmen Polletta, H. E. Smith et al., "High-redshift QSOs in the swire survey and the $z \sim 3$ QSO luminosity function," *Astrophysical Journal*, vol. 675, no. 1, pp. 49–70, 2008.
- [61] D. E. Gruber, J. L. Matteson, L. E. Peterson, and G. V. Jung, "The spectrum of diffuse cosmic hard X-rays measured with HEAO 1," *Astrophysical Journal*, vol. 520, no. 1, pp. 124–129, 1999.
- [62] A. De Luca and S. Molendi, "The 2–8 keV cosmic X-ray background spectrum as observed with XMM-Newton," *Astronomy and Astrophysics*, vol. 419, no. 3, pp. 837–848, 2004.
- [63] E. Churazov, R. Sunyaev, M. Revnivtsev et al., "INTEGRAL observations of the cosmic X-ray background in the 5–100 keV range via occultation by the Earth," *Astronomy and Astrophysics*, vol. 467, no. 2, pp. 529–540, 2007.
- [64] M. Ajello, J. Greiner, G. Sato et al., "Cosmic X-Ray background and Earth albedo spectra with swift bat," *Astrophysical Journal*, vol. 689, no. 2, pp. 666–677, 2008.
- [65] S. F. Hönig and T. Beckert, "Active galactic nuclei dust tori at low and high luminosities," *Monthly Notices of the Royal Astronomical Society*, vol. 380, no. 3, pp. 1172–1176, 2007.
- [66] J. Tueller, R. F. Mushotzky, S. Barthelmy et al., "Swift bat survey of AGNs," *Astrophysical Journal*, vol. 681, no. 1, pp. 113–127, 2008.
- [67] R. Krivonos, M. Revnivtsev, A. Lutovinov, S. Sazonov, E. Churazov, and R. Sunyaev, "INTEGRAL/IBIS all-sky survey in hard X-rays," *Astronomy and Astrophysics*, vol. 475, no. 2, pp. 775–784, 2007.
- [68] A. Malizia, J. B. Stephen, L. Bassani, A. J. Bird, F. Panessa, and P. Ubertini, "The fraction of Compton-thick sources in an INTEGRAL complete AGN sample," *Monthly Notices of the Royal Astronomical Society*, vol. 399, pp. 944–951, 2009.

- [69] F. A. Harrison et al., “The nuclear spectroscopic telescope array (NuSTAR),” in *Proceedings of the Space Telescopes and Instrumentation: Ultraviolet to Gamma Ray*, vol. 7732 of *Proceedings of SPIE*, July 2010.
- [70] P. Tozzi, R. Gilli, V. Mainieri et al., “X-ray spectral properties of active galactic nuclei in the Chandra Deep Field South,” *Astronomy and Astrophysics*, vol. 451, no. 2, pp. 457–474, 2006.
- [71] M. Schmidt, “3C 273: a star-like object with large red-shift,” *Nature*, vol. 197, no. 4872, p. 1040, 1963.
- [72] K. I. Kellermann, R. Sramek, M. Schmidt, D. B. Shaffer, and R. Green, “VLA observations of objects in the Palomar Bright Quasar Survey,” *Astronomical Journal*, vol. 98, pp. 1195–1207, 1989.
- [73] M. C. Begelman, R. D. Blandford, and M. J. Rees, “Theory of extragalactic radio sources,” *Reviews of Modern Physics*, vol. 56, no. 2, pp. 255–351, 1984.
- [74] D. B. Sanders, E. S. Phinney, G. Neugebauer, B. T. Soifer, and K. Matthews, “Continuum energy distribution of quasars—shapes and origins,” *Astrophysical Journal*, vol. 347, pp. 29–51, 1989.
- [75] E. Treister, C. N. Cardamone, K. Schawinski et al., “Heavily obscured agn in star-forming galaxies at $z \approx 2$,” *Astrophysical Journal*, vol. 706, no. 1, pp. 535–552, 2009.
- [76] P. Ranalli, A. Comastri, and G. Setti, “The 2–10 keV luminosity as a star formation rate indicator,” *Astronomy and Astrophysics*, vol. 399, no. 1, pp. 39–50, 2003.
- [77] D. M. Alexander, R. R. Chary, A. Pope et al., “Reliable identification of compton-thick quasars at $z \approx 2$: spitzer mid-infrared spectroscopy of HDF-oMD49,” *Astrophysical Journal*, vol. 687, no. 2, pp. 835–847, 2008.
- [78] F. Fiore et al., “Chasing highly obscured QSOs in the COSMOS field,” *The Astrophysical Journal*, vol. 693, pp. 447–462, 2009.
- [79] B. Yencho, A. J. Barger, L. Trouille, and L. M. Winter, “The optx project. II. hard X-ray luminosity functions of active galactic nuclei for $z \lesssim 5$,” *Astrophysical Journal*, vol. 698, no. 1, pp. 380–396, 2009.
- [80] R. Della Ceca, A. Caccianiga, P. Severgnini et al., “The cosmological properties of AGN in the XMM-Newton Hard Bright Survey,” *Astronomy and Astrophysics*, vol. 487, no. 1, pp. 119–130, 2008.
- [81] J. A. Baldwin, M. M. Phillips, and R. Terlevich, “Classification parameters for the emission-line spectra of extragalactic objects,” *Astronomical Society of the Pacific*, vol. 93, pp. 5–19, 1981.
- [82] A. D. Montero-Dorta, D. J. Croton, R. Yan et al., “The DEEP2 Galaxy Redshift Survey: the red sequence AGN fraction and its environment and redshift dependence,” *Monthly Notices of the Royal Astronomical Society*, vol. 392, no. 1, pp. 125–134, 2009.
- [83] M. Brusa, R. Gilli, F. Civano, A. Comastri, R. Fiore, and C. Vignali, “Identification of (high-redshift) AGN with WFXT: lessons from COSMOS and CDFS,” *Memorie della Societa Astronomica Italiana Supplementi*, vol. 17, p. 106, 2011.
- [84] J. D. Silverman, P. J. Green, W. A. Barkhouse et al., “The luminosity function of X-ray-selected active galactic nuclei: evolution of supermassive black holes at high redshift,” *Astrophysical Journal*, vol. 679, no. 1, pp. 118–139, 2008.
- [85] J. Ebrero, F. J. Carrera, M. J. Page et al., “The XMM-Newton serendipitous survey VI. The X-ray luminosity function,” *Astronomy and Astrophysics*, vol. 493, no. 1, pp. 55–69, 2009.
- [86] J. Aird, K. Nandra, E. S. Laird et al., “The evolution of the hard X-ray luminosity function of AGN,” *Monthly Notices of the Royal Astronomical Society*, vol. 401, pp. 2531–2551, 2010.
- [87] M. Brusa, A. Comastri, R. Gilli et al., “High-redshift quasars in the COSMOS survey: the space density of $z > 3$ X-ray selected QSOs,” *Astrophysical Journal*, vol. 693, pp. 8–22, 2009.
- [88] X. Fan, C. L. Carilli, and B. Keating, “Observational constraints on cosmic reionization,” *Annual Review of Astronomy and Astrophysics*, vol. 44, pp. 415–462, 2006.
- [89] M. Elvis, T. Maccararo, A. S. Wilson et al., “Seyfert galaxies as X-ray sources,” *Monthly Notices of the Royal Astronomical Society*, vol. 183, pp. 129–157, 1978.
- [90] F. E. Bauer, D. M. Alexander, W. N. Brandt et al., “The fall of active galactic nuclei and the rise of star-forming galaxies: a close look at the Chandra Deep Field X-ray number counts,” *Astronomical Journal*, vol. 128, no. 5, pp. 2048–2065, 2004.
- [91] E. Treister, C. Megan Urry, E. Chatzichristou et al., “Obscured active galactic nuclei and the X-RAY, optical, and far-infrared number counts of active galactic nuclei in the goods fields,” *Astrophysical Journal*, vol. 616, no. 1, pp. 123–135, 2004.
- [92] S. Sazonov, M. Revnivtsev, R. Krivonos, E. Churazov, and R. Sunyaev, “Hard X-ray luminosity function and absorption distribution of nearby AGN: INTEGRAL all-sky survey,” *Astronomy and Astrophysics*, vol. 462, no. 1, pp. 57–66, 2007.
- [93] G. T. Richards, M. Lacy, L. J. Storrie-Lombardi et al., “Spectral energy distributions and multiwavelength selection of type 1 quasars,” *Astrophysical Journal Supplement Series*, vol. 166, no. 2, pp. 470–497, 2006.
- [94] M. Haas, U. Klaas, S. A. H. Müller et al., “The ISO view of Palomar-Green quasars,” *Astronomy and Astrophysics*, vol. 402, no. 1, pp. 87–111, 2003.
- [95] E. Treister, C. M. Urry, J. Van Dуйne et al., “Spitzer number counts of active galactic nuclei in the goods fields,” *Astrophysical Journal*, vol. 640, no. 2, pp. 603–611, 2006.
- [96] E. Glikman, D. J. Helfand, and R. L. White, “A near-infrared spectral template for quasars,” *Astrophysical Journal*, vol. 640, no. 2, pp. 579–591, 2006.
- [97] M. Nenkova, Z. Ivezić, and M. Elitzur, “Dust emission from active galactic nuclei,” *Astrophysical Journal*, vol. 570, no. 1, pp. L9–L12, 2002.
- [98] E. Treister, J. H. Krolik, and C. Dullemond, “Measuring the fraction of obscured quasars by the infrared luminosity of unobscured quasars,” *Astrophysical Journal*, vol. 679, no. 1, pp. 140–148, 2008.
- [99] C. N. Cardamone, C. M. Urry, M. Damen et al., “Mid-infrared properties and color selection for X-ray-detected active galactic nuclei in the MUSYC Extended Chandra Deep Field-South,” *Astrophysical Journal*, vol. 680, no. 1, pp. 130–142, 2008.
- [100] M. Schmidt, “Space distribution and luminosity functions of Quasi-Stellar radio sources,” *Astrophysical Journal*, vol. 151, p. 393, 1968.
- [101] H. L. Marshall, “The evolution of optically selected quasars with Z less than 2.2 and B less than 20,” *Astrophysical Journal*, vol. 299, pp. 109–121, 1985.
- [102] G. Mathez, “Evolution of the luminosity function of quasars—a model with constant density and luminosity evolution,” *Astronomy & Astrophysics*, vol. 53, pp. 15–21, 1976.
- [103] G. Hasinger, T. Miyaji, and M. Schmidt, “Luminosity-dependent evolution of soft X-ray selected AGN: new Chandra and XMM-Newton surveys,” *Astronomy and Astrophysics*, vol. 441, no. 2, pp. 417–434, 2005.
- [104] Y. Ueda, M. Akiyama, K. Ohta, and T. Miyaji, “Cosmological evolution of the hard X-ray active galactic nucleus luminosity function and the origin of the hard X-ray background,” *Astrophysical Journal*, vol. 598, no. 2, pp. 886–908, 2003.

- [105] A. J. Barger, L. L. Cowie, R. F. Mushotzky et al., “The cosmic evolution of hard X-ray-selected active galactic nuclei,” *Astronomical Journal*, vol. 129, no. 2, pp. 578–609, 2005.
- [106] S. Cristiani, D. M. Alexander, F. Bauer et al., “The space density of high-redshift QSOs in the Great Observatories Origins Deep Survey,” *Astrophysical Journal*, vol. 600, no. 2, pp. L119–L122, 2004.
- [107] X. Fan, V. K. Narayanan, R. H. Lupton et al., “A survey of $z > 5.8$ quasars in the sloan digital sky survey. I. Discovery of three new quasars and the spatial density of luminous quasars at $z \sim 6$,” *Astronomical Journal*, vol. 122, no. 6, pp. 2833–2849, 2001.
- [108] C. J. Willott, P. Delorme, C. Reyl e et al., “The Canada-France high- z quasar survey: nine new quasars and the luminosity function at redshift 6,” *Astronomical Journal*, vol. 139, no. 3, pp. 906–918, 2010.
- [109] F. Civano et al., “The population of high-redshift active galactic nuclei in the Chandra-COSMOS survey,” *Astrophysical Journal*, vol. 741, p. 91, 2011.
- [110] R. Maiolino and G. H. Rieke, “Low-luminosity and obscured seyfert nuclei in nearby galaxies,” *Astrophysical Journal*, vol. 454, no. 1, pp. 95–105, 1995.
- [111] D. R. Ballantyne, J. E. Everett, and N. Murray, “Connecting galaxy evolution, star formation, and the cosmic X-ray background,” *Astrophysical Journal*, vol. 639, no. 2, pp. 740–752, 2006.
- [112] A. T. Steffen, A. J. Barger, L. L. Cowie, R. F. Mushotzky, and Y. Yang, “The changing active galactic nucleus population,” *Astrophysical Journal*, vol. 596, no. 1, pp. L23–L26, 2003.
- [113] C. Simpson, “The luminosity dependence of the type 1 active galactic nucleus fraction,” *Monthly Notices of the Royal Astronomical Society*, vol. 360, no. 2, pp. 565–572, 2005.
- [114] A. Akylas and I. Georgantopoulos, “Can photo-ionisation explain the decreasing fraction of X-ray obscured AGN with luminosity?” *Astronomy and Astrophysics*, vol. 479, no. 3, pp. 735–740, 2008.
- [115] E. Treister, S. Virani, E. Gawiser et al., “Optical spectroscopy of X-ray sources in the extended Chandra deep field south,” *Astrophysical Journal*, vol. 693, pp. 1713–1727, 2009.
- [116] R. Barvainis, “Hot dust and the near-infrared bump in the continuum spectra of quasars and active galactic nuclei,” *Astrophysical Journal*, vol. 320, pp. 537–544, 1987.
- [117] F. L. A. Franca, F. Fiore, A. Comastri et al., “The HEL-LAS2XMM survey. VII. The hard X-ray luminosity function of AGNs up to $z = 4$: More absorbed AGNs at low luminosities and high redshifts,” *Astrophysical Journal*, vol. 635, no. 2, pp. 864–879, 2005.
- [118] G. Hasinger, “Absorption properties and evolution of active galactic nuclei,” *Astronomy and Astrophysics*, vol. 490, no. 3, pp. 905–922, 2008.
- [119] A. Akylas, I. Georgantopoulos, A. Georgakakis, S. Kitsionas, and E. Hatziminaoglou, “XMM-Newton and Chandra measurements of the AGN intrinsic absorption: dependence on luminosity and redshift,” *Astronomy and Astrophysics*, vol. 459, no. 3, pp. 693–701, 2006.
- [120] D. Lutz, R. Maiolino, H. W. W. Spoon, and A. F. M. Moorwood, “The relation between AGN hard X-ray emission and mid-infrared continuum from ISO spectra: scatter and unification aspects,” *Astronomy and Astrophysics*, vol. 418, no. 2, pp. 465–473, 2004.
- [121] J. J. Bock, K. A. Marsh, M. E. Ressler, and M. W. Werner, “High-resolution mid-infrared imaging of the nucleus of NGC 1068,” *Astrophysical Journal*, vol. 504, no. 1, pp. L5–L10, 1998.
- [122] J. T. Radoński, R. K. Piña, C. Packham et al., “Resolved mid-infrared emission in the narrow-line region of NGC 4151,” *Astrophysical Journal*, vol. 587, no. 1, pp. 117–122, 2003.
- [123] W. Saunders, M. Rowan-Robinson, A. Lawrence et al., “The 60-micron and far-infrared luminosity functions of IRAS galaxies,” *Monthly Notices of the Royal Astronomical Society*, vol. 242, pp. 318–337, 1990.
- [124] K. Iwasawa, D. B. Sanders, A. S. Evans, N. Trentham, G. Miniutti, and H. W. W. Spoon, “Fe K emission in the ultra-luminous infrared galaxy Arp 220,” *Monthly Notices of the Royal Astronomical Society*, vol. 357, pp. 565–571, 2005.
- [125] C. K. Seyfert, “Nuclear emission in spiral nebulae,” *Astrophysical Journal*, vol. 97, p. 28, 1943.
- [126] G. Risaliti, R. Maiolino, and M. Salvati, “The distribution of absorbing column densities among Seyfert 2 galaxies,” *Astrophysical Journal*, vol. 522, no. 1, pp. 157–164, 1999.
- [127] J. B. de Veny, W. H. Osborn, and K. Janes, “A catalogue of quasars,” *Astronomical Society of the Pacific*, vol. 83, p. 611, 1971.
- [128] G. Matt, A. C. Fabian, M. Guainazzi, K. Iwasawa, L. Bassani, and G. Malaguti, “The X-ray spectra of compton-thick Seyfert 2 galaxies as seen by BeppoSAX,” *Monthly Notices of the Royal Astronomical Society*, vol. 318, pp. 173–179, 2000.
- [129] M. H. K. de Grijp, G. K. Miley, J. Lub, and T. De Jong, “Infrared Seyferts: a new population of active galaxies?” *Nature*, vol. 314, no. 6008, pp. 240–242, 1985.
- [130] C. Winkler, T. J. L. Courvoisier, G. Di Cocco et al., “The INTEGRAL mission,” *Astronomy and Astrophysics*, vol. 411, no. 1, pp. L1–L6, 2003.
- [131] N. Gehrels, G. Chincarini, P. Giommi et al., “The Swift gamma-ray burst mission,” *Astrophysical Journal*, vol. 611, no. 2, pp. 1005–1020, 2004.
- [132] P. Ubertini, F. Lebrun, G. Di Cocco et al., “IBIS: the imager on-board INTEGRAL,” *Astronomy and Astrophysics*, vol. 411, no. 1, pp. L131–L139, 2003.
- [133] P. Vignati, S. Molendi, G. Matt et al., “BeppoSAX unveils the nuclear component in NGC 6240,” *Astronomy and Astrophysics*, vol. 349, no. 2, pp. L57–L60, 1999.
- [134] N. A. Levenson, T. M. Heckman, J. H. Krolik, K. A. Weaver, and P. T. Zycki, “Penetrating the deep cover of compton-thick active galactic nuclei,” *Astrophysical Journal*, vol. 648, no. 1, pp. 111–127, 2006.
- [135] G. Ghisellini, F. Haardt, and G. Matt, “The contribution of the obscuring torus to the X-ray spectrum of Seyfert galaxies—a test for the unification model,” *Monthly Notices of the Royal Astronomical Society*, vol. 267, p. 743, 1994.
- [136] D. M. Alexander, W. N. Brandt, A. E. Hornschemeier et al., “The Chandra deep field north survey. VI. The nature of the optically faint X-ray source population,” *Astronomical Journal*, vol. 122, no. 5, pp. 2156–2176, 2001.
- [137] A. J. Barger, L. L. Cowie, P. Capak et al., “Optical and infrared properties of the 2 Ms Chandra Deep Field North X-ray sources,” *Astronomical Journal*, vol. 126, no. 2, pp. 632–665, 2003.
- [138] V. Mainieri, D. Rigopoulou, I. Lehmann et al., “Submillimetre detection of a high-redshift type 2 QSO,” *Monthly Notices of the Royal Astronomical Society*, vol. 356, no. 4, pp. 1571–1575, 2005.
- [139] I. Georgantopoulos, A. Akylas, A. Georgakakis, and M. Rowan-Robinson, “The Compton-thick AGN in the Chandra Deep Field North,” *Astronomy and Astrophysics*, vol. 507, no. 2, pp. 747–756, 2009.
- [140] E. Daddi, D. M. Alexander, M. Dickinson et al., “Multiwavelength study of massive galaxies at $z \sim 2$. II. Widespread

- compton-thick active galactic nuclei and the concurrent growth of black holes and bulges,” *Astrophysical Journal*, vol. 670, no. 1, pp. 173–189, 2007.
- [141] F. Fiore, A. Grazian, P. Santini et al., “Unveiling obscured accretion in the Chandra Deep Field-South,” *Astrophysical Journal*, vol. 672, no. 1, pp. 94–101, 2008.
- [142] I. Georgantopoulos, A. Georgakakis, M. Rowan-Robinson, and E. Rovilos, “Searching for mid-IR obscured AGN in the Chandra deep field North,” *Astronomy and Astrophysics*, vol. 484, no. 3, pp. 671–678, 2008.
- [143] J. L. Donley, G. H. Rieke, P. G. Pérez-González, and G. Barro, “Spitzer’s contribution to the AGN population,” *Astrophysical Journal*, vol. 687, no. 1, pp. 111–132, 2008.
- [144] A. Pope, R. S. Bussmann, A. Dey et al., “The nature of faint Spitzer-selected dust-obscured galaxies,” *Astrophysical Journal*, vol. 689, no. 1, pp. 127–133, 2008.
- [145] A. Georgakakis, M. Rowan-Robinson, K. Nandra, J. Digby-North, P. G. Pérez-González, and G. Barro, “Infrared excess sources: compton thick QSOs, low-luminosity seyferts or starbursts?” *Monthly Notices of the Royal Astronomical Society*, vol. 406, no. 1, pp. 420–433, 2010.
- [146] M. Salvato, G. Hasinger, O. Ilbert et al., “Photometric redshift and classification for the XMM-cosmos sources,” *Astrophysical Journal*, vol. 690, no. 2, pp. 1250–1263, 2009.
- [147] C. N. Cardamone, P. G. Van Dokkum, C. M. Urry et al., “The multiwavelength Survey by Yale-Chile (MUSYC): deep medium-band optical imaging and high-quality 32-band photometric redshifts in the ECDF-S,” *Astrophysical Journal Supplement Series*, vol. 189, no. 2, pp. 270–285, 2010.
- [148] P. F. Hopkins, L. Hernquist, T. J. Cox, and D. Kerbs, “A cosmological framework for the co-evolution of quasars, supermassive black holes, and elliptical galaxies. i. galaxy mergers and quasar activity,” *Astrophysical Journal Supplement Series*, vol. 175, no. 2, pp. 356–389, 2008.
- [149] J. S. Kartaltepe, D. B. Sanders, E. Le Floch et al., “A multiwavelength study of a sample of 70 μm selected galaxies in the cosmos field. I. Spectral energy distributions and luminosities,” *Astrophysical Journal*, vol. 709, no. 2, pp. 572–596, 2010.
- [150] E. Treister, C. M. Urry, K. Schawinski, C. N. Cardamone, and D. B. Sanders, “Heavily obscured active galactic nuclei in high-redshift luminous infrared galaxies,” *Astrophysical Journal*, vol. 722, no. 2, pp. L238–L243, 2010.
- [151] B. Luo, F. E. Bauer, W. N. Brandt et al., “The Chandra deep field-south survey: 2 Ms source catalogs,” *Astrophysical Journal Supplement Series*, vol. 179, no. 1, pp. 19–36, 2008.
- [152] E. Treister, K. Schawinski, M. Volonteri, P. Natarajan, and E. Gawiser, “Black hole growth in the early Universe is self-regulated and largely hidden from view,” *Nature*, vol. 474, no. 7351, pp. 356–358, 2011.
- [153] R. J. Bouwens, G. D. Illingworth, J. P. Blakeslee, and M. Franx, “Galaxies at $z \sim 6$: the UV luminosity function and luminosity density from 506 HUDF, HUDF parallel ACS field, and goods i-dropouts,” *Astrophysical Journal*, vol. 653, no. 1, pp. 53–85, 2006.
- [154] F. Fiore, S. Puccetti, and S. Mathur, “Demography of high redshift,” In press.
- [155] C. J. Willott, “No evidence of obscured, accreting black holes in most $z = 6$ star-forming galaxies,” *Astrophysical Journal Letters*, vol. 742, no. 1, 2011.
- [156] L. L. Cowie, A. J. Barger, and G. Hasinger, “The faintest X-ray sources from $z = 0-8$,” In press.
- [157] Y. Ueda, S. Eguchi, Y. Terashima et al., “Suzaku observations of active galactic nuclei detected in the SWIFT BAT survey: discovery of a ”new type” of buried supermassive black holes,” *Astrophysical Journal*, vol. 664, no. 2 II, pp. L79–L82, 2007.
- [158] R. Gilli, J. Su, C. Norman et al., “A Compton-thick active galactic nucleus at $z \sim 5$ in the 4 Ms Chandra deep field south,” *The Astrophysical Journal Letters*, vol. 730, no. 2, p. L28, 2011.
- [159] M. Volonteri and N. Y. Gnedin, “Relative role of stars and quasars in cosmic reionization,” *Astrophysical Journal*, vol. 703, no. 2, pp. 2113–2117, 2009.
- [160] C. A. Faucher-Giguère, A. Lidz, L. Hernquist, and M. Zaldarriaga, “Evolution of the intergalactic opacity: implications for the ionizing background, cosmic star formation, and quasar activity,” *Astrophysical Journal*, vol. 688, no. 1, pp. 85–107, 2008.
- [161] P. Salucci, E. Szuszkiewicz, P. Monaco, and L. Danese, “Mass function of dormant black holes and the evolution of active galactic nuclei,” *Monthly Notices of the Royal Astronomical Society*, vol. 307, no. 3, pp. 637–644, 1999.
- [162] Q. Yu and S. Tremaine, “Observational constraints on growth of massive black holes,” *Monthly Notices of the Royal Astronomical Society*, vol. 335, no. 4, pp. 965–976, 2002.
- [163] A. Marconi, G. Risaliti, R. Gilli, L. K. Hunt, R. Maiolino, and M. Salvati, “Local supermassive black holes, relics of active galactic nuclei and the X-ray background,” *Monthly Notices of the Royal Astronomical Society*, vol. 351, no. 1, pp. 169–185, 2004.
- [164] F. Shankar, P. Salucci, G. L. Granato, G. De Zotti, and L. Danese, “Supermassive black hole demography: the match between the local and accreted mass functions,” *Monthly Notices of the Royal Astronomical Society*, vol. 354, no. 4, pp. 1020–1030, 2004.
- [165] P. Natarajan and E. Treister, “Is there an upper limit to black hole masses?” *Monthly Notices of the Royal Astronomical Society*, vol. 393, no. 3, pp. 838–845, 2009.
- [166] F. Shankar, D. H. Weinberg, and J. Miralda-Escudé, “Self-consistent models of the AGN and black hole populations: duty cycles, accretion rates, and the mean radiative efficiency,” *Astrophysical Journal*, vol. 690, no. 1, pp. 20–41, 2009.
- [167] M. Vestergaard and B. M. Peterson, “Determining central black hole masses in distant active galaxies and quasars. II. Improved optical and UV scaling relationships,” *Astrophysical Journal*, vol. 641, no. 2, pp. 689–709, 2006.
- [168] R. J. McLure and J. S. Dunlop, “The cosmological evolution of quasar black hole masses,” *Monthly Notices of the Royal Astronomical Society*, vol. 352, no. 4, pp. 1390–1404, 2004.
- [169] M. Vestergaard, “Early growth and efficient accretion of massive black holes at high redshift,” *Astrophysical Journal*, vol. 601, no. 2, pp. 676–691, 2004.
- [170] B. C. Kelly, M. Vestergaard, X. Fan, P. Hopkins, L. Hernquist, and A. Siemiginowska, “Constraints on black hole growth, quasar lifetimes, and Eddington ratio distributions from the SDSS broad-line quasar black hole mass function,” *Astrophysical Journal*, vol. 719, no. 2, pp. 1315–1334, 2010.
- [171] J. A. Kollmeier, C. A. Unken, C. S. Kochanek et al., “Black hole masses and eddington ratios at $0.3 < z < 4$,” *Astrophysical Journal*, vol. 648, no. 1, pp. 128–139, 2006.
- [172] J. R. Trump et al., “Observational limits on type 1 active galactic nucleus accretion rate in COSMOS,” *The Astrophysical Journal*, vol. 700, pp. 49–55, 2009.
- [173] P. Natarajan, “The formation and evolution of massive black hole seeds in the early universe,” *Bulletin of the Astronomical Society of India*, vol. 39, no. 1, pp. 145–161, 2011.

- [174] G. Lodato and P. Natarajan, “The mass function of high-redshift seed black holes,” *Monthly Notices of the Royal Astronomical Society*, vol. 377, no. 1, pp. L64–L68, 2007.
- [175] D. R. Ballantyne, A. R. Draper, K. K. Madsen, J. R. Rigby, and E. Treister, “Lifting the veil on obscured accretion: active galactic nuclei number counts and survey strategies for imaging hard X-ray missions,” *Astrophysical Journal*, vol. 736, no. 1, p. 56, 2011.
- [176] T. Takahashi, K. Mitsuda, R. Kelley et al., “The ASTRO-H mission,” in *Proceedings of the Space Telescopes and Instrumentation: Ultraviolet to Gamma Ray*, vol. 7732 of *Proceedings of SPIE*, 2010.
- [177] L. Yao, E. R. Seaquist, N. Kuno, and L. Dunne, “CO molecular gas in infrared-luminous galaxies,” *Astrophysical Journal*, vol. 588, no. 2, pp. 771–791, 2003.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

