

Transient neurological symptoms in the older population: report of a prospective cohort study—the Medical Research Council Cognitive Function and Ageing Study (CFAS)

Nahal Mavaddat,¹ George M Savva,² Daniel S Lasserson,³ Matthew F Giles,⁴ Carol Brayne,⁵ Jonathan Mant¹

To cite: Mavaddat N, Savva GM, Lasserson DS, *et al*. Transient neurological symptoms in the older population: report of a prospective cohort study—the Medical Research Council Cognitive Function and Ageing Study (CFAS). *BMJ Open* 2013;**3**:e003195. doi:10.1136/bmjopen-2013-003195

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-003195>).

Received 9 May 2013
Revised 11 June 2013
Accepted 12 June 2013

For numbered affiliations see end of article.

Correspondence to
Dr Nahal Mavaddat;
nm212@medschl.cam.ac.uk

ABSTRACT

Objective: Transient ischaemic attack (TIA) is a recognised risk factor for stroke in the older population requiring timely assessment and treatment by a specialist. The need for such TIA services is driven by the epidemiology of transient neurological symptoms, which may not be caused by TIA. We report prevalence and incidence of transient neurological symptoms in a large UK cohort study of older people.

Design: Longitudinal cohort study

Setting: The Medical Research Council Cognitive Function and Ageing Study (CFAS) is a population representative study based on six centres across England and Wales.

Participants: Random samples of people in their 65th year were obtained from Family Health Service Authority lists. The participation rate was 80% (n=13 004). Interview at baseline included questions about stroke and three transient neurological symptoms, repeated in a subsample after 2 years. Patients were flagged for mortality.

Main outcome measures: Prevalence and 2-year incidence of transient neurological symptoms.

Results: In 11 903 participants without a history of stroke, 271 (2.3%) reported transient problems with speech, 872 (7.6%) with sight and 596 (5.1%) weakness in a limb with 1456 (12.7%) reporting at least one symptom. Of those reinterviewed (n=6748), 675 (9.8%) reported at least one symptom over 2 years.

Conclusions: Lifetime prevalence and incidence of transient neurological symptoms in people aged 65 years and over is high and is substantially greater than the incidence of TIA in hospital-based and population-based studies. These high rates of transient neurological symptoms in the community in the older population should be considered when planning TIA services.

ARTICLE SUMMARY

Article focus

- Prompt initiation of secondary prevention following transient ischaemic attack (TIA) is associated with up to 80% reduction in risk of subsequent stroke.
- Many people presenting to specialist TIA clinics with transient neurological symptoms do not have TIA.
- The prevalence and incidence of transient neurological symptoms (as opposed to TIA) in older age groups is unknown.

Key messages

- In a large multicentred community-based study representative of the older population in the UK, we found a high prevalence and incidence of transient neurological symptoms, significantly greater than that of TIAs in hospital-based and population-based studies.
- These high rates of transient neurological symptoms in the community in the older population should be considered when planning TIA services.

Strengths and limitations of this study

- Previous studies of incidence and prevalence of transient neurological symptoms have mostly been conducted in populations not representative of older age groups where TIA is most common and few have been undertaken in UK populations.
- The incidence of transient neurological symptoms in the study was determined in people without severe cognitive impairment, and may be an underestimation, since cognitive impairment can be a manifestation of vascular disease. The wording of questions to identify transient neurological symptoms may have picked up some people who had symptoms lasting more than 24 h.

INTRODUCTION

Transient ischaemic attack (TIA) is an established and powerful risk factor for stroke. Eight per cent of patients who have TIA suffer from stroke within 7 days, many within 48 h.^{1 2} Immediate specialist assessment and treatment is associated with substantial reductions in this early risk of stroke,³ and is recommended for people with suspected TIA.⁴

As awareness grows of the urgency of early management of TIA among both primary care practitioners and the general public,⁵ more patients with symptoms suggestive of TIA are likely to present to primary and secondary care services. Need for specialist services will be driven by incidence of symptoms that might represent TIA (ie, transient neurological symptoms) rather than the epidemiology of TIA per se. Currently about 48 000 probable or definite TIAs (transient neurological symptoms lasting less than 24 h of likely vascular aetiology) and 43 000 minor strokes (stroke events causing minimal or no neurological deficits) are managed as outpatients every year in England, but it is not clear to what extent this might be the tip of the iceberg.⁶ Data are available worldwide on the prevalence and incidence of TIA.^{7–11} The epidemiology in the community of transient neurological symptoms (ie, neurological symptoms of sudden onset of which TIA is a subset) is however less well defined. Previous studies of incidence and prevalence of transient neurological symptoms have mostly been conducted in populations not representative of older age groups where TIA is most common.^{7 11–16} Very few have been undertaken in UK populations. While incidence of transient neurological symptoms will drive the need for specialist TIA services, prevalence of transient neurological symptoms in the community is also important. TIA is associated with a long-term increase in risk of stroke,^{17 18} so there may be value in diagnosing ‘old’ events that have not presented to medical services in order to target secondary prevention. Hence, the importance of determining both the incidence and prevalence of transient neurological symptoms in the community.

This study reports prevalence and incidence of three common transient neurological symptoms (limb weakness, loss of speech and disturbance of vision) in a population-based multicentred cohort study in the UK (England and Wales) in those aged 65 years and over, the Medical Research Council Cognitive Function and Ageing Study (CFAS).¹⁹

METHODS

The Medical Research Council CFAS is a population-representative study of individuals aged 65 years and over. The study began in 1991 and was designed to determine the incidence of dementia in the older population. It has six centres across England and Wales chosen to represent the national variation of urban–rural mix, socioeconomic deprivation and rates of chronic disease.¹⁹ Five of these with identical study designs (Oxford, Nottingham, Cambridgeshire, Gwynedd and Newcastle) are used in

the present investigation. The sixth centre (Liverpool) used a different design and is therefore excluded. Random samples of people in their 65th year and above were obtained from Family Health Service Authority lists (agency responsible for maintaining registers of general practice populations at that time). The sample was stratified by age (65–74, 75 years and over) and equal numbers were randomly selected from these groups with the aim of recruiting 2500 to each centre. All study centres obtained ethical approval from local research committees (REC Ref: 05/MRE05/37). Full details of methodology are available elsewhere.¹⁹

Eligible participants (or their proxies where appropriate) provided informed consent. Trained interviewers undertook baseline interviews in the participants’ homes, including sociodemographic characteristics, cognitive function and disease history, including previous stroke, coronary heart disease and diabetes (full details at <http://www.cfas.ac.uk>). On the basis of baseline screening, the study sample was divided into two groups at baseline: people without cognitive impairment and a group consisting of those with cognitive impairment plus a stratified subsample of those without cognitive impairment. The first group underwent no further assessment at baseline and was ‘rescreened’ after 2 years. The second group underwent a further detailed cognitive assessment at baseline, but was not followed up for transient neurological symptoms. The estimates of prevalence of transient neurological symptoms are calculated for all people at baseline, while the incidence estimates use only the first group. All participants were flagged with the National Health Service (NHS) Central Register. Deaths and underlying causes of death attributed to stroke (International Classification of Diseases (ICD) codes 430–438) were notified to the study.

Prevalence of transient neurological symptoms

At baseline all participants were asked “Have you ever experienced sudden problems with:

1. speech, which got better after a day?
2. weakness in the arms or legs, which got better after a day?
3. sight, which got better after a day?”

Social class was determined using the Registrar General’s Occupational Classification. Cognitive status was determined using the Mini-Mental State Examination (MMSE)²⁰ and the Automated Geriatric Examination for Computer Assisted Taxonomy (AGECAT).²¹

Incidence of transient neurological symptoms

All those in the ‘rescreen’ arm were asked at 2-year follow-up if they had experienced each transient neurological symptom in the past 2 years.

ANALYSIS

All analyses were carried out using STATA V.11. Inverse probability weights were used throughout to ensure that

Transient Neurological symptoms in the older population in the MRC-CFAS

the sample was representative of the target population. Weights were estimated with logistic regression using presence in each phase of the study as an outcome and taking into account oversampling of over-75s at baseline. Weights for incidence calculations adjust for attrition based on baseline characteristics and stratified selection into the assessment arm as appropriate. Baseline prevalence of each transient neurological symptom was calculated for age-specific strata and as a weighted percentage to provide a population estimate of prevalence for people aged 65 years and over. Associations with gender, age, social class (manual (IIIb, IV and V) and non-manual (I, II and IIIa)) and cardiovascular comorbidity were explored using logistic regression models adjusting for all other factors and cognitive function based on MMSE by score (less than 18, 18–21, 22–25 and 26–30).

Two-year incidence of transient neurological symptoms was estimated using weighted percentages of those reporting any of the symptoms during follow-up. Weights were adjusted for refusals, dropouts and for non-reassessment of people with cognitive impairment. Calculation of attrition weights for incident transient neurological symptoms excluded those who died between baseline and follow-up. These estimates are therefore applicable to the population over 65 years without severe cognitive impairment or stroke at baseline surviving at least 2 years.

RESULTS

The participation rate in the CFAS was 80% (13 004/16 258). For the prevalence analysis, participants who had a stroke at baseline (963, 7.4%) or for whom baseline information about stroke was missing (138, 1.1%) were excluded, leaving 11 903 participants. In total, 2283 of the original participants were not allocated to the 'rescreen' arm at 2 years; 754 died, 1973 declined to participate and 145 were lost to follow-up, leaving 6748 (76% response rate of potential participants) for the incidence analysis.

Table 1 shows the demographic features of participants in the CFAS, studied in this analysis at baseline. Forty per cent of participants were men and 60% were women. 12.3% of all participants reported at least one transient neurological symptom; 2.3% reported transient loss of speech, 5.1% transient weakness and 7.6% transient loss of sight.

Table 2 shows prevalence data with weighted percentages and ORs (adjusted for age, sex, social class, cognition and cardiovascular morbidities) of reported transient neurological symptoms at baseline by demographic factors and comorbidities. In total, 12.8% of men and 12% of women reported at least one of the three transient neurological symptoms of loss of speech, loss of sight or weakness. There was no significant association of gender with reporting at least one transient neurological symptom, but significantly lower odds of reporting a transient neurological symptom in the over

Table 1 Demographics of participants in the Cognitive Function and Aging Study (CFAS) by sex, age, social class and comorbidity (data are counted in percentages)

	All (n=11903)		Males (n=4689)		Females (n=7214)	
	N	Per cent*	N	Per cent*	N	Per cent*
	Age (years)					
65–74	5980	58.8	2635	64.5	3345	55.0
75–84	4624	32.1	1716	29.6	2908	33.8
85+	1299	9.1	338	5.9	961	11.2
Social class						
I	558	4.8	225	4.8	333	4.8
II	2988	26.1	1213	26.3	1775	25.9
IIIa	1337	11.6	456	9.9	881	12.8
IIIb	4295	37.7	1865	40.8	2430	35.5
IV	1727	14.9	650	13.9	1077	15.7
V	561	4.9	196	4.3	365	5.3
Comorbidity						
Angina	1923	16.3	878	15.3	1045	11.1
Diabetes	672	5.5	310	6.4	362	4.9
Heart attack	1139	9.5	632	13.6	507	6.8

*Weighted percentages.

85 age group. The odds of reporting any transient neurological symptom were higher in those in manual compared to non-manual social classes. The presence of cardiovascular morbidities of angina and heart attack were also significantly associated with the odds of having at least one of the transient neurological symptoms.

There was no difference between genders with regard to reporting individual transient neurological symptoms of loss of speech, loss of sight or weakness. However, there was a significantly lower odds of reporting transient visual loss in those aged over 85 years, and an increased odds of reporting transient symptoms of weakness and loss of sight in manual compared to non-manual social classes. The presence of cardiovascular comorbidities of angina and heart attack were associated with higher odds of all three symptoms.

Table 3 shows 2-year incidence of transient neurological symptoms in respondents attending the 'rescreen'. A total of 9.8% of participants reported at least one transient neurological symptom over the past 2 years, with the highest incidence of reported transient neurological symptoms being due to loss of sight followed by weakness, with loss of speech being the least frequently reported transient neurological symptom. The incidence of each of the transient neurological symptoms was highest in the 75–84 age group and lowest in the over 85 age group.

DISCUSSION

Our findings suggest that transient neurological symptoms are common in the older population in England

Table 2 Distribution and adjusted odds of transient neurological symptoms in those aged 65 years or over in the Cognitive Function and Aging Study (CFAS) by sex, social class and comorbidity (data are counted in percentages)

	Loss of speech			Weakness			Loss of sight			At least one symptom		
	N	Per cent*	OR† (95% CI)	N	Per cent*	OR† (95% CI)	N	Per cent*	OR† (95% CI)	N	Per cent*	OR† (95% CI)
Gender												
Male	119	2.5	1.0	257	5.5	1.0	347	7.6	1.0	598	12.8	1.0
Female	152	2.2	0.9 (0.7 to 1.2)	339	4.8	0.9 (0.7 to 1.0)	525	7.6	1.0 (0.9 to 1.2)	858	12.0	1.0 (0.9 to 1.1)
Age (years)												
65–74	137	2.3	1.0	302	5.1	1.0	480	8.1	1.0	768	12.9	1.0
75–84	104	2.3	0.7 (0.7 to 1.2)	229	5.0	0.9 (0.7 to 1.1)	327	7.2	0.9 (0.7 to 1.1)	553	12.0	0.9 (0.8 to 1.0)
85+	30	2.5	0.8 (0.5 to 1.3)	65	5.4	0.9 (0.6 to 1.2)	65	5.4	0.6 (0.5 to 0.8)	134	10.4	0.7 (0.6 to 0.9)
Social class‡												
Non-manual	96	5.8	1.0	63	11.0	1.0	295	17.9	1.0	497	10.2	1.0
Manual	164	7.2	1.1 (0.9 to 1.5)	390	17.8	1.4 (1.2 to 1.7)	556	25.7	1.4 (1.2 to 1.6)	920	14.1	1.4 (1.2 to 1.6)
Comorbidity												
Angina (n=1538)	64	4.3	1.5 (1.0 to 2.2)	176	11.7	2.5 (2.0 to 3.2)	160	10.6	1.3 (1.0 to 1.6)	312	20.7	1.7 (1.4 to 1.9)
Diabetes (n=672)	28	4.0	1.4 (0.9 to 2.2)	40	6.1	1.0 (0.7 to 1.4)	66	10.1	1.3 (0.9 to 1.7)	102	15.2	1.1 (0.9 to 1.4)
Heart attack (n=1139)	52	4.7	1.8 (1.2 to 2.7)	131	10.5	1.4 (1.2 to 1.9)	119	11.5	1.4 (1.1 to 1.8)	233	12.4	1.4 (1.2 to 1.8)

*Weighted percentages.

†Adjusted for age, sex, social class, cognition and cardiovascular comorbidities.

‡Reference non-manual (I, II and IIIa) compared to manual social class (IIIb, IV and V).

156 participants had missing data on the speech questionnaire, 155 on the weakness and 154 on the sight question.

Transient Neurological symptoms in the older population in the MRC-CFAS

Table 3 Two-year incidence of transient neurological symptoms in the Cognitive Function and Aging Study (CFAS) in those without stroke at baseline at 2-year screening assessment, N at 2 years was 6748 (data are in N (%))

Transient neurological symptoms	Age (years)							
	All		65–74 (N=3657)		75–84 (N=2618)		85+ (N=473)	
	N	Per cent*	N	Per cent*	N	Per cent*	N	Per cent*
Loss of speech	149	2.1	55	1.5	84	3.3	10	2.0
Weakness	239	3.7	117	3.4	106	4.1	16	3.4
Loss of sight	410	6.3	214	6.3	170	6.6	26	4.5
At least one symptom	675	9.8	327	9.3	304	11.6	44	7.8

*Weighted percentage.

98 participants had missing data on the speech questionnaire, 97 on the weakness and 105 on the sight question.

and Wales, with at least 12% of people aged 65 years and over having experienced a transient neurological symptom of the arm or leg, speech or vision, which gets better after a day, and approximately 5% having experienced at least one such symptom over the course of a year. The commonest of these was transient symptoms of vision, followed by limb weakness. Those aged over 85 years reported lower rates of transient neurological symptoms, predominantly due to less frequent reporting of transient visual loss. Problems of memory and recall may have contributed to under-reporting, and higher mortality rates associated with neurological incidents to underestimation of rates of transient neurological symptoms in this age group. Chronic visual problems may have also potentially masked transient visual losses in the oldest-old population. Incidence of at least one transient neurological symptom in the CFAS population was approximately 2.6 times greater than that of confirmed TIA presenting to medical services in the Oxford Vascular (OxVASC) study in people aged over 85 years (approximately 15/1000/year), eight times greater in those aged 75–84 years (7/1000/year) and approximately 15 times greater in those aged 65–74 years (3/1000 population/year).¹⁰

The prevalence and incidence of transient neurological symptoms in the CFAS was somewhat higher than those found by questionnaires used in other studies internationally (mostly conducted in younger age groups), but comparable to those of Wilkinson *et al* in an over 60 age group in a US-based study.^{7 12–14 16} Online supplementary table S4 shows the prevalence of sudden onset of neurological symptoms of weakness in a limb or of loss of speech or sight in previous population and community studies.^{7 12–14 16} Questions used to elicit transient neurological symptoms in the CFAS are likely to have captured greater numbers of transient neurological events in the population compared with those seeking more specific vascular symptoms, or to the few studies defining more precisely the onset, offset and timing of the event.^{7 12–14} While some respondents will have been describing true transient ischaemic events, it is likely that many did not have a true TIA. Validated measures to determine the presence of previous TIA are limited, and most studies of transient neurological

symptoms in questionnaires overestimate true transient ischaemic attacks.^{13 14 22} Wilkinson *et al*¹⁴ suggest that around 10% of transient symptoms reported in a questionnaire are subsequently diagnosed as TIA after a neurologist's assessment. Self-reported transient neurological symptoms in other community-based studies have low positive predictive value, with higher values only in studies in outpatient populations (over 70%).^{13 14 22–24} This suggests that the use of screening to identify possible past transient ischaemic attacks may generate unnecessary extra strain on primary care and secondary TIA services with limited benefit.

Patients presenting with transient neurological symptoms present considerable diagnostic dilemmas for primary care practitioners reflected in low rates of TIA diagnosis confirmation in patients referred to UK TIA clinics (over 50% being for non-TIA causes).^{6 25 26} Many symptoms of TIA are non-specific and occur in non-vascular syndromes. Dizziness, ophthalmological problems, migraine, epilepsy, nerve entrapment or psychological states are the commonest non-TIA diagnoses.^{25–29} Often, however, no diagnosis can be determined.^{25 26} Scales such as the ABCD² help determine the urgency of TIA referral, but do not distinguish TIA from non-TIA symptoms. Wilkinson *et al*¹⁴ reported transient loss of speech and loss of sight to be more reliable than limb weakness for neurologist diagnosis of TIA, while Hart *et al*⁷ found the strongest and most consistent relationship with subsequent stroke to be loss of power in an arm. Other studies suggest that the presence of certain symptoms such as headache, dizziness, loss of consciousness, memory loss, blurred vision, generalised weakness, pain in limbs and seizures make the diagnosis of TIA less likely.^{14 29 30}

Study limitations

The incidence of transient neurological symptoms was determined in people without severe cognitive impairment, and may be an underestimate since cognitive impairment can be a manifestation of vascular disease.^{31 32} Only three transient neurological symptoms were reported in this study. Some symptoms of TIA (eg, posterior circulation or pure sensory symptoms) were not sought. The wording of questions to identify transient

Transient Neurological symptoms in the older population in the MRC-CFAS

neurological symptoms may have picked up some people who had symptoms lasting more than 24 h. Such patients are nevertheless likely to draw on TIA services for further assessment. In addition, it was not possible to determine whether different transient neurological symptoms in a participant occurred simultaneously or separately in time. The problem of recall bias in these self-reports of transient symptoms needs to be considered. Questions were designed to be easily comprehensible to the older population. However, the need to consider multiple criteria when responding to a question, for example, 'sudden onset' and 'less than a day', may have led to difficulties in interpretation of the transient neurological symptoms questions for some participants. Finally, baseline and 2-year follow-up were carried out in the 1990s and changes in the rates of vascular events may have occurred since then. While the age-specific incidence of stroke appears to be declining, it is not clear that the same is true of TIA.^{10 33 34} This study is more recent than many studies estimating rates of transient neurological symptoms in the population, many of which have been carried out during the 1970s or 1980s.^{12–14}

CONCLUSION

In a large multicentred community-based study, highly representative of the older population in the UK and conducted in the age group where TIAs are most common, we found a high prevalence and incidence of transient neurological symptoms. The incidence of such symptoms in the community is significantly greater than that of TIAs. This highlights the need for adequate provision of TIA services and the potential importance of the development of valid diagnostic tools to assist the general practitioner and hospital doctor in better triage of people presenting with transient neurological symptoms.

Author affiliations

¹Primary Care Unit, Department of Public Health and Primary Care, University of Cambridge, Strangeways Laboratory, Worts Causeway, Cambridge, UK

²School of Nursing Sciences, University of East Anglia, Norwich Research Park, Norwich, UK

³Department of Primary Care Health Sciences, University of Oxford, Oxford, UK

⁴Stroke Prevention Research Unit, Department of Clinical Neurology, John Radcliffe Hospital, NIHR Biomedical Research Centre, Oxford University, Oxford, UK

⁵Department of Public Health and Primary Care, University of Cambridge, Institute of Public Health, Cambridge, UK

Acknowledgements We would like to thank the MRC CFAS study group for data collection and management. We are also grateful to all respondents, their families and their primary care teams from across the country for their participation in the MRC CFAS.

Contributors NM and GMS contributed to analysis of data. All authors contributed to study design and intellectual input.

Funding The MRC CFAS has had funding by major awards from the Medical Research Council (MRC/G9901400) and Department of Health. NM was funded by a National Institute for Health Research (NIHR) Walport Clinical Lectureship in Primary Care Research.

Competing interests None.

Patient consent Obtained.

Ethics approval Multicentre REC (Ref: 05/MRE05/37).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data can be shared through application. For further information please refer to the application form on the website <http://www.cfas.ac.uk>.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 3.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/3.0/>

REFERENCES

- Coull AJ, Lovett JK, Rothwell PM. Population based study of early risk of stroke after transient ischaemic attack or minor stroke: implications for public education and organisation of services. *BMJ* 2004;328:326.
- Rothwell PM, Giles MF, Flossmann E, *et al*. A simple score (ABCD) to identify individuals at high early risk of stroke after transient ischaemic attack. *Lancet* 2005;366:29–36.
- Rothwell PM, Giles MF, Chandratheva A, *et al*. Effect of urgent treatment of transient ischaemic attack and minor stroke on early recurrent stroke (EXPRESS study): a prospective population-based sequential comparison. *Lancet* 2007;370:1432–42.
- Department of Health. National Stroke Strategy. 2007.
- Department of Health. Stroke: Act F.A.S.T. awareness campaign. 2009.
- Giles MF, Rothwell PM. Substantial underestimation of the need for outpatient services for TIA and minor stroke. *Age Ageing* 2007;36:676–80.
- Hart CL, Hole DJ, Smith GD. The relation between questions indicating transient ischaemic attack and stroke in 20 years of follow up in men and women in the Renfrew/Paisley Study. *J Epidemiol Community Health* 2001;55:653–6.
- Mant J, Wade DT, Winner S. Health care needs assessment: stroke. In: Stevens A, Raftery J, Mant J, Simpson S, eds. *Health care needs assessment: the epidemiologically based needs assessment reviews*. 2nd edn. Oxford: Radcliffe Medical Press, 2004: 141–244.
- Orlandi G, Gelli A, Fanucchi S, *et al*. Prevalence of stroke and transient ischaemic attack in the elderly population of an Italian rural community. *Eur J Epidemiol* 2003;18:879–82.
- Rothwell PM, Coull AJ, Silver LE, *et al*. Population-based study of event-rate, incidence, case fatality, and mortality for all acute vascular events in all arterial territories (Oxford Vascular Study). *Lancet* 2005;366:1773–83.
- Bos MJ, van Rijn MJ, Witteman JC, *et al*. Incidence and prognosis of transient neurological attacks. *JAMA* 2007;298:2877–85.
- Mules J, Henderson M, Kuller L. A population survey of symptoms suggestive of transient ischemic attacks. *Stroke* 1971;2:114–27.
- Toole JF, Lefkowitz DS, Chambless LE, *et al*. Self-reported transient ischemic attack and stroke symptoms: methods and baseline prevalence. The ARIC Study, 1987–1989. *Am J Epidemiol* 1996;144:849–56.
- Wilkinson WE, Heyman A, Burch JG, *et al*. Use of a self-administered questionnaire for detection of transient cerebral ischemic attacks: I. Survey of elderly persons living in retirement facilities. *Ann Neurol* 1979;6:40–6.
- Bots ML, van der Wilk EC, Koudstaal PJ, *et al*. Transient neurological attacks in the general population. Prevalence, risk factors, and clinical relevance. *Stroke* 1997;28:768–73.
- Howard VJ, McClure LA, Meschia JF, *et al*. High prevalence of stroke symptoms among persons without a diagnosis of stroke or transient ischemic attack in a general population: the REasons for Geographic And Racial Differences in Stroke (REGARDS) study. *Arch Intern Med* 2006;166:1952–8.
- van Wijk I, Kappelle LJ, van GJ, *et al*. Long-term survival and vascular event risk after transient ischaemic attack or minor ischaemic stroke: a cohort study. *Lancet* 2005;365: 2098–104.
- Clark TG, Murphy MF, Rothwell PM. Long term risks of stroke, myocardial infarction, and vascular death in "low risk" patients with a non-recent transient ischaemic attack. *J Neurol Neurosurg Psychiatry* 2003;74:577–80.

Transient Neurological symptoms in the older population in the MRC-CFAS

19. Brayne C, McCracken C, Matthews FE. Cohort profile: the Medical Research Council Cognitive Function and Ageing Study (CFAS). *Int J Epidemiol* 2006;35:1140–5.
20. Folstein MF, Folstein SE, McHugh PR. Mini-mental state. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–98.
21. Copeland JR, Dewey ME, Griffiths-Jones HM. A computerized psychiatric diagnostic system and case nomenclature for elderly subjects: GMS and AGE-CAT. *Psychol Med* 1986;16:89–99.
22. Lasserson DS. Initial management of suspected transient cerebral ischaemia and stroke in primary care: implications of recent research. *Postgrad Med J* 2009;85:422–7.
23. Jones WJ, Williams LS, Meschia JF. Validating the Questionnaire for Verifying Stroke-Free Status (QVSFS) by neurological history and examination. *Stroke* 2001;32:2232–6.
24. Wilson A, Potter J, Taub N, *et al*. The effectiveness of a modified version of the Wilkinson questionnaire in screening for TIA and minor stroke in the United Kingdom. *Age Ageing* 2005;34:30–5.
25. Murray S, Bashir K, Lees KR, *et al*. Epidemiological aspects of referral to TIA clinics in Glasgow. *Scott Med J* 2007;52:4–8.
26. Martin PJ, Young G, Enevoldson TP, *et al*. Overdiagnosis of TIA and minor stroke: experience at a regional neurovascular clinic. *QJM* 1997;90:759–63.
27. Boysen G, Jensen G, Schnohr P. Frequency of focal cerebral transient ischemic attacks during a 12 month period. The Copenhagen city heart study. *Stroke* 1979;10:533–4.
28. Johnson SE, Skre H. Transient cerebral ischemic attacks in the young and middle aged. A population study. *Stroke* 1986;17:662–6.
29. Amort M, Fluri F, Schafer J, *et al*. Transient ischemic attack versus transient ischemic attack mimics: frequency, clinical characteristics and outcome. *Cerebrovasc Dis* 2011;32:57–64.
30. Dawson J, Lamb KE, Quinn TJ, *et al*. A recognition tool for transient ischaemic attack. *QJM* 2009;102:43–9.
31. De Haan EH, Nys GM, van Zandvoort MJ. Cognitive function following stroke and vascular cognitive impairment. *Curr Opin Neurol* 2006;19:559–64.
32. Selnes OA, Vinters HV. Vascular cognitive impairment. *Nat Clin Pract Neurol* 2006;2:538–47.
33. Dennis MS, Bamford JM, Sandercock PA, *et al*. Incidence of transient ischemic attacks in Oxfordshire, England. *Stroke* 1989;20:333–9.
34. Rothwell PM, Coull AJ, Giles MF, *et al*. Change in stroke incidence, mortality, case-fatality, severity, and risk factors in Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet* 2004;363:1925–33.

BMJ Open

Transient neurological symptoms in the older population: report of a prospective cohort study—the Medical Research Council Cognitive Function and Ageing Study (CFAS)

Nahal Mavaddat, George M Savva, Daniel S Lasserson, et al.

BMJ Open 2013 3:

doi: [10.1136/bmjopen-2013-003195](https://doi.org/10.1136/bmjopen-2013-003195)

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/3/7/e003195.full.html>

These include:

Data Supplement

"Supplementary Data"

<http://bmjopen.bmj.com/content/suppl/2013/07/22/bmjopen-2013-003195.DC1.html>

References

This article cites 31 articles, 16 of which can be accessed free at:

<http://bmjopen.bmj.com/content/3/7/e003195.full.html#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 3.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Cardiovascular medicine](#) (227 articles)

[Epidemiology](#) (589 articles)

[Neurology](#) (128 articles)

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>