

UNDERSTANDING THE DISEASE



Understanding renal recovery

Max Bell^{1*} , Lakhmir S. Chawla² and R. Wald³

© 2017 The Author(s). This article is an open access publication

The traditional view of recovery following acute kidney injury (AKI) was that patients who successfully overcame the severe illness underlying their AKI would experience a full restoration of their pre-morbid kidney function. Recent work has suggested that patients who survived an episode of AKI are at ongoing risk of adverse outcomes. These include de novo chronic kidney disease (CKD), progression of pre-existing CKD, end-stage kidney disease (ESRD), and death [1–4]. With a greater appreciation of the fact that AKI survivors often have a challenging clinical course, healthcare providers and researchers must establish practical definitions to cover the full spectrum of post-AKI kidney outcomes.

While ESRD reflects the extreme end of the CKD spectrum and fortunately affects a minority of AKI survivors, progressive non-dialysis requiring CKD is a far more frequent outcome following an episode of AKI. The incidence of progressive CKD in AKI survivors may be underestimated partly because kidney function is not routinely monitored in structured follow-up programs [5]. This is further limited by the existence of variable definitions for kidney recovery. Finally, even when serial assessments of kidney function are available, reliance on creatinine as a marker of glomerular filtration rate (GFR) risks overestimating renal function shortly after an acute illness during which muscle wasting might have occurred. Prowle and co-workers demonstrated that a potential 135% increase in CKD diagnoses when adjusting for the confounding effect of prolonged major illness on the serum creatinine concentration [6].

Data from large registries have shed some light on the risk of CKD following critical illness that was complicated by AKI. In a cohort of 130,134 critically ill patients

from the Swedish Intensive Care Register from 2005 to 2011, followed for 1–7 years, the relative risks of de novo CKD and ESRD after an episode of AKI (patients with pre-admission CKD excluded) were 7- and 22-fold higher as compared to patients who did not experience AKI [7]. The incidence of CKD was 6.0% (95% CI 5.1–7.0) at 1 year and 10.5% (95% CI 8.5–13.0) at 5 years, with ESRD occurring in 0.44% (95% CI 0.18–0.24) and 1.8% (95% CI 1.6–1.9) at corresponding time points. This confirms data from prior studies that have demonstrated an association between AKI and subsequent CKD, whereby CKD develops in 20–40% of AKI survivors [8–10].

Numerous factors impact the trajectory of kidney function after an episode of AKI. Prompt recovery of AKI anticipates a lower likelihood of CKD as compared to a slower recovery [11]. In a cohort of 17,000 patients, the authors observed five patterns of recovery following an initial episode of AKI: early reversal, with kidney function sustained to discharge (26.6%); no reversal at all (26.5%); late reversal after day 7 (9.7%); early reversal with relapse/relapses but ultimate recovery (22.5%); and relapse without recovery (14.7%) [11]. Schiffel and Fischer demonstrated that 26% of AKI survivors with non-recovery of kidney function at discharge improved renal function and 10.7% returned to a normalized estimated GFR; all changes took place within the first year [12]. A study of hospitalized patients described a return to baseline creatinine in 92.5%, partial recovery in 7%, and no recovery in 0.6% of patients 3 months after AKI. Patients with RIFLE F had significantly lower rates of recovery ($p < 0.001$) [13].

The Acute Disease Quality Initiative (ADQI) group has developed consensus definitions for renal recovery after AKI (Fig. 1). The new proposed definitions differentiate rapid recovery from delayed recovery and provide a framework for staging the post-AKI/pre-CKD period. Once validated, this proposed scheme could help

*Correspondence: max.bell@sil.se¹ Perioperative Medicine and Intensive Care, Karolinska University Hospital, Stockholm, Sweden

Full author information is available at the end of the article




Fig. 1 Hypothetical trajectories of acute kidney disease (AKD). AKD follows on from acute kidney injury (AKI) in those patients who do not fully recover within 7 days. The trajectory of AKD can take many forms largely depending on the severity of the initial AKI episode. Here, a series of hypothetical scenarios representing typical trajectories of the AKI–AKD continuum are depicted. Stage 3 AKI might slowly improve to stage 2 AKI and then progress to AKD (1). Stage 1 AKI might progress to stage 3, then improve rapidly to stage 1 AKI before progressing to stage 1 AKD (2). An episode of persistent AKI (>48 h) might be followed by a period of sustained reversal (*asterisks*), then a second episode of AKI (*double dagger*) leading to AKD (3). Stage 2 AKI might rapidly reverse (4). Subacute AKD might occur wherein the first 7 days are marked with slowly worsening renal function that does not technically meet the criteria for AKI, and progress to Stage 3 AKD (5). This trajectory can be seen in patients treated with chronic nephrotoxic medications (e.g., with aminoglycosides) Modified from Acute Dialysis Quality Initiative 16; www.adqi.org

standardize the nomenclature of kidney recovery and enable the design of robust trials that examine kidney recovery as an outcome in trials testing various candidate interventions [14].

The optimal follow-up of AKI survivors remains controversial. In a cohort of patients discharged from hospital after an episode of dialysis-requiring AKI, Harel et al. showed that only a minority of patients visited a nephrologist in the weeks following their hospitalization; however, there was an association between visiting a nephrologist and improved survival [15]. Though it might seem intuitive to extend early nephrology follow-up to all survivors of non-dialysis requiring AKI, the high incidence of hospital-associated AKI would make this impracticable. Moreover, many AKI survivors will have a benign clinical course suggesting that more targeted selection of high-risk patients is required. At present, there is limited information to predict adverse outcomes in AKI survivors and a biomarker that could reliably predict such outcomes would be of high value to clinicians. Dedicated clinics for the follow-up of AKI survivors have been established at some centers [16]. An ongoing multicenter pilot

trial in Canada is randomizing survivors of AKI (KDIGO stage 2 or higher) to follow-up in a dedicated post-AKI clinic for 1 year as compared to usual care [Clinicaltrials.gov NCT02483039]. As we await the findings of this trial more research is needed to clarify the most impactful components of care delivery in such clinics.

In the meantime we suggest that patients with severe AKI, KDIGO stage 3 and certainly those who received renal replacement therapy, receive targeted follow-up with a nephrologist within 30 days of hospital discharge with further follow-up individualized to the patient's needs. Such follow-up should comprise a reassessment of kidney function/kidney damage, attention to blood pressure control and cardiovascular risk factors, medication reconciliation, and patient education regarding the implications of a prior AKI episode.

Author details

¹ Perioperative Medicine and Intensive Care, Karolinska University Hospital, Stockholm, Sweden. ² Veterans Affairs Medical Center, Washington DC, USA. ³ Division of Nephrology, St. Michael's Hospital and the University of Toronto, Toronto, Canada.

Compliance with ethical standards

Conflicts of interest

None.

Open Access This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Received: 3 February 2017 Accepted: 10 March 2017

Published online: 23 March 2017

References

- Chawla LS, Eggers PW, Star RA, Kimmel PL (2014) Acute kidney injury and chronic kidney disease as interconnected syndromes. *N Engl J Med* 371:58–66
- Nisula S, Kaukonen KM, Vaara ST, Korhonen AM, Poukkanen M, Karlsson S, Haapio M, Inkinen O, Parviainen I, Suojaranta-Ylinen R, Laurila JJ, Tenhunen J, Reinikainen M, Ala-Kokko T, Ruokonen E, Kuitunen A, Pettila V, FINNAKI Study Group (2013) Incidence, risk factors and 90-day mortality of patients with acute kidney injury in Finnish intensive care units: the FINNAKI study. *Intensive Care Med* 39:420–428
- Rimes-Stigare C, Frumento P, Bottai M, Martensson J, Martling CR, Bell M (2015) Long-term mortality and risk factors for development of end-stage renal disease in critically ill patients with and without chronic kidney disease. *Crit Care* 19:383
- Wald R, Quinn RR, Luo J, Li P, Scales DC, Mamdani MM, Ray JG, University of Toronto Acute Kidney Injury Research Group (2009) Chronic dialysis and death among survivors of acute kidney injury requiring dialysis. *JAMA* 302:1179–1185
- Siew ED, Peterson JF, Eden SK, Hung AM, Speroff T, Ikizler TA, Matheny ME (2012) Outpatient nephrology referral rates after acute kidney injury. *J Am Soc Nephrol* 23:305–312
- Prowle JR, Kolic I, Purdell-Lewis J, Taylor R, Pearse RM, Kirwan CJ (2014) Serum creatinine changes associated with critical illness and detection of persistent renal dysfunction after AKI. *Clin J Am Soc Nephrol* 9:1015–1023
- Rimes-Stigare C, Frumento P, Bottai M, Martensson J, Martling CR, Walther SM, Karlstrom G, Bell M (2015) Evolution of chronic renal impairment and long-term mortality after de novo acute kidney injury in the critically ill; a Swedish multi-centre cohort study. *Crit Care* 19:221
- Amdur RL, Chawla LS, Amodeo S, Kimmel PL, Palant CE (2009) Outcomes following diagnosis of acute renal failure in U.S. veterans: focus on acute tubular necrosis. *Kidney Int* 76:1089–1097
- Bucaloiu ID, Kirchner HL, Norfolk ER, Hartle JE 2nd, Perkins RM (2012) Increased risk of death and de novo chronic kidney disease following reversible acute kidney injury. *Kidney Int* 81:477–485
- Goldstein SL, Jaber BL, Faubel S, Chawla LS, Acute Kidney Injury Advisory Group of American Society of Nephrology (2013) AKI transition of care: a potential opportunity to detect and prevent CKD. *Clin J Am Soc Nephrol* 8:476–483
- Kellum JA, Sileanu FE, Bihorac A, Hoste EA, Chawla LS (2016) Recovery after acute kidney injury. *Am J Respir Crit Care Med*. doi:10.1164/rccm.201604-0799OC
- Schiff H, Fischer R (2008) Five-year outcomes of severe acute kidney injury requiring renal replacement therapy. *Nephrol Dial Transplant* 23:2235–2241
- Ali T, Khan I, Simpson W, Prescott G, Townend J, Smith W, Macleod A (2007) Incidence and outcomes in acute kidney injury: a comprehensive population-based study. *J Am Soc Nephrol* 18:1292–1298
- Chawla LS, Bellomo R, Bihorac A, Goldstein SL, Siew ED, Bagshaw SM, Bittleman D, Cruz D, Endre Z, Fitzgerald RL, Forni L, Kane-Gill SL, Hoste E, Koyner J, Liu KD, Macedo E, Mehta R, Murray P, Nadim M, Ostermann M, Palevsky PM, Pannu N, Rosner M, Wald R, Zarbock A, Ronco C, Kellum JA, Acute Disease Quality Initiative Workgroup 16 (2017) Acute kidney disease and renal recovery: consensus report of the Acute Disease Quality Initiative (ADQI) 16 Workgroup. *Nat Rev Nephrol*. doi:10.1038/nrneph.2017.2
- Harel Z, Wald R, Bargman JM, Mamdani M, Etchells E, Garg AX, Ray JG, Luo J, Li P, Quinn RR, Forster A, Perl J, Bell CM (2013) Nephrologist follow-up improves all-cause mortality of severe acute kidney injury survivors. *Kidney Int* 83:901–908
- Hemmett J, Er L, Chiu HH, Cheung C, Djurdjev O, Levin A (2015) Time to revisit the problem of CIN? The low incidence of acute kidney injury with and without contrast in hospitalized patients: an observational cohort study. *Can J Kidney Health Dis* 2:38