



Inositol Pyrophosphates: Energetic, Omnipresent and Versatile Signalling Molecules

Akruti Shah^{1,2}, Shubhra Ganguli^{1,2}, Jayraj Sen^{1,2} and Rashna Bhandari^{1*}

Abstract | Inositol pyrophosphates (PP-IPs) are a class of energy-rich signalling molecules found in all eukaryotic cells. These are derivatives of inositol that contain one or more diphosphate (or pyrophosphate) groups in addition to monophosphates. The more abundant and best studied PP-IPs are diphosphoinositol pentakisphosphate (IP₇) and bis-diphosphoinositol tetrakisphosphate (IP₈). These molecules can influence protein function by two mechanisms: binding and pyrophosphorylation. The former involves the specific interaction of a particular inositol pyrophosphate with a binding site on a protein, while the latter is a unique attribute of inositol pyrophosphates, wherein the β-phosphate moiety is transferred from a PP-IP to a pre-phosphorylated serine residue in a protein to generate pyrophosphoserine. Both these events can result in changes in the target protein's activity, localisation or its interaction with other partners. As a consequence of their ubiquitous presence in all eukaryotic organisms and all cell types examined till date, and their ability to modify protein function, PP-IPs have been found to participate in a wide range of metabolic, developmental, and signalling pathways. This review highlights many of the known functions of PP-IPs in the context of their temporal and spatial distribution in eukaryotic cells.

Keywords: 5-Diphosphoinositol pentakisphosphate (IP₇), Bis-diphosphoinositol tetrakisphosphate (IP₈), Inositol hexakisphosphate (IP₆), IP₆ kinase (IP6K), Inositol phosphate

1 Introduction

Myo-inositol, a stereoisomer of cyclohexanehexol with one axial and five equatorial hydroxyl groups, is a component of biomolecules found in all forms of life.¹ In eukaryotic cells, the phosphorylated derivatives of *myo*-inositol include lipid phosphatidyl inositols and water-soluble inositol polyphosphates.¹ Inositol hexakisphosphate (IP₆), at a concentration range of 10–100 μM in yeast and animal cells and 500 μM in slime moulds, is the most abundant inositol polyphosphate in eukaryotes.^{2, 3} In the early 1990s, a distinct subclass of inositol polyphosphates containing 'high energy' pyrophosphate groups was identified in slime mould and mammalian cells.^{4–6} Since then several studies have characterized these inositol pyrophosphates (PP-IPs) and their functions (for

reviews see^{2, 7–23}). The major PP-IPs, diphosphoinositol pentakisphosphate, PP-IP₅ (or IP₇) and bis-diphosphoinositol tetrakisphosphate, [PP]₂-IP₄ (or IP₈), are derived by the addition of phosphate groups to pre-existing monophosphates on IP₆ (Fig. 1). IP₇ is the most abundant PP-IP, and its concentration ranges from 0.5 to 1.3 μM in yeast and mammalian cells.² IP₈ is present at much lower levels in most organisms, ranging from undetectable to approximately 50% of IP₇ levels in budding yeast²⁴ and some mammalian cell lines.^{2, 25}

The pathway of synthesis of inositol polyphosphates has been characterized in yeast, slime moulds, plants and animals. The simplest anabolic pathway, characterized in the yeast *Saccharomyces cerevisiae* (Fig. 1), involves the release

A. Shah, S. Ganguli and J. Sen contributed equally.
¹ Laboratory of Cell Signalling, Centre for DNA Fingerprinting and Diagnostics, Hyderabad, Telangana, India. ² Graduate Studies, Manipal University, Manipal, Karnataka, India.
*rashna@cdfd.org.in

REVIEW
ARTICLE

of $I(1,4,5)P_3$ from $PI(4,5)P_2$ by phospholipase C (PLC), subsequent phosphorylation of IP_3 to IP_4 and IP_5 by inositol polyphosphate multikinase (IPMK or *Ipk2*), and the conversion of IP_5 to IP_6 by the IP_5 2-kinase, *Ipk1*.²⁶ Inositol pyrophosphates are synthesized by two classes of enzymes, IP_6 kinases and PP- IP_5 kinases. IP_6 kinases, identified by the Snyder group, convert IP_6 to 5PP- IP_5 (also called 5- IP_7).²⁷ *S. cerevisiae* have a single IP_6 kinase, called *Kcs1*, whereas mammals have three isoforms, *IP6K1*, *IP6K2* and *IP6K3*. PP- IP_5 kinases, identified independently by the York and Shears laboratories, convert 5PP- IP_5 to 1,5[PP]₂- IP_4 (IP_8).^{28, 29} These kinases can also synthesize an alternative form of IP_7 (1PP- IP_5 , also called 1- IP_7) from IP_6 .^{30, 31} Yeast has a single PP- IP_5 kinase (*Vip1* in *S. cerevisiae*), and mammals have two

isoforms, *PPIP5K1* and *PPIP5K2*.²⁹ 1- IP_7 is the minor IP_7 isoform under normal conditions.^{2, 32} PP- IP_4 and [PP]₂- IP_3 are derived from IP_5 by the action of IP_6 kinases and PP- IP_5 kinases.^{2, 33} Diphosphoinositol polyphosphate phosphohydrolase, or DIPP (*Ddp1* in *S. cerevisiae*) hydrolyzes the pyrophosphate moiety of inositol pyrophosphates, rapidly degrading IP_8 to IP_6 ³⁴ (Fig. 1). Interestingly, the *PPIP5Ks* also possess a phosphatase domain which selectively cleaves the 1-position β -phosphate of 1- IP_7 and IP_8 , thus targeting the products of the kinase domain.^{35, 36} A recent study identified another yeast phosphatase, *Siw14*, that specifically targets the 5-position β -phosphate of PP- IPs ³⁷ (Fig. 1).

As PP- IPs are found in all eukaryotic organisms, they display many conserved and divergent




Figure 1: The pathway of synthesis of inositol pyrophosphates. The simplest pathway in yeast starts with the formation of IP_3 from $PI(4,5)P_2$ by the action of phospholipase C (PLC). Subsequent phosphorylation by *Ipk2* (IPMK in mammals) converts IP_3 to IP_4 and IP_5 . *Ipk1* (IP_5 -2K in mammals) converts IP_5 to IP_6 . *Kcs1* (IP6Ks in mammals) phosphorylates IP_6 to 5PP- IP_5 (or 5- IP_7). *Vip1* (PPIP5Ks in mammals) acts on IP_6 to form 1PP- IP_5 (or 1- IP_7) and on 5- IP_7 to form 1,5[PP]₂- IP_4 (or IP_8). *Kcs1* can also convert 1- IP_7 to IP_8 . 5PP- IP_4 and 1,5[PP]₂- IP_3 are synthesised from IP_5 by the action of *Kcs1* and *Vip1*. IP_6 kinases prefer IP_6 over IP_5 due to their higher affinity towards the former.¹⁰² In yeast, the minor inositol pyrophosphates, 5PP- IP_4 and 1,5[PP]₂- IP_3 are detected only on deletion of *Ipk1*.³⁰ DIPP (diphosphoinositol polyphosphate phosphohydrolase), which has five isoforms in mammals and a single isoform, *Ddp1*, in yeast, hydrolyses diphosphate groups on IP_7 and IP_8 to form IP_6 , and on PP- IP_4 and [PP]₂- IP_3 to form IP_5 .^{2, 131} *Siw14*, an inositol pyrophosphate phosphatase in yeast, preferentially cleaves the C5 β -phosphate on PP- IPs .³⁷ The yeast enzymes are depicted in purple, and mammalian enzymes are depicted in green and are bracketed. The undetermined inositol pyrophosphate structure is represented with an interrogation mark. *Myo*-inositol contains five equatorial (parallel to the axis) and one axial (perpendicular to the axis) hydroxyl groups. Carbon atoms on the *myo*-inositol ring are numbered on the structures of $PI(4,5)P_2$ and IP_6 .

functions in yeast, plants and mammals. These small molecules participate in a wide array of cellular and organismal processes ranging from apoptosis and DNA repair to energy homeostasis and blood clotting. Inositol pyrophosphates regulate protein function via two molecular mechanisms, (a) protein binding and (b) protein pyrophosphorylation (Fig. 2). The inositol pyrophosphates 1-IP₇ and 5-IP₇ show isomer-specific binding to proteins to regulate their function.^{38–41} Conversely, any inositol pyrophosphate may act as a phosphate donor, transferring its β-phosphate to a pre-phosphorylated serine to form pyrophosphoserine, bringing about protein pyrophosphorylation^{42, 43} (Fig. 2). Pyrophosphorylation is an enzyme-independent reaction, requiring only the inositol pyrophosphate donor, the pre-phosphorylated protein acceptor, and divalent cations such as Mg²⁺. The acceptor serines are pre-phosphorylated by a protein kinase, usually CK1 or CK2, and occur in acidic serine sequence motifs, i.e. one or more Ser with neighbouring Glu/Asp residues, i.e. one or more Ser with neighbouring Glu/Asp residues. IP₇-mediated pyrophosphorylation of specific proteins can regulate glycolysis and rRNA synthesis in yeast,^{44, 45} and viral particle release and dynein motor driven retrograde trafficking in mammalian cells.^{46, 47} Specific serine phosphatases such as PP1 and PP2C cannot remove the pyrophosphate group,^{42, 48} but

alkaline phosphatase can dephosphorylate serine,⁴⁸ suggesting that pyrophosphorylation is a reversible modification with many potential roles in cell signalling.

2 Temporal Regulation of Inositol Pyrophosphates

2.1 Detection and Measurement of PP-IPs

The biochemical determination of PP-IP levels presents several technical challenges, and till date there are no antibody-based or colourimetry-based assays to detect these molecules. Most measurements have relied on monitoring radiolabelled [³H]inositol or [³²P]phosphate after strong anion exchange HPLC-based resolution to separate individual inositol phosphates. The Mayr laboratory developed a post-HPLC column derivatization method that allows spectrophotometric detection of unlabelled inositol polyphosphates,⁴⁹ and was also able to resolve 1-IP₇ and 5-IP₇.³⁰ A new method reported recently employs a monosaccharide binding resin (CarboPac™) to resolve 1-IP₇ and 5-IP₇ and has estimated that 1-IP₇ constitutes less than 2% of total IP₇ in a human colon cancer cell line.³² All HPLC-based resolution methods however require deproteinization of the sample using perchloric acid or




Figure 2: Mechanism of regulation of protein function by PP-IPs. PP-IPs modulate protein function by two mechanisms: **a** direct binding to proteins in which the positively charged binding pockets formed by Lys/Arg residues (shown in *blue*) electrostatically neutralise the high negative charge of the PP-IP molecule,¹²⁵ and **b** protein pyrophosphorylation, which involves the non-enzymatic transfer of the β-phosphate from a PP-IP to a pre-phosphorylated Ser residue (shown in *mauve*) surrounded by Asp/Glu residues (shown in *red*).

trichloroacetic acid, and employ acidic buffers to elute inositol phosphates from chromatography columns. These acidic conditions lead to hydrolysis of the β -phosphate moiety of PP-IPs, and their likely underestimation in cells and tissues.⁵⁰ A simple method developed by the Saiardi laboratory employing high-percentage polyacrylamide gels to resolve IP₆, IP₇, and IP₈ avoids the use of low pH buffers, thereby improving the efficiency and ease of PP-IP detection. The sensitivity of PP-IP detection was also improved by staining gels with the fluorescent dye 4',6-diamidino-2-phenylindole (DAPI),⁵⁰ and by using titanium oxide (TiO₂) beads to enrich for these phosphate containing molecules in cell and tissue extracts.²⁵ These newer methods have suggested that earlier estimates of PP-IP levels may be off the mark.³ Nevertheless, while studies over the past twenty years employing HPLC methods may have underestimated the absolute levels of PP-IPs, the information on an increase or decrease in the concentration of these molecules, or changes in enzyme activity, is likely to be accurate. This section therefore presents a summary of our current knowledge of the temporal changes in PP-IP levels under different conditions.

While classical second messenger molecules like cAMP display acute changes in response to extracellular signals, most measurements suggest that PP-IP levels change only marginally under certain conditions. The levels of PP-IPs are tightly regulated and these molecules display a high turnover rate, with studies showing that the IP₇ pool can turn over ten times every 40 min,⁵¹ suggesting that pathways for the synthesis and utilisation of IP₇ are constantly active in the cell. Several studies show that IP₇ or IP₈ levels vary in response to physiological stresses, during different phases of the cell cycle, and over the course of development and ageing.

2.2 PP-IPs Fluctuate in Response to Stress

IP₇ and IP₈ are involved in the cell's response to different physiological stresses in mammals, yeast, and plants. In mammalian cells, hyper-osmotic stress can trigger an acute 25-fold increase in the level of IP₈,⁵² and thermal stress can increase IP₈ levels 3–4 fold.⁵³ PPIP5K1, one of the mammalian enzymes responsible for IP₈ synthesis from 5-IP₇, is activated fourfold upon osmotic stress.²⁹ Initial studies using protein kinase inhibitors suggested that the ERK/MEK kinase pathway was responsible for these changes in IP₈,^{52, 53} but later data revealed that these kinase inhibitors actually

acted via off-target effects on the cellular AMP/ATP ratio⁵⁴ (discussed later). Exposure of cells to apoptosis inducing drugs such as cisplatin or staurosporine has also been shown to elevate the levels of IP₇ and to a lesser extent IP₈⁵⁵ (discussed in detail later). Interestingly, exposure of neutrophils to nicotine or cigarette-smoke extract decreases IP₇ levels, suggesting that IP₇ plays a role in the pathogenesis of tobacco-induced chronic obstructive pulmonary disease.⁵⁶

In budding yeast, the effect of phosphate starvation on inositol pyrophosphate levels has yielded conflicting results. One study observed an increase in cellular IP₇ concentration during limiting extracellular phosphate levels,⁵⁷ whereas others revealed a decrease in IP₇ levels following phosphate starvation.^{58, 59} PP-IP levels in yeast decrease rapidly upon treatment with hydrogen peroxide, as a consequence of inhibition of Kcs1 enzyme activity via oxidation of a cysteine residue.²⁴ Yeast lacking Kcs1 demonstrate reduced cell death upon exposure to H₂O₂²⁴ but are sensitive to several other stresses, including osmotic and thermal stresses.⁶⁰ Yeast carrying deletions for both *kcs1* and *vip1* genes have no PP-IPs and show no changes in transcription in response to osmotic, heat or oxidative stress, suggesting that PP-IPs are required for a cell to adapt in order to survive these stresses.⁶¹

Plant seeds are the most abundant source of IP₆ (also known as phytic acid), which serves as a store for phosphorus, inositol, and cations required by the seedling during germination. Recently, two groups independently demonstrated the presence of IP₇ and IP₈ in plant seeds and vegetative tissues.^{62, 63} These studies also identified two Vip1 orthologues in Arabidopsis that catalyse the formation of IP₈ from IP₇. Although no IP6K orthologue has been identified in plants, it was speculated that an unknown enzyme activity is responsible for plant IP₇ synthesis.⁶³ It is also likely that the IP₇ and IP₈ isomers found in plants are different from those occurring in yeast and mammals.⁶³ The plant stress hormone abscisic acid was shown to cause a twofold increase in the levels of both IP₇ and IP₈, whereas treatment with the plant defence hormone **jasmonate** led to a sustained twofold increase in IP₈ but did not affect IP₇ levels.⁶³ Further probing the specific function of IP₈ in jasmonate signalling, this study suggested that IP₈ binds the F-box protein COI1 which is part of an E3 ligase complex responsible for proteasomal degradation of the transcriptional repressor protein JAZ. The combined binding of jasmonate and IP₈ to the COI1-JAZ jasmonate coreceptor

Jasmonate: A lipid based hormone that regulates plant growth and is required for plant defence responses against herbivores and environmental stress

complex facilitates JAZ degradation, thereby permitting the expression of jasmonate responsive genes involved in plant defences against insect herbivores and fungal pathogens.

2.3 PP-IP Changes During the Cell Cycle

PP-IP levels have also been shown to fluctuate during the cell cycle in both yeast and mammals. *S. cerevisiae* cells can be arrested in the G1 phase of the cell cycle by treatment with the α -factor mating pheromone.⁶⁴ IP₇ and IP₈ levels increase by approximately twofold between 30 to 60 min after release from this arrest, in the time that corresponds to synchronised entry of the cells into the S phase.⁶⁵ The levels of the PP-IPs decrease again during the G2/M phase. The activity of Kcs1 was shown to mirror these changes in IP₇ and IP₈ levels, suggesting that signalling pathways operating during the cell cycle may post-translationally modify Kcs1 to regulate its activity.⁶⁵ Mammalian cells can be arrested in the G₀ phase of the cell cycle by allowing them to grow to confluence, and they can be arrested in early mitosis by treatment with nocodazole, which interferes with microtubule polymerization and prevents the formation of the mitotic spindle.⁶⁶ When rat mammary tumour cells were synchronised by both these methods, IP₇ was found to be twice as high in the G1 phase as compared with the other phases of the cell cycle where the basal levels of this PP-IP are approximately 0.6 μ M.⁶⁷ The functional significance of these changes in PP-IP levels during the cell cycle is still unclear, but it is likely that one or more cell cycle modulating factors may be regulated by IP₇ binding or pyrophosphorylation.

2.4 PP-IPs in Development and Ageing

Unlike most other eukaryotes, the slime moulds *Dictyostelium discoideum* and *Polysphondylium pallidum* display an abundance of PP-IPs, and are also the only documented organisms in which IP₈ is more abundant than IP₇.^{3, 68} Recent measurements using TiO₂ beads and polyacrylamide gel electrophoresis estimate that vegetative stage *D. discoideum* contain IP₇ at 60 μ M and IP₈ at 180 μ M,³ and that IP₈ levels go up approximately threefold during starvation induced development, when these amoebae aggregate, form a multicellular “slug”, and eventually develop into a fruiting body. The chemoattractant cAMP released by *D. discoideum* during starvation has been shown to lead to a rapid and sustained three- to fourfold increase in IP₇ and IP₈ levels.⁶⁹ IP₇ competes with the lipid inositide PI(3,4,5)P₃

to bind the PH domain of the protein Crac, and interferes with Crac translocation to the plasma membrane. As Crac translocation is required for cAMP-dependent chemotaxis, it was postulated that IP₇ is a negative regulator of chemotaxis, modulating the sensitivity of cells to cAMP stimulation.

IP₇ levels have been shown to increase in mice as they age.³⁹ While both IP₆ and IP₇ are higher in hepatocytes derived from 10-month-old as compared with 2-month-old mice, the IP₇/IP₆ ratio increases more than twofold in older mice, and correlates with a decrease in insulin sensitivity. Like *D. discoideum* Crac, the PH domain of mammalian AKT can bind IP₇, which competes with PI(3,4,5)P₃ binding to inhibit membrane translocation and activation of AKT in response to insulin. *Ip6k1*^{-/-} mice with reduced levels of IP₇ and upregulated AKT signalling are thus more insulin sensitive as they age. Another study demonstrated increased IP₇ production in ageing bone marrow derived mesenchymal stem cells as compared to young cells.⁷⁰ Consequently, there is decreased AKT phosphorylation and activation in older cells, leading to increased sensitivity to hypoxic injury with age.

3 The Functions of PP-IPs in Different Cellular Compartments

Technical limitations imposed by currently available methods for the detection of PP-IPs have translated into the lack of any convincing data on the subcellular compartmentalisation of PP-IPs. However, there are several studies on the tissue distribution and subcellular localisation of the IP6Ks and PPIP5Ks, which suggest that IP₇ and IP₈ are likely to be found in a variety of tissues and in all cell compartments. In *S. cerevisiae*, both PP-IP kinases, Kcs1 and Vip1 are located predominantly in the cytoplasm (<http://yeastgfp.yeast-genome.org/>), but several studies have revealed important functions for PP-IPs in various subcellular compartments including the nucleus and vacuoles. In mice, *Ip6k1* and *Ip6k2* mRNA are expressed at varying levels in all tissues,²⁷ whereas IP6K3 is highly expressed in the cerebellum.^{71, 72} *Ip6k1* mRNA shows highest expression in testes,²⁷ correlating with spermatogenesis failure observed in *Ip6k1* knockout mice.⁷³ Although protein overexpression studies show IP6K1 located in the cytoplasm and nucleus, IP6K2 mainly in the nucleus, and IP6K3 localised predominantly in the cytoplasm,⁷¹ such analyses are plagued with anomalies arising from high levels of unregulated protein expression, and often do not reflect the

Chemotaxis: Phenomenon of movement of a cell or an organism in response to a chemical stimulus.

behaviour of the endogenous protein. *Ppip5k1* mRNA is expressed in many human tissues, with greater abundance observed in skeletal muscle, heart and brain.⁷⁴ Overexpressed PPIP5K1 and PPIP5K2 localise mainly to the cytoplasm,^{29, 74} but PPIP5K2 has a nuclear localisation signal that is absent in PPIP5K1, enabling its translocation to the nucleus in a phosphorylation-dependent manner.⁷⁵ Both the PPIP5K isoforms possess a lipid inositide binding domain distinct from their kinase domains, and agonist-stimulated production of PI(3,4,5)P₃ can lead to translocation of PPIP5K1 from the cytoplasm to the plasma membrane.^{41, 76}

As suggested by the ubiquitous expression and localisation of the kinases responsible for their synthesis, PP-IPs have been shown to participate in a myriad functions in many different tissues and subcellular compartments. This section describes several functions of these small molecules in different locations within a cell (Fig. 3).

3.1 Plasma Membrane

One of the most intriguing yet controversial functions of IP₇ is its ability to compete with the lipid inositide PI(3,4,5)P₃ for binding to PH domains in specific proteins, and thereby displacing these proteins from the plasma membrane. The first demonstration that IP₇ can compete with PI(3,4,5)P₃ to bind a PH domain was in *Dictyostelium*⁶⁹ (discussed earlier). This study also demonstrated that IP₇ binds the PH domain of the mammalian kinase AKT with an affinity comparable to I(1,3,4,5)P₄, which mimics the PI(3,4,5)P₃ head group. A subsequent study went on to show that the IP₇-AKT interaction is physiologically relevant, as *Ip6k1*^{-/-} mice show higher AKT activity, leading to increased insulin sensitivity and reduced weight gain in response to a high-fat diet.³⁹ It has also been shown that enhanced PI(3,4,5)P₃-dependent AKT activation in neutrophils derived from *Ip6k1*^{-/-} mice leads to greater phagocytic and bactericidal ability in these cells.⁴⁰ Although one report showed that




Figure 3: Representation of cellular functions of inositol pyrophosphates in an eukaryotic cell. **1** IP₇ physiologically inhibits AKT signalling by competitively binding to the PH domain of AKT and thus preventing it from binding to PI(3,4,5)P₃ (PIP3). IP₇ and IP6K2 promote, **2** apoptosis and **3** formation of autophagosomes. **4** PP-IPs regulate the cellular levels of ATP by their action on the glycolysis pathway and mitochondrial membrane potential. **5** IP₇ is responsible for maintaining endoplasmic reticulum morphology in yeast cells. **6** IP₇-mediated pyrophosphorylation regulates dynein binding to membranes and thereby influences Golgi morphology. **7** PP-IPs positively regulate the synthesis of polyphosphates in yeast vacuoles. **8** In pancreatic β cells, IP₇ upregulates insulin secretion by increasing the readily releasable pool of insulin granules docked at the plasma membrane. **9** IP₇ inhibits the synaptic exocytotic pathway in neurons. In the nucleus, PP-IPs are responsible for **10** telomere length maintenance in yeast, **11** DNA repair via the homologous recombination (HR) and nucleotide excision repair (NER) pathways, and **12** epigenetic modifications that influence chromatin remodelling to control global transcription. **13** In yeast, IP₇ influences ribosome biogenesis by regulating nucleolar rDNA transcription.

IP₇ is not able to displace PI(3,4,5)P₃ from AKT in an in vitro binding assay,⁷⁷ it was later demonstrated that non-hydrolysable analogues of 5-IP₇ or 1-IP₇ can inhibit AKT activation even in the presence of PI(3,4,5)P₃.⁷⁸ A recent study revealed that chemically synthesized 5-IP₇ released into mammalian cells is able to induce translocation of AKT from the plasma membrane to the cytoplasm.⁷⁹ It has been shown that AKT and other PH domains bind 5-IP₇ and IP₆ with higher affinity as compared with 1-IP₇ or IP₈.⁴¹ This study also suggested that PI(3,4,5)P₃ dependent recruitment of PIP5K1 to the plasma membrane would deplete subplasmalemmal 5-IP₇ and IP₆ by using them as substrates, thus providing positive feedback to PI(3,4,5)P₃ binding by PH domains at the plasma membrane. Although the structural conundrum of how the AKT PH domain which specifically binds PI(3,4,5)P₃ over PI(4,5)P₂ can bind 5-IP₇ better than IP₆ remains unresolved, it is likely that binding of specific PP-IPs to different PH domains is a conserved mechanism by which these molecules regulate signalling at the plasma membrane.

3.2 Cytoplasm

Studies in yeast and mammalian cells have revealed many functions for PP-IPs in the cytoplasm, including key roles in energy metabolism, apoptosis and autophagy.

3.2.1 Energy Metabolism PP-IPs have been described as ‘energy sensors’ and ‘metabolic messengers’. This is because PP-IP levels can both reflect and regulate the cellular levels of ATP. The IP6Ks have a very low affinity for ATP, with a Km of approximately 1 mM, which lies within the range of cellular ATP concentrations. Consequently, fluctuations in ATP levels correlate with changes in the intracellular concentration of IP₇. Lowering the cellular ATP concentration by treatment with sodium azide, oligomycin, or certain kinase inhibitors has been shown to significantly reduce the levels of PP-IPs.^{54, 80} Interestingly, at a low ATP/ADP ratio, IP6Ks can switch to being ADP phosphotransferases, transferring the 1-phosphate from IP₆ to ADP to generate I(2,3,4,5,6)P₅,⁸¹ depleting cytosolic IP₆ and perhaps further lowering IP₇ synthesis. It is possible that the dual enzymatic activity of IP6Ks allows them to function as cellular adenylate energy sensors, converting IP₆ to IP₇ or IP₅ under high or low energy conditions respectively, so that these products may transduce information on the cellular energy

status to regulate different metabolic and signalling pathways.

PP-IPs in turn affect ATP levels by regulating glycolysis. *S. cerevisiae* lacking Kcs1 have a higher cellular ATP concentration than wild-type yeast.⁴⁴ It was shown that IP₇ produced by Kcs1 pyrophosphorylates the major glycolytic transcription factor Gcr1, which is known to interact with Gcr2. In *kcs1Δ* cells, reduced pyrophosphorylation of Gcr1 increases its interaction with Gcr2, thereby increasing **glycolytic flux**.

In mammals, IP6K1 has been shown to influence fat accumulation by regulating adipocyte energy metabolism.⁸² The AMP-activated protein kinase (AMPK) signalling pathway augments the transformation of white adipose tissue to beige, enhancing energy expenditure in the form of heat, and correlating inversely with type II diabetes and fat-induced obesity. IP₆ causes stimulatory phosphorylation of AMPK, thereby activating the pathway that leads to browning of white adipose tissue. Specific deletion of IP6K1 in mouse adipocytes led to increased thermogenic energy expenditure in these cells, presumably due to increased availability of IP₆. It was suggested that this phenomenon, coupled with increased insulin sensitivity due to AKT activation (described earlier), leads to reduced weight gain when *Ip6k1*^{-/-} mice are provided a high-fat diet.

3.2.2 Apoptosis 5-IP₇ produced by IP6K2 has been shown to upregulate apoptotic signalling pathways in many mammalian cells and tissues. Overexpression of either of the three IP6K isoforms in various cancer cell lines under normal as well as stress conditions leads to increased cell death.^{55, 83, 84} However, siRNA mediated depletion of only IP6K2 and not IP6K1 or IP6K3 abrogated cell death, suggesting that IP6K2 is the only isoform that is physiologically capable of inducing apoptosis.^{55, 85} Work by the Lindner group revealed that overexpressed IP6K2 in untreated cancerous or non-cancerous cells is mainly cytoplasmic and upon apoptotic induction it translocates to the nucleus.⁸⁴ IP6K2 has a nuclear localization signal, the removal of which keeps the protein in the cytoplasm, lowering its pro-apoptotic effect. The Snyder group showed that overexpressed IP6K2 translocates to the mitochondria upon cytotoxic stress and co-localizes with the pro-apoptotic protein Bax.⁵⁵ Overexpression of catalytically inactive IP6K2 did not promote stress-induced apoptosis, indicating that the pro-apoptotic function of IP6K2 relies on IP₇ synthesis. Indeed, the levels of IP₇ were found

Glycolytic flux: Rate at which molecules proceed through the glycolysis pathway.

Apoptosis: Regulated or programmed cell death seen in multicellular organisms.

to be higher in cells undergoing cytotoxic stress as compared with untreated cells. IP6K2 has been shown to promote apoptosis by acting on several signalling pathways. IP₇ binding to the PH domain of AKT inhibits AKT-dependent prosurvival signalling,⁶⁹ indirectly enhancing apoptosis.^{55, 86} IP6K2 also influences apoptotic signalling via the transcription factor p53, which is known to activate both pro- and anti-apoptotic genes. Kinase activity independent direct binding of IP6K2 to p53 augments apoptosis by downregulating expression of p21, which is responsible for stressed cells choosing cell cycle arrest instead of apoptosis.⁸⁷ IP6K2 also binds TTI1, a protein that forms part of the TTT co-chaperone complex that stabilises the protein kinases DNAPKcs and ATM.⁸⁸ IP₇ synthesized by IP6K2 binds the protein kinase CK2 and promotes its phosphorylation of the TTT complex, which in turn stabilises DNAPKcs and ATM, leading to increased phosphorylation of p53 and activation of cell death. In another study, IP6K2 binding to TRAF2 (TNF receptor associated factor) prevents TAK1 (transforming growth factor β -activated kinase) phosphorylation and subsequent NF κ B activation, sensitizing cells to TNF- α induced apoptosis.⁸⁹ The heat shock protein HSP90 has been shown to bind IP6K2 and inhibit its catalytic activity, thereby protecting cells from apoptosis and augmenting cell survival.⁹⁰ The importance of IP6K2 in promoting apoptosis is also reflected in phenotypes observed in *Ip6k2* knockout mice.⁹¹ Chronic exposure to the UV mimetic carcinogen 4-nitroquinoline 1-oxide (4NQO) led to a fourfold increased incidence of invasive squamous cell carcinoma of the oral cavity and oesophagus in *Ip6k2*^{-/-} mice as compared with their wild-type littermates. This was attributed at least in part to resistance to cell death in tissues lacking IP6K2. Whole genome expression profiling of tissues from *Ip6k2*^{-/-} as compared with *Ip6k2*^{+/+} mice showed upregulation of certain oncogenes and downregulation of tumour suppressor genes, suggesting that other pathways regulated by IP6K2 also impinge upon survival versus death signals to influence carcinogenesis.

3.2.3 Autophagy IP6Ks in yeast and mammals have also been shown to participate in autophagy, a process by which cytoplasmic proteins and organelles are degraded. Autophagy is initiated with the formation of the phagophore, a double membrane that begins to engulf macromolecules and organelles in the cytoplasm.⁹² The phagophore membrane closes to form an autophagosome,

which fuses with a lysosome, leading to degradation and recycling of the engulfed components. Budding yeast deficient in Kcs1 display undetectable levels of PP-IPs and exhibit a reduction in the number and size of autophagosomes formed upon nitrogen starvation.⁹³ *kcs1 Δ* yeast show mislocalisation of phagophore assembly sites to the plasma membrane, suggesting that Kcs1 is required for normal autophagosome formation. Overexpression of IP6Ks in mammalian cells led to an increase in the number of stress-induced autophagosomes as compared with control cells, and reduced expression of IP6Ks using RNA interference suppressed autophagosome formation.⁹⁴ Expression of an inactive IP6K had no effect, revealing that autophagosome formation is IP₇ dependent. The evolutionarily conserved protein kinase mTOR is a master regulator of cell growth and metabolism, and negatively regulates autophagy. In human cells, IP6K expression levels showed an inverse correlation with mTOR activity, suggesting that IP₇ may promote autophagy via the mTOR signalling pathway.⁹⁴

3.3 Membrane Bound Organelles

PP-IPs are not membrane permeant, and there is little evidence to suggest that PP-IP synthesising enzymes are localised within membrane-bound organelles in mammalian cells. However, changes in intracellular PP-IP levels have been shown to impact normal function in several membrane bound organelles, suggesting that these molecules may be synthesized within these organelles, may act on the cytoplasmic face of the membrane, or may be transported across membranes by yet unknown mechanisms.

3.3.1 Mitochondria Yeast devoid of Kcs1 and mouse embryonic fibroblasts (MEFs) lacking IP6K1 have dysfunctional mitochondria, which display a significantly lower oxygen consumption as compared with their wild-type counterparts.⁴⁴ Without IP6K1, mitochondria are unable to generate the membrane potential required for ATP production, and are also deficient in the respiratory chain protein cytochrome c oxidase. Despite these mitochondrial defects, *kcs1 Δ* yeast show higher ATP levels because of increased glycolytic flux (discussed earlier). However, *kcs1 Δ* yeast shows a decreased growth rate as compared with wild-type cells, which could be partly attributed to defects in the bio-synthesis of major macromolecules like fatty acids and nucleotides that depend on enzymes of the mitochondria.⁴⁴

3.3.2 Endoplasmic Reticulum The endoplasmic reticulum (ER) is generally nondescript and parallel to the plasma membrane in wild-type budding yeast, whereas in *kcs1Δ* cells, it appears hypertrophied and perpendicular to the plasma membrane.⁹⁵ This suggests that PP-IPs have a role to play in ER function. However, there was no apparent defect in the secretion of newly synthesised proteins into the medium by *kcs1Δ* yeast, suggesting that ER functions remain largely unaffected in the absence of significant levels of PP-IPs. There are no studies examining whether PP-IPs influence ER structure or function in mammalian cells.

3.3.3 Golgi Our recent studies show that *Ip6k1^{-/-}* MEFs display a fragmented Golgi morphology when compared with *Ip6k1^{+/+}* MEFs.⁴⁷ This phenotype was restored by the expression of catalytically active IP6K1 but not by the inactive protein, suggesting that PP-IPs are essential to maintain Golgi morphology. The pericentriolar position of the Golgi apparatus in mammalian cells is dependent on the cytoplasmic motor protein dynein, and the Golgi appears fragmented if dynein function is compromised.⁹⁶ Further analysis showed that PP-IPs regulate dynein binding to membranes, and thereby influence Golgi morphology (discussed later).

3.3.4 Lysosomes and Related Organelles The number and distribution of lysosomes marked by the protein LAMP2 (lysosome associated membrane protein 2) appears unaltered in *Ip6k1^{-/-}* MEFs (Chanduri and Bhandari, unpublished). However, PP-IPs have an important function in yeast vacuoles, which are lysosome-like organelles.⁹⁷ Yeast vacuoles are the site of synthesis and storage of inorganic polyphosphates (polyP), which are linear chains of orthophosphate moieties linked via phosphoanhydride bonds.⁹⁸ PolyP are found in all life forms and are involved in diverse physiological functions like energy metabolism, transcription regulation, and blood clotting.⁹⁸ An intriguing link was established between PP-IPs and polyP, when it was observed that *kcs1Δ* yeast also have substantially reduced levels of polyP.⁹⁹ Our work showed that the link between PP-IPs and polyP levels is also conserved in mammals.¹⁰⁰ *Ip6k1^{-/-}* mice have reduced levels of polyP in platelet dense granules, which are lysosome-related organelles. Consequently, these mice display delayed clotting time and altered clot architecture, making them less susceptible to

thromboembolism. Elegant structural and biochemical studies from the Mayer laboratory uncovered the molecular link between PP-IPs and polyP. The first study identified that the yeast vacuolar transport chaperone (VTC) complex generates polyP from ATP, with the subunit VTC4 responsible for enzyme activity.¹⁰¹ A recent study determined that multiple subunits of the VTC complex, including VTC4, contain an SPX domain, which has a positively charged surface that can specifically bind inositol polyphosphates.⁵⁹ Binding of 5-IP₇ to this domain enhances VTC-dependent polyP synthesis in isolated vacuoles, suggesting that PP-IPs modulate polyP synthesis by allosteric regulation of the polyP polymerase. As the levels of 5-IP₇ have been shown to decrease upon phosphate starvation in yeast,^{58, 59} it was speculated that PP-IPs may act as sensors of inorganic phosphate (Pi) levels in the cytoplasm and communicate Pi fluctuations to SPX domains, which then interact with various proteins to regulate Pi uptake, transport and storage.⁵⁹

3.3.5 Vesicles Several studies have investigated the role of PP-IPs in vesicle trafficking processes. Budding yeast lacking *Kcs1* show abnormally small and fragmented vacuoles,¹⁰² which reflect a defect in the endocytosis pathway.⁹⁵ In these cells, the endosomes formed upon endocytosis fail to fuse with vacuoles and accumulate as large multilamellar endosomal intermediates. The exact mechanism by which PP-IPs regulate endocytosis remains unclear. PP-IPs synthesized by IP6K1 are essential to ensure normal insulin secretion in mammals. *Ip6k1^{-/-}* mice have lower serum insulin levels as compared with their wild-type littermates,⁷³ and in pancreatic β cells, 5-IP₇ synthesized by IP6K1 upregulates insulin secretion.¹⁰³ 5-IP₇ increases the readily releasable pool of insulin containing granules which are docked at the plasma membrane, ready for the first phase of exocytosis upon glucose uptake by β cells.¹⁰³ IP6K1 can also influence synaptic vesicle exocytosis. IP6K1 acts independent of its catalytic activity to bind GRAB, a guanine-nucleotide exchange factor (GEF) for the small G-protein Rab3A, which is a negative regulator of synaptic vesicle exocytosis.¹⁰⁴ IP6K1 binding to GRAB inhibits the activity of Rab3A to augment neurotransmitter release from synaptic vesicles. In contrast, a recent study showed that 5-IP₇ synthesized by IP6K1 can inhibit the synaptic exocytotic pathway by interacting with synaptotagmin, which is a calcium-sensing

Thromboembolism: Obstruction of a blood vessel by a blood clot.

Metastasis: Spread of cancer cells from one part of the body to another part via the blood or lymphatic system.

protein in the synaptic vesicle membrane and promotes vesicle fusion.¹⁰⁵ 5-IP₇ (but not 1-IP₇ or IP₆) binds and restrains synaptotagmin in a fusion-incompetent conformation to inhibit neurotransmitter release. These contrasting studies suggest that PP-IPs and their kinases can regulate exocytosis via multiple non-overlapping pathways in different cell types and tissues.

In addition to regulating vesicle fusion events, PP-IPs have also been shown to influence vesicle transport along microtubules. In the first demonstration of how serine pyrophosphorylation can regulate cellular processes, the Saiardi laboratory showed that pyrophosphorylation of the β subunit of the adaptor protein complex AP3 can regulate the release of HIV-1 virus like particles from mammalian cells.⁴⁶ 5-IP₇-mediated pyrophosphorylation of AP3B1 was shown to inhibit its interaction with the microtubule plus end-directed kinesin motor protein Kif3A, thereby lowering kinesin-dependent intracellular trafficking of HIV1 virus like-particles. Recent work from our laboratory has shown that 5-IP₇-mediated protein pyrophosphorylation can also regulate dynein-motor driven microtubule minus-end directed vesicle trafficking.⁴⁷ We found that pyrophosphorylation of Ser51 on the dynein-intermediate chain (IC) is important for its interaction with the p150^{Glued} subunit of the dynactin complex, and is required to recruit dynein to vesicles. *Ip6k1*^{-/-} MEFs have reduced interaction between dynein IC and p150^{Glued}, and as a consequence have defects in several dynein-dependent trafficking processes, including endosomal sorting of transferrin and Golgi morphology.

Dysplasia: Enlargement of an organ or a tissue due to excessive proliferation of cells of an abnormal type.

Cytoskeleton: A network of protein filaments and tubules in the cytoplasm of cells, which gives them shape.

3.4 Actin Cytoskeleton

The role of PP-IPs in actin cytoskeleton regulation was first identified by studies in the fission yeast, *Schizosaccharomyces pombe*, which possesses a Vip1 orthologue named Asp1.²⁸ *Asp* (*arp*, *sop*, *profilin* interactor) was originally identified as a high-copy suppressor of a mutation in *Arp3*, which is part of the actin-related protein complex Arp2/3, an essential component of the actin cytoskeleton in yeast.¹⁰⁶ *asp1* Δ yeast exhibit temperature sensitive morphological defects and defects in polarised growth due to a disorganised actin cytoskeleton. Later, when Vip1 and Asp1 were found to possess inositol pyrophosphate synthesis activity, it was observed that the kinase activity of Asp1 was essential to maintain the normal rod-shaped morphology of fission yeast.²⁸

Remodelling of the actin cytoskeleton is essential for cell migration and invasion associated with carcinogenesis. Recent studies have shed light on the role of IP₇ in the promotion of tumour growth and metastasis using in vitro and in vivo approaches.^{107, 108} The Snyder group showed that gene deletion of *Ip6k2* in HCT116 human colorectal cancer cells and *Ip6k2* knock-down in breast and lung cancer cell lines led to a reduction in focal adhesion kinase (FAK) phosphorylation, correlating with reduced cell spreading and cell-matrix adhesion.¹⁰⁷ Epithelial to mesenchymal transition properties including cell migration and invasion were also significantly reduced in these cells. Subcutaneous xenograft of *Ip6k2* knockout HCT116 cells in immune-compromised mice led to tumours that were smaller in size as compared with those formed by parent HCT116 cells. This study also worked out the molecular mechanism underlying the influence of IP6K2 on actin cytoskeleton remodelling. Liver Kinase B1 (LKB1), which suppresses FAK phosphorylation dependent migration, invasion and metastasis, is localised in the cytosol and is activated upon phosphorylation by PKC-zeta. IP6K2 binds LKB1 and generates IP₇ which through an unknown mechanism leads to reduced PKC-zeta dependent LKB1 phosphorylation, causing its nuclear sequestration and inactivation, thus increasing FAK activity to facilitate actin remodelling. Recent work from our laboratory showed that IP6K1 also promotes cell migration and invasion.¹⁰⁸ Knockdown of IP6K1 expression in cancer cells leads to a reduction in cell migration, invasion and anchorage-independent growth. *Ip6k1* knockout mice fed with the oral carcinogen 4NQO showed reduced progression from epithelial dysplasia to invasive carcinoma in the upper aerodigestive tract as compared with their wild-type littermates, showing that IP6K1 is also required to promote cell invasion in vivo. This phenotype of resistance to aerodigestive tract carcinoma seen in *Ip6k1* knockout mice is in direct contrast to the outcome of 4NQO treatment observed in *Ip6k2* knockout mice (described earlier), which showed a higher incidence of carcinoma.⁹¹ These contrasting observations suggest that in aerodigestive tract epithelial cells, IP6K1 is responsible for promoting carcinogenesis, whereas the predominant function of IP6K2 is to prevent transformation by promoting apoptosis.

It was recently shown that the third IP₆ kinase isoform, IP6K3, also influences the actin cytoskeleton. This study examined the effect of the loss of *Ip6k3* in specialised neurons in the cerebellum

called Purkinje cells, which express high levels of IP6K3.⁷² Behavioural tests revealed that *Ip6k3* knockout mice manifest defects in motor learning and coordination. These mice showed abnormalities in **synapse** number and structure of Purkinje cells. The shape of Purkinje cell **dendritic spines** is regulated by the arrangement and attachment of cytoskeletal elements including actin and actin-regulating proteins such as spectrins and adducins.^{109, 110} Spectrin is a structural protein that forms a mesh on the cytoplasmic face of the plasma membrane and adducin binds spectrin to promote its association with filamentous actin.¹¹¹ IP6K3, but not IP6K1 or IP6K2, was shown to bind spectrin and adducin, and cells lacking IP6K3 showed reduced spectrin-adducin interaction.⁷² Catalytically active and inactive forms of IP6K3 can promote adducin binding to spectrin, providing another example of IP₆ kinases functioning as structural scaffolds independent of their ability to synthesize PP-IPs.

3.5 Nucleus

The work of several laboratories has shed light on the importance of PP-IPs in the regulation of essential housekeeping functions in the nucleus, including the maintenance of genome integrity and regulation of transcription.

3.5.1 Genome Integrity PP-IPs have been shown to participate in many processes that are responsible for maintaining the integrity of the eukaryotic genome. Telomeres are protein–DNA structures present at the ends of linear eukaryotic chromosomes to protect them against degradation. The repetitive DNA sequences present in telomeres shorten with each cell division and telomere shortening correlates with ageing. Two independent studies showed that PP-IPs play a role in maintaining the length of telomeres in budding yeast. Using yeast mutants deleted for various inositol phosphate biosynthetic enzymes, it was shown that the loss of PP-IPs led to telomere lengthening, whereas their overproduction led to shortening of telomeres.¹¹² Telomere lengthening in *kcs1Δ* yeast could be reversed by expression of active but not inactive Kcs1, indicating that PP-IP synthesis by this enzyme is necessary to maintain normal telomere length. In another study, it was observed that yeast lacking PP-IPs are resistant to treatment with wortmannin and caffeine, inhibitors of phosphoinositide 3-kinase (PI3K) and PI3K-related protein kinases, which are known regulators of telomere length.¹¹³ Yeast lacking

Kcs1 showed longer than normal telomeres, but interestingly, yeast lacking the IP₆ synthesizing enzyme *Ipk1* showed shortening of telomeres. This was attributed to the high levels of inositol pyrophosphates PP-IP₄ and [PP]₂-IP₃ synthesized from IP₅ in these cells, so that the total PP-IP component of *ipk1Δ* yeast is actually higher than wild-type yeast. This data showed that any PP-IP can act to maintain telomere length, suggesting that perhaps protein(s) involved in telomere length maintenance are pyrophosphorylation targets of PP-IPs.

PP-IPs have been shown to participate in two key DNA repair pathways, homologous recombination (HR) mediated repair and nucleotide excision repair (NER). The yeast IP₆ kinase Kcs1 (kinase C suppressor-1) was initially identified in a genetic screen for second site mutations that suppress the hyperrecombination phenotype observed in yeast carrying a mutant form of protein kinase C (*Pkc*).¹¹⁴ Subsequently, it was shown that the inositol pyrophosphate synthesis activity of Kcs1 is essential to support hyperrecombination in *pkc* mutant yeast.¹¹⁵ Our data revealed that the DNA recombination promoting function of PP-IPs is also conserved in mammalian cells.¹¹⁶ When MEFs were allowed to recover from DNA damage induced by the replication stress-inducer hydroxyurea or the radiomimetic antibiotic neocarzinostatin, cells lacking IP6K1 showed delayed entry into the next phase of the cell cycle, and ultimately underwent cell death. The DNA repair markers, γ -H2AX, BLM helicase, and Rad51 were recruited to the sites of DNA damage in *Ip6k1*^{-/-} MEFs, suggesting that HR is initiated in these cells. However, these markers persisted for a longer time in *Ip6k1*^{-/-} MEFs, and the DNA breaks were not repaired, indicating that repair is incomplete in these cells. Expression of active but not inactive IP6K1 could reverse this phenotype, showing that 5-IP₇ is required to support HR in mammals. It was recently reported that IP6K1 also promotes NER in an enzyme activity dependent manner.¹¹⁷ IP6K1 was shown to interact with damage-specific DNA binding protein 1 (DDB1), which is part of the Cullin-RING ubiquitin ligase CRL4 complex, an E3 ubiquitin ligase that initiates NER.¹¹⁸ IP6K1 binding to CRL4 promotes its interaction with the COP9 signalosome (CSN) to keep the E3 ligase inactive. UV exposure leads to dissociation of DDB1 from IP6K1, allowing the synthesis of 5-IP₇, which acts as a transducer for NER by promoting the dissociation of CRL4 from CSN.

Synapse: Junction between two nerve cells, which consists of a small gap across which impulses pass by diffusion of a neurotransmitter.

Dendritic spines: Small protrusion from a neuron's dendrite which receives signal from a single axon at the synapse.

3.5.2 Chromatin Remodelling Two independent studies highlighted the role of PP-IPs in regulating epigenetic modifications in yeast and mammals. In budding yeast, PP-IPs were shown to play a critical role in regulating the environmental stress response. The yeast strain *kcs1Δvip1Δ*, which is incapable of producing any PP-IPs, showed little to no transcriptional response to heat, osmotic, or oxidative stress.⁶¹ These cells displayed a decrease in stress dependent histone deacetylation brought about by the HDAC complex Rpd3L. A putative inositol phosphate binding site was identified on the catalytic subunit Rpd3, suggesting that PP-IP binding may directly activate Rpd3L to regulate the global transcription response to environmental stress. Another study identified that mouse IP6K1 interacts with the histone demethylase JMJD2C (Jumonji domain containing 2C).¹¹⁹ Cells lacking IP6K1 showed a global reduction in the levels of trimethylation on histone H3 lysine 9 (H3K9me3), and a concomitant increase in H3K9 acetylation. Overexpression of active but not inactive IP6K1 led to dissociation of JMJD2C from chromatin and a consequent increase in H3K9me3 levels. It was suggested that IP₇ acts on one or more chromatin associated proteins to lower JMJD2C recruitment to chromatin.

3.5.3 Transcription Regulation Studies in budding yeast have shown that PP-IPs control transcription to regulate major metabolic pathways. The Pho80–Pho85 cyclin-cyclin dependent kinase (CDK) complex is responsible for regulating the phosphate (Pi) responsive (PHO) pathway by phosphorylating the transcription factor Pho4 to promote its cytoplasmic accumulation. Under Pi starvation conditions, the CDK inhibitor Pho81 lowers the kinase activity of Pho80–Pho85, leading to dephosphorylation and nuclear translocation of Pho4 to trigger the transcription of PHO genes.^{120,121} O’Shea and colleagues found that PP-IPs produced by Vip1 during phosphate starvation lowered Pho4 phosphorylation by the Pho80–Pho85–Pho81 complex.⁵⁷ They further determined that 1-IP₇ binding to Pho81 triggers a structural change that occludes the binding of Pho4 to the Pho85 kinase active site, but that 5-IP₇ does not have this effect,³⁸ highlighting how PP-IP binding-mediated regulation of protein function is specific to individual PP-IP isoforms. Another example of transcription regulation by 1-IP₇ was seen in the mammalian innate immune response.¹²² 1-IP₇ but not 5-IP₇ increased phosphorylation and activation of the transcription factor IRF3, which is responsible

for production of the cytokine IFN β upon viral infection. Interestingly, a non-hydrolysable analogue of 1-IP₇ could not recapitulate this effect, suggesting that 1-IP₇-mediated specific pyrophosphorylation may be involved in this pathway. This study provides the only hint that PP-IP-mediated protein pyrophosphorylation may also be stereo-selective towards a particular isoform.

Yeast lacking Kcs1 exhibit inositol auxotrophy and decreased intracellular inositol levels.¹²³ This is due to reduced transcription of the *INO1* gene which encodes *myo*-inositol-3-phosphate synthase, the enzyme that converts glucose-6-phosphate to inositol-3-phosphate, the rate limiting step of *de novo* inositol synthesis in eukaryotes. Inositol depletion in the growth medium led to increased Kcs1 protein levels, and PP-IPs synthesized by Kcs1 were found to be essential for the upregulation of *INO1* transcription under these conditions. As expression of mouse IP6K1 in *kcs1Δ* yeast rescued their inositol auxotrophy,¹²⁴ it was expected that PP-IP dependent regulation of inositol synthesis would be conserved in mammals. However, MEFs lacking IP6K1 exhibited an unexpected increase in mammalian *Ino1* transcription, and a corresponding increase in inositol levels as compared with wild-type MEFs.¹²⁴ This study further demonstrated that IP6K1 is localised to the nucleus by binding to the lipid phosphatidic acid, and that PP-IP synthesis by IP6K1 increases methylation of *Ino1* DNA to reduce its transcription. The phenomenon of *Ino1* transcription regulation by PP-IPs is a rare example of evolutionary divergence between yeast and mammals with regard to the influence of PP-IPs on a specific metabolic pathway.

Studies in our laboratory have shown that PP-IPs regulate rRNA transcription by RNA polymerase I in budding yeast.⁴⁵ *kcs1Δ* yeast exhibit reduced protein synthesis due to a decrease in ribosome biogenesis, which in turn is attributed to a decrease in rRNA levels. Although there was no defect in the recruitment of RNA polymerase I on rDNA, the rate of transcription elongation was reduced in *kcs1Δ* as compared with wild-type yeast. We identified that 5-IP₇ can pyrophosphorylate three subunits of the RNA polymerase I complex, and suggest that this modification may be essential to maintain normal transcription elongation by this polymerase. A recent study by the Fiedler group has shown that several potential IP₇ pyrophosphorylation targets are nucleolar proteins associated with RNA polymerase I,

suggesting that there may be additional proteins on which IP₇ acts to regulate rRNA synthesis.¹²⁵

4 Perspective on the Future

Although PP-IPs were identified more than 20 years ago, there are still only a handful of researchers attempting to uncover the physiological functions of these unique molecules. The main reason for the lack of popularity faced by these molecules is that their study remains technically challenging for most cell biologists. There are no commercially available kits for the detection and measurement of PP-IPs, no fluorescently tagged or radiolabelled PP-IPs readily available for use in protein binding or pyrophosphorylation assays, and no easy methods to detect pyrophosphorylated proteins. Recent advances by the Fiedler, Jessen and Potter laboratories are likely to help in surmounting these challenges.^{78, 79, 126–129} Highly pure PP-IPs with β-phosphate moieties at specific carbon atoms have been synthesized by these groups, including non-hydrolysable analogues that are stable in cells and can bind target proteins but not pyrophosphorylate them.^{47, 78, 130} A system for intracellular delivery and photouncaging of chemically synthesized PP-IPs has recently been developed,⁷⁹ which promises to open up new methods to study the functions of these molecules in different subcellular compartments. The use of IP₇ as an affinity reagent revealed two different classes of interacting proteins, depending on the absence or presence of the divalent cation Mg²⁺ during the interaction, representing IP₇ binding or pyrophosphorylation targets respectively.¹²⁵ These latest advances promise a bright future for PP-IPs, with the hope that availability of new tools and information on novel PP-IP target proteins will draw new researchers into examining whether these versatile small molecules can regulate the protein or pathway of their interest.

Acknowledgements

The authors thank Manasa Chanduri for valuable feedback and inputs. AS and JS are recipients of Junior Research Fellowships of the University Grants Commission, Government of India, and SG is a recipient of a Junior Research Fellowship from the Centre for DNA Fingerprinting and Diagnostics (CDFD). RB acknowledges funding from the Wellcome Trust/Department of Biotechnology India Alliance (500020/Z/09/Z), Human Frontier Science Program (RGP0025/2016) and core funds from the CDFD.

Received: 5 November 2016. Accepted: 16 November 2016
Published online: 28 February 2017

Open Access

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Michell RH (2008) Inositol derivatives: evolution and functions. *Nat Rev Mol Cell Biol* 9:151–161
2. Wundenberg T, Mayr GW (2012) Synthesis and biological actions of diphosphoinositol phosphates (inositol pyrophosphates), regulators of cell homeostasis. *Biol Chem* 393:979–998
3. Pisani F, Livermore T, Rose G, Chubb JR, Gaspari M, Saiardi A (2014) Analysis of *Dictyostelium discoideum* inositol pyrophosphate metabolism by gel electrophoresis. *PLoS One* 9:e85533
4. Mayr GW, Radenberg T, Thiel U, Vogel G, Stephens LR (1992) Phosphoinositol diphosphates: non-enzymic formation in vitro and occurrence in vivo in the cellular slime mold *Dictyostelium*. *Carbohydr Res* 234:247–262
5. Stephens L, Radenberg T, Thiel U, Vogel G, Khoo KH, Dell A et al (1993) The detection, purification, structural characterization, and metabolism of diphosphoinositol pentakisphosphate(s) and bisdiphosphoinositol tetrakisphosphate(s). *J Biol Chem* 268:4009–4015
6. Menniti FS, Miller RN, Putney JW Jr, Shears SB (1993) Turnover of inositol polyphosphate pyrophosphates in pancreatoma cells. *J Biol Chem* 268:3850–3856
7. Bennett M, Onnebo SM, Azevedo C, Saiardi A (2006) Inositol pyrophosphates: metabolism and signaling. *Cell Mol Life Sci* 63:552–564
8. Shears SB (2007) Understanding the biological significance of diphosphoinositol polyphosphates ('inositol pyrophosphates'). *Biochem Soc Symp* 74:211–221
9. Barker CJ, Illies C, Gaboardi GC, Berggren PO (2009) Inositol pyrophosphates: structure, enzymology and function. *Cell Mol Life Sci* 66:3851–3871
10. Burton A, Hu X, Saiardi A (2009) Are inositol pyrophosphates signalling molecules? *J Cell Physiol* 220:8–15
11. Shears SB (2009) Diphosphoinositol polyphosphates: metabolic messengers? *Mol Pharmacol* 76:236–252
12. Tsui MM, York JD (2010) Roles of inositol phosphates and inositol pyrophosphates in development, cell signaling and nuclear processes. *Adv Enzyme Regul* 50:324–337

13. Chakraborty A, Kim S, Snyder SH (2011) Inositol pyrophosphates as mammalian cell signals. *Sci Signal* 4:1
14. Azevedo C, Sziogyarto Z, Saiardi A (2011) The signaling role of inositol hexakisphosphate kinases (IP6Ks). *Adv Enzyme Regul* 51:74–82
15. Shears SB, Gokhale NA, Wang H, Zaremba A (2011) Diphosphoinositol polyphosphates: what are the mechanisms? *Adv Enzyme Regul* 51:13–25
16. Saiardi A (2012) Cell signalling by inositol pyrophosphates. *Subcell Biochem* 59:413–443
17. Saiardi A (2012) How inositol pyrophosphates control cellular phosphate homeostasis? *Adv Biol Regul* 52:351–359
18. Wilson MS, Livermore TM, Saiardi A (2013) Inositol pyrophosphates: between signalling and metabolism. *Biochem J* 452:369–379
19. Barker CJ, Berggren PO (2013) New horizons in cellular regulation by inositol polyphosphates: insights from the pancreatic beta-cell. *Pharmacol Rev* 65:641–669
20. Koldobskiy MA, Snyder SH (2011) Inositol pyrophosphates in cell death and life. *Cell Cycle* 10:568–570
21. Thomas MP, Potter BV (2014) The enzymes of human diphosphoinositol polyphosphate metabolism. *FEBS J* 281:14–33
22. Shears SB (2015) Inositol pyrophosphates: why so many phosphates? *Adv Biol Regul* 57:203–216
23. Thota SG, Bhandari R (2015) The emerging roles of inositol pyrophosphates in eukaryotic cell physiology. *J Biosci* 40:593–605
24. Onnebo SM, Saiardi A (2009) Inositol pyrophosphates modulate hydrogen peroxide signalling. *Biochem J* 423:109–118
25. Wilson MS, Bulley SJ, Pisani F, Irvine RF, Saiardi A (2015) A novel method for the purification of inositol phosphates from biological samples reveals that no phytate is present in human plasma or urine. *Open Biol* 5:150014
26. Irvine RF, Schell MJ (2001) Back in the water: the return of the inositol phosphates. *Nat Rev Mol Cell Biol* 2:327–338
27. Saiardi A, Erdjument-Bromage H, Snowman AM, Tempst P, Snyder SH (1999) Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher inositol polyphosphate kinases. *Curr Biol* 9:1323–1326
28. Mulugu S, Bai W, Fridy PC, Bastidas RJ, Otto JC, Dollins DE et al (2007) A conserved family of enzymes that phosphorylate inositol hexakisphosphate. *Science* 316:106–109
29. Choi JH, Williams J, Cho J, Falck JR, Shears SB (2007) Purification, sequencing, and molecular identification of a mammalian PP-InsP5 kinase that is activated when cells are exposed to hyperosmotic stress. *J Biol Chem* 282:30763–30775
30. Lin H, Fridy PC, Ribeiro AA, Choi JH, Barma DK, Vogel G et al (2009) Structural analysis and detection of biological inositol pyrophosphates reveal that the family of VIP/diphosphoinositol pentakisphosphate kinases are 1/3-kinases. *J Biol Chem* 284:1863–1872
31. Wang H, Falck JR, Hall TM, Shears SB (2012) Structural basis for an inositol pyrophosphate kinase surmounting phosphate crowding. *Nat Chem Biol* 8:111–116
32. Gu C, Wilson MS, Jessen HJ, Saiardi A, Shears SB (2016) Inositol pyrophosphate profiling of two HCT116 cell lines uncovers variation in InsP8 levels. *PLoS One* 11:e0165286
33. Draskovic P, Saiardi A, Bhandari R, Burton A, Ilc G, Kovacevic M et al (2008) Inositol hexakisphosphate kinase products contain diphosphate and triphosphate groups. *Chem Biol* 15:274–286
34. Safrany ST, Caffrey JJ, Yang X, Bembenek ME, Moyer MB, Burkhart WA et al (1998) A novel context for the 'MutT' module, a guardian of cell integrity, in a diphosphoinositol polyphosphate phosphohydrolase. *EMBO J* 17:6599–6607
35. Wang H, Nair VS, Holland AA, Capolicchio S, Jessen HJ, Johnson MK et al (2015) Asp1 from *Schizosaccharomyces pombe* binds a [2Fe–2S](2+) cluster which inhibits inositol pyrophosphate 1-phosphatase activity. *Biochemistry* 54:6462–6474
36. Shears SB, Baughman BM, Gu C, Nair VS, Wang H (2016) The significance of the 1-kinase/1-phosphatase activities of the PPIP5K family. *Adv Biol Regul* 63:96–106
37. Steidle EA, Chong LS, Wu M, Crooke E, Fiedler D, Resnick AC et al (2016) A novel inositol pyrophosphate phosphatase in *Saccharomyces cerevisiae*: Siw14 protein selectively cleaves the beta-phosphate from 5-diphosphoinositol pentakisphosphate (5PP-IP5). *J Biol Chem* 291:6772–6783
38. Lee YS, Huang K, Quioco FA, O'Shea EK (2008) Molecular basis of cyclin-CDK-CKI regulation by reversible binding of an inositol pyrophosphate. *Nat Chem Biol* 4:25–32
39. Chakraborty A, Koldobskiy MA, Bello NT, Maxwell M, Potter JJ, Juluri KR et al (2010) Inositol pyrophosphates inhibit Akt signaling, thereby regulating insulin sensitivity and weight gain. *Cell* 143:897–910
40. Prasad A, Jia Y, Chakraborty A, Li Y, Jain SK, Zhong J et al (2011) Inositol hexakisphosphate kinase 1 regulates neutrophil function in innate immunity by inhibiting phosphatidylinositol-(3,4,5)-trisphosphate signaling. *Nat Immunol* 12:752–760
41. Gokhale NA, Zaremba A, Janoshazi AK, Weaver JD, Shears SB (2013) PPIP5K1 modulates ligand competition between diphosphoinositol polyphosphates and PtdIns(3,4,5)P3 for polyphosphoinositide-binding domains. *Biochem J* 453:413–426
42. Bhandari R, Saiardi A, Ahmadibeni Y, Snowman AM, Resnick AC, Kristiansen TZ et al (2007) Protein pyrophosphorylation by inositol pyrophosphates is a posttranslational event. *Proc Natl Acad Sci USA* 104:15305–15310

43. Saiardi A, Bhandari R, Resnick AC, Snowman AM, Snyder SH (2004) Phosphorylation of proteins by inositol pyrophosphates. *Science* 306:2101–2105
44. Szijgyarto Z, Garedeu A, Azevedo C, Saiardi A (2011) Influence of inositol pyrophosphates on cellular energy dynamics. *Science* 334:802–805
45. Thota SG, Unnikannan CP, Thampatty SR, Manorama R, Bhandari R (2015) Inositol pyrophosphates regulate RNA polymerase I-mediated rRNA transcription in *Saccharomyces cerevisiae*. *Biochem J* 466:105–114
46. Azevedo C, Burton A, Ruiz-Mateos E, Marsh M, Saiardi A (2009) Inositol pyrophosphate mediated pyrophosphorylation of AP3B1 regulates HIV-1 Gag release. *Proc Natl Acad Sci USA* 106:21161–21166
47. Chanduri M, Rai A, Malla AB, Wu M, Fiedler D, Malik R et al (2016) Inositol hexakisphosphate kinase 1 (IP6K1) activity is required for cytoplasmic dynein-driven transport. *Biochem J* 473:3031–3047
48. Yates LM, Fiedler D (2015) Establishing the stability and reversibility of protein pyrophosphorylation with synthetic peptides. *ChemBioChem* 16:415–423
49. Mayr GW (1988) A novel metal-dye detection system permits picomolar-range h.p.l.c. analysis of inositol polyphosphates from non-radioactively labelled cell or tissue specimens. *Biochem J* 254:585–591
50. Losito O, Szijgyarto Z, Resnick AC, Saiardi A (2009) Inositol pyrophosphates and their unique metabolic complexity: analysis by gel electrophoresis. *PLoS One* 4:e5580
51. Glennon MC, Shears SB (1993) Turnover of inositol pentakisphosphates, inositol hexakisphosphate and diphosphoinositol polyphosphates in primary cultured hepatocytes. *Biochem J* 293:583–590
52. Pesesse X, Choi K, Zhang T, Shears SB (2004) Signaling by higher inositol polyphosphates. Synthesis of bisdiphosphoinositol tetrakisphosphate (“InsP8”) is selectively activated by hyperosmotic stress. *J Biol Chem* 279:43378–43381
53. Choi K, Mollapour E, Shears SB (2005) Signal transduction during environmental stress: InsP(8) operates within highly restricted contexts. *Cell Signal* 17:1533–1541
54. Choi K, Mollapour E, Choi JH, Shears SB (2008) Cellular energetic status supervises the synthesis of bisdiphosphoinositol tetrakisphosphate independently of AMP-activated protein kinase. *Mol Pharmacol* 74:527–536
55. Nagata E, Luo HR, Saiardi A, Bae BI, Suzuki N, Snyder SH (2005) Inositol hexakisphosphate kinase-2, a physiologic mediator of cell death. *J Biol Chem* 280:1634–1640
56. Xu Y, Li H, Bajrami B, Kwak H, Cao S, Liu P et al (2013) Cigarette smoke (CS) and nicotine delay neutrophil spontaneous death via suppressing production of diphosphoinositol pentakisphosphate. *Proc Natl Acad Sci USA* 110:7726–7731
57. Lee YS, Mulugu S, York JD, O’Shea EK (2007) Regulation of a cyclin-CDK-CDK inhibitor complex by inositol pyrophosphates. *Science* 316:109–112
58. Lonetti A, Szijgyarto Z, Bosch D, Loss O, Azevedo C, Saiardi A (2011) Identification of an evolutionarily conserved family of inorganic polyphosphate endopolyphosphatases. *J Biol Chem* 286:31966–31974
59. Wild R, Gerasimaite R, Jung JY, Truffault V, Pavlovic I, Schmidt A et al (2016) Control of eukaryotic phosphate homeostasis by inositol polyphosphate sensor domains. *Science* 352:986–990
60. Dubois E, Scherens B, Vierendeels F, Ho MM, Messenguy F, Shears SB (2002) In *Saccharomyces cerevisiae*, the inositol polyphosphate kinase activity of Kcs1p is required for resistance to salt stress, cell wall integrity, and vacuolar morphogenesis. *J Biol Chem* 277:23755–23763
61. Worley J, Luo X, Capaldi AP (2013) Inositol pyrophosphates regulate cell growth and the environmental stress response by activating the HDAC Rpd3L. *Cell Rep* 3:1476–1482
62. Desai M, Rangarajan P, Donahue JL, Williams SP, Land ES, Mandal MK et al (2014) Two inositol hexakisphosphate kinases drive inositol pyrophosphate synthesis in plants. *Plant J* 80:642–653
63. Laha D, Johnen P, Azevedo C, Dynowski M, Weiss M, Capolicchio S et al (2015) VIH2 regulates the synthesis of inositol pyrophosphate InsP8 and jasmonate-dependent defenses in Arabidopsis. *Plant Cell* 27:1082–1097
64. Bucking-Throm E, Duntze W, Hartwell LH, Manney TR (1973) Reversible arrest of haploid yeast cells in the initiation of DNA synthesis by a diffusible sex factor. *Exp Cell Res* 76:99–110
65. Banfic H, Bedalov A, York JD, Visnjic D (2013) Inositol pyrophosphates modulate S phase progression after pheromone-induced arrest in *Saccharomyces cerevisiae*. *J Biol Chem* 288:1717–1725
66. Rosner M, Schipany K, Hengstschlager M (2013) Merging high-quality biochemical fractionation with a refined flow cytometry approach to monitor nucleocytoplasmic protein expression throughout the unperturbed mammalian cell cycle. *Nat Protoc* 8:602–626
67. Barker CJ, Wright J, Hughes PJ, Kirk CJ, Michell RH (2004) Complex changes in cellular inositol phosphate complement accompany transit through the cell cycle. *Biochem J* 380:465–473
68. Laussmann T, Pikzack C, Thiel U, Mayr GW, Vogel G (2000) Diphospho-myo-inositol phosphates during the life cycle of Dictyostelium and Polysphondylium. *Eur J Biochem* 267:2447–2451
69. Luo HR, Huang YE, Chen JC, Saiardi A, Iijima M, Ye K et al (2003) Inositol pyrophosphates mediate chemotaxis in Dictyostelium via pleckstrin homology domain-PtdIns(3,4,5)P3 interactions. *Cell* 114:559–572

70. Zhang Z, Zhao C, Liu B, Liang D, Qin X, Li X et al (2014) Inositol pyrophosphates mediate the effects of aging on bone marrow mesenchymal stem cells by inhibiting Akt signaling. *Stem Cell Res Ther* 5:33
71. Saiardi A, Nagata E, Luo HR, Snowman AM, Snyder SH (2001) Identification and characterization of a novel inositol hexakisphosphate kinase. *J Biol Chem* 276:39179–39185
72. Fu C, Xu J, Li RJ, Crawford JA, Khan AB, Ma TM et al (2015) Inositol hexakisphosphate kinase-3 regulates the morphology and synapse formation of cerebellar purkinje cells via spectrin/adducin. *J Neurosci* 35:11056–11067
73. Bhandari R, Juluri KR, Resnick AC, Snyder SH (2008) Gene deletion of inositol hexakisphosphate kinase 1 reveals inositol pyrophosphate regulation of insulin secretion, growth, and spermiogenesis. *Proc Natl Acad Sci USA* 105:2349–2353
74. Fridy PC, Otto JC, Dollins DE, York JD (2007) Cloning and characterization of two human VIP1-like inositol hexakisphosphate and diphosphoinositol pentakisphosphate kinases. *J Biol Chem* 282:30754–30762
75. Yong ST, Nguyen HN, Choi JH, Bortner CD, Williams J, Pulloor NK et al (2015) Identification of a functional nuclear translocation sequence in hPPIP5K2. *BMC Cell Biol* 16:17
76. Gokhale NA, Zaremba A, Shears SB (2011) Receptor-dependent compartmentalization of PPIP5K1, a kinase with a cryptic polyphosphoinositide binding domain. *Biochem J* 434:415–426
77. Downes CP, Gray A, Fairservice A, Safrany ST, Batty IH, Fleming I (2005) The regulation of membrane to cytosol partitioning of signalling proteins by phosphoinositides and their soluble headgroups. *Biochem Soc Trans* 33:1303–1307
78. Wu M, Dul BE, Trevisan AJ, Fiedler D (2013) Synthesis and characterization of non-hydrolysable diphosphoinositol polyphosphate second messengers. *Chem Sci* 4:405–410
79. Pavlovic I, Thakor DT, Vargas JR, McKinlay CJ, Hauke S, Anstaett P et al (2016) Cellular delivery and photochemical release of a caged inositol-pyrophosphate induces PH-domain translocation in cellulo. *Nat Commun* 7:10622
80. Nagel A, Barker CJ, Berggren PO, Illies C (2010) Diphosphoinositol polyphosphates and energy metabolism: assay for ATP/ADP ratio. *Methods Mol Biol* 645:123–131
81. Wundenberg T, Grabinski N, Lin H, Mayr GW (2014) Discovery of InsP6-kinases as InsP6 dephosphorylating enzymes provides a new mechanism of cytosolic InsP6 degradation driven by the cellular ATP/ADP ratio. *Biochem J* 462:173–184
82. Zhu Q, Ghoshal S, Rodrigues A, Gao S, Asterian A, Kamenecka TM et al (2016) Adipocyte-specific deletion of Ip6k1 reduces diet-induced obesity by enhancing AMPK-mediated thermogenesis. *J Clin Invest* 126:4273–4288
83. Morrison BH, Bauer JA, Hu J, Grane RW, Ozdemir AM, Chawla-Sarkar M et al (2002) Inositol hexakisphosphate kinase 2 sensitizes ovarian carcinoma cells to multiple cancer therapeutics. *Oncogene* 21:1882–1889
84. Morrison BH, Tang Z, Jacobs BS, Bauer JA, Lindner DJ (2005) Apo2L/TRAIL induction and nuclear translocation of inositol hexakisphosphate kinase 2 during IFN-beta-induced apoptosis in ovarian carcinoma. *Biochem J* 385:595–603
85. Morrison BH, Bauer JA, Kalvakolanu DV, Lindner DJ (2001) Inositol hexakisphosphate kinase 2 mediates growth suppressive and apoptotic effects of interferon-beta in ovarian carcinoma cells. *J Biol Chem* 276:24965–24970
86. Nagata E, Saiardi A, Tsukamoto H, Okada Y, Itoh Y, Satoh T et al (2011) Inositol hexakisphosphate kinases induce cell death in Huntington disease. *J Biol Chem* 286:26680–26686
87. Koldobskiy MA, Chakraborty A, Werner JK Jr, Snowman AM, Juluri KR, Vandiver MS et al (2010) p53-mediated apoptosis requires inositol hexakisphosphate kinase-2. *Proc Natl Acad Sci USA* 107:20947–20951
88. Rao F, Cha J, Xu J, Xu R, Vandiver MS, Tyagi R et al (2014) Inositol pyrophosphates mediate the DNA-PK/ATM-p53 cell death pathway by regulating CK2 phosphorylation of Tti1/Tel2. *Mol Cell* 54:119–132
89. Morrison BH, Bauer JA, Lupica JA, Tang Z, Schmidt H, DiDonato JA et al (2007) Effect of inositol hexakisphosphate kinase 2 on transforming growth factor beta-activated kinase 1 and NF-kappaB activation. *J Biol Chem* 282:15349–15356
90. Chakraborty A, Koldobskiy MA, Sixt KM, Juluri KR, Mustafa AK, Snowman AM et al (2008) HSP90 regulates cell survival via inositol hexakisphosphate kinase-2. *Proc Natl Acad Sci USA* 105:1134–1139
91. Morrison BH, Haney R, Lamarre E, Drazba J, Prestwich GD, Lindner DJ (2009) Gene deletion of inositol hexakisphosphate kinase 2 predisposes to aerodigestive tract carcinoma. *Oncogene* 28:2383–2392
92. Kaur J, Debnath J (2015) Autophagy at the crossroads of catabolism and anabolism. *Nat Rev Mol Cell Biol* 16:461–472
93. Taylor R Jr, Chen PH, Chou CC, Patel J, Jin SV (2012) KCS1 deletion in *Saccharomyces cerevisiae* leads to a defect in translocation of autophagic proteins and reduces autophagosome formation. *Autophagy* 8:1300–1311
94. Nagata E, Saiardi A, Tsukamoto H, Satoh T, Itoh Y, Itoh J et al (2010) Inositol hexakisphosphate kinases promote autophagy. *Int J Biochem Cell Biol* 42:2065–2071
95. Saiardi A, Sciambi C, McCaffery JM, Wendland B, Snyder SH (2002) Inositol pyrophosphates regulate endocytic trafficking. *Proc Natl Acad Sci USA* 99:14206–14211

96. Cortesy-Theulaz I, Pauloin A, Pfeffer SR (1992) Cytoplasmic dynein participates in the centrosomal localization of the Golgi complex. *J Cell Biol* 118:1333–1345
97. Li SC, Kane PM (2009) The yeast lysosome-like vacuole: endpoint and crossroads. *Biochim Biophys Acta* 1793:650–663
98. Azevedo C, Saiardi A (2014) Functions of inorganic polyphosphates in eukaryotic cells: a coat of many colours. *Biochem Soc Trans* 42:98–102
99. Auesukaree C, Tochio H, Shirakawa M, Kaneko Y, Harashima S (2005) Plc1p, Arg82p, and Kcs1p, enzymes involved in inositol pyrophosphate synthesis, are essential for phosphate regulation and polyphosphate accumulation in *Saccharomyces cerevisiae*. *J Biol Chem* 280:25127–25133
100. Ghosh S, Shukla D, Suman K, Lakshmi BJ, Manorama R, Kumar S et al (2013) Inositol hexakisphosphate kinase 1 maintains hemostasis in mice by regulating platelet polyphosphate levels. *Blood* 122:1478–1486
101. Hothorn M, Neumann H, Lenherr ED, Wehner M, Rybin V, Hassa PO et al (2009) Catalytic core of a membrane-associated eukaryotic polyphosphate polymerase. *Science* 324:513–516
102. Saiardi A, Caffrey JJ, Snyder SH, Shears SB (2000) The inositol hexakisphosphate kinase family. Catalytic flexibility and function in yeast vacuole biogenesis. *J Biol Chem* 275:24686–24692
103. Illies C, Gromada J, Fiume R, Leibiger B, Yu J, Juhl K et al (2007) Requirement of inositol pyrophosphates for full exocytotic capacity in pancreatic cells. *Science* 318:1299–1302
104. Luo HR, Saiardi A, Nagata E, Ye K, Yu H, Jung TS et al (2001) GRAB: a physiologic guanine nucleotide exchange factor for Rab3A, which interacts with inositol hexakisphosphate kinase. *Neuron* 31:439–451
105. Lee T-S, Lee J-Y, Kyung JW, Yang Y, Park SJ, Lee S et al (2016) Inositol pyrophosphates inhibit synaptotagmin-dependent exocytosis. *Proc Natl Acad Sci USA* 113:8314–8319
106. Feoktistova A, McCollum D, Ohi R, Gould KL (1999) Identification and characterization of *Schizosaccharomyces pombe* asp1(+), a gene that interacts with mutations in the Arp2/3 complex and actin. *Genetics* 152:895–908
107. Rao F, Xu J, Fu C, Cha JY, Gadalla MM, Xu R et al (2015) Inositol pyrophosphates promote tumor growth and metastasis by antagonizing liver kinase B1. *Proc Natl Acad Sci USA* 112:1773–1778
108. Jadav RS, Kumar D, Buwa N, Ganguli S, Thampatty SR, Balasubramanian N et al (2016) Deletion of inositol hexakisphosphate kinase 1 (IP6K1) reduces cell migration and invasion, conferring protection from aerodigestive tract carcinoma in mice. *Cell Signal* 28:1124–1136
109. Landis DM, Reese TS (1983) Cytoplasmic organization in cerebellar dendritic spines. *J Cell Biol* 97:1169–1178
110. Matsuoka Y, Li X, Bennett V (1998) Adducin is an in vivo substrate for protein kinase C: phosphorylation in the MARCKS-related domain inhibits activity in promoting spectrin–actin complexes and occurs in many cells, including dendritic spines of neurons. *J Cell Biol* 142:485–497
111. Gardner K, Bennett V (1987) Modulation of spectrin–actin assembly by erythrocyte adducin. *Nature* 328:359–362
112. York SJ, Armbruster BN, Greenwell P, Petes TD, York JD (2005) Inositol diphosphate signaling regulates telomere length. *J Biol Chem* 280:4264–4269
113. Saiardi A, Resnick AC, Snowman AM, Wendland B, Snyder SH (2005) Inositol pyrophosphates regulate cell death and telomere length through phosphoinositide 3-kinase-related protein kinases. *Proc Natl Acad Sci USA* 102:1911–1914
114. Huang KN, Symington LS (1995) Suppressors of a *Saccharomyces cerevisiae* pkc1 mutation identify alleles of the phosphatase gene PTC1 and of a novel gene encoding a putative basic leucine zipper protein. *Genetics* 141:1275–1285
115. Luo HR, Saiardi A, Yu H, Nagata E, Ye K, Snyder SH (2002) Inositol pyrophosphates are required for DNA hyperrecombination in protein kinase c1 mutant yeast. *Biochemistry* 41:2509–2515
116. Jadav RS, Chanduri MV, Sengupta S, Bhandari R (2013) Inositol pyrophosphate synthesis by inositol hexakisphosphate kinase 1 is required for homologous recombination repair. *J Biol Chem* 288:3312–3321
117. Rao F, Xu J, Khan AB, Gadalla MM, Cha JY, Xu R et al (2014) Inositol hexakisphosphate kinase-1 mediates assembly/disassembly of the CRL4-signalosome complex to regulate DNA repair and cell death. *Proc Natl Acad Sci USA* 111:16005–16010
118. Li J, Wang QE, Zhu Q, El-Mahdy MA, Wani G, Praetorius-Ibba M et al (2006) DNA damage binding protein component DDB1 participates in nucleotide excision repair through DDB2 DNA-binding and cullin 4A ubiquitin ligase activity. *Cancer Res* 66:8590–8597
119. Burton A, Azevedo C, Andreassi C, Riccio A, Saiardi A (2013) Inositol pyrophosphates regulate JMJD2C-dependent histone demethylation. *Proc Natl Acad Sci USA* 110:18970–18975
120. Lenburg ME, O'Shea EK (1996) Signaling phosphate starvation. *Trends Biochem Sci* 21:383–387
121. Carroll AS, O'Shea EK (2002) Pho85 and signaling environmental conditions. *Trends Biochem Sci* 27:87–93
122. Pulloor NK, Nair S, Kostic AD, Bist P, Weaver JD, Riley AM et al (2014) Human genome-wide RNAi screen identifies an essential role for inositol pyrophosphates in Type-I interferon response. *PLoS Pathog* 10:e1003981
123. Ye C, Bandara WM, Greenberg ML (2013) Regulation of inositol metabolism is fine-tuned by inositol

- pyrophosphates in *Saccharomyces cerevisiae*. *J Biol Chem* 288:24898–24908
124. Yu W, Ye C, Greenberg ML (2016) Inositol hexakisphosphate kinase 1 (IP6K1) regulates inositol synthesis in mammalian cells. *J Biol Chem* 291:10437–10444
 125. Wu M, Chong LS, Perlman DH, Resnick AC, Fiedler D (2016) Inositol polyphosphates intersect with signaling and metabolic networks via two distinct mechanisms. *Proc Natl Acad Sci USA* 113:E6757–E6765
 126. Conway JH, Fiedler D (2015) An affinity reagent for the recognition of pyrophosphorylated peptides. *Angew Chem Int Ed Engl* 54:3941–3945
 127. Marmelstein AM, Yates LM, Conway JH, Fiedler D (2014) Chemical pyrophosphorylation of functionally diverse peptides. *J Am Chem Soc* 136:108–111
 128. Capolicchio S, Wang H, Thakor DT, Shears SB, Jensen HJ (2014) Synthesis of densely phosphorylated bis-1,5-diphospho-myo-inositol tetrakisphosphate and its enantiomer by bidirectional P-anhydride formation. *Angew Chem Int Ed Engl* 53:9508–9511
 129. Riley AM, Wang H, Shears SB (2015) BV, L.P. Synthetic tools for studying the chemical biology of InsP8. *Chem Commun (Camb)* 51:12605–12608
 130. Wang H, Godage HY, Riley AM, Weaver JD, Shears SB, Potter BV (2014) Synthetic inositol phosphate analogs reveal that PPIP5K2 has a surface-mounted substrate capture site that is a target for drug discovery. *Chem Biol* 21:689–699
 131. Kilari RS, Weaver JD, Shears SB, Safrany ST (2013) Understanding inositol pyrophosphate metabolism and function: kinetic characterization of the DIPPs. *FEBS Lett* 587:3464–3470



Akruti Shah completed her integrated M.Sc. in Medical Biotechnology from ARIBAS, affiliated with Sardar Patel University, Anand, Gujarat, in 2014. She is currently a graduate student in the Laboratory of Cell Signalling, Centre for DNA

Fingerprinting and Diagnostics, Hyderabad, under the guidance of Dr. Rashna Bhandari. Her research primarily focuses on elucidating the role of IP6K1 in the formation and maintenance of mRNA-protein granules called processing bodies.



Shubhra Ganguli completed her B.Sc. in Biochemistry from Sri Venkateswara College, University of Delhi, in 2012, and M.Sc. in Biotechnology from the Maharaja Sayajirao University of Baroda, Gujarat, in 2014. She is currently pursuing a Ph.D. in the

Laboratory of Cell Signalling at the Centre for DNA Fingerprinting and Diagnostics, Hyderabad, under the guidance of Dr. Rashna Bhandari. Her doctoral research is directed at understanding the role played by inositol pyrophosphates in regulating DNA repair pathways in mammalian cells.



Jayraj Sen received his B.Sc. degree in Biotechnology from the Government PG College, Guna, affiliated to Jiwaji University, Gwalior, Madhya Pradesh, in 2011, and M.Sc. in Biochemistry from Devi Ahilya Vishwavidyalaya, Indore, Madhya Pradesh,

in 2014. He is currently pursuing a Ph.D. in the Laboratory of Cell Signalling, under the guidance of Dr. Rashna Bhandari at the Centre for DNA Fingerprinting and Diagnostics, Hyderabad. His research goal is to investigate the role of IP6K1 in glandular epithelia.



Dr. Rashna Bhandari received her B.Sc. (Hons) degree in Human Biology from the All India Institute of Medical Sciences, Delhi. She obtained her M.Sc. and Ph.D. degrees in Biological Sciences from the Indian Institute of Science, Bengaluru,

where she worked with Dr. Sandhya Visweswariah on signal transduction by the membrane-bound guanylyl cyclase, GCC, which is involved in maintaining fluid and ion homeostasis across the intestinal membrane. As a post-doctoral fellow, she worked in the laboratory of Dr. John Kuriyan at the University of California at Berkeley, USA, where she studied the structure–function relationship of JAK-STAT proteins and the EGF receptor. Later, she moved to the Johns Hopkins University School of Medicine in Baltimore, USA, where she worked with Dr. Solomon Snyder to discover that inositol pyrophosphates can influence protein function by serine pyrophosphorylation. In 2008, she joined the Centre for DNA Fingerprinting and Diagnostics, Hyderabad, as a staff scientist, to head the Laboratory of Cell Signalling. Her laboratory uses budding yeast, mammalian cell lines and mice as model systems to investigate cellular pathways and physiological functions regulated by IP₆ kinases and inositol pyrophosphates.