

# *In vitro* selection of transgenic sugarcane callus utilizing a plant gene encoding a mutant form of acetolactate synthase

Christell van der Vyver · Tobie Conradie ·  
Jens Kossmann · James Lloyd

Received: 7 September 2012 / Accepted: 18 January 2013 / Published online: 5 February 2013 / Editor: J. Foster  
© The Author(s) 2013. This article is published with open access at Springerlink.com

**Abstract** Selection genes are routinely used in plant genetic transformation protocols to ensure the survival of transformed cells by limiting the regeneration of non-transgenic cells. In order to find alternatives to the use of antibiotics as selection agents, we followed a targeted approach utilizing a plant gene, encoding a mutant form of the enzyme acetolactate synthase, to convey resistance to herbicides. The sensitivity of sugarcane callus (*Saccharum* spp. hybrids, cv. NCo310) to a number of herbicides from the sulfonylurea and imidazolinone classes was tested. Callus growth was most affected by sulfonylurea herbicides, particularly 3.6 µg/l chlorsulfuron. Herbicide-resistant transgenic sugarcane plants containing mutant forms of a tobacco acetolactate synthase (*als*) gene were obtained following biolistic transformation. Post-bombardment, putative transgenic callus was selectively proliferated on MS medium containing 3 mg/l 2,4-dichlorophenoxyacetic acid (2,4-D), 20 g/l sucrose, 0.5 g/l casein, and 3.6 µg/l chlorsulfuron. Plant regeneration and rooting was done on MS medium lacking 2,4-D under similar selection conditions. Thirty vigorously growing putative transgenic plants were successfully *ex vitro*-acclimatized and established under glasshouse conditions. Glasshouse spraying of putative transgenic plants with 100 mg/l chlorsulfuron dramatically decreased the amount of non-transgenic plants that had escaped the *in vitro* selection regime. PCR analysis showed that six surviving plants were *als*-positive and that five of these expressed the mutant *als* gene. This report is the first to describe a selection system for sugarcane transformation that uses a selectable marker gene of plant origin targeted by a sulfonylurea herbicide.

**Keywords** Acetolactate synthase · Acetolactate synthase mutation · Sugarcane transformation · Sulfonylurea herbicide

## Introduction

Since the early 1990s of the last century, much progress has been made in the biotechnological manipulation of sugarcane. Both *in vitro* regeneration and genetic transformation were established and are now routinely used (Franks and Birch 1991; Bower and Birch 1992). These technologies, however, depend heavily on the antibiotic selection of *in vitro* genetically transformed cells. In the past, antibiotic and herbicide systems that were predominantly used as selectable marker genes in the process of transgenic sugarcane production included the antibiotic *npII* (neomycin phosphotransferase) and *hpt* (hygromycin phosphotransferase) genes, as well as the *bar* (phosphinotricin acetyltransferase) gene (Enriquez-Obregon et al. 1998; Leibbrandt and Snyman 2003; Joyce et al. 2010). To date, only one other selection system, namely, mannose in combination with the *Escherichia coli* phosphomannose isomerase gene, has been tested in sugarcane (Chengalrayan et al. 2001; Jain et al. 2007). However, the escape rate for this *in vitro* selection system was more than 40%, with 15 of 34 putative transgenic clones being false positives. Since the existing selection systems for use in sugarcane transformation are predominantly based on antibiotic resistance, there is an urgent need for the development of alternative selection systems that might be more acceptable to producers and consumers. Even though some of these antibiotic selectable markers have been declared safe to use in transgenic crops by the US Environmental Protection Agency (1994), they are still not acceptable to a large portion of consumers (Franks 1999; Chern et al. 2002). Developing additional

C. van der Vyver (✉) · T. Conradie · J. Kossmann · J. Lloyd  
Institute for Plant Biotechnology, Department of Genetics,  
University of Stellenbosch, Stellenbosch 7602, South Africa  
e-mail: cvdv@sun.ac.za

alternative selection systems for specific crops will also allow for repeated transformations where more than one selection system is needed for multiple gene transfer in gene stacking approaches into a particular plant species.

In recent years, alternative selection systems have been developed, including the so-called positive selection systems that can be used to supply the putative transgenic plants cells with a metabolic advantage over non-transformed cells (Joersbo and Okkels 1996). In addition, plant genes containing amino acid substitutions leading to decreased herbicide binding have been used for selection (Li et al. 1992; Ogawa et al. 2008; Endo et al. 2012). One such plant gene is acetolactate synthase (ALS) targeted by a large number of herbicides including members of sulfonylurea, imidazolinone, triazolopyrimidine, pyrimidinylthiobenzoate, and sulfonyl-aminocarbonyl-triazolinone (LaRossa and Schloss 1984; Mazur and Falco 1989; Duggleby and Pang 2000). ALS is the first enzyme in the biosynthetic pathway for the branched-chain amino acids valine, leucine, and isoleucine (Chaleff and Mauvais 1984; Yadav et al. 1986; Singh and Shaner 1995). The enzyme is further nuclear-encoded and targeted to the chloroplast (Corbett and Tardif 2006). It catalyzes the condensation of two pyruvate molecules to acetolactate as well as the condensation of pyruvate and ketobutyrate to acetoxybutyrate (LaRossa and Schloss 1984; Schloss et al. 1985; Yadav et al. 1986; Singh and Shaner 1995). Herbicides act on the catalytic region of ALS as competitive inhibitors for pyruvate (Chaleff and Mauvais 1984; McCourt et al. 2006), resulting in the inactivation of the enzyme and the subsequent halting of branched amino acid synthesis that leads to plant death.

Mutations in the *als* gene of *E. coli* (Yadav et al. 1986), *Saccharomyces cerevisiae* (Falco and Dumas 1985; Yadav et al. 1986), and plants (Chaleff and Mauvais 1984; Chaleff and Ray 1984; Lee et al. 1988; Kochevenko and Willmitzer 2003; Khruangchan et al. 2011; Schnell et al. 2012) confer resistance to herbicides. Single nucleotide changes in the gene result in an amino acid substitution, rendering the enzyme less sensitive to inhibition. When reviewed in 2010, there were 22 substitutions at seven sites across *als* genes leading to resistance (Powles and Yu 2010). The most prominent mutation is Pro-197 in which 11 amino acid substitutions affect herbicide resistance in almost 50 plant species (Warwick et al. 2008). Pro-197 mutations result mainly in sulfonylurea resistance, whereas mutations at Ala-205, Asp-376, Trp-574, Ser-653, or Gly-654 cause resistance against imidazolinone and, to a lesser extent, sulfonylurea herbicides in plants such as canola, rice, sugar beet, maize, wheat, and cotton (Sebastian et al. 1989; Swanson et al. 1989; Newhouse et al. 1991, 1992; Rajasekaran et al. 1996; Wright and Penner 1998; Bae et al. 2002; Laplante et al. 2009). So far, only one mutation (Ala-559) in the *als*

gene has been identified in field-grown sugarcane conferring tolerance to the herbicide imazapyr (Punyadee et al. 2007; Khruangchan et al. 2011).

A few of these *als* gene mutations in combination with herbicide use have been developed into *in vitro* plant selection systems (Lee et al. 2007; Sundar and Sakthivel 2008). Commercially available genetically modified crops with the selectable marker gene *als* are available for plant species such as carnation, cotton, canola, soybean, and maize (Miki and McHugh 2004). Previously, a mutant ALS allele from *Arabidopsis* (*csr1-1*) was used as a selectable marker in a rice transformation experiment (Li et al. 1992). Since the transgenic rice was both fertile and showed high resistance to chlorsulfuron, ALS inhibition and protein transport to the chloroplast are conserved between dicots and monocots, demonstrating that the gene can therefore be used as an effective selectable marker in monocots. Applicability for various crops was also confirmed by two studies which used a mutated form of the rice *als* gene as a selectable marker in rice and wheat genetic transformation (Ogawa et al. 2008; Endo et al. 2012). However, none of these *als* mutations have yet been tested and developed for herbicide selection in transgenic sugarcane production.

To date, sugarcane has only been genetically transformed for herbicide resistance with the *pat* (phosphinothricin acetyltransferase) gene, conferring resistance to phosphinothricin (Gallo-Meagher and Irvine 1996; Falco et al. 2000; Leibbrandt and Snyman 2003). Mutation breeding was also used to produce imidazolinone-resistant sugarcane which is effective against a wide spectrum of grass and broadleaf weeds (Koch et al. 2012). In this study, we therefore determined the sensitivity of sugarcane callus and plantlets to herbicides from the sulfonylurea and imidazolinone herbicide classes and report on the effectiveness of a mutated ALS as an *in vitro* selection system for the recovery of putative transgenic sugarcane callus.

## Materials and Methods

*Plant material and establishment of in vitro cultures.* In order to produce embryonic callus, the basal part of sugarcane (*Saccharum* spp. hybrids) cultivar NCo310 leaf roll, just above the apical meristem, was excised, cut into 2-mm-thick transverse sections, and placed on MS medium (Murashige and Skoog 1962; Highveld Biological, Johannesburg, South Africa) containing 2% (*w/v*) sucrose, 3 mg/l 2,4-dichlorophenoxyacetic acid (2,4-D), 0.5 g casein, and 0.22% (*w/v*) gelrite (Sigma-Aldrich, Johannesburg, South Africa), pH 6.0 (MS3). This was incubated for 6 wk at 24°C either in the dark or, if otherwise indicated, under a light regime of 16-h light/8-h dark.

**Herbicide response curves.** The sensitivity of sugarcane callus toward chlorsulfuron, nicosulfuron, rimsulfuron, and the imidazolinone herbicides imazapyr, imazethapyr, and imazaquin was examined (Sigma-Aldrich). A known amount of sugarcane callus (0.25 g) was placed on basic MS3 medium (control) and compared to the callus placed on MS3 medium containing different concentrations of a single herbicide (between 3 and 500  $\mu\text{g/l}$ ). The sensitivity of the sugarcane callus to these herbicides was monitored for the following 15 wk with sub-culturing onto fresh medium every 2 wk. Callus embryogenicity and cell death were visually monitored and callus biomass determined by the increased weight of the starting material after 4 wk. The influence of various parameters, such as growth room light intensities, absence of light, and the presence/absence of casein in the growth medium, was also visually monitored and noted.

Sugarcane plantlets, regenerated from embryogenic callus, were kept at 25°C in natural light in the glasshouse and further sprayed with the six different herbicides at concentrations between 28 and 180 mg/l to determine *ex vitro* sugarcane sensitivity toward the different herbicides tested.

**Isolation of tobacco mutant *als* gene.** Seeds of the *rchl 6.6* chlorosulfuron-resistant tobacco line (Kochevenko and Willmitzer 2003) were germinated and allowed to grow for 4 wk. RNA was extracted from mature plant leaves using the method of Malnoy et al. (2001) and Hu et al. (2002). cDNA was synthesized from total RNA using the Fermentas RevertAid™ H First-strand cDNA synthesis kit according to the manufacturer's specifications (Fermentas, Pretoria, South Africa).

The primers designed for the tobacco *als* coding sequence (accession no. X07645) were ALS FW (5'-AAGGATCCTTCGTCTCTCACT-3') and ALS REV (5'-AACCTAGGAGCTCTGTAGCAC-3') for PCR amplification from the cDNA template. These primers included the *Bam*HI restriction sites on both ends of the PCR product. PCR was performed using the *pfu* DNA polymerase enzyme (Fermentas) according to the manufacturer's specifications. The 2-kb *als* PCR product was purified, sequenced, and ligated into the *Bam*HI site of the pUBI510 plant expression vector under control of the maize ubiquitin promoter and nos terminator (pUBI-*als*).

**Biolistic gene transfer, selection, and regeneration of plants.** Embryogenic callus explants were placed on an osmoticum medium consisting of the basic MS3 medium and 0.2 M of each sorbitol and mannitol (Sigma Aldrich) 4 h prior to and after bombardment (Vain et al. 1993).

Sugarcane callus was transformed using a biolistic particle delivery system as previously described by Franks and Birch (1991), with some modifications. Sterilized tungsten particles (M10; Bio-Rad, Hercules, CA) were mixed with

10  $\mu\text{l}$  plasmid DNA (1  $\mu\text{g}/\mu\text{l}$ ), 50  $\mu\text{l}$  2.5 M  $\text{CaCl}_2$ , and 20  $\mu\text{l}$  0.1 M spermidine (Sigma Aldrich). Particle inflow gun was constructed locally and was used for all biolistic DNA delivery. For bombardment, 5  $\mu\text{l}$  of the particle suspension was placed into the center of a 1-mm<sup>2</sup> metal grid above the explants. Target tissue was placed 16.5 cm below the particle source and the chamber was evacuated to a pressure of 80 kPa before the particles were discharged.

Bombarded tissue was maintained on MS3 medium for 3 d before selection started on predetermined concentrations of chlorsulfuron for 8–10 wk in the dark with sub-culturing onto fresh medium every 2 wk. Tissue was regenerated on herbicide-containing MS medium, lacking 2,4-D, in the light for 8 wk or until roots formed. Calli were regenerated into plantlets on rooting media containing chlorsulfuron to further eliminate non-transgenic clones. Plantlets, 5 cm high with roots, were *ex vitro*-acclimatized in soil-containing pots in the glasshouse. Surviving 2-wk-old plantlets were sprayed with predetermined concentrations of herbicide (100 mg/l chlorsulfuron) to reduce the number of non-transgenic plants that escaped the *in vitro* selection regime.

**PCR analysis of putative transgenic plants.** Putative transformed plants were analyzed for the presence of the *als* transgene using PCR where one primer binds in the ubiquitin promoter region (Ubi-exp: 5'-ATACGCTATTT ATTGCTTGG-3') and the second primer in the *als* gene (ALS REV). Approximately 200 ng of the genomic DNA template was used for a 50- $\mu\text{l}$  PCR reaction. Genomic DNA was isolated according to the manufacturer's instruction of the Fermentas GeneJet Plant Genomic DNA purification kit (Fermentas).


For gene expression analysis, 500 ng of total RNA was extracted from putative transgenic sugarcane leaf material by the method of Malnoy et al. (2001) and Hu et al. (2002). cDNA was synthesized from total RNA with the Fermentas RevertAid™ H First-strand synthesis cDNA kit according to the manufacturer's specifications (Fermentas). Tobacco *als* cDNA-specific primers (forward: 5'-CAATGGGAGGA TCGG TTCTA-3'; reverse: 5'-CAAGTATGGCCCAGG AGTGT-3') and sugarcane actin cDNA primers, as internal controls (forward: 5'-TCACACTTTCTACAATGAGCT-3'; reverse: 5'-GATATCCACATC ACACTT CAT-3'), were used for amplification.

## Results

**Herbicide sensitivity.** Sugarcane callus was placed on different concentrations of sulfonylurea and imidazolinone herbicides. Those tested included the sulfonylurea herbicides—chlorsulfuron, nicosulfuron, and rimsulfuron—and

the imidazolinone herbicides—imazapyr, imazethapyr, and imazaquin. Herbicide concentrations varied between 3 and 500  $\mu\text{g/l}$ . The absence/presence of light had little influence on herbicide activity (Figs. 1 and 2). Sugarcane callus placed in direct light showed severe signs of stress due to the production of phenolic compounds in the cells (visible as a purple coloration in the cells). However, this was not related to the presence of herbicide in the growth medium (Fig. 1). Furthermore, the presence or absence of casein in the growth medium did not significantly influence the sensitivity of the sugarcane callus toward the herbicides (data not shown). Although imidazolinone-type herbicides had very little effect on the growth of sugarcane callus (data not shown), inhibition was found on medium containing chlorsulfuron and rimsulfuron at 3.6 and 200  $\mu\text{g/l}$ , respectively (Table 1 and Figs. 1 and 2). Callus placed on the control MS3 medium showed on average a 10.5-fold increase in fresh weight over 4 wk (0.25–2.63 g), while that placed on chlorsulfuron or rimsulfuron showed no, or at most, a 5-fold increase in weight over 4 wk, respectively (Fig. 2). Prolonged exposure of the callus (more than 12 wk) to the different herbicides did not increase the sensitivity response already obtained during the first 8 wk; rather, it tended to reduce the effect of the initial exposure. Based on visual observations and weight measurements, inhibition of callus growth was most obvious between 6 and 8 wk after callus initiation on chlorsulfuron- or rimsulfuron-containing media (Figs. 1 and 2). Although the exposure of callus to any of the six herbicides did not result in cell death, it did cause loss of callus embryogenesis and a reduction in biomass.

Sugarcane plantlets were further exposed to different herbicide concentrations between 28 and 180 mg/l, and



**Figure 1.** Sugarcane callus growth on basic MS3 medium (control) compared to the medium supplemented with different concentrations of chlorsulfuron, placed under filtered light. Observations shown were recorded after 5 wk on chlorsulfuron. *Square* indicates treatment (3.6  $\mu\text{g/l}$  chlorsulfuron) resulting in the most severe callus growth inhibition.


these showed very low sensitivity to the imidazolinone class of herbicides. Sugarcane plantlets started dying on 100 mg/l of chlorsulfuron and rimsulfuron after 4 wk (Fig. 3). Plant death could, however, be accelerated to 3 wk post-exposure when the herbicide concentration was increased to 140 mg/l.

*ALS isolation and transfer into sugarcane callus.* The mutant *ALS SurB* isoform (*alsb*) was isolated from tobacco cDNA (Kochevenko and Willmitzer 2003). The PCR product was sequenced and a single nucleotide change at base pair 1,719 (TGG to TTG), which results in a Trp-573-Leu amino acid substitution, was confirmed. The insert was ligated in a “sense” orientation with respect to the promoter into the *Bam*HI restriction site of the pUBI 510 plant expression vector, resulting in the vector pUBI-als.

The plant expression vector pUBI-als was bombarded into embryogenic sugarcane callus using 12 plates. Each plate contained a 2.5-cm diameter circle of callus closely stacked together. Bombarded calli were placed on medium containing 3.6  $\mu\text{g/l}$  chlorsulfuron as a selection agent (amount as determined by a herbicide kill curve). Bombarded callus was placed in the dark for 8 wk with sub-culturing onto fresh medium after 2, 4, and 6 wk. Following selection, putative transformed calli were transferred to the light on a regeneration medium lacking 2,4-D. After 8 wk on chlorsulfuron selection, up to 20% of the bombarded calli survived the treatment and showed active growth with a healthy yellow color compared to non-surviving dormant and grayish calli. The herbicide had little effect on the rooting ability of the putative transgenic clones. Regenerated plantlets were transferred to pots containing soil in the greenhouse for *ex vitro* acclimatization. After 2 wk, 30 putative transgenic plantlets were established in the glasshouse. These plants were sprayed with 100 mg/l chlorsulfuron, which reduced the number of plantlets by ~70%, to eight surviving plants. The surviving putative transgenic plants were tested for the presence of the *alsb* gene *via* PCR. Six lines were found to be positively transgenic for the maize ubiquitin promoter (*Ubi-1*)–*alsb* fusion gene (Fig. 3). In five of these transgenic lines, the tobacco *als* gene transcript was verified by reverse transcription polymerase chain reaction (Fig. 4).

## Discussion

In this study, a mutated *als* gene was successfully used for the first time as a selectable marker in combination with the herbicide chlorsulfuron as a selection agent, resulting in the production of transgenic sugarcane lines resistant against the herbicide. The first step in our study toward developing the ALS selection system in sugarcane was to determine the sensitivity of the sugarcane callus toward a range of ALS



**Figure 2.** Sugarcane callus growth on basic MS3 medium (control) compared to the medium supplemented with different concentrations of chlorsulfuron (A–D) and rimsulfuron (E), placed in the dark. Observations were done after 5 wk (A), 10 wk (B, E), and 15 wk (C) on

herbicide. **D.** A close-up view of the callus growth after 10 wk on 0 (control), 3.6, and 20 µg/l chlorsulfuron-containing medium. *Arrows* indicated treatments resulting in the most severe callus growth inhibition on chlorsulfuron (3.6 µg/l) or rimsulfuron (200 µg/l).

targeting herbicides. Callus biomass proliferation was significantly inhibited, up to 10-fold, when exposed to the different herbicides, in particular chlorsulfuron. However, the growth response did not always correlate with an increased herbicide concentration. For example, chlorsulfuron was the most effective herbicide to inhibit callus growth at a concentration of 3.6 µg/l (10 nM), while higher concentrations, such as 50 µg/l chlorsulfuron, did not affect growth at all. A similar low concentration of chlorsulfuron (10 nM) has previously been found to effectively inhibit transgenic rice protoplast growth (Li et al. 1992). On the other hand, chlorsulfuron-resistant maize calli were selected on 50 nM chlorsulfuron, while soybean embryos were only sensitive to ALS herbicides (mix of primisulfuron and prosulfuron)

when a concentration as high as 183 mg/l was applied (Fromm et al. 1990; Rao et al. 2009).

In our study, not all of the herbicides affected sugarcane callus growth. Imadazolinone-type herbicides had very little or no effect on the growth of Nco310 sugarcane callus. This is in contrast to a recent study done with N19 sugarcane callus where regeneration was suppressed when callus was exposed to imazapyr concentrations as low as 0.08 µM (21 µg/l; Koch et al. 2012). The concentrations of imazapyr, imazethapyr, or imazaquin as high as 500 µg/l scarcely affected sugarcane callus biomass and embryogenicity and also did not result in cell death. This is similar to the result found by Heering et al. (1992), where wheat calli were less sensitive to 100 µM imazapyr, in contrast to the whole

**Table 1.** Sugarcane callus growth on MS3 medium supplemented with different concentrations of chlorsulfuron relative to the control medium without supplementation


Environment	Phenotype <sup>z</sup>	Chlorsulfuron concentrations <sup>y</sup>																	
		Control			3.6 µg/l			20 µg/l			50 µg/l			200 µg/l			500 µg/l		
		5	10	15	5	10	15	5	10	15	5	10	15	5	10	15	5	10	15
Filtered light	Biomass	0	0	0	-2	-3	-2	1	0	-1	0	-2	-1	1	0	0	1	-2	-2
	Embryogenesis	2	1	0	-1	-1	0.5	1	-1	0	1	-1	0	2	1	0	1	-1	0
	Cell death	0	0	2	0	0	1	0	1	0	0	0	0	0	0	1	0	0	0
Dark	Biomass	0	0	0	-2	-2	-2	-1	-3	-1	0	0	0	0	0	0	0	1	0
	Embryogenesis	1	1	1	0	-3	0	0	-2	0	1	1	1	1	1	1	1	1	1
	Cell death	0	0	0	0	1	1.5	0	2	1	0	0	0	0	1	0	0	0	0
Light	Biomass	0	0	0	0	-2	1	0	-1	-2	0	-1	0	0	-1	0	0	-1	-1
	Embryogenesis	1	1	0	1	-1	0	1	1	0	1	1	0	1	1	0	1	1	0
	Cell death	0	0	1	0	0	0	0	0	2	0	0	1	0	1	1	0	0	0
Filtered light, no casein	Biomass	0	0		0.5	1		-0.5	-2		0	0		0	0.5		0.5	1	
	Embryogenesis	1	0		1	0		1	0		1	0		1	0		1	0	
	Cell death	0	1		0	1		2	1		2	1		0	1		1	1	
Dark, no casein	Biomass	0		0	-0.5		1	0		0	0		0	0		-1	0		0
	Embryogenesis	2		1	1		1	1		1	1		1	2		0	2		1
	Cell death	0		0	2		0	1.5		0	2		0	0		0	1		0

<sup>z</sup> Observations were recorded after 5, 10, and 15 wk on chlorsulfuron. Biomass: control=0, >1 more than control, <1 less than control; Embryogenesis: no embryos=0, >1=visible embryos, <1 loss of cell viability; Cell death: no cell death=0, >1=visible cell death  
<sup>y</sup> Values in *italics* indicate herbicide concentrations where callus survival was the most affected

wheat plant. Therefore, the effectiveness of herbicide concentrations on growth differs significantly across plant tissues and plant species.


The ALS selection system was less effective in sugarcane when compared to other well-established systems, such as the *nptII* gene and use of geneticin. In the past, when the geneticin/*nptII* system was applied for selecting transgenic

sugarcane calli, 1–5% of the total bombarded calli would survive with almost no escapes (Bower and Birch 1992; Snyman et al. 1996). In our chlorsulfuron/*als* system, however, up to 20% of the bombarded calli survived after 8 wk of *in vitro* selection with insufficient new growth to easily differentiate between transgenic and non-transgenic calli. Also, from the surviving calli, 30 putative transgenic plants



**Figure 3.** Analysis of putative transgenic plants which survived the *in vitro* selection regime. *A*, PCR amplification of a fragment of the fused ubiquitin promoter (*ubi*) and mutant acetolactate synthase *B* (*alsb*) gene from gDNA extracted from putative transformed pUBI-*als* sugarcane plants. Lane *M*, λDNA digested with *Pst*I; *H*<sub>2</sub>*O*, water control; *C* control Nco310 sugarcane cultivar, *Pos* positive control, pUBI-*als*

vector. Lines 1, 3, 5–8 were positive transgenic plants. Lines 2 and 4 were negative. *B*, Four weeks after putative transgenic plants were sprayed with 100 mg/l of chlorsulfuron in the glasshouse. Lines 1, 3, 5–8 survived the herbicide spraying. Lines 2 and 4 died after chlorsulfuron spraying (white arrows). Lines correspond to the lanes indicated in the PCR analysis.



**Figure 4.** PCR amplification using sugarcane cDNA as the template to determine gene expression. (A) Internal standard as amplified with sugarcane actin primers. (B) Expression of the tobacco *als* gene in

sugarcane leaves. Lane M,  $\lambda$ DNA digested with *Pst*I. *H<sub>2</sub>O*, water control; C, control Nco310 sugarcane cultivar. Lanes 1–5 represent five transgenic sugarcane clones containing the tobacco *als* gene insert.

regenerated, of which six contained the *als* gene, an escape rate of 20%. Escapes could be dramatically reduced in our study by ~70% with *ex vitro* herbicide spraying. Similar to the escape rate for sugarcane, the ALS selectable marker system applied in wheat had an escape rate of 7.7–60% of surviving embryos not containing the *als* selectable marker gene (Ogawa et al. 2008). Therefore, based on the escape rate of the chlorsulfuron/*als* system reported here, it seems to be comparable to ALS systems developed for other crops, but less efficient than the geneticin/*npt*II system in sugarcane.

A disadvantage of sulfonylurea and imidazolinone herbicide application for selection might be the lack of induced cell death. In this study, exposure of callus to these herbicides only resulted in biomass reduction and loss of embryogenesis. A similar result has been reported for the mannose selection system in sugarcane where different concentrations of mannose inhibited growth by 50–80%, but could not completely suppress callus growth (Jain et al. 2007). Also, wheat callus exposed to imazapyr only showed 50% growth inhibition after 70 d. (Heering et al. 1992).

Our results have further shown that light was not required to ensure herbicide activity. Since callus cultured under light turned purple due to excessive accumulation of anthocyanin pigments as a result of light stress, application of the selection system should preferably be done in the dark. In the dark, coloration changes included a change from a healthy pale yellow to a grayish white, and shades in between. Very long exposure to the different herbicides beyond 6–8 wk also did not result in greater growth inhibition of sugarcane callus or cell death as callus seems to overcome the inhibitory growth effect of the herbicides.

Overall, in this study, the expression of the *als* gene, together with chlorsulfuron as a selection agent, was successfully used to produce transgenic sugarcane plants. The *in vitro* selection system was complimented and further improved by a single *ex vitro* herbicide spray application of surviving plantlets. Since a large variation exists in the sensitivity of plant tissues to different herbicides, in the future, it may be useful to also test additional herbicide classes, such as pyrimidinylcarboxylates, which also target the *als* gene, in sugarcane. In general,

applying ALS in combination with sulfonylurea or imidazolinone herbicides for the selection of transgenic plants has several advantages. Firstly, these herbicides have little or no known toxic side effects on animals and no measurable mutagenic properties (Levitt 1983; Powles and Yu 2010). In addition, the use of a mutated *als* gene is less likely to prompt concerns regarding food safety in GM plants because a native *als* gene exists in all plants. The mutated *als* gene can also act both as an *in vitro* selectable marker gene during sugarcane genetic transformation as well as a genetic insertion event that conveys resistance to herbicides in the field during crop production. In this regard, a sugarcane crop with a herbicide resistance trait will supply much needed protection against weeds during crop production. It has been shown that weeds can reduce sugarcane yields by more than 40%, greatly increasing the cost of production (Millhollon 1992, 1995). ALS-inhibiting herbicides are also used for weed control in the South African sugar industry, and these ALS-inhibiting herbicides account for about 17.5% of the total global herbicide market (Green 2007). Introduction of a herbicide-resistant sugarcane would not only provide a useful selection system but would ultimately also be beneficial to sugarcane producers.

**Acknowledgments** We would like to thank Prof. Lothar Willmitzer (Max Planck Institute of Molecular Plant Physiology, Germany) for providing us with seeds from the *rchl* 6.6 chlorosulfuron-resistant tobacco line. This project was funded by Biosafety, South Africa ([www.biosafety.org.za](http://www.biosafety.org.za)).

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- Bae C, Young-Ill L, Yong-Pyo L, Yong-Won S, Do-Jin L, Deuk-Chum Y, Hyo-Yeon L (2002) Selection of herbicide tolerant cell lines from  $\gamma$ -ray-irradiated cell cultures in rice (*Oryza sativa* L. cv. Ilpumbyeo). *J Plant Biotechnol* 4:123–127
- Bower R, Birch R (1992) Transgenic sugarcane plants via micro-projectile bombardment. *Plant J* 2:409–416

- Chaleff RS, Mauvais CJ (1984) Acetolactate synthase is the site of action of two sulfonylurea herbicides in higher plants. *Science* 224:1443–1445
- Chaleff RS, Ray TB (1984) Herbicide resistant mutant from tobacco cell cultures. *Science* 223:1148–1161
- Chengalrayan K, Gallo-Meagher M, English RG (2001) Novel selection agents for sugarcane transformation. *Soil Crop Sci Soc Florida Proc* 60:81–87
- Chern WS, Rickertsen K, Tsuboi N, Fu TT (2002) Consumer acceptance and willingness to pay for genetically modified vegetable oil and salmon: a multiple-country assessment. *AgBioForum: J Agrobiotechnol Manag Economics* 5:105–112. <http://www.agbioforum.org>. Accessed 1 November 2012
- Corbett CL, Tardif FJ (2006) Detection of resistance to acetolactate synthase inhibitors in weeds with emphasis on DNA-based techniques: a review. *Pest Manag Sci* 62:584–597
- Duggleby RG, Pang SS (2000) Acetohydroxyacid synthase. *J Biochem Mol Biol* 33:1–36
- Endo M, Shimizu T, Toki S (2012) Selection of transgenic rice plants using a herbicide tolerant form of the *acetolactate synthase* gene. *Methods Mol Biol* 847:59–66
- Enriquez-Obregon GA, Nazquez PR, Prieto SD, Riva-Gustavo AD, Selmann HG (1998) Herbicide resistant sugarcane (*Saccharum officinarum* L.) plants by *Agrobacterium*-mediated transformation. *Planta* 206:20–27
- Falco MC, Neto AT, Ulian EC (2000) Transformation and expression of a gene for herbicide resistance in a Brazilian sugarcane. *Plant Cell Rep* 19:1188–1194
- Falco SC, Dumas KS (1985) Genetic analysis of mutants of *Saccharomyces cerevisiae* resistant to the herbicide sulfometuron methyl. *Genetics* 1099:21–35
- Franks JR (1999) The status and prospects for genetically modified crops in Europe. *Food Policy* 24:565–584
- Franks T, Birch RG (1991) Gene transfer into intact sugarcane callus using microprojectile bombardment. *Aus J Plant Physiol* 18:471–480
- Fromm ME, Morrish F, Armsoren C, Williams R, Thomas J, Klein TM (1990) Inheritance and expression of chimeric genes in the progeny of transgenic maize plants. *Biotechnology* 8:833–839
- Gallo-Meagher M, Irvine JM (1996) Herbicide resistant transgenic sugarcane plants containing the bar gene. *Crop Sci* 36:1367–1374
- Green JM (2007) Review of glyphosate and ALS-inhibiting herbicide crop resistance and resistant weed management. *Weed Technol* 21:547–558
- Heering DC, Guenzi AC, Peeper TF, Claypool PL (1992) Growth response of wheat (*Triticum aestivum*) callus to imazapyr and *in vitro* selection of resistance. *Weed Sci* 40:174–179
- Hu CG, Honda C, Kita M, Zhang Z, Tsuda T, Moriguchi T (2002) A simple protocol for RNA isolation from fruit trees containing high levels of polysaccharides and polyphenol compounds. *Plant Mol Biol Rep* 20:69/1–69/7
- Jain M, Chengalrayan K, Abouzid A, Gallo M (2007) Prospecting the utility of a PMI/mannose selection system for the recovery of transgenic sugarcane (*Saccharum* spp. hybrid) plants. *Plant Cell Rep* 26:581–590
- Joersbo M, Okkels FT (1996) A novel principle for selection of transgenic plant cells: positive selection. *Plant Cell Rep* 16:219–221
- Joyce P, Kuwahata M, Turner N, Lakshmanan P (2010) Selection system and co-cultivation medium are important determinants of *Agrobacterium*-mediated transformation of sugarcane. *Plant Cell Rep* 29:173–183
- Khruangchan N, Chatchawankanphanich O, Pornprom T (2011) *Acetolactate synthase* gene mutation conferring imazapyr tolerance in sugarcane clone. *NRCT J* 41:115–127
- Koch AC, Ramgareeb S, Rutherford RS, Snyman SJ, Watt MP (2012) An *in vitro* mutagenesis protocol for the production of sugarcane tolerant to the herbicide imazapyr. *In Vitro Cell Dev Biol—Plant* 48:417–427. doi:10.1007/s11627-012-9448-x
- Kochevenko A, Willmitzer L (2003) Chimeric RNA/DNA oligonucleotide-based site-specific modification of the tobacco *acetolactate synthase* gene. *Plant Physiol* 132:174–184
- LaRossa RA, Schloss JV (1984) The sulfonylurea herbicide sulfometuron methyl is an extremely potent and selective inhibitor of acetolactate synthase in *Salmonell typhimurium*. *J Biol Chem* 259:8753–8757
- Laplante J, Rajcan I, Tardif FJ (2009) Multiple allelic forms of acetohydroxyacid synthase are responsible for herbicide resistance in *Setaria viridis*. *Theor Appl Genet* 119:577–586
- Lee K, Yang K, Kang K, Kang S, Lee N, Back K (2007) Use of *Myxococcus xanthus* protoporphyrinogen oxidase as a selectable marker for transformation of rice. *Pest Biochem Physiol* 88:31–35
- Lee KY, Townsend J, Tepperman J, Black M, Chui CF, Mazur B, Dunsmuir P, Bedbrook J (1988) The molecular basis of sulfonylurea herbicide resistance in tobacco. *EMBO J* 7:1241–1248
- Leibbrandt NB, Snyman SJ (2003) Stability of gene expression and agronomic performance of a transgenic herbicide-resistant sugarcane line in South Africa. *Crop Sci* 43:671–678
- Levitt G (1983) Sulfonylureas, new herbicides. In: Miyamoto J, Keraney PC (eds) *Pesticide chemistry, human welfare and the environment*. Oxford University Press, New York, pp 243–250
- Li Z, Hayashimoto A, Murai N (1992) A sulfonylurea herbicide resistance gene from *Arabidopsis thaliana* as a new selectable marker for production of fertile transgenic rice plants. *Plant Physiol* 100:662–668
- Malnoy M, Reynoird JP, Mourgues F, Chevreau E, Simoneau P (2001) A method for isolating total RNA from pear leaves. *Plant Mol Biol Rep* 16:69
- Mazur BJ, Falco SC (1989) The development of herbicide resistant crops. *Annu Rev Plant Physiol Plat Mol Biol* 40:441–470
- McCourt JA, Pang SS, King-Scott J, Guddat LW, Duggleby RG (2006) Herbicide-binding sites revealed in the structure of plant acetohydroxyacid synthase. *Proc Natl Acad Sci U S A* 103:569–573
- Miki B, McHugh S (2004) Selectable marker genes in transgenic plants: applications, alternatives and biosafety. *J Biotechnol* 107:193–232
- Millhollon RW (1992) Effect of itchgrass (*Rottboellia cochinchinensis*) interference on growth and yield of sugarcane (*Saccharum* spp. hybrids). *Weed Sci* 40:48–53
- Millhollon RW (1995) Growth and yield of sugarcane as affected by johnsongrass (*Sorghum halepense*) interference. *J Am Soc Sugarcane Technol* 15:32–40
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 15:473–497
- Newhouse KE, Singh BK, Shaner DL, Stidham M (1991) Mutations in corn (*Zea mays*) conferring resistance to imidazolinones. *Theor Appl Genet* 83:65–70
- Newhouse KE, Smith W, Starret M, Schaefer T, Singh BK (1992) Tolerance to imidazolinone herbicides in wheat. *Plant Physiol* 100:882–886
- Ogawa T, Kawahigashi H, Toki S, Handa H (2008) Efficient transformation of wheat by using a mutated rice *acetolactate synthase* gene as a selectable marker. *Plant Cell Rep* 27:1325–1331
- Powles SB, Yu Q (2010) Evolution in action: plants resistant to herbicides. *Annu Rev Plant Biol* 61:317–347
- Punyadee P, Thongros M, Pornprom T (2007) Biochemical mechanism of resistance to imazapyr in sugarcane cell selections. *Thai J Agric Sci* 40:133–141
- Rajasekaran K, Grula JW, Anderson DM (1996) Selection and characterization of mutant cotton (*Gossypium hirsutum* L.) cell lines



- resistant to sulfonylurea and imidazolinone herbicides. *Plant Sci* 199:115–124
- Rao SS, Mamodou L, McConnell M, Polisetty R, Kwanyuen P, Hildebrand D (2009) Non-antibiotic selection systems for soybean somatic embryos: the lysine analog aminoethyl-cysteine as a selection agent. *BMC Biotechnol* 9:94
- Sebastian SA, Fader MG, Ulrich FJ, Forney RD, Chaleff SR (1989) Semidominant soybean mutation for resistance to sulfonylurea herbicides. *Crop Sci* 199:115–124
- Schloss JV, Van Dyk DE, Vasta JF, Kutny RM (1985) Purification and properties of *Salmonella typhimurium* acetolactate synthase isozyme II from *Escherichia coli* HB101/pDU9. *Biochemistry* 24:4952–4959
- Schnell J, Labbe H, Kovinich N, Manabe Y, Miki B (2012) Comparability of imazapyr-resistant *Arabidopsis* created by transgenesis and mutagenesis. *Transgenic Res* 21:1255–1264. doi:10.1007/s112480129597z
- Singh BK, Shaner DL (1995) Biosynthesis of branched amino acids: from test tube to field. *Plant Cell* 7:935–944
- Snyman SJ, Meyer GM, Carson DL, Botha FC (1996) Establishment of embryogenic callus and transient gene expression in selected sugarcane varieties. *S Afr J Bot* 62:151–154
- Sundar IK, Sakthivel N (2008) Advances in selectable marker genes for plant transformation. *J Plant Physiol* 165:1698–1716
- Swanson EB, Hergesell MJ, Arnoldo M, Sippel DW, Wong RSC (1989) Microspore mutagenesis and selection: canola plants with field tolerance to imidazolinones. *Theor Appl Genet* 78:525–530
- Environmental Protection Agency US (1994) Neomycin phosphotransferase: II. Tolerance exemption. *Fed Regist* 56:4935
- Vain P, McMullen MD, Finer JJ (1993) Osmotic treatment enhances particle bombardment-mediated transient and stable transformation of maize. *Plant Cell Rep* 12:84–88
- Warwick SI, Xu R, Sauder C, Beckie HJ (2008) Acetolactate synthase target-site mutations and single nucleotide polymorphism genotyping in ALS-resistant kochia (*Kochia scoparia*). *Weed Sci* 56:797–806
- Wright T, Penner D (1998) Cell selection and inheritance of imidazolinone resistance in sugar beet (*Beta vulgaris*). *Theor Appl Genet* 96:612–620
- Yadav N, McDevitt RE, Benard S, Falco SC (1986) Single amino acid substitutions in the enzyme acetolactate synthase confer resistance to the herbicide sulfometuron methyl. *Proc Natl Acad Sci U S A* 83:4418–4422