



What is the role of nephrologists and nurses of the dialysis department in providing fertility care to CKD patients? A questionnaire study among care providers

Gaby F. van Ek¹ · Esmée M. Krouwel¹ · Melianthe P. J. Nicolai¹ ·
Brenda L. Den Oudsten² · Marjolein E. M. Den Ouden³ · Sandra W. M. Dieben⁴ ·
Hein Putter⁵ · Rob C. M. Pelger¹ · Henk W. Elzevier¹

Received: 2 January 2017 / Accepted: 20 March 2017 / Published online: 29 March 2017
© The Author(s) 2017. This article is an open access publication

Abstract

Purpose This study evaluated current fertility care for CKD patients by assessing the perspectives of nephrologists and nurses in the dialysis department.

Methods Two different surveys were distributed for this cross-sectional study among Dutch nephrologists ($N = 312$) and dialysis nurses ($N = 1211$).

Results Response rates were 50.9% (nephrologists) and 45.4% (nurses). Guidelines on fertility care were present in the departments of 9.0% of the nephrologists and 15.6% of the nurses. 61.7% of the nephrologists and 23.6% of the nurses informed $\geq 50\%$ of their patients on potential changes in fertility due to a decline in renal function. Fertility subjects discussed by nephrologists included “wish to have children” (91.2%), “risk of pregnancy for patients’ health” (85.8%), and “inheritance of the disease” (81.4%). Barriers withholding nurses from discussing FD were based on “the age of the patient” (62.6%), “insufficient training” (55.2%), and “language and ethnicity” (51.6%).

29.2% of the nurses felt competent in discussing fertility, 8.3% had sufficient knowledge about fertility, and 75.7% needed to expand their knowledge. More knowledge and competence were associated with providing fertility health care ($p < 0.01$).

Conclusions In most nephrology departments, the guidelines to appoint which care provider should provide fertility care to CKD patients are absent. Fertility counseling is routinely provided by most nephrologists, nurses often skip this part of care mainly due to insufficiencies in self-imposed competence and knowledge and barriers based on cultural diversity. The outcomes identified a need for fertility guidelines in the nephrology department and training and education for nurses on providing fertility care.

Keywords Chronic kidney disease · Fertility care · Practice patterns · Questionnaires · Renal care providers

Introduction

Chronic kidney disease (CKD) is associated with a decrease in reproductive function [1–4]. Although fertility disorders (FD) are common in both male and female patients, the precise etiology is largely unknown. A major component is a disturbance in the hypothalamic–pituitary axis caused by CKD, resulting in menstrual irregularities, anovulation, and infertility in female patients [1]. For example, over 94% of female patients receiving dialysis experience menstrual irregularities [5]. If pregnancy does occur, patients are highly at risk for further deterioration of their renal function, pre-eclampsia, and the need for blood transfusions [1, 6, 7]. In addition, pregnancy and possible transfusions may complicate future renal transplantation as immunological sensitization might be induced [1]. The

✉ Gaby F. van Ek
G.F.van_Ek@lumc.nl

¹ Department of Urology, Leiden University Medical Center, Albinusdreef 2, PO Box 9600, 2300 RC Leiden, The Netherlands

² Department of Medical and Clinical Psychology and Centre of Research on Psychological and Somatic Disorders, Tilburg University, Tilburg, The Netherlands

³ Saxion, University of Applied Sciences, Enschede, The Netherlands

⁴ Department of Gynecology, Leiden University Medical Center, Leiden, The Netherlands

⁵ Department of Medical Statistics, Leiden University Medical Center, Leiden, The Netherlands

fetus is also at risk for intrauterine growth restrictions, along with multiple complications associated with preterm delivery [1, 6, 7]. In male CKD patients, besides disturbances in the hypothalamic–pituitary axis, oxidative stress and uremia both contribute to FD by decreasing testicular volume and impairment of spermatogenesis [2, 3].

Where little is documented on fertility management and counseling in male CKD patients, several studies exist on the management of fertility in female CKD patients [1, 8]. Counseling prior to conception is necessary for all female patients of reproductive age due to the complexity of fertility in CKD [1, 8]. Every disease stage requires different counseling, and adequate information on fertility is especially needed for those patients receiving renal replacement therapy [6]. Although FD are common during dialysis, prevention of unwanted pregnancy is necessary since recent innovations in dialysis induce hormonal normalization and thereby increase pregnancy rates up to 75% [1, 6, 7]. If pregnancy does occur, early confirmation is of great importance as close monitoring of renal function, patients' health, and the pregnancy itself is essential [1, 6, 7, 9].

Adequate fertility care, including counseling, management, and referral to a physician specialized in fertility, determines whether the pregnancy and maternal and fetal outcomes are successful [1]. A multidisciplinary approach is fundamental for success in this part of renal health care [1, 8]. Unfortunately, little is known about current fertility care for CKD patients provided by renal care providers as this subject is often ignored in research. No information is available on the current format of this part of renal care or the availability of guidelines appointing which care providers are accountable for addressing fertility issues. Since the nephrologist is a major renal care provider throughout all stages of CKD, it is most likely that the provision of fertility care depends on this renal care provider. However, considering the nephrologist's increasing workload and a limited time available during consultation, nurses might play an important role in the future for providing fertility care for CKD patients. Nurses working in the dialysis unit could especially contribute to this part of renal care since they often have frequent and intensive one-on-one contact with their patients.

The aim of this multidisciplinary study was to examine current fertility care for CKD patients from the perspectives of the nephrologists and nurses working in the dialysis unit. Guidelines appointing who should provide fertility care within nephrology departments were focused on, as were practice patterns of nephrologists and nurses regarding informing, discussing, counseling on fertility, and referral to a physician specialized in fertility. In addition, the evaluation among nurses addressed possible barriers toward discussing fertility care and tools needed to provide it. Nurses' competence, knowledge, and education regarding fertility

in CKD patients were evaluated, as well as how these factors influence fertility care currently provided by them.

Methods

Study design

Data for this cross-sectional study were collected among nephrologists and nurses using two separate surveys. All practicing Dutch nephrologists ($n = 318$) who were member of the Dutch Federation of Nephrology (Nefrovisie) were requested to participate in the survey. A total of 312 questionnaires were sent to nephrologists' home addresses; six obtained addresses were out of date. Non-respondents received a reminder letter two and/or four months after the initial mailing.

For the survey among nurses, all Dutch dialysis centers (both in and outside of the hospital) were approached for participation ($n = 63$). Thirty-four centers (54.0%) agreed on participation, and all their employed dialysis nurses and nurses specialized in nephrology received a questionnaire ($n = 1171$) at their work address. An e-mail was sent to non-responding centers received 2 and/or 4 months after the initial invitation as a reminder. Participating centers received a motivational e-mail after 4 months asking them to motivate their staff on returning the questionnaire. In addition, 40 questionnaires were handed out during an informative meeting for nurses specialized in nephrology resulting in a total of 1211 distributed questionnaires.

Instrument design

The authors developed both questionnaires as validated instruments for assessing the aims of the studies were not available. The questionnaires were derived from instruments used in previous studies evaluating healthcare in medical departments [10–13]. Contents were based on issues described in the literature and additional topics identified by the authors. The classification of renal failure was according to K/DOQI clinical guidelines [9]. Both questionnaires were part of surveys that evaluated the discussion of sexual dysfunction as well. The items on sexual dysfunction were studied separately [14]. The questionnaire used in the nephrologist study was pilot-tested for content, length, and comprehensiveness by nephrologists and nephrology residents from the Leiden University Medical Centre ($n = 7$). No modifications were made as no remarks were made. The first sheet of the questionnaire contained demographic questions and an opt-out possibility. The (remaining) ten items assessed:

- Guidelines on fertility care within the nephrology department.
- Nephrologists' practice patterns regarding informing, discussing, and counseling on fertility, including referral to physicians specialized in fertility.
- Fertility subjects addressed by the nephrologist.

The nurses' questionnaire was pilot-tested at the Leiden University Medical Centre as well. Twenty-three nurses working in the dialysis department inspected layout, linguistics, and comprehensiveness of items. No comments were made, so the final questionnaire was identical. Demographic questions and an opt-out possibility were placed on the first sheet; the other ten questions assessed:

- Guidelines on fertility care within the dialysis or nephrology department.
- Nurses' practice patterns regarding informing, discussing, and counseling on fertility with dialysis patients.
- Competence of the nurse in providing fertility care to dialysis patients.
- Possible barriers to discussing fertility with dialysis patients.
- Current knowledge and education on fertility in dialysis patients, including influence on fertility care currently provided by the nurse.
- Tools that could assist in providing fertility care to dialysis patients.

Data analysis

Data were analyzed using IBM SPSS statistics 23 (SPSS Inc., Chicago, IL, USA). Descriptive statistics were used to describe demographic information and the answers given by the respondents. Bivariate associations were calculated using the Cochran–Armitage trend test. Two-sided p values of <0.05 were considered statistically significant.

Ethical considerations

Since no patients or interventions were involved during both surveys, no formal ethical approval was needed in the Netherlands.

Results

Participation

In the nephrologist survey, a total of 159 of the 312 questionnaires sent were returned (response rate 50.9%). Thirty-eight respondents were not willing to participate, and reasons for not participating were, for instance, “lack of time”

($n = 28$), “not practicing” ($n = 5$), and “not interested” ($n = 2$). Fifteen respondents were excluded since they were specialized in pediatric nephrology. A total of 113 (36.2%) questionnaires were analyzed.

Of the 1211 questionnaires sent to the nurses, 550 were returned (response rate 45.4%). Twenty-three nurses declined participation, three questionnaires were excluded because less than 50% was filled in, and one respondent was excluded for not working as a nurse. Reasons not to participate were, for example, “no interest” ($n = 7$), “insufficient time” ($n = 4$), and “not enough experience” ($n = 2$). Including the 23 questionnaires from the pilot study, a total of 546 surveys were used for analyses. Demographic information of the respondents of both surveys is displayed in Table 1.

Guidelines on providing fertility care

Both surveys assessed if clear guidelines were present within nephrology departments regarding which care provider is accountable for discussing patients' fertility. Ten nephrologists (9.0%) answered positively on the question if clear guidelines were present. Seven (6.3%) noted that their guidelines indicated the nephrologist was accountable for advising CKD patients on fertility matters, and one nephrologist (0.9%) noted the gynecologist was responsible. Of the nurses, 84 (15.6%) were aware of guidelines within their department regarding the accountability for discussing patients' fertility. Thirty-one nurses (6.2%) stated these guidelines pointed out their own group of professionals as accountable. Other answers given concerning accountability were the “nephrologist” ($n = 22$, 4.4%), “nephrologist and nurse together” ($n = 17$, 3.4%), and “nephrologist, nurse and nephrology social worker” ($n = 5$, 1.0%).

Providing information

Nephrologists and nurses were asked how often they informed their dialysis patients about the association between a decline in renal function, dialysis, and reduced fertility. Their answers on this question are listed in Table 2. More experienced nurses and nurses who were aware of guidelines regarding the accountability for discussing patients' fertility informed their patients more often on reduced fertility (Linear-by-Linear Association, $p = 0.004$ resp. $p < 0.001$). No association was found between experience or awareness of guidelines and providing information among nephrologists.

Discussion of fertility by the nephrologist

A majority of the nephrologists ($n = 96$, 88.1%) advised their female patients with renal failure not to become pregnant. Fifty-eight (62.4%) did so when their patients reached CKD stage 4 and 12.9% ($n = 12$) discouraged

Table 1 Demographic characteristics

	Nurses (<i>n</i> = 546)	<i>n</i> ^a (%)	Nephrologists (<i>n</i> = 113)	<i>n</i> ^a (%)
Gender			Gender	
Male		55 (10.1)	Male	70 (61.9)
Female		491 (89.9)	Female	43 (38.1)
Age (years)			Age (years)	
Median 47.0 (range 22–65)		542 (99.3)	Median 47.0 (range 33–62)	113 (100.0)
Mean 44.8			Mean 47.16	
Position			Position	
Dialysis nurse		491 (89.9)	Nephrologist	111 (98.2)
Nurse in dialysis registration training		16 (3.0)	Resident	2 (1.8)
Team leader dialysis department		17 (3.2)		
Nurse specialized in nephrology		19 (3.5)		
Other ^b		22 (3.9)		
Years of employment			Years of employment	
<1 year		13 (2.4)	<1 year	1 (0.9)
1–2 years		24 (4.4)	1–2 years	2 (1.8)
3–5 years		66 (12.1)	3–5 years	13 (11.5)
6–10 years		134 (24.5)	6–10 years	28 (24.8)
11–15 years		106 (19.4)	11–15 years	19 (16.8)
>15 years		203 (37.2)	>15 years	50 (44.2)
Clinical setting			Clinical setting	
University hospital		64 (11.7)	University hospital	27 (23.9)
District general teaching hospital		214 (39.2)	District general teaching hospital	54 (47.8)
District general hospital		184 (33.7)	District general hospital	30 (26.5)
Tertiary and district general hospital		6 (1.1)	Tertiary and district general hospital	1 (0.9)
Dialysis centre, outside the hospital		94 (17.2)	Dialysis centre, outside the hospital	6 (5.3)

^a *n* may differ due to multiple answers that could be given to questions or because the questions were not answered consistently, some were skipped or forgotten

^b Includes, e.g., diabetic nurse, pre-dialysis nurse, quality officer

Table 2 Providing information on fertility

How often do you inform your (dialysis) patients that a decline in renal function and dialysis are associated with reduced fertility?	Never <i>n</i> (%)	<50% of the cases <i>n</i> (%)	In 50% of the cases <i>n</i> (%)	>50% of the cases <i>n</i> (%)	Always <i>n</i> (%)
Nephrologists (<i>n</i> = 110)	12 (10.9)	30 (27.3)	16 (14.5)	27 (24.5)	25 (22.7)
Nurses (<i>n</i> = 537)	190 (35.3)	221 (41.0)	41 (7.6)	47 (8.7)	40 (7.4)

n differs because the questions were not answered consistently, and some were skipped or forgotten

pregnancy in stage 5. Other nephrologists advised their patients to avoid pregnancy during stage 1 (*n* = 2, 2.2%), stage 2 (*n* = 1, 1.1%), and stage 3 (*n* = 20, 21.5%) of CKD. Besides advising their female patients to prevent pregnancy before renal transplantation, nephrologists discussed several other subjects regarding fertility with their female patients. These included: wish to have children (*n* = 103, 91.2%), health risks of pregnancy (*n* = 97, 85.8%), and inheritance of the disease (*n* = 92, 81.4%). With their male patients, they also discussed their wish to have children (*n* = 85, 75.2%) and inheritance of the

disease (*n* = 76, 67.3%). On top of that, almost three-quarter of nephrologists (*n* = 84, 74.3%) inquired after erectile dysfunction. With regards to referring CKD patients to a physician specialized in fertility, on average the nephrologists referred 14.5% (SD 27.0) of female patients and 9.7% (SD 22.5) of male patients.

Discussion of fertility by the nurse

Nurses were asked how often they discussed fertility with male and/or female patients and in how many cases the

partner was present when they did. Answers are listed in Table 3.

Nurses with more years of experience discussed fertility more often with male (Linear-by-Linear Association, $p = 0.001$) and female patients (Linear-by-Linear Association, $p = 0.001$). On the question “What could be helpful for the discussion of fertility”, 86.0% ($n = 467$) answered informative brochures, 55.6% ($n = 302$) thought training to discuss fertility would be helpful, and 15.1% ($n = 82$) recommended posters about fertility in the waiting room. Almost 45% ($n = 242$, 44.6%) felt that the nephrologist should initiate the conversation on fertility so nurses could then refer to that conversation. Self-reported answers to this question were, for example, “more privacy or private appointments to discuss fertility” ($n = 22$, 4.4%), “more knowledge” ($n = 12$, 2.4%), and “referral to a physician” ($n = 4$, 0.8%).

Barriers experienced by the nurse

The survey contained a list of possible barriers that might prevent nurses from discussing fertility. They were asked to write down to which extent they agreed with the barriers. The barriers are listed in Table 4. Barriers most agreed on by the nurses were based on, “age of the patient” (62.6%), “insufficient training” (55.2%), “language and ethnicity” (51.9%), “insufficient knowledge” (49.4%) and “culture or religion” (47.1%).

Knowledge and competence of the nurse

On the question “Is the subject fertility in CKD patients addressed during your training to become a nurse?” two-thirds of nurses ($n = 351$) answered affirmatively. Next,

Table 3 Discussion of fertility by the nurse

	Never n (%)	Seldom n (%)	Routinely n (%)	Often n (%)
Male patients	284 (53.2)	218 (40.8)	27 (5.1)	5 (0.9)
Female patients	235 (44.0)	253 (47.4)	37 (6.9)	9 (1.7)
	Never n (%)	<50% of the cases n (%)	≥50% of the cases ^a n (%)	Always n (%)
How often is the partner present?	359 (72.2)	79 (15.9)	37 (7.4)	22 (4.4)

n differs because the questions were not answered consistently, and some were skipped or forgotten

^a ≥50% of the cases contains the answers “in 50% of the cases” and “more than 50% of the cases”

Table 4 Barriers for nurses not to discuss fertility

	Agree n (%) ^a	Indecisive n (%)	Disagree n (%) ^b
Age of the patient	333 (62.6)	114 (21.4)	85 (16.0)
Insufficient training	295 (55.2)	146 (27.3)	93 (17.4)
Barriers based on language and ethnicity	275 (51.6)	178 (33.4)	80 (15.0)
Insufficient knowledge	265 (49.4)	164 (30.6)	107 (20.0)
Barriers based on culture or religion	251 (47.1)	175 (32.8)	107 (20.1)
Could not find a suitable moment	230 (43.4)	158 (29.8)	142 (26.8)
Patients do not bring up fertility spontaneously	219 (41.1)	170 (31.9)	144 (27.0)
Presence of a third person	209 (39.5)	148 (28.0)	172 (32.5)
Patient is too ill to discuss fertility	147 (27.5)	199 (37.3)	188 (35.2)
I feel uncomfortable to discuss fertility	135 (25.4)	170 (32.0)	227 (42.7)
No connection with the patient	124 (23.3)	151 (28.4)	257 (48.3)
Afraid to offend the patient	107 (20.1)	138 (25.9)	288 (54.0)
No referral options	106 (19.9)	143 (27.0)	285 (53.4)
Patient is not ready to discuss fertility	104 (19.5)	244 (45.9)	184 (34.6)
Age difference between yourself and the patient	95 (17.8)	119 (22.3)	319 (59.8)
Someone else is accountable for discussing fertility	84 (15.8)	186 (34.9)	263 (49.3)
Changed fertility is not a problem for the patient	57 (10.7)	216 (40.6)	259 (48.7)
Insufficient time	50 (9.4)	117 (22.0)	366 (68.7)

n differs because the questions were not answered consistently, and some were skipped or forgotten

^a Agree contains the answers “totally agree” and “agree”

^b Disagree contains the answers “totally disagree” and “disagree”

the nurses were asked to rate their own level of knowledge regarding fertility in CKD patients. Forty-seven nurses (8.7%) stated to have no knowledge at all, 254 (47.0%) answered “a little” and 194 (35.9%) had some knowledge. Forty-five nurses (8.3%) rated their own knowledge as sufficient and one nurse (0.2%) answered in having a lot of knowledge. Nearly 30% of the nurses ($n = 156$, 29.2%) felt competent to discuss fertility with their patients, three-quarters ($n = 407$, 75.7%) stated to be in need of expanding their knowledge. Nurses with less knowledge, who did not feel competent to discuss fertility or who were in need of expanding their knowledge, informed their patients less often on fertility. Furthermore, these nurses discussed fertility issues less often with their patients and noted “insufficient training” as a barrier to discuss fertility more often than other nurses. The p values ranged from <0.001 up to 0.01.

Discussion

This analysis was the first to evaluate the extent of fertility care and information provided by nephrologists and nurses working in the dialysis unit for CKD patients. The study identified the absence of clear guidelines on this important part of renal health care in the majority of the Dutch nephrology and dialysis departments. Despite the lack of guidelines that appoint a health care professional to provide fertility care, the study shows that most nephrologists routinely discuss fertility during consultation. Unfortunately, this trend is not present throughout the whole nephrology department; nurses working in the dialysis unit who were unaware of guidelines or who had less work experience often skipped discussing fertility. However, the nurses are willing to provide this part of renal care in cooperation with the nephrologist. A suggestion was made that the nephrologists could initiate the subject of fertility enabling the nurses to continue the conversation during their own consultation with their patients. Still, multiple reasons exist that restrain nurses from assessing this subject. Besides the age of the patient, major barriers that withhold nurses from discussing fertility with their CKD patients derive from diversity in language, ethnicity, culture, and religion. Dealing with these factors in a diverse patient population requires certain skills, knowledge, and attitudes defined as cultural competence [15]. Cultural competence is becoming ever more important due to increasing cultural diversity among patients and care providers [15, 16]. This is especially so when providing fertility care, as sensitive issues need to be explained and discussed.

This study revealed the importance of training and nurses' self-imposed knowledge and feelings of competence in providing fertility care. Unfortunately, study

outcomes also indicated that these components were not self-evident among nurses of the dialysis department. This lack of training, self-imposed knowledge, and competence might be explained when focusing on the current educational system. Even though almost 70% of the nurses received education on fertility in CKD during their nurse training, a pressing need for additional training in discussing fertility exists. These lacuna in competence, knowledge and training regarding fertility issues is a problem not confined to the nurses working in nephrology; similar challenges arise among nurses in other medical departments as well [17, 18].

Strengths and limitations

This study was a pioneer in assessing fertility care provided by renal care providers for CKD patients. In this light, a limitation of the study is the lack of formal comparison as no previous study of this nature has been conducted. Response bias may have occurred due to low responses in both surveys. However, this may also be interpreted as a lack of interest or knowledge regarding the subject. In addition, no validated questionnaires exist that assess the study aims. For this reason, two questionnaires were developed, which have not been formally validated. Validation of these two surveys was not performed as they will not be reused. Nevertheless, the authors attempted to develop reliable, literature-based and pilot-tested instruments. Finally, response bias may have occurred as socially desirable answers may have been given. As a result, answers might have been over or underestimated.

Recommendations for practice

In order to achieve a cooperative fertility care system in CKD for both nephrologists and nurses, study results suggest several adjustments need to be made which involve both care providers. First, clear guidelines should be formulated within all nephrology departments to ensure multidisciplinary fertility care is provided. These guidelines should highlight the roles in providing fertility care of both nephrologists and nurses of the dialysis department, so any uncertainties regarding for instance accountability will be diminished. At the request of nurses themselves, informative brochures on fertility should be made available within nephrology departments and dialysis units to hand out to patients. In order to establish a functional system, nurses' competence and knowledge regarding discussing fertility, especially in culturally diverse populations, should be improved. Renal care providers' competence and skills regarding discussing fertility

are of great importance as it is their ethical responsibility to explain the complexity of fertility in CKD to their patients [19]. Literature suggests that improvement of competence in discussing fertility and providing fertility care in culturally diverse populations could be achieved by providing adequate training; a recommendation confirmed by more than half of the nurses in this survey [15, 16, 20]. Discretion is called when providing fertility counseling as some patients could find being advised not to become pregnant by their physician as traumatizing [8, 21]. The lack of privacy is experienced by some of the nurses; this could call for the facilitation of a private and scheduled consultation to discuss a delicate subject such as fertility. In addition, CKD patients might benefit from an adequate referral system to physicians specialized in fertility [21, 22]. Wiles et al. [22] showed that in the UK referral to specialized pre-pregnancy counseling renal clinics resulted in highly satisfied female CKD patients. Awareness should be raised among renal care providers on the importance of referral to physicians specialized in fertility as currently only a small percentage of the CKD patients are referred. Finally, to be able to enhance current fertility care according to patients' wishes, more research should be performed to determine their needs and preferences regarding this part of renal healthcare. Especially among male CKD patients more research is needed, as little is known about the preferable format, type and timing of fertility care for men.

Conclusion

Guidelines that appoint who should provide fertility care to CKD patients are absent in the majority of Dutch nephrology departments. Current fertility care is not provided in a cooperative manner: most of the nephrologists assessed fertility routinely during consultation; less experienced nurses working in dialysis units often skipped this part of renal care. Besides a patients' age, a lack of

competence, insufficient knowledge and training, unawareness of guidelines, and barriers based on cultural diversity are reasons for nurses to omit providing fertility care. To achieve a cooperative fertility care system in CKD for both nephrologists and nurses, results emphasize the need for guidelines on this part of renal care in Dutch nephrology and dialysis departments and the provision of training to improve of nurses' competence regarding cultural diversity and fertility. Finally, more research should be performed to determine patients' needs and preferences regarding this important part of renal healthcare, especially among male patients.

Acknowledgements The authors thank all Dutch nephrologists and nurses who took the time to fill out the questionnaire. Linguistic supervision was performed by Emma Horton, MD. Furthermore, the authors would like to thank Stijn van der Struijs for his assistance with the data collection.

Funding This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Appendix 1: Questionnaire nephrologist

Reason not to participate:

- No interested
- No time
- Not enough experience
- No enhancement possible in this area.
- Retired
- Other:.....

For each question chose the most suitable answer. Please do not skip any questions. Thank you in advance for your effort.

Demographics

1. What is your gender?
 - Male
 - Female
2. What is your age? _____ years
3. What is your current position at work?
 - Nephrologist
 - Resident
 - Other: _____
4. Time of practice in nephrology (residency included)?
 - 0-11 months
 - 1-2 years
 - 3-5 years
 - 6-10 years
 - 11-15 years
 - > 15 years
5. Type of clinic/practice?
 - Tertiary referral hospital (or university hospital)
 - General teaching hospital
 - District general hospital
 - Tertiary and district general hospital
 - Dialysis centre, outside of a hospital
 - Other: _____

Fertility in chronic kidney disease patients

Definition according to the KDOQI*: patients suffering from chronic kidney disease and above the age of 16. This includes with or without renal replacement therapy.

Table 10. Stages of Chronic Kidney Disease

Stage	Description	GFR (mL/min/1.73 m ²)
1	Kidney damage with normal or ↑ GFR	>90
2	Kidney damage with mild ↓ GFR	60–89
3	Moderate ↓ GFR	30–59
4	Severe ↓ GFR	15–29
5	Kidney failure	<15 (or dialysis)

Chronic kidney disease is defined as either kidney damage or GFR <60 mL/min/1.73 m² for ≥3 months. Kidney damage is defined as pathologic abnormalities or markers of damage, including abnormalities in blood or urine tests or imaging studies.

1. How often do you inform your patients of reproductive age that a decline in renal function is associated with reduced fertility?
 - Never/almost never
 - In less than half of the cases
 - In half of the cases
 - In more than half of the cases
 - Almost always/Always

2. Do you advise your female patients with renal failure not to become pregnant?
- Yes
 - No
3. If yes, in which stage of disease?
- Stage 1
 - Stage 2
 - Stage 3
 - Stage 4
 - Stage 5
4. Are you aware of clear guidelines within your department regarding which care provider is accountable for discussing patients' fertility?
- Yes, that is _____
 - No
 - Unaware
5. Does your hospital offer specialized pre-conceptive counselling for chronic kidney disease patients?
- Yes
 - No
 - Unaware
6. What percentage of your female patients of reproductive age have you referred to a care provider specialized in fertility?
_____%
7. What percentage of your male patients of reproductive age have you referred to a care provider specialized in fertility?
_____%
8. When you discuss fertility with a female patient, which subject(s) do you discuss? (multiple answers possible)
- Wish to have children
 - Menopausal complaints
 - Risk of early termination of pregnancy
 - Inheritance of the disease
 - Risk of pregnancy for patients' health
 - Option to wait with a pregnancy until after kidney transplantation
 - Possibility of oocyte cryopreservation
 - Fear of malformations in their child
 - Other, _____
9. When you discuss fertility with a male patient, which subject(s) do you discuss? (multiple answers possible)
- Wish to have children
 - Presence of erectile dysfunction
 - Ability of ejaculation
 - Cryopreservation of sperm
 - Inheritance of the disease
 - Fear of malformations in their child
 - Other, _____
10. If a female transplant recipient wishes to get pregnant, which subject(s) do you discuss? (multiple answers possible)
- Chance of a successful pregnancy
 - Pregnancy in the first years after RTx is not possible
 - Possible decline in function of donor kidney due to pregnancy
 - Risk of early termination of pregnancy
 - Inheritance of the disease
 - Other, _____

Thank you very much for completing the survey!

Appendix 2: Questionnaire dialysis nurses and nurses specialized in nephrology

Reason not to participate:

- No interested
- No time
- Not enough experience
- No enhancement possible in this area.
- Retired
- Other:.....

Choose at each question the most suitable answer. Please do not skip any questions. Thank you in advance for your effort.

Demographics

1. What is your gender?
 - Male
 - Female
2. What is your age? _____ years
3. What is your current position at work?
 - Dialysis Nurse
 - Nurse in Dialysis registration training
 - Team leader dialysis department
 - Nurse specialised in Nephrology
 - Other: _____
4. Years of employment?

<input type="checkbox"/> 0-11 months	<input type="checkbox"/> 6-10 years
<input type="checkbox"/> 1-2 years	<input type="checkbox"/> 11-15 years
<input type="checkbox"/> 3-5 years	<input type="checkbox"/> > 15 years
5. Type of clinic/practice?
 - University Hospital
 - District general teaching hospital
 - District general hospital
 - Tertiary and district general hospital
 - Dialysis centre, outside of a hospital
 - Other: _____

Fertility in chronic kidney disease patients

Definition: patients suffering from chronic kidney disease who receive any form of renal replacement therapy and are above the age of 16.

1. How often do you inform your patients of reproductive age that dialysis is associated with reduced fertility?
 - Never/ Almost never
 - In less than half of the cases
 - In half of the cases
 - In more than half of the cases
 - Almost always/ Always
2. How do you rate your own knowledge on fertility?
 - No knowledge at all
 - Not a lot
 - Some knowledge
 - Sufficient knowledge
 - A lot of knowledge

3. Is the subject fertility in CKD patients addressed during your training to become a nurse?

- Yes
- No

4. Are you in need of extending your knowledge on the discussing of sexual dysfunction?

- Yes
- No

5. Do you feel competent to discuss fertility with your patients?

- Yes
- No

6. Are you aware of clear guidelines within your department regarding which care provider is accountable for discussing patients' fertility?

- Yes, that is _____
- No
- Unaware

7. How often is the partner of the patient present while discussing fertility?

- Never
- In less than half of the cases
- In half of the cases
- In more than half of the cases
- Always

8. What could be helpful for the discussion of fertility? (multiple answers possible)

- Informative brochures
- Training to discuss fertility
- Posters about fertility in the waiting room
- The nephrologist should initiate the conversation on fertility, so that I could refer to this conversation
- Other, _____

9. How often do you discuss fertility with:

	Never	Seldom	Routinely	Often
- Male patients	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
- Female patients	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Possible barriers towards discussing fertility are listed below. To which extent are the barriers applicable to you? Please give only one answer for each barrier.

	Totally disagree	Disagree	Slightly disagree/ Slightly agree	Agree	Totally agree
Insufficient time	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Insufficient knowledge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Insufficient training	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Someone else is accountable for discussing fertility	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Patient is not ready to discuss fertility	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Changed fertility is not a problem for the patient	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Patient is too ill to discuss fertility	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Barriers based on culture or religion	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Barriers based on language or ethnicity	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Age of the patient	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Age difference between yourself and the patient	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Presence of a third person	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Afraid to offend the patient	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
No connection with the patient	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Could not find a suitable moment	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I feel uncomfortable to discuss fertility	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Patients do not bring up fertility spontaneously	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
No referral options	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Thank you very much for completing the survey!

References

- Hladunewich M, Schatell D (2016) Intensive dialysis and pregnancy. *Hemodial Int* 20(3):339–348. doi:10.1111/hdi.12420
- Shiraishi K, Shimabukuro T, Naito K (2008) Effects of hemodialysis on testicular volume and oxidative stress in humans. *J Urol* 180(2):644–650
- Lessan-Pezeshki M, Ghazizadeh S (2008) Sexual and reproductive function in end-stage renal disease and effect of kidney transplantation. *Asian J Androl* 10(3):441–446. doi:10.1111/j.1745-7262.2008.00348.x
- Hill NR, Fatoba ST, Oke JL, Hirst JA, O'Callaghan CA, Lasserson DS, Hobbs FD (2016) Global prevalence of chronic kidney disease—a systematic review and meta-analysis. *PLoS ONE* 11(7):e0158765. doi:10.1371/journal.pone.0158765
- Douglas NC, Shah M, Sauer MV (2007) Fertility and reproductive disorders in female solid organ transplant recipients. *Semin Perinatol* 31(6):332–338. doi:10.1053/j.semperi.2007.09.002
- Todros GBP, Anne C, Valentina C, Elena V, Rossella A, Maria Chiara D, Salvatore B, Tullia (2010) Pregnancy in dialysis patients: is the evidence strong enough to lead us to change our counseling policy? *Clin J Am Soc Nephrol*. doi:10.2215/CJN.05660809
- Yang LY, Thia EW, Tan LK (2010) Obstetric outcomes in women with end-stage renal disease on chronic dialysis: a review. *Obstet Med* 3(2):48–53. doi:10.1258/om.2010.10000110.1258/om.2010.100001 (Epub 2010 Jun 3)
- Tong A, Jesudason S, Craig JC, Winkelmayr WC (2015) Perspectives on pregnancy in women with chronic kidney disease: systematic review of qualitative studies. *Nephrol Dial Transplant* 30(4):652–661. doi:10.1093/ndt/gfu378
- National Kidney Foundation (2002) K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 39 (2 Suppl 1):S1–266
- Krouwel EM, Nicolai MP, vS-vT AQ, Putter H, Osanto S, Pelger RC, Elzevier HW (2015) Addressing changed sexual functioning in cancer patients: a cross-sectional survey among Dutch oncology nurses. *Eur J Oncol Nurs* 19(6):707–715
- Nicolai MP, Liem SS, Both S, Pelger RC, Putter H, Schali MJ, Elzevier HW (2013) What do cardiologists know about the effects of cardiovascular agents on sexual function? A survey among Dutch cardiologists. Part I. *Neth Heart J* 21(12):540–544. doi:10.1007/s12471-013-0471-2
- van Ek GF, Krouwel EM, Nicolai MP, Bouwsma H, Ringers J, Putter H, Pelger RC, Elzevier HW (2015) Discussing Sexual Dysfunction with Chronic Kidney Disease Patients: practice Patterns in the Office of the Nephrologist. *J Sex Med* 12(12):2350–2363. doi:10.1111/jsm.1306210.1111/jsm.13062 (Epub 2015 Dec 3)

13. Van Ek GF KE, Van der Veen E, Nicolai MP, Van den Ouden BL, Ringers J, Putter H, Pelger RC, Elzevier HW (2017) The discussion of sexual dysfunction before and after kidney transplantation from the perspective of the renal transplant surgeon. *Progress in Transplantation* Accepted for publication
14. van Ek GF, Krouwel EM, Nicolai MP, Bouwsma H, Ringers J, Putter H, Pelger RC, Elzevier HW (2015) Discussing sexual dysfunction with chronic kidney disease patients: practice patterns in the office of the nephrologist. *J Sex Med* 12(12):2350–2363
15. Calvillo E, Clark L, Ballantyne JE, Pacquiao D, Purnell LD, Villarruel AM (2009) Cultural competency in baccalaureate nursing education. *J Transcult Nurs* 20(2):137–145. doi:[10.1177/1043659608330354](https://doi.org/10.1177/1043659608330354)
16. Kai J, Beavan J, Faull C, Dodson L, Gill P, Beighton A (2007) Professional uncertainty and disempowerment responding to ethnic diversity in health care: a qualitative study. *PLoS Med* 4(11):e323. doi:[10.1371/journal.pmed.0040323](https://doi.org/10.1371/journal.pmed.0040323)
17. Krouwel EM, Nicolai MP, Jeanne van Steijn-van Tol AQM, Putter H, Osanto S, Pelger RCM, Elzevier HW (2016) Fertility preservation counselling in Dutch oncology practice: Are Nurses ready to assist physicians?. *Eur J Cancer Care* (in press)
18. Ussher JM, Cummings J, Dryden A, Perz J (2017) Talking about fertility in the context of cancer: health care professional perspectives. *Eur J Cancer Care* 25(1):99–111. doi:[10.1111/ecc.12379](https://doi.org/10.1111/ecc.12379)
19. Lopez LF, Martinez CJ, Castaneda DA, Hernandez AC, Perez HC, Lozano E (2014) Pregnancy and kidney transplantation, triple hazard? Current concepts and algorithm for approach of preconception and perinatal care of the patient with kidney transplantation. *Transpl Proc* 46(9):3027–3031. doi:[10.1016/j.transproceed.2014.07.013](https://doi.org/10.1016/j.transproceed.2014.07.013)
20. Fleckman JM, Dal Corso M, Ramirez S, Begaliev M, Johnson CC (2015) Intercultural competency in public health: a call for action to incorporate training into public health education. *Front Public Health* 3:210. doi:[10.3389/fpubh.2015.00210](https://doi.org/10.3389/fpubh.2015.00210)
21. Tong A, Brown MA, Winkelmayer WC, Craig JC, Jesudason S (2015) Perspectives on pregnancy in women with CKD: a semi-structured interview study. *Am J Kidney Dis* 66(6):951–961
22. Wiles KS, Bramham K, Vais A, Harding KR, Chowdhury P, Taylor CJ, Nelson-Piercy C (2015) Pre-pregnancy counselling for women with chronic kidney disease: a retrospective analysis of nine years' experience. *BMC Nephrol* 16:28. doi:[10.1186/s12882-015-0024-6](https://doi.org/10.1186/s12882-015-0024-6)