


Progress toward cariporide analogs for sodium-proton exchange inhibition

ORGN 653

Jessica Moore¹, Mark Graves II¹, Jacob A Vervynckt, jacob.vervynckt@wku.edu¹, William Harley², Fredric Gorin², and Hasan Palandoken, hasan.palandoken@wku.edu¹. (1) Department of Chemistry, Western Kentucky University, 1906 College Heights Boulevard, Bowling Green, KY 42101, (2) Center for Neuroscience, University of California, Davis, 1515 Newton Court, Davis, CA 95616

The sodium proton exchanger (NHE) is particularly important in maintaining the intracellular pH in human heart and brain. Under anaerobic conditions (i.e., ischemia), a shift from oxidative to nonoxidative glycolysis occurs. The resultant decrease in the intracellular pH activates NHE, which increases the intracellular sodium, initiating the sequence of physiological events that lead to cell death. Thus, there has been great interest in the development of compounds that inhibit NHE. Indeed, potent NHE inhibitors are available. However, a fundamental impediment to the field is the delivery of these compounds to poorly vascularized tissues during the early phases of ischemia when NHE inhibition is most beneficial. We have synthesized analogs of cariporide, a potent (e.g., nanomolar IC₅₀ activity) NHE inhibitor, to address these temporal and delivery challenges. The preparation and biological activities of our cariporide analogs will be discussed.



[Materials, Devices and Switches, Metal-Mediated Reactions, Asymmetric Reactions, Total Synthesis, Biologically-Related Molecules and Processes](#)

7:00 PM-9:00 PM, Wednesday, April 9, 2008 Morial Convention Center -- La Louisiane, Blrm. C, Poster

[Sci-Mix](#)

8:00 PM-10:00 PM, Monday, April 7, 2008 Morial Convention Center -- Hall A, Sci-Mix

[Division of Organic Chemistry](#)

[The 235th ACS National Meeting, New Orleans, LA, April 6-10, 2008](#)