

Hypoxia induces connexin 43 dysregulation by modulating matrix metalloproteinases via MAPK signaling

Xianghong Wu · Wen Huang · Gang Luo · Laval Andy Alain

Received: 18 March 2013 / Accepted: 23 August 2013 / Published online: 4 September 2013
© The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract Connexin 43 (Cx43) is a major structural protein found in the gap junctions of the ventricular myocardium and a major determinant of its electrical properties. The effects of matrix metalloproteinases (MMPs), the mitogen-activated protein kinase (MAPK) signaling pathway, transcription factor NF- κ B, and activator protein-1 (AP-1)/c-Jun on the regulation of Cx43 gene expression in H9c2 cardiomyocytes were assessed. The MAPK signaling pathway (MEK/ERK1/2 and PI3K) and transcription factors NF- κ B and AP-1/c-Jun were inhibited, then Cx43 expression was assessed using Western blot analysis, and MMP-9 activity was assessed using gelatin zymography. Hypoxia decreased the Cx43 protein level by approximately 30–50 %. Doxycycline (10 μ g/mL), an inhibitor of MMP, markedly attenuated the hypoxia-induced downregulation of Cx43 protein expression at 6 h. The hypoxia-induced decrease in Cx43 protein expression was significantly reversed by U0126 (10 μ M), a MEK/ERK1/2 inhibitor, at 6 and 12 h; LY294002 (30 μ M), a PI3K inhibitor, downregulated Cx43 expression. Hypoxia-induced MMP-9 activation was inhibited by treatment with LY294002, U0126, and, most especially, U0126. JSH-23 (30 μ M), an NF- κ B inhibitor, and SP600125 (10 μ M), an AP-1/c-Jun inhibitor, attenuated the loss of Cx43. These results suggest that MAPK signaling and the activities NF- κ B and MMPs play an important roles in the regulation of Cx43 expression.

Keywords Connexin 43 · Matrix metalloproteinases · Mitogen-activated protein kinases · Hypoxia

Abbreviations

Cx43	Connexin 43
MMPs	Matrix metalloproteinases
MAPK	Mitogen-activated protein kinases
AP-1	Activator protein-1
ECM	Extracellular matrix
PI3K	Phosphoinositide-3 kinase
NF- κ B	Kappa-light-chain enhancer of activated B cells
LPS	Lipopolysaccharide

Introduction

Gap junctions, which are composed of channel-forming integral membrane proteins known as connexins, mediate cell–cell communication in almost all tissues [1]. Connexin 43 (Cx43) is a major structural protein found in the gap junctions of the ventricular myocardium and a major determinant of myocardial electrical properties [2]. Cx43 dysfunction in cardiomyocytes may contribute to the pathogenesis of ventricular arrhythmias. Transferring the Cx43 gene to pigs with anterior infarction reduces ventricular tachycardia in the border zone of the healed scar [3].

Matrix metalloproteinases (MMPs) reportedly play an important role in the degradation of the extracellular matrix (ECM) [4]. The degradation of the ECM by MMPs is involved in the pathogenesis of cardiovascular diseases, including atherosclerosis and myocardial infarction [5, 6]. MMP expression is regulated at the transcription level by the activation of various transcription factors, such as

X. Wu · G. Luo · L. A. Alain
Department of Cardiology, First Affiliated Hospital, Guangxi Medical University, Nanning, People's Republic of China

W. Huang (✉)
Department of Neurology, First Affiliated Hospital, Guangxi Medical University, Nanning 530021, People's Republic of China
e-mail: hwen1229@163.com

activator protein (AP)-1 and nuclear factor kappa-light-chain enhancer of activated B cells (NF- κ B) [7–9]. MMP-9 is expressed within minutes of myocardial ischemia [10]. MMPs are also capable of mediating Cx43 cleavage. Myocardial ischemia-induced reduction in Cx43 expression is also attenuated in MMP-7^{-/-} mice [11]. Our previously reported study found that lipopolysaccharide (LPS)-induced MMP-2 activation is affected by NF- κ B-dependent Ras-MEK1/2 signaling [12]. Serine/threonine protein kinase B (PKB/Akt)—a major downstream effector of activated phosphoinositide-3 kinase (PI3K) and the ERK1/2 pathway—influences the transcription of MMP-9 [13]. HIV-1 induces the dysregulation of tight junction proteins by modulating MMPs via the ERK and Akt signaling [14, 15]. But, are MMPs and mitogen-activated protein kinase (MAPK) signaling involved in the regulation of gap junctions in cardiomyocytes, as they are in tight junctions? This present study was performed to determine if MMPs and MAPK signaling are involved in the hypoxia-induced regulation of Cx43. This study was performed to help to better understand the mechanisms involved in the development of ventricular arrhythmia following myocardial ischemia, as well as provide data that could be used to pharmacologically prevent ventricular arrhythmia in patients with a history of myocardial infarction.

Methods

Cell culture

Rat H9C2 cardiomyocytes were obtained from the Chinese Academy of Sciences (Beijing, China) and cultured in Dulbecco's modified Eagle's medium (DMEM; Sigma, St. Louis, MO, USA) supplemented with 10 % fetal bovine serum (FBS; Hyclone) and penicillin plus streptomycin (100 U/mL and 100 μ g/mL, respectively). All cultures were maintained at 37 °C in a 5 % CO₂ atmosphere.

Hypoxic conditions and chemical treatment

Cells were incubated under hypoxic conditions to mimic *in vivo* myocardial ischemia. Hypoxic conditions were achieved in an incubator containing a gaseous mixture of 94 % N₂, 5 % CO₂, and 1 % O₂ (v/v).

Cells were incubated overnight in serum-free essential medium before treatment with the indicated agents. In brief, H9C2 cardiomyocytes were cultured in 6-well plates until 90 % confluence, then treated under hypoxic conditions in the presence or absence of the following inhibitors: MMP inhibitor doxycycline (10 μ g/mL; Sigma); PI3K inhibitor LY294002 (30 μ M; Sigma); and MEK/ERK1/2 inhibitor U0126 (10 μ M; Sigma). In specific experiments,

H9C2 cells were treated with the NF- κ B inhibitor JSH 23 (30 μ M; Santa Cruz) or AP-1/c-Jun inhibitor SP600125 (10 μ M; Sigma) for 6 or 12 h. Control cells were cultured in complete medium under normoxic conditions. After treatment, cells were harvested and assessed using Western blot analysis, and the culture medium was assayed for MMP-9 activity.

Cell viability assay

Relative cell viability was determined using 125 mg/mL MTT (3-[4,5-dimethylthiazolyl-2]-2,5-diphenyltetrazolium bromide in phosphate-buffered saline) [16], which was added after treatment. Cells were incubated in MTT for 3 h at 37 °C and solubilized in dimethyl formamide (50 %; v/v) and sodium dodecyl sulfate (SDS) (20 %; w/v), and then absorbance was measured at 570 nm.

Western blot analysis

Proteins from the H9C2 cells that were cultured in the 6-well plates were extracted using RIPA lysis buffer (Santa Cruz Biotechnology) and centrifuged at 15,000 g for 15 min. The supernatants were collected, and protein concentrations were determined using the BCM Protein Assay Kit (Thermo Scientific Pierce). An aliquot of 30 μ g proteins from each sample was separated on 10 % Tris-HCl SDS-polyacrylamide gels, transferred onto nitrocellulose membranes, incubated with 3 % skim milk in Tris-buffered saline solution for 1 h, incubated overnight with the respective antibodies at 4 °C, and finally incubated with the peroxidase-conjugated secondary antibody at room temperature for 120 min. To visualize the proteins, the immunoblots were analyzed using the ECL Western blotting detection kit (Thermo Scientific Pierce). Rabbit anti-rat Cx43 antibodies were purchased from Abcam. Anti-ERK1/2, phosphorylated ERK1/2 (p-ERK1/2), PI3K, and phosphorylated PI3K (p-PI3K) antibodies were purchased from cell signaling. HRP-GAPDH was purchased from Kangcheng Biotechnology. All secondary antibodies were purchased from Santa Cruz Biotechnology.

MMP-9 activity assay

MMP-9 activity was assessed by gelatin zymography [12] using premade 10 % polyacrylamide gels containing 0.1 % gelatin and 10 μ L serum-free media from the treated cultures; these procedures were performed according to the instructions provided by the manufacturer (Invitrogen). Briefly, cells in serum-free medium were pretreated with 30 μ M LY294002 or 10 μ M U0126 for 2 h, followed by coexposure to hypoxia and specific inhibitors for 6 or 12 h. After electrophoresis, the gel was removed and incubated

with Renaturing Buffer for 30 min at room temperature with gentle agitation, then equilibrated overnight with Developpin Buffer. Bands were visualized by staining for 30–60 min with 0.1 % Coomassie R-250 in 40 % ethanol and 10 % acetic acid, followed by destaining for 2 h at room temperature with a solution containing 10 % ethanol and 7.5 % acetic acid.

MMP-9 promoter activity assays

H9C2 cells in 24-well plates were transfected with 0.5 μ g pGL3 MMP-9 using lipofectin (Invitrogen) as the transfection reagent. To normalize to transfection efficiency, cells were cotransfected with 0.05 μ g pRL-TK construct (Promega), which encodes for *Renilla* luciferase. Firefly and *Renilla* luciferase activities were determined using the Dual-Luciferase Reporter Assay System (Promega), as previously described [13].

Statistical analysis

The intensities of the bands corresponding to specific proteins were determined using image J software. Routine statistical analyses were completed using SPSS 15.0. Values are presented as the mean \pm SEM. One-way ANOVA was used to evaluate between-group differences, followed by the Tukey test. In this study, $p < 0.05$ is considered significant.

Results

MMPs and MAPK signaling are involved in Cx43 regulation

Rat H9C2 cardiomyocytes are shown in Fig. 1a. None of the chemical treatments affected cell viability at the concentrations that were used in the present study (Fig. 1b).

MMPs are involved in the cleavage of Cx43. Administering MMP-2 to normal rats decreases Cx43 expression by 40 % [17]. To investigate the effects of MMPs on the expression of Cx43, H9C2 cells were treated with doxycycline, an inhibitor of MMPs. As illustrated in Fig. 2, hypoxia decreased the total Cx43 protein level by approximately 30–50 %. Doxycycline markedly attenuated the hypoxia-induced downregulation of total Cx43 protein expression at both 6 and 12 h, but most especially at 6 h. Our data indicate that MMPs are involved in the regulation of Cx43 protein expression, which is consistent with previous studies [17].

Our previously reported data indicate that the Akt and ERK1/2 pathways influence MMP-9 expression [14, 15]. To determine if MAPK signaling is involved in the

regulation of Cx43 expression, the inhibitors LY294002 and U0126 were used to block PI3K and the MEK/ERK signaling pathway, respectively. The hypoxia-induced reduction in Cx43 protein expression was significantly reversed by U0126 at 6 and 12 h (Fig. 2a–c). Interestingly, the PI3K inhibitor LY294002 did not protect against hypoxia-induced alterations in Cx43. On the contrary, LY294002 acts synergistically with hypoxia to down-regulate Cx43 protein expression.

The polyclonal Cx43 antibody may recognize several forms of phosphorylated Cx43 (p-Cx43). As shown in Fig. 2, lower bands correspond to low-molecular-weight proteins that are weakly phosphorylated, and upper bands correspond to high-molecular-weight proteins that are strongly phosphorylated [18]. Notably, MAPK signaling affected Cx43 phosphorylation. The ratio of p-Cx43 to total Cx43 proteins (T-Cx43) was significantly higher following exposure to U0126 and hypoxia in comparison with cells that were treated with hypoxia alone; however, a weaker phosphorylated form of Cx43 was induced with LY294002 under hypoxia (Fig. 2d–e). Meanwhile, p-ERK1/2 and p-PI3K levels were markedly upregulated in response to hypoxia and downregulated in response to U0126 and LY294002 (Fig. 2f–g).

ERK1/2 signaling regulates Cx43 expression by modulating MMP-9 enzyme activity

To determine if MMP-9 enzyme activity is involved in MAPK signaling and the regulation of Cx43, LY294002 (PI3K inhibitor) and U0126 (MEK/ERK inhibitor) were applied, respectively, and MMP-9 activity was measured using zymography. The zymography media were obtained from the same cultures used in the Western blot analysis. As shown in Fig. 3, hypoxia-induced MMP-9 activation was inhibited by treatment with LY294002 and U0126 at 6 and 12 h, especially U0126.

NF- κ B and AP-1/c-Jun are involved in the regulation of Cx43

JNK/SAPK inhibitors significantly reversed the decrease in Cx32 and Cx43 protein expression [19]. To further evaluate the roles of the transcription factors NF- κ B and AP-1/c-Jun in the regulation of Cx43, H9C2 cells were cultured in the presence of specific inhibitors of NF- κ B or AP-1/c-Jun and total Cx43 protein expression was assessed by Western blot analysis. As indicated in Fig. 4a, Cx43 expression decreased following hypoxia at 6 and 12 h. However, this effect was notably attenuated by both NF- κ B and AP-1/c-Jun inhibitors, JSH-23 and SP600125, respectively (Fig. 4b–c). This indicates that NF- κ B and AP-1/c-Jun contribute to the regulation of Cx43 expression under

Fig. 1 a Photomicrographs of the H9C2 cardiomyocytes. **b** Inhibitors on cell viability determined with MTT assays. H9C2 cells were treated with inhibitors for 12 h at different concentrations (doxycycline: 1, 5, 10, 20, and 30 $\mu\text{g/ml}$; LY294002: 5, 15, 30, 60, and 120 μM ; U0126: 5, 10, 15, 30, and 60 μM ; JSH-23: 15, 30, 60, 120, and 240 μM ; SP600125: 1, 5, 10, 20, and 30 μM)





Fig. 2 a Showing Western Blot band of Cx43 with treatment of hypoxia in the absence or presence of MMPs inhibitor doxycycline (10 $\mu\text{g/ml}$) or PI3K inhibitor LY294002 (30 μM) or MEK/ERK 1/2 inhibitor U0126 (10 μM) at 6 h and 12 h. **b–g** The quantified results are depicted in the form of bar graphs. Cx43 protein expression with treatment for 6 h (**b**) and 12 h (**c**). Hypoxia decreased the total Cx43 protein level by approximately 30–50 %. Doxycycline or U0126 markedly attenuated hypoxia-induced downregulation of Cx43 protein expression. LY294002 acts synergistically with hypoxia to downregulate Cx43 protein expression. The ratio of P-Cx43 to T-Cx43 is shown in (**d–e**). p-ERK1/2 and p-PI3K levels in response to U0126 and LY294002 were shown in **f–g**. Results represent the mean \pm SEM of 3 separate experiments. (*Asterisk*) Significantly different as compared to the control. (*Hash*) Data in the groups co-treated with hypoxia and specific inhibitor are significantly different from those treated with hypoxia alone






Fig. 3 **a** Showing Zymography band of Cx43 with treatment of hypoxia in the absence or presence of LY294002 (30 μ M) or U0126 (10 μ M) at 6 h and 12 h. **b–c** The quantified results are depicted in the form of bar graphs. Cx43 protein expression with treatment for 6 h (**b**) and 12 h (**c**). Results represent the mean \pm SEM of 3 separate

experiments. Significantly different as compared to the control. (*Hash*) Data in the groups co-treated with hypoxia and specific inhibitor are significantly different from those treated with hypoxia alone

hypoxic conditions. Both JSH-23 and SP600125 demonstrated the most obvious effects on Cx43 expression at 6 h. Interestingly, both JSH-23 and SP600125 shifted Cx43 expression to its phosphorylated forms under hypoxic conditions (Fig. 4d–e).

Our previous study indicated that the transcription factors NF- κ B and AP-1 are involved in the regulation of MMP-9 promoter activity [13]. To determine the roles that the NF- κ B and AP-1 pathways play in modulating MMP-9 activity under hypoxic conditions, MMP-9 promoter activity was assessed following treatment with JSH-23 and SP600125. As shown in Fig. 4f, both JSH-23 and SP600125 protected against hypoxia-induced MMP-9 promoter upregulation.

Discussion


Primary cardiac myocytes are widely used to investigate cellular and molecular changes in cardiac studies; however, the major disadvantage of primary cardiac myocytes is the high number of neonatal animals that are required. The H9C2 rat cardiomyocyte cell line is a reliable alternative for some studies. H9C2 rat cardiomyocytes demonstrate many similarities to primary cardiomyocytes, including membrane morphology, G-protein signaling expression, and electrophysiological properties [20, 21], and were used in all of our experiments.

It is widely accepted that oxidative stress plays a central role in the induction of pathogenic cell responses. The activation of the redox-regulated ERK1/2 and Akt signaling cascades has been observed in atherosclerosis and myocardial ischemia [22]. NF- κ B acts broadly on genes

that affect cell survival, differentiation, and proliferation. NF- κ B is also associated with cancer, autoimmune diseases, and cardiovascular disease [23]. The contribution of NF- κ B to cardiovascular disease, like atherosclerosis, is most easily understood in the context of chronic inflammation, wherein proinflammatory cytokines drive the activation of NF- κ B that in turn drives target gene expression [24]. Previous studies report that the NF- κ B inhibitor BAY11-7082 and the PI3K inhibitor LY294002 both significantly reverse the effects of Toll-like receptors via the ligand-mediated downregulation of Cx43 expression [25]. The nuclear transcription factor AP-1 can combine with many genes and, subsequently, regulate the transcription of these target genes. AP-1 also facilitates the transcription and expression of Cx43 in myometrial cells. Blocking the AP-1 sites that bind to the Cx43 promoter can neutralize the corticotropin-releasing hormone-induced upregulation of Cx43 [26]. Cx43 mRNA expression increases in cardiac myocytes following the addition of amphetamine, and these effects are completely attenuated by the AP-1/c-Jun inhibitor SP600125 [27]. Angiotensin II induces Cx43 expression in vascular smooth muscle cells via the activation of transcription factor AP-1 [28].

To assess the roles of MMPs and MAPK signaling in the regulation of Cx43 gene expression in H9c2 cardiomyocytes, the MMP inhibitor doxycycline, MEK/ERK 1/2 inhibitor U0126, and PI3K inhibitor LY294002 were used in the present study. In agreement with previous data, the inhibition of MMP activity attenuated the hypoxia-induced downregulation of Cx43 protein expression in H9c2 cardiomyocytes (Fig. 2a–d). One novel finding of the present study is that ERK1/2 and PI3K/Akt signaling act differently on Cx43 expression. Inhibiting ERK1/2

Fig. 4 a Showing Western Blot band of Cx43 with treatment of hypoxia in the absence or presence of JSH-23 (30 μ M) or SP600125 (10 μ M) at 6 h and 12 h. **b–f** The quantified results are depicted in the form of bar graphs. Cx43 protein expression with treatment for 6 h (b) and 12 h (c). Hypoxia induced a decrease in Cx43 expression, however, this effect was notably attenuated by both NF- κ B and AP-1/c-Jun inhibitors, JSH-23 and SP600125, respectively. The ratio of P-Cx43 to T-Cx43 is shown in (d–e). MMP-9 promoter activity assessed with Dual-Luciferase Reporter Assay System was shown in f. Results represent the mean \pm SEM of three separate experiments. (*Asterisk*) Significantly different as compared to the control. (*Hash*) data in the groups co-treated with hypoxia and specific inhibitor are significantly different from those treated with hypoxia alone



signaling attenuated the hypoxia-induced downregulation of Cx43 protein expression, however, blocking PI3K/Akt enhanced the loss of Cx43 in cardiomyocytes (Fig. 2a–c). MAPK signaling probably acts on Cx43 expression by modulating MMP-9 enzymes, as indicated in Fig. 3. Our previously reported study indicated that transcription factors NF- κ B and AP-1 are involved in the regulation of MMP-9 promoter activity by mutating the binding sites of NF- κ B and AP-1 [13]. To further study the functions of transcription factors under hypoxic conditions, JSH-23 and SP600125 were used to block the NF- κ B and AP-1/c-Jun pathways. As shown in Fig. 4a–c, the administration of JSH-23 and SP600125 attenuated the hypoxia-induced loss of Cx43 while also protecting against MMP-9 promoter

upregulation (Fig. 4f). This is a good indication that MMP activity, MEK/ERK1/2 and AP-1/c-Jun signaling, and NF- κ B play an important roles in regulation of Cx43 expression in H9c2 cardiomyocytes, as summarized in Fig. 5.

The Cx43 expression level depends on the time of oxygen deprivation. Cultured atrial cells exposed to hypoxia for 24 and 48 h demonstrate 75 and 90 % reductions in the Cx43 protein level, respectively [29]. In the present study, exposure to hypoxia for 6 and 12 h caused a decrease in the Cx43 level by approximately 30–50 % (Figs. 2b–c and 4b–c).

Studies indicate that phosphatases may regulate connexin during cardiac ischemia [30]. Connexin phosphorylation is associated with assembly, stability, and other channel




Fig. 5 Schematic diagram of the MAPK signaling pathways involved in regulation of Cx43 expression by modulating MMP-9 enzyme. The regulation of Cx43 expression was studied by blocking the pathways of PI3K/Akt or MEK/ERK 1/2 NF-κB or AP-1/c-Jun and MMPs,

respectively, with specific inhibitors. The signaling cascades, transcription factors NF-κB and AP-1/c-Jun, and MMPs activity appear to be critical factors in hypoxia-induced disruption of gap junction integrity

properties in gap junctions [31]. p-Cx43 is the major constituent of gap junctions; however, dephosphorylation of Cx43 occurs during hypoxia and ischemia and is associated with p-ERK1/2 levels [32]. Our results are partly consistent with studies in the literature indicating that MAPK signaling affects p-Cx43. Coexposure to hypoxia and U0126, JSH-23, or SP600125 shifts Cx43 expression to p-Cx43; however, the dephosphorylation of Cx43 is induced by the addition LY294002 (Figs. 2d–e, 4d–e).

Nevertheless, it should be noted that these data were collected by a primary study investigating the connection between MAPK signaling, MMP activity, and gap junctions. Further studies should be performed to investigate the exact mechanisms that regulate Cx43, especially the phosphorylated and dephosphorylated forms of Cx43. This will lead to better understanding of the working mechanisms involved in MAPK signaling, MMP, and Cx43 expression.

In summary, exposure to hypoxia decreases Cx43 protein expression in H9c2 cardiomyocytes, inhibits MMP activity, and blocks ERK1/2 signaling, thereby attenuating the hypoxia-induced downregulation of Cx43. Transcription factors NF-κB and AP-1/c-Jun participate in the regulation of hypoxia-induced Cx43 expression. MAPK signaling acts on Cx43 expression, most likely by modulating MMP enzymes.

Acknowledgments This work was supported by the National Nature Science Foundation of China (Grant No. 81360057, 81371333, 81160152) and Guangxi Nature Science Foundation (2013GXNSFAA 019184, 2013GXNSFCA019013).

Disclosures The authors have nothing to disclose.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Kumar NM, Gilula NB (1996) The gap junction communication channel. *Cell* 84:381–388
- Howarth FC, Nowotny N, Zilahi E, El Haj MA, Lei M (2007) Altered expression of gap junction connexin proteins may partly underlie heart rhythm disturbances in the streptozotocin-induced diabetic rat heart. *Mol Cell Biochem* 305:145–151
- Greener ID, Sasano T, Wan X, Igarashi T, Strom M, Rosenbaum DS et al (2012) Connexin43 gene transfer reduces ventricular tachycardia susceptibility after myocardial infarction. *J Am Coll Cardiol* 60:1103–1110
- Garcia-Touchard A, Henry TD, Sangiorgi G, Spagnoli LG, Mauriello A, Conover C et al (2005) Extracellular proteases in atherosclerosis and restenosis. *Arterioscler Thromb Vasc Biol* 25:1119–1127
- Siefert SA, Sarkar R (2012) Matrix metalloproteinases in vascular physiology and disease. *Vascular* 20:210–216
- Koenig GC, Rowe RG, Day SM, Sabeh F, Atkinson JJ, Cooke KR et al (2012) MT1-MMP-dependent remodeling of cardiac extracellular matrix structure and function following myocardial infarction. *Am J Pathol* 180:1863–1878
- Cho HJ, Kang JH, Kwak JY, Lee TS, Lee IS, Park NG et al (2007) Ascofuranone suppresses PMA-mediated matrix metalloproteinase-9 gene activation through the Ras/Raf/MEK/ERK- and Ap1-dependent mechanisms. *Carcinogenesis* 28:1104–1110
- Byun HJ, Hong IK, Kim E, Jin YJ, Jeoung DI, Hahn JH et al (2006) A splice variant of CD99 increases motility and MMP-9 expression of human breast cancer cells through the AKT-, ERK-, and JNK-dependent AP-1 activation signaling pathways. *J Biol Chem* 281:34833–34847
- Wang HH, Hsieh HL, Wu CY, Sun CC, Yang CM (2009) Oxidized low-density lipoprotein induces matrix metalloproteinase-9 expression via a p42/p44 and JNK-dependent AP-1 pathway in brain astrocytes. *Glia* 57:24–38
- Etoh T, Joffs C, Deschamps AM, Davis J, Dowdy K, Hendrick J et al (2001) Myocardial and interstitial matrix metalloproteinase activity after acute myocardial infarction in pigs. *Am J Physiol Heart Circ Physiol* 281:H987–H994
- Lindsey ML, Escobar GP, Mukherjee R, Goshorn DK, Sheats NJ, Bruce JA et al (2006) Matrix metalloproteinase-7 affects connexin-43 levels, electrical conduction, and survival after myocardial infarction. *Circulation* 113:2919–2928

12. Wu X, Li L (2012) Rosiglitazone suppresses lipopolysaccharide-induced matrix metalloproteinase-2 activity in rat aortic endothelial cells via Ras-MEK1/2 signaling. *Int J Cardiol* 158:54–58
13. Huang W, Andras IE, Rha GB, Hennig B, Toborek M (2011) PPARalpha and PPARgamma protect against HIV-1-induced MMP-9 overexpression via caveolae-associated ERK and Akt signaling. *FASEB J* 25:3979–3988
14. Huang W, Eum SY, Andras IE, Hennig B, Toborek M (2009) PPARalpha and PPARgamma attenuate HIV-induced dysregulation of tight junction proteins by modulations of matrix metalloproteinase and proteasome activities. *FASEB J* 23:1596–1606
15. Huang W, Rha GB, Chen L, Seelbach MJ, Zhang B, Andras IE et al (2010) Inhibition of telomerase activity alters tight junction protein expression and induces transendothelial migration of HIV-1-infected cells. *Am J Physiol Heart Circ Physiol* 298:H1136–H1145
16. Lu HS, Chen HP, Wang S, Yu HH, Huang XS, Huang QR et al (2012) Hypoxic preconditioning up-regulates DJ-1 protein expression in rat heart-derived H9c2 cells through the activation of extracellular-regulated kinase 1/2 pathway. *Mol Cell Biochem* 370:231–240
17. Mohammad G, Kowluru RA (2011) Novel role of mitochondrial matrix metalloproteinase-2 in the development of diabetic retinopathy. *Invest Ophthalmol Vis Sci* 52:3832–3841
18. Glukhov AV, Fedorov VV, Kalish PW, Ravikumar VK, Lou Q, Janks D et al (2012) Conduction remodeling in human end-stage nonischemic left ventricular cardiomyopathy. *Circulation* 125:1835–1847
19. Lin FL, Chang CI, Chuang KP, Wang CY, Liu HJ (2010) Advanced glycation end products down-regulate gap junctions in human hepatoma SKHep 1 cells via the activation of Src-dependent ERK1/2 and JNK/SAPK/AP1 signaling pathways. *Journal of agricultural and food chemistry*. 58:8636–8642
20. Hescheler J, Meyer R, Plant S, Krautwurst D, Rosenthal W, Schultz G (1991) Morphological, biochemical, and electrophysiological characterization of a clonal cell (H9c2) line from rat heart. *Circ Res* 69:1476–1486
21. Sipido KR, Marban E (1991) L-type calcium channels, potassium channels, and novel nonspecific cation channels in a clonal muscle cell line derived from embryonic rat ventricle. *Circ Res* 69:1487–1499
22. Weissenberg A, Sydlik U, Peuschel H, Schroeder P, Schneider M, Schins RP et al (2010) Reactive oxygen species as mediators of membrane-dependent signaling induced by ultrafine particles. *Free Radical Biol Med* 49:597–605
23. Hayden MS, Ghosh S (2012) NF-kappaB, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev* 26:203–234
24. Lawrence T (2009) The nuclear factor NF-kappaB pathway in inflammation. *Cold Spring Harb Perspect Biol*. 1:a001651
25. Zhao Y, Riviuccio MA, Lutz S, Scemes E, Brosnan CF (2006) The TLR3 ligand polyI: c downregulates connexin 43 expression and function in astrocytes by a mechanism involving the NF-kappaB and PI3 kinase pathways. *Glia*. 54:775–785
26. Mitchell JA, Lye SJ (2005) Differential activation of the connexin 43 promoter by dimers of activator protein-1 transcription factors in myometrial cells. *Endocrinology* 146:2048–2054
27. Shyu KG, Wang BW, Yang YH, Tsai SC, Lin S, Lee CC (2004) Amphetamine activates connexin43 gene expression in cultured neonatal rat cardiomyocytes through JNK and AP-1 pathway. *Cardiovasc Res* 63:98–108
28. Jia G, Cheng G, Gangahar DM, Agrawal DK (2008) Involvement of connexin 43 in angiotensin II-induced migration and proliferation of saphenous vein smooth muscle cells via the MAPK-AP-1 signaling pathway. *J Mol Cell Cardiol* 44:882–890
29. Severino A, Narducci ML, Pedicino D, Pazzano V, Giglio AF, Biasucci LM et al (2012) Reversible atrial gap junction remodeling during hypoxia/reoxygenation and ischemia: a possible arrhythmogenic substrate for atrial fibrillation. *Gen Physiol Biophys* 31:439–448
30. Beardslee MA, Lerner DL, Tadros PN, Laing JG, Beyer EC, Yamada KA et al (2000) Dephosphorylation and intracellular redistribution of ventricular connexin43 during electrical uncoupling induced by ischemia. *Circ Res* 87:656–662
31. Marquez-Rosado L, Solan JL, Dunn CA, Norris RP, Lampe PD (2012) Connexin43 phosphorylation in brain, cardiac, endothelial and epithelial tissues. *Biochim Biophys Acta* 1818:1985–1992
32. Turner MS, Haywood GA, Andreka P, You L, Martin PE, Evans WH et al (2004) Reversible connexin 43 dephosphorylation during hypoxia and reoxygenation is linked to cellular ATP levels. *Circ Res* 95:726–733