

Regulation of membrane excitability: a convergence on voltage-gated sodium conductance

Wei-Hsiang Lin · Richard A. Baines

Received: 17 January 2014 / Accepted: 11 March 2014 / Published online: 29 March 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract The voltage-gated sodium channel (Na_v) plays a key role in regulation of neuronal excitability. Aberrant regulation of Na_v expression and/or function can result in an imbalance in neuronal activity which can progress to epilepsy. Regulation of Na_v activity is achieved by coordination of a multitude of mechanisms including RNA alternative splicing and translational repression. Understanding of these regulatory mechanisms is complicated by extensive genetic redundancy: the mammalian genome encodes ten Na_v s. By contrast, the genome of the fruitfly, *Drosophila melanogaster*, contains just one Na_v homologue, encoded by *paralytic* (*DmNa_v*). Analysis of splicing in *DmNa_v* shows variants exhibit distinct gating properties including varying magnitudes of persistent sodium current (I_{NaP}). Splicing by Pasilla, an identified RNA splicing factor, alters I_{NaP} magnitude as part of an activity-dependent mechanism. Enhanced I_{NaP} promotes membrane hyperexcitability that is associated with seizure-like behaviour in *Drosophila*. Nova-2, a mammalian Pasilla homologue, has also been linked to splicing of Na_v s and, moreover, mouse gene knockouts display seizure-like behaviour.

Expression level of Na_v s is also regulated through a mechanism of translational repression in both flies and mammals. The translational repressor Pumilio (Pum) can bind to *Na_v* transcripts and repress the normal process of translation, thus regulating sodium current (I_{Na}) density in neurons. Pum2-deficient mice exhibit spontaneous EEG abnormalities. Taken together, aberrant regulation of Na_v function and/or expression is often epileptogenic. As such, a better understanding of regulation of membrane excitability through RNA alternative splicing and translational repression of Na_v s should provide new leads to treat epilepsy.

Keywords Excitability · *Drosophila* · Epilepsy · *Paralytic* · Splicing · Translational repression

Introduction

The regulation of neuronal excitability—primarily the ability to maintain action potential firing within physiological constraints—is an important mechanism for maintenance of neuronal stability [1]. Without such regulation, chronic changes in levels of synaptic excitation have the potential to destabilise neural circuits leading to an imbalance in neuronal activity. One consequence of activity imbalance is seizure, which if recurrent is termed epilepsy [2]. The voltage-gated sodium channel (Na_v) plays a key role in the regulation of neuronal excitability because its activation results in action potential firing. It is perhaps, therefore, not surprising that many mechanisms that act to stabilise neuronal activity do so through modifying the activity of this class of ion channel [1, 3–7].

Ten genes (*SCN1A-SCN11A*), encoding pore-forming α -subunits, are present in mammals [8]. This relatively high number is, however, insufficient to support the wide diversity of Na_v kinetics reported in the nervous system. Diversity of signalling is critically reliant on additional mechanisms such as alternative splicing, RNA editing, and protein modification (i.e., phosphorylation) [9–12, 4]. However, whilst the importance of posttranscriptional and posttranslational modifications is appreciated for refining activity of channel subtypes, understanding of the mechanisms that neurons employ to determine which form of Na_v to express remains rudimentary. In contrast to mammals, the genome of the fruitfly *Drosophila melanogaster* contains only one Na_v channel homologue: encoded by *paralytic* (*DmNa_v*) [13, 5]. The lack of redundancy, coupled with a high degree of structural and functional homology, makes *DmNa_v* an advantageous model with which to study the role of this ion channel family [14,

W.-H. Lin · R. A. Baines (✉)
Faculty of Life Sciences, University of Manchester, Oxford Road,
Manchester, UK
e-mail: Richard.Baines@manchester.ac.uk

15]. In this review, we use $DmNa_v$ as a model to summarise recent findings relating to how neurons generate diversity in Na_v channel activity and to stabilise neuronal circuit function when faced with changing levels of synaptic excitation.

Alternative splicing generates diversity in Na_v channel activity

Alternative splicing involves the substitution, removal, and/or inclusion of exonic sequences within a pre-messenger RNA (mRNA) to produce transcripts encoding related protein isoforms [9]. Estimates indicate that ~95 % of human genes are alternatively spliced [16, 17]. Variant transcripts of $DmNa_v$, first reported by Loughney et al., (1989), were among the first evidence for the existence of alternative splicing of this family of gene products. Subsequent studies in *Drosophila*, *Musca*, and cockroach have identified 15 alternatively spliced exons [18, 19, 14, 20, 15]. Importantly, alternative splicing of exons is replicated in mammalian Na_v channels [21–23]. Spliced exons are conserved across evolutionarily diverged species, strongly indicative of fundamental physiological importance.

A recent structure-function study has described the effects to $DmNa_v$ channel kinetics of alternative splicing [15]. Of the 15 known splice decisions, two splice events are mutually exclusive incorporating one of either a pair of exons (*C/D* and *K/L*). Both exon pairs are membrane spanning, contributing to domains IIS4–5 and IIIS3–4, respectively. The remaining 11 spliced exons (*J*, 7, 8, *I*, *A*, *B*, *E*, *F*, 22, *H*, 23) are independent and cytoplasmic. Heterologous expression of $DmNa_v$ splice variants in *Xenopus* oocytes shows that such splicing imparts specific attributes to channel kinetics. For example, inclusion of exon *F* results in a hyperpolarising shift in activation kinetics, indicative of increased excitability for those neurons that express *F*-containing variants. By contrast, inclusion of exons *J* and *E* results in a depolarising shift of activation voltages which are predicted to reduce neuron excitability. On the other hand, channels expressing exon *H* inactivate at more depolarised voltages, predicted to make neurons more excitable. Finally, the choice to include mutually exclusive exons *K* or *L* markedly affects the magnitude of the persistent current (I_{NaP}) that arises from incomplete inactivation of the channel [24, 5, 25]. Inclusion of exon *K* results in a smaller I_{NaP} relative to that observed from expression of transcripts that contain exon *L*, in otherwise identical channels. Increasing I_{NaP} leads to an increased frequency of action potential firing [26, 5]. Figure 1 summarises the known splicing events of $DmNa_v$, and the effect on channel kinetics and/or I_{NaP} is summarised in Table 1. Of course, the caveat to heterologous expression is that the nature of the cell membrane of the cell type used may influence the kinetics of expressed channels compared to expression, in this instance, in *Drosophila* neurons [27]. Attempts to express $DmNa_v$

variants in *Drosophila* neurons, using the well-characterised GAL4/UAS system, has repeatedly failed to produce functional channels, for unknown reasons (Lin and Baines, unpublished observations).

Both embryonic and adult *Drosophila* CNS expresses a wide diversity of $DmNa_v$ splice forms. However, the profile of splicing differs between these two stages. This is indicative that different Na_v properties are required at each stage and that these differences are achieved through splicing. Differences of spliced exons expressed in these two stages include a greater usage of exon *J* (89 %) but not of exon *F* (10 %) in adults and vice versa in embryos (10 % exon *J* and 78 % exon *F*) [14, 15]. However, the physiological significance of these differences is not clear. It is interesting to note that $DmNa_v$ transcripts which lack a majority of common cytoplasmically located spliced exons result in channels with shifted activation and inactivation kinetics towards hyperpolarised and depolarised voltages, respectively, and which also exhibit a much larger I_{NaP} . These properties are predicted to make neurons highly excitable [15]. Similarly, analysis of splicing of Na_v in other insects shows that it is important for functional properties of the expressed channel. For example, exclusion of optional exon *B* (located at the linker between the domains I and II, but not equivalent to exon *B* in *Drosophila*) in cockroach sodium channels ($BgNa_v$) potentiates the amplitude of the fast-activating and inactivating I_{Na} transient current (I_{NaT}), which is likely to increase cell excitability (Table 1) [28]. Indeed, an emerging theme is that splicing in of optional exons primarily reduces channel activity and hence, membrane excitability, in order to suit the requirements of neural signalling.

Splicing in intracellular coding regions of mammalian Na_v s can also result in changes to channel activity. For example, the human $Na_v1.3$ (*SCN3A*) alternative spliced exon 12, which encodes an intracellular loop between domains I and II, results in the generation of multiple isoforms. By using multiple splice donor sites in exon 12, four different variants are produced: 12v1, 12v2, 12v3, and 12v4. The variant 12v4, when compared to 12v2, seemingly increases membrane excitability by shifting activation kinetics of the expressed I_{Na} . By contrast, inactivation kinetics showed a shift toward hyperpolarising potentials for 12v1 over 12v3, indicative that expressing 12v1 might be expected to decrease membrane excitability (Table 1) [29, 30]. Taken together, Na_v gating properties can be determined by the inclusion of exons to alter membrane excitability. However, details of how inclusion of specific spliced exons change gating of the affected channels remains to be determined. A possible mechanism for altering channel kinetics is the phosphorylation state of the channel [30, 31]. Analysis of the amino acid sequence of human $Na_v1.3$ splice variants revealed the presence of two additional phosphorylation sites (protein kinase C on Ser^{631/632} and casein kinase II on Ser⁶⁴⁶) in 12v3 and 12v4 that are absent from other variants [30]. Changing membrane excitability




Fig. 1 Schematic of the predicted topology of the voltage-gated sodium channel showing approximate locations of *Drosophila* spliced exons. Cytoplasmic *DmNav_v* exons J, 7, 8, I, A, B, E, F, 22, H, 23 are optional, while exons C/D and K/L are mutually exclusive. *DmNav_v* exon 8 is conserved in human *Nav_s* as mutually exclusive spliced exons 5A and 5N (6A and 6N in *hNav_v.1.2* and *hNav_v.1.3* due to different exon numbering in the consensus gene sequence), and identical residues are shown in black boxes. Exon 5A and 5N of *hNav_v.1.1* differ by 3 amino acids, shown in grey boxes in the 5N sequence. Mutually exclusive *DmNav_v* spliced exon L and cockroach *BgNav_v* exon G1 are identical and are conserved in

human: exon 18A of *hNav_v.1.1* and *hNav_v.1.6*. *DmNav_v* exons K and L differ by 16/41 residues (shown in grey boxes in the exon K sequence). Inclusion of *BgNav_v* exon G3 and *hNav_v.1.6* exon 18N generated a truncated channel. Exon 12 of *hNav_v.1.3* is located in the intracellular loop between domains I and II. By using different splice donor sites in exon 12, four spliced variants, 12v1, 12v2, 12v3, and 12v4 can be generated. The amino acid sequences are obtained as follows: *DmNav_v*, 8, K, and L [15]; *hNav_v*, 5A and 5N [61]; *hNav_v.1.6* 18A and 18N [23]; *BgNav_v*, G1, G2, and G3 [20]; *hNav_v.1.3* 12v1, 12v2, 12v3, and 12v4 [30]

Table 1 Summary of spliced *Nav_v* exons that are known to affect channel kinetics. Predicted influence on neuron excitability due to splicing are stated, increased (↑), decreased (↓), or complex (?)

Channel	Exon	Expression system	Predicted effect on cell excitability (by changing)	References
<i>DmNav_v</i>	J, E	<i>Xenopus</i> oocytes	↓ (act →)	[15]
<i>DmNav_v</i>	F	<i>Xenopus</i> oocytes	↑ (act ←)	[15]
<i>DmNav_v</i>	H	<i>Xenopus</i> oocytes	↑ (inact →)	[15]
<i>DmNav_v</i>	K	<i>Xenopus</i> oocytes	↓ (<i>I_{NaP}</i> amplitude ↓)	[15]
<i>DmNav_v</i>	L	<i>Xenopus</i> oocytes	↑ (<i>I_{NaP}</i> amplitude ↑)	[15]
<i>hNav_v.1.1</i>	5A	HEK293T	↑ (<i>I_{NaP}</i> amplitude ↑, inact →)	[61]
<i>hNav_v.1.1</i>	5N	HEK293T	↓ (<i>I_{NaP}</i> amplitude ↓, inact ←)	[61]
<i>hNav_v.1.3</i>	12v1	<i>Xenopus</i> oocytes	↓ (inact ←)	[30]
<i>hNav_v.1.3</i>	12v2	<i>Xenopus</i> oocytes	↓ (act →)	[30]
<i>hNav_v.1.3</i>	12v3	<i>Xenopus</i> oocytes	↑ (inact →)	[30]
<i>hNav_v.1.3</i>	12v4	<i>Xenopus</i> oocytes	↑ (act ←)	[30]
<i>BgNav_v</i>	B	<i>Xenopus</i> oocytes	↑ (<i>I_{NaT}</i> amplitude ↑)	[28]
<i>BgNav_v</i>	G1	<i>Xenopus</i> oocytes	? (<i>I_{NaT}</i> amplitude ↓, act ←, inact →)	[20]
<i>BgNav_v</i>	G2	<i>Xenopus</i> oocytes	? (<i>I_{NaT}</i> amplitude ↑, act →, inact ←)	[20]

Specific changes observed to channel kinetics are as follows: depolarising (→) or hyperpolarising (←) shifts in activation (act) or inactivation (inact) and/or increased (↑) or decreased (↓) transient (*I_{NaT}*) or persistent sodium current (*I_{NaP}*) amplitude

through phosphorylation in the I-II linker of Na_v may influence current amplitude without significantly affecting gating properties [11, 32–36].

Persistent Na current and membrane excitability

The persistent Na current (I_{NaP}) has been identified to play critical roles in regulating membrane excitability [37]. Moreover, numerous point mutations in human Na_vs , identified in patients with epilepsy, potentiate this component of the voltage-gated Na current (I_{Na}) [26]. Interestingly, I_{NaP} is also a primary target of some clinically used antiepileptic drugs, including phenytoin, valproic acid, and lamotrigine. [38–40]. It is significant, therefore, that the magnitude of this current can be altered through alternative splicing. However, our understanding of the molecular machinery that regulates splicing of Na_vs is poor. This is unfortunate because a fuller understanding may offer new leads for antiepileptic drug design.


In early behavioural screens of *Drosophila*, different single-gene mutations were identified that induce a seizure-like phenotype when flies are exposed to strong sensory stimuli. Following a mechanical shock, such as vortexing or harsh-tapping of the culture vial, bang-sensitive (bs) mutant flies exhibit a stereotyped sequence of seizure-like spasms, followed by a period of paralysis, and then a second recovery seizure-like phase that precedes a more complete recovery (Fig. 2) [41, 42]. Despite the evolutionary distance, the resemblance in epileptiform activity between fly and humans and the response to clinical antiepileptic drugs make bs mutants an accepted model for studying epilepsy [42–49]. One such bs mutant—*slamdance* (*sda*)—which has a deficiency of aminopeptidase N, exhibits increased seizure-like activity in

response to electrical stimulation in the larval stage. Detailed electrophysiology shows that I_{NaP} is significantly increased in central motoneurons in this mutant [48]. A molecular analysis reveals that splicing of *DmNa_v* is similarly altered in the *sda* mutation to favour inclusion of exon *L* at the expense of exon *K* [25]. As previously described (see above), inclusion of exon *L* results in channels that, when expressed in *Xenopus* oocytes, exhibit a larger I_{NaP} [15]. Seizure-like behaviour, in response to electric shock, along with the increased I_{NaP} and increased inclusion of *L* isoform are all reversed by feeding larvae with antiepileptic drugs (AEDs) including phenytoin (Phy) and gabapentin (Gbp) [25, 48]. Thus, a better understanding of how I_{NaP} is regulated, particularly through splicing, may be beneficial for epilepsy therapy.

Activity-dependent alternative splicing regulation of I_{NaP} expression

Seizures can be induced in both mammals and flies through ingestion of proconvulsants such as picrotoxin (PTx) [50, 46]. PTx elicits seizure through antagonism of the GABA_A receptor-suppressing synaptic inhibition [51, 52]. Remarkably, we showed that enhancement of synaptic activity in wild-type larvae, through ingestion of PTx, is sufficient to increase inclusion of exon *L* in *DmNa_v*, increasing I_{NaP} as a consequence and inducing a bang-sensitive phenotype. Conversely, seizure activity can be rescued via enhancing synaptic inhibition in *sda* through ingestion of GABA [25]. Both manipulations suggest that the ‘decision’ to splice either exon *L* or *K* is dictated by neuronal activity: i.e. activity-dependent. Increasing synaptic excitatory input results in greater inclusion of exon *L*, which in turn increases I_{NaP} and membrane excitability [25]. Increased excitability would be

Fig. 2 *Drosophila* bang-sensitive mutant behaviour. Brief vortexing (~10 s) of the culture vial, containing bang-sensitive mutant flies, induces a stereotyped sequence of seizure-like spasms, followed by a period of paralysis, and then a recovery seizure-like phase that precedes a normal but refractory phase followed ultimately by a complete recovery



predicted to further increase inclusion of exon *L* up to a maximum of 100 % (which is observed in *sda* and other bs mutants). Such a self-reinforcing cycle provides a plausible, although untested, explanation of the clinical observation in which untreated seizures beget seizures, i.e. paroxysmal activity has the potential to promote susceptibility to further seizures [53, 54].

Splicing of *Na_v* transcripts in response to activity provides an important mechanism for inducing changes in excitability. Because of the complexity of the mammalian CNS, with its larger number of expressed *Na_v*s, the extent of splicing and its functional consequences are not well understood. However, the high degree of homology between *DmNa_v* and its mammalian counterparts allows us to use the former to guide future studies in mammals. *Drosophila* exon K/L (located in homology domain III S3–4) is conserved in the homologous domain III from insect to mammal [20, 18, 23], although the outcome of splicing differs. Splicing at this location in cockroach produces three mutually exclusive transcripts that contain spliced exons *G1*, *G2*, or *G3*. *G3* contains a stop codon and generates a nonfunctional channel, whereas *G1* and *G2* result in channels that differ in peak *I_{NaT}* amplitude, gating properties (Table 1) and sensitivity to deltamethrin, a pyrethroid insecticide [20]. In mammals, this same region is also spliced in *Na_v1.1* and *Na_v1.6*—resulting in the inclusion of exons *18A* or *18N* [21–23]. Exon *18A* predominates in adult brain and *18N* in embryo and nonneuronal tissues. Similar to the cockroach exon *G3*, mammalian exon *18N* contains a stop codon and generates a truncated channel. These truncated *Na_v* channels that contain only the first two domains, express mainly in nonneuronal tissues, and are hypothesised to be a ‘fail-safe’ mechanism to prevent the expression of functional *Na_v*s in nonexcitable cells [55, 23, 21, 22].

A second splicing event in mammalian *Na_v*s is noteworthy because it occurs at the equivalent S3–4 region of homologous domain I. Similar to *DmNa_v*, splicing at exon 5 in *Na_v1.1* is mutually exclusive with the choice of either exons *5A* or *5N* (again for adult and neonatal). Alternative splicing in this region is also observed in *Na_v1.2*, *1.3*, *1.6* and *Na_v1.7* in both human and mouse [56–60]. In human *Na_v1.1*, three amino acids differ between exon *5A* and *5N*; however, the channels exhibit distinct gating properties. Heterologous expression of human *Na_v1.1-5N*, in HEK293T cells, produces channels which exhibit more rapid inactivation and reduced *I_{NaP}* compared to *Na_v1.1-5A*. Whilst much needs to be learnt about this splice event, these results suggest that splicing at this location is sufficient to confer changes in neuronal excitability (Table 1) [61]. Intriguingly, inclusion of neonatal exon *6N* is increased in both *Na_v1.2* and *Na_v1.3* following electrical or kainate-induced seizure in adult rat hippocampus [62, 63], perhaps indicative that splicing may similarly be activity-regulated in mammals, as it is in the fly.

Pasilla/Nova, critical factors involved in activity-dependent alternative splicing

A screen of RNA-binding proteins in *Drosophila* first identified Pasilla (Ps) to be sufficient to regulate splicing of mutually exclusive exons *K* and *L* in *DmNa_v* [64]. The inclusion of exon *K* is significantly increased to 50 % in a *ps* loss-of-function mutant indicating that the presence of Ps is necessary for the inclusion of exon *L* [15]. Loss of one copy of *ps* is also sufficient to rescue the bs-associated seizure behaviour of *sda* mutants and, moreover, to also prevent PTx-induced seizure in WT background (Fig. 3) [25]. These data suggest that Ps is required for the underlying activity-dependent splicing mechanism. Ps, which contains a K-homology (KH) RNA-binding domain [65, 66], encodes the *Drosophila* homologue of the human neuro-oncological ventral antigen 1 and 2 (Nova-1 and Nova-2, respectively) proteins [67, 68]. Nova-1 and Nova-2 are expressed to high levels in brain, however, in largely nonoverlapping patterns [69–71]. By recognising YCAY-motifs, located either in introns or 3’UTRs of target




Fig. 3 Pasilla is required for activity-dependent inclusion of exon *L* of *DmNa_v*. Prolonged mean recovery time to electroshock of third instar larvae (i.e. increased severity of seizure) is observed in both slamdance (*sda*) mutants and picrotoxin (PTx)-fed WT flies. Analysis of splicing of *DmNa_v* in whole CNS of such larvae shows that inclusion of exon *L* increased to ~100 %. In *sda*, loss of one copy of *pasilla* (*sda*^{+/-}, *ps*^{+/-}) is sufficient to decrease the inclusion of exon *L* and to rescue seizure-like behaviour. Similarly, removal of one copy of *ps* in WT larvae (*ps*^{+/-}) diminishes PTx-induced seizure, as well as inclusion of exon *L*. Data are taken from [25]

transcripts, Nova1/2 regulate neuronal alternative splicing and also mediate transportation of some target transcripts between the nucleus and cytoplasm [68, 72, 73]. Splicing in at least 17 ion channel genes, including $Na_v s$, is predicated to be regulated by Nova-2 [74, 75]. Significantly, overexpression of Nova-2 in HEK293 cells results in an increase in the $Na_v 1.1-5N$ splice variant [75]. In support of this, *Nova-2* and $Na_v 1.1-5N$ transcript abundance are upregulated in temporal neocortical tissue of mesial temporal lobe epilepsy patients. [75]. The relationship between Nova expression and epilepsy has been further examined by EEG recordings in *Nova-2*^{+/-} heterozygous mice (*Nova-2*^{-/-} mice die within 2–3 weeks of birth). Perturbing Nova steady-state levels in *Nova-2*^{+/-} heterozygous mice gives rise to cortical hyperexcitability and also to spontaneous generalised seizure discharge [73]. Moreover, Nova localization shifts from primarily nuclear to cytoplasmic within 2–4 h after pilocarpine-induced seizure [73]. Taken together, these findings strongly implicate perturbation of Nova-2 function contributes to epileptogenesis. The corollary would be that manipulation of Nova activity might be antiepileptic. The conservation of function between Ps and Nova offers the exciting opportunity to utilise *Drosophila* to rapidly identify molecules that might influence Nova function.

Sodium channel expression and homeostasis

Control of neuron excitability is known to be achieved at the genomic level through transcriptional regulation of Na_v channel genes [1, 3, 76, 6]. In *Drosophila* CNS, the regulation of the voltage-gated sodium current (I_{Na}) can be achieved through activity-dependent alteration of *DmNa_v* mRNA level [77, 4, 5, 78]. Removal of excitatory synaptic inputs to motoneurons, achieved by expressing tetanus toxin light chain in all central neurons, significantly increased *DmNa_v* transcript abundance and also the magnitude of I_{Na} in motoneurons. On the other hand, enhanced excitatory synaptic release, achieved by increasing cAMP level in the CNS, decreased both mRNA level and I_{Na} [77, 5]. This homeostatic mechanism is ideally suited to allow membrane excitability to track the degree of synaptic excitation to which a neuron is exposed (i.e. neuronal homeostasis).

Mammalian neurons (e.g. rat) exhibit the same type of activity-dependent homeostasis of membrane excitability [3]. Deprivation of synaptic excitation in cortical neuron cultures, achieved by chronically blocking glutamatergic signalling, resulted in increased $Na_v 1.6$ mRNA expression, I_{Na} , and membrane excitability [79]. The underlying mechanism of this homeostatic regulation, in both flies and mammals, requires the protein Pumilio (Pum) [79, 80, 5, 78]. Pumilio is a member of the Pum and FBF (PuF) RNA-binding protein family [81, 82] and is evolutionarily conserved in many species including yeast (*Saccharomyces cerevisiae*),

C. elegans, *Drosophila*, *Anopheles*, zebrafish, *Xenopus*, mouse, and human [82, 83].

In the fly CNS, activity-dependent increase in Pum level results in the translational repression of *DmNa_v* transcripts, reducing I_{Na} and membrane excitability [5, 78]. This mechanism is dynamic, such that decreasing levels of synaptic excitation results in decreased Pum level, increased *DmNa_v* transcript abundance, and potentiation of membrane excitability. In rat cortical neurons, the level of Pum was similarly observed to be activity-dependent, mirroring the mechanism observed in the fly [79, 84]. Pum is able to repress translation through binding a specific motif—termed Nanos response element (NRE) [85]—present in both *DmNa_v* and rat $Na_v 1.6$ transcripts [79, 78]. Once Pum is bound to a transcript, cofactors Nanos [86] and brain tumour [86] are recruited to form a quaternary RNA-protein complex that causes transcript deadenylation [87] and consequently repression of translation. An 8-nucleotide core motif UGUA(A/U/C)AUA [88] of the NRE is sufficient for the binding of Pum to *DmNa_v* transcripts [78], and this motif exists in about 10 % of all *Drosophila* transcripts [88]. Notably, those 10 % of transcripts were only interrogated for NREs present in the 3'UTR region; however, Pum binds to the NRE located in the 3' end of the open reading frame (ORF) in both *DmNa_v* and rat $Na_v 1.6$ [79, 78]. Therefore, there might be many more Pum targets yet to be identified.

In a genome-wide screening of transcripts associated with the RNA-binding region of Pum, more than 1,000 distinct mRNAs were identified [88]. This suggests that Pum is broadly involved in posttranscriptional regulation of many genes. Indeed, in addition to regulating translation of $Na_v s$, Pum has also been implicated to regulate dendritogenesis [89, 90], expression of glutamate receptors [91], and aspects of memory and learning in higher brain centres [92]. Behaviour training of long-term memory (LTM) produced by spaced training (ten training sessions with a 15-min rest interval between each session), compared to anaesthesia-resistant memory (ARM) produced by massed training (ten training sessions without rest intervals), resulted in *pum* mRNA upregulation. Pum mutant flies also showed defects in LTM formation. [92]. Pum regulates NMJ morphology via negative regulation of the translational factor eIF-4E expression by directly binding to an NRE in the 3'UTR of the *eIF-4E* transcript [90]. Pum loss-of-function mutants show enhanced expression of eIF-4E and upregulated GluRIIA expression and increased frequency of spontaneous neurotransmitter release [91]. Thus, Pum is seemingly central to many aspects of CNS function, not least of which is homeostatic control of neuronal excitability. In this regard, it is significant that in mouse, Pum2 deficiency leads to spontaneous EEG abnormalities and lower seizure thresholds to the proconvulsant pentylentetrazole [93]. Similar to Pum, the $Na_v 1.6$ transcript is upregulated in *CELF4* (CUGBP, ELAV-like family

member 4) deficient mice [94]. CELF4 is similarly a brain-specific neuronal RNA-binding protein and binds to the 3'UTR of *Nav1.6*. Because mammalian *Nav1.6* is the primary determinant of action potential initiation and main contributor of I_{NaP} in excitatory neurons, upregulated *Nav1.6* mRNA results in increasing neuronal excitability [95]. Consequently, *CELF4* deficient mice exhibit both convulsive and nonconvulsive (absence-like) seizures and also have a lower seizure threshold [94, 96]. These findings demonstrate that understanding the regulation of I_{Na} or I_{NaP} via RNA-binding proteins is a potentially important approach for epilepsy therapy.

Summary and outlook

Voltage-gated sodium channels are important determinants for controlling membrane excitability. Regulation of Na_v activity is achieved, at least in part, by coordination of RNA alternative splicing and translational repression of *Nav* transcripts (Fig. 4). When one considers additional mechanisms of regulation of Na_v channel activity, including RNA editing [28], phosphorylation [33, 32, 11, 4], trafficking [97, 98], and degradation [98–100], it becomes clear that these channels are subject to both considerable and diverse regulation consistent with the high level of channel diversity observable in the multitude of neuron types in the human brain. The utilisation of model systems, including *Drosophila*, offers the significant opportunity to rapidly progress understanding in these and related areas.

A particular area where *Drosophila* is already making a contribution to understanding epilepsy is through modelling human *Nav* point mutations. A variety of techniques now exist to allow such mutations to be ‘knocked-in’ to *DmNav*. Sun et al. [101] recently reported a *Nav1.1* (K1270T) knock-in that recapitulates a mutation associated with genetic epilepsy with

febrile seizures plus (GEFS+). Electrophysiological analysis shows this to be a gain-of-function mutation that results in a hyperpolarizing shift in the deactivation potential for I_{NaP} . This approach not only serves to validate the genetic basis of human disease, but also provides a sensitised genetic background for high-throughput, low cost, screens to identify novel compounds that have antiepileptic properties. Identification of novel targets, such as splicing regulators, can also be quickly developed as the basis of screens with the potential advantage of identifying antiepileptic compounds which interact with nontraditional targets. By far the most common targets of currently used AEDs are ion channels and, whilst these offer effective therapeutic targets, there might be much to be gained from identifying additional targets which would facilitate combinatorial therapy. Combinations of AEDs are showing promise for the treatment of intractable epilepsy [102].

Use of *Drosophila* (and other simple model systems) also offers the prospect of exploring the mechanistic basis of epileptogenesis from understanding how small seizures may lead to larger seizures to providing novel approaches to prevent epilepsy from progressing, even when an epilepsy-associated mutation is present. For example, we recently reported that the presence of phenytoin, during embryogenesis when the CNS first forms neural circuits, prevents the normal seizure phenotype characteristic of the *Drosophila sda* mutant [48]. The inference from this study is that early intervention may be beneficial in blocking epileptogenesis by preventing activity-dependent feedback mechanisms that we spotlight in this review. The finding of conservation of regulatory mechanisms between insects such as *Drosophila* and mammals validates the use of simpler model organisms to provide better understanding of *Nav* regulation in humans with an obvious benefit of novel therapies for epilepsy.




Fig. 4 Membrane excitability is regulated by activity-dependent RNA alternative splicing and translational repression of voltage-gated sodium channel transcripts. Control of membrane excitability through Na_v activity is achieved by regulation of channel kinetics, current density (I_{Na}), and magnitude of persistent Na current (I_{NaP}). RNA alternative splicing results in splice variants which exhibit different channel gating properties including activation and inactivation kinetics and I_{NaP} . Splicing is

regulated, in part, by Pasilla in *Drosophila* and in humans by its homologue, Nova. In *Drosophila*, increased synaptic excitation results in increased I_{NaP} which in turn feeds back to further increase synaptic excitation. This self-reinforcing cycle likely further increases I_{NaP} (dashed line) leading to seizure. Current density of Na_v can be regulated through a mechanism of translational repression of *Nav* transcripts via Pumilio and possibly CELF4

Acknowledgments We thank Dr. Stephanie Schorge (UCL, UK) for commenting on the manuscript. We also thank members of the Baines group for helpful comments and advice. Research in the Baines lab is supported by both the BBSRC (BB/J005002/1) and MRC (MR/J009180/1).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Turrigiano GG, Nelson SB (1998) Thinking globally, acting locally: AMPA receptor turnover and synaptic strength. *Neuron* 21(5):933–935
- Catterall WA (2000) From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron* 26(1):13–25
- Desai NS, Rutherford LC, Turrigiano GG (1999) Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nature neuroscience* 2(6):515–520. doi:10.1038/9165
- Baines RA (2003) Postsynaptic protein kinase A reduces neuronal excitability in response to increased synaptic excitation in the *Drosophila* CNS. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 23(25):8664–8672
- Mee CJ, Pym EC, Moffat KG, Baines RA (2004) Regulation of neuronal excitability through pumilio-dependent control of a sodium channel gene. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 24(40):8695–8703. doi:10.1523/JNEUROSCI.2282-04.2004
- Marder E, Goaillard JM (2006) Variability, compensation and homeostasis in neuron and network function. *Nature reviews Neuroscience* 7(7):563–574. doi:10.1038/nrn1949
- Schulz DJ, Baines RA, Hempel CM, Li L, Liss B, Misonou H (2006) Cellular excitability and the regulation of functional neuronal identity: from gene expression to neuromodulation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 26(41):10362–10367. doi:10.1523/JNEUROSCI.3194-06.2006
- Goldin AL (2001) Resurgence of sodium channel research. *Annual review of physiology* 63:871–894. doi:10.1146/annurev.physiol.63.1.871
- Li Q, Lee JA, Black DL (2007) Neuronal regulation of alternative pre-mRNA splicing. *Nature reviews Neuroscience* 8(11):819–831. doi:10.1038/nrn2237
- Dong K (2007) Insect sodium channels and insecticide resistance. *Invertebrate neuroscience : IN* 7(1):17–30. doi:10.1007/s10158-006-0036-9
- Smith RD, Goldin AL (1996) Phosphorylation of brain sodium channels in the I–II linker modulates channel function in *Xenopus* oocytes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 16(6):1965–1974
- Smith RD, Goldin AL (2000) Potentiation of rat brain sodium channel currents by PKA in *Xenopus* oocytes involves the I–II linker. *American journal of physiology Cell physiology* 278(4):C638–C645
- Feng G, Deak P, Chopra M, Hall LM (1995) Cloning and functional analysis of TipE, a novel membrane protein that enhances *Drosophila* para sodium channel function. *Cell* 82(6):1001–1011
- Olson RO, Liu Z, Nomura Y, Song W, Dong K (2008) Molecular and functional characterization of voltage-gated sodium channel variants from *Drosophila melanogaster*. *Insect biochemistry and molecular biology* 38(5):604–610. doi:10.1016/j.ibmb.2008.01.003
- Lin WH, Wright DE, Muraro NI, Baines RA (2009) Alternative splicing in the voltage-gated sodium channel DmNav regulates activation, inactivation, and persistent current. *Journal of neurophysiology* 102(3):1994–2006. doi:10.1152/jn.00613.2009
- Pan Q, Shai O, Lee LJ, Frey BJ, Blencowe BJ (2008) Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nature genetics* 40(12):1413–1415. doi:10.1038/ng.259
- Wang ET, Sandberg R, Luo S, Khrebtkova I, Zhang L, Mayr C, Kingsmore SF, Schroth GP, Burge CB (2008) Alternative isoform regulation in human tissue transcriptomes. *Nature* 456(7221):470–476. doi:10.1038/nature07509
- Davies TG, Field LM, Usherwood PN, Williamson MS (2007) A comparative study of voltage-gated sodium channels in the Insecta: implications for pyrethroid resistance in Anopheline and other Neopteran species. *Insect molecular biology* 16(3):361–375. doi:10.1111/j.1365-2583.2007.00733.x
- Lee SH, Ingles PJ, Knipple DC, Soderlund DM (2002) Developmental regulation of alternative exon usage in the house fly *Vssc1* sodium channel gene. *Invertebrate neuroscience : IN* 4(3):125–133. doi:10.1007/s10158-001-0014-1
- Tan J, Liu Z, Nomura Y, Goldin AL, Dong K (2002) Alternative splicing of an insect sodium channel gene generates pharmacologically distinct sodium channels. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 22(13):5300–5309. doi:20026551
- Diss JK, Archer SN, Hirano J, Fraser SP, Djamgoz MB (2001) Expression profiles of voltage-gated Na(+) channel alpha-subunit genes in rat and human prostate cancer cell lines. *Prostate* 48(3):165–178. doi:10.1002/pros.1095
- Oh Y, Waxman SG (1998) Novel splice variants of the voltage-sensitive sodium channel alpha subunit. *Neuroreport* 9(7):1267–1272
- Plummer NW, McBurney MW, Meisler MH (1997) Alternative splicing of the sodium channel SCN8A predicts a truncated two-domain protein in fetal brain and non-neuronal cells. *The Journal of biological chemistry* 272(38):24008–24015
- Kiss T (2008) Persistent Na-channels: origin and function. A review. *Acta biologica Hungarica* 59(Suppl):1–12. doi:10.1556/ABiol.59.2008.Suppl.1
- Lin WH, Gunay C, Marley R, Prinz AA, Baines RA (2012) Activity-dependent alternative splicing increases persistent sodium current and promotes seizure. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32(21):7267–7277. doi:10.1523/JNEUROSCI.6042-11.2012
- Stafstrom CE (2007) Persistent sodium current and its role in epilepsy. *Epilepsy currents / American Epilepsy Society* 7(1):15–22. doi:10.1111/j.1535-7511.2007.00156.x
- Escayg A, Goldin AL (2010) Sodium channel SCN1A and epilepsy: mutations and mechanisms. *Epilepsia* 51(9):1650–1658. doi:10.1111/j.1528-1167.2010.02640.x
- Song W, Liu Z, Tan J, Nomura Y, Dong K (2004) RNA editing generates tissue-specific sodium channels with distinct gating properties. *The Journal of biological chemistry* 279(31):32554–32561. doi:10.1074/jbc.M402392200
- Dietrich PS, McGivern JG, Delgado SG, Koch BD, Eglén RM, Hunter JC, Sangameswaran L (1998) Functional analysis of a voltage-gated sodium channel and its splice variant from rat dorsal root ganglia. *Journal of neurochemistry* 70(6):2262–2272
- Thimmapaya R, Neelands T, Niforatos W, Davis-Taber RA, Choi W, Putman CB, Kroeger PE, Packer J, Gopalakrishnan M, Faltynek CR, Surowy CS, Scott VE (2005) Distribution and functional characterization of human Nav1.3 splice variants. *The European journal of neuroscience* 22(1):1–9. doi:10.1111/j.1460-9568.2005.04155.x
- Chatelier A, Dahllund L, Eriksson A, Krupp J, Chahine M (2008) Biophysical properties of human Na v1.7 splice variants and their regulation by protein kinase A. *Journal of neurophysiology* 99(5):2241–2250. doi:10.1152/jn.01350.2007

32. Gershon E, Weigl L, Lotan I, Schreibley W, Dascal N (1992) Protein kinase A reduces voltage-dependent Na⁺ current in *Xenopus* oocytes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 12(10):3743–3752
33. Li M, West JW, Lai Y, Scheuer T, Catterall WA (1992) Functional modulation of brain sodium channels by cAMP-dependent phosphorylation. *Neuron* 8(6):1151–1159
34. Murphy BJ, Rossie S, De Jongh KS, Catterall WA (1993) Identification of the sites of selective phosphorylation and dephosphorylation of the rat brain Na⁺ channel alpha subunit by cAMP-dependent protein kinase and phosphoprotein phosphatases. *The Journal of biological chemistry* 268(36):27355–27362
35. Smith RD, Goldin AL (1997) Phosphorylation at a single site in the rat brain sodium channel is necessary and sufficient for current reduction by protein kinase A. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 17(16):6086–6093
36. Cantrell AR, Smith RD, Goldin AL, Scheuer T, Catterall WA (1997) Dopaminergic modulation of sodium current in hippocampal neurons via cAMP-dependent phosphorylation of specific sites in the sodium channel alpha subunit. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 17(19):7330–7338
37. Yue C, Remy S, Su H, Beck H, Yaari Y (2005) Proximal persistent Na⁺ channels drive spike afterdepolarizations and associated bursting in adult CA1 pyramidal cells. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 25(42):9704–9720. doi:10.1523/JNEUROSCI.1621-05.2005
38. Chao TI, Alzheimer C (1995) Effects of phenytoin on the persistent Na⁺ current of mammalian CNS neurones. *Neuroreport* 6(13):1778–1780
39. Taverna S, Mantegazza M, Franceschetti S, Avanzini G (1998) Valproate selectively reduces the persistent fraction of Na⁺ current in neocortical neurons. *Epilepsy research* 32(1–2):304–308
40. Spadoni F, Hainsworth AH, Mercuri NB, Caputi L, Martella G, Lavaroni F, Bernardi G, Stefani A (2002) Lamotrigine derivatives and riluzole inhibit INa, P in cortical neurons. *Neuroreport* 13(9):1167–1170
41. Ganetzky B, Wu CF (1982) *Drosophila* mutants with opposing effects on nerve excitability: genetic and spatial interactions in repetitive firing. *Journal of neurophysiology* 47(3):501–514
42. Lee J, Wu CF (2002) Electroconvulsive seizure behavior in *Drosophila*: analysis of the physiological repertoire underlying a stereotyped action pattern in bang-sensitive mutants. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 22(24):11065–11079
43. Kuebler D, Tanouye M (2002) Anticonvulsant valproate reduces seizure-susceptibility in mutant *Drosophila*. *Brain research* 958(1):36–42
44. Reynolds ER, Stauffer EA, Feeney L, Rojahn E, Jacobs B, McKeever C (2004) Treatment with the antiepileptic drugs phenytoin and gabapentin ameliorates seizure and paralysis of *Drosophila* bang-sensitive mutants. *Journal of neurobiology* 58(4):503–513. doi:10.1002/neu.10297
45. Tan JS, Lin F, Tanouye MA (2004) Potassium bromide, an anticonvulsant, is effective at alleviating seizures in the *Drosophila* bang-sensitive mutant bang senseless. *Brain research* 1020(1–2):45–52. doi:10.1016/j.brainres.2004.05.111
46. Stilwell GE, Saraswati S, Littleton JT, Chouinard SW (2006) Development of a *Drosophila* seizure model for in vivo high-throughput drug screening. *The European journal of neuroscience* 24(8):2211–2222. doi:10.1111/j.1460-9568.2006.05075.x
47. Song J, Tanouye MA (2008) From bench to drug: human seizure modeling using *Drosophila*. *Progress in neurobiology* 84(2):182–191. doi:10.1016/j.pneurobio.2007.10.006
48. Marley R, Baines RA (2011) Increased persistent Na⁺ current contributes to seizure in the slamdance bang-sensitive *Drosophila* mutant. *Journal of neurophysiology* 106(1):18–29. doi:10.1152/jn.00808.2010
49. Parker L, Padilla M, Du Y, Dong K, Tanouye MA (2011) *Drosophila* as a model for epilepsy: bss is a gain-of-function mutation in the para sodium channel gene that leads to seizures. *Genetics* 187(2):523–534. doi:10.1534/genetics.110.123299
50. White HS (2003) Preclinical development of antiepileptic drugs: past, present, and future directions. *Epilepsia* 44(Suppl 7):2–8
51. Constanti A (1978) The “mixed” effect of picrotoxin on the GABA dose/conductance relation recorded from lobster muscle. *Neuropharmacology* 17(3):159–167
52. Jones-Davis DM, Macdonald RL (2003) GABA(A) receptor function and pharmacology in epilepsy and status epilepticus. *Current opinion in pharmacology* 3(1):12–18
53. Reynolds EH (1987) Early treatment and prognosis of epilepsy. *Epilepsia* 28(2):97–106
54. Chadwick DW (2008) The treatment of the first seizure: the benefits. *Epilepsia* 49(Suppl 1):26–28. doi:10.1111/j.1528-1167.2008.01446.x
55. Plummer NW, Galt J, Jones JM, Burgess DL, Sprunger LK, Kohrman DC, Meisler MH (1998) Exon organization, coding sequence, physical mapping, and polymorphic intragenic markers for the human neuronal sodium channel gene SCN8A. *Genomics* 54(2):287–296. doi:10.1006/geno.1998.5550
56. Sarao R, Gupta SK, Auld VJ, Dunn RJ (1991) Developmentally regulated alternative RNA splicing of rat brain sodium channel mRNAs. *Nucleic acids research* 19(20):5673–5679
57. Yarowsky PJ, Krueger BK, Olson CE, Clevinger EC, Koos RD (1991) Brain and heart sodium channel subtype mRNA expression in rat cerebral cortex. *Proc Natl Acad Sci U S A* 88(21):9453–9457
58. Gustafson TA, Clevinger EC, O’Neill TJ, Yarowsky PJ, Krueger BK (1993) Mutually exclusive exon splicing of type III brain sodium channel alpha subunit RNA generates developmentally regulated isoforms in rat brain. *The Journal of biological chemistry* 268(25):18648–18653
59. Raymond CK, Castle J, Garrett-Engle P, Armour CD, Kan Z, Tsinoremas N, Johnson JM (2004) Expression of alternatively spliced sodium channel alpha-subunit genes. Unique splicing patterns are observed in dorsal root ganglia. *The Journal of biological chemistry* 279(44):46234–46241. doi:10.1074/jbc.M406387200
60. Kasai N, Fukushima K, Ueki Y, Prasad S, Nosakowski J, Sugata K, Sugata A, Nishizaki K, Meyer NC, Smith RJ (2001) Genomic structures of SCN2A and SCN3A—candidate genes for deafness at the DFNA16 locus. *Gene* 264(1):113–122
61. Fletcher EV, Kullmann DM, Schorge S (2011) Alternative splicing modulates inactivation of type 1 voltage-gated sodium channels by toggling an amino acid in the first S3–S4 linker. *The Journal of biological chemistry* 286(42):36700–36708. doi:10.1074/jbc.M111.250225
62. Aronica E, Yankaya B, Troost D, van Vliet EA, Lopes da Silva FH, Gorter JA (2001) Induction of neonatal sodium channel II and III alpha-isoform mRNAs in neurons and microglia after status epilepticus in the rat hippocampus. *The European journal of neuroscience* 13(6):1261–1266
63. Gastaldi M, Bartolomei F, Massacrier A, Planells R, Robaglia-Schlupp A, Cau P (1997) Increase in mRNAs encoding neonatal II and III sodium channel alpha-isoforms during kainate-induced seizures in adult rat hippocampus. *Brain research Molecular brain research* 44(2):179–190
64. Park JW, Parisky K, Celotto AM, Reenan RA, Graveley BR (2004) Identification of alternative splicing regulators by RNA interference in *Drosophila*. *Proc Natl Acad Sci U S A* 101(45):15974–15979. doi:10.1073/pnas.0407004101
65. Siomi H, Matunis MJ, Michael WM, Dreyfuss G (1993) The pre-mRNA binding K protein contains a novel evolutionarily conserved motif. *Nucleic acids research* 21(5):1193–1198

66. Grishin NV (2001) KH domain: one motif, two folds. *Nucleic acids research* 29(3):638–643
67. Ule J, Jensen KB, Ruggiu M, Mele A, Ule A, Darnell RB (2003) CLIP identifies Nova-regulated RNA networks in the brain. *Science* 302(5648):1212–1215. doi:10.1126/science.1090095
68. Ule J, Stefani G, Mele A, Ruggiu M, Wang X, Taneri B, Gaasterland T, Blencowe BJ, Darnell RB (2006) An RNA map predicting Nova-dependent splicing regulation. *Nature* 444(7119):580–586. doi:10.1038/nature05304
69. Buckanovich RJ, Posner JB, Darnell RB (1993) Nova, the paraneoplastic Ri antigen, is homologous to an RNA-binding protein and is specifically expressed in the developing motor system. *Neuron* 11(4):657–672
70. Yang YY, Yin GL, Darnell RB (1998) The neuronal RNA-binding protein Nova-2 is implicated as the autoantigen targeted in POMA patients with dementia. *Proc Natl Acad Sci U S A* 95(22):13254–13259
71. Seshiah P, Miller B, Myat MM, Andrew DJ (2001) pasilla, the Drosophila homologue of the human Nova-1 and Nova-2 proteins, is required for normal secretion in the salivary gland. *Developmental biology* 239(2):309–322. doi:10.1006/dbio.2001.0429
72. Racca C, Gardiol A, Eom T, Ule J, Triller A, Darnell RB (2010) The neuronal splicing factor Nova co-localizes with target RNAs in the dendrite. *Frontiers in neural circuits* 4:5. doi:10.3389/neuro.04.005.2010
73. Eom T, Zhang C, Wang H, Lay K, Fak J, Noebels JL, Darnell RB (2013) NOVA-dependent regulation of cryptic NMD exons controls synaptic protein levels after seizure. *eLife* 2:e00178. doi:10.7554/eLife.00178
74. Ule J, Ule A, Spencer J, Williams A, Hu JS, Cline M, Wang H, Clark T, Fraser C, Ruggiu M, Zeeberg BR, Kane D, Weinstein JN, Blume J, Darnell RB (2005) Nova regulates brain-specific splicing to shape the synapse. *Nature genetics* 37(8):844–852. doi:10.1038/ng1610
75. Heinzen EL, Yoon W, Tate SK, Sen A, Wood NW, Sisodiya SM, Goldstein DB (2007) Nova2 interacts with a cis-acting polymorphism to influence the proportions of drug-responsive splice variants of SCN1A. *American journal of human genetics* 80(5):876–883. doi:10.1086/516650
76. Zhang W, Linden DJ (2003) The other side of the engram: experience-driven changes in neuronal intrinsic excitability. *Nature reviews Neuroscience* 4(11):885–900. doi:10.1038/nrn1248
77. Baines RA, Uhler JP, Thompson A, Sweeney ST, Bate M (2001) Altered electrical properties in Drosophila neurons developing without synaptic transmission. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 21(5):1523–1531
78. Muraro NI, Weston AJ, Gerber AP, Luschnig S, Moffat KG, Baines RA (2008) Pumilio binds para mRNA and requires Nanos and Brat to regulate sodium current in Drosophila motoneurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 28(9):2099–2109. doi:10.1523/JNEUROSCI.5092-07.2008
79. Driscoll HE, Muraro NI, He M, Baines RA (2013) Pumilio-2 regulates translation of Nav1.6 to mediate homeostasis of membrane excitability. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 33(23):9644–9654. doi:10.1523/JNEUROSCI.0921-13.2013
80. Schweers BA, Walters KJ, Stern M (2002) The Drosophila melanogaster translational repressor pumilio regulates neuronal excitability. *Genetics* 161(3):1177–1185
81. Zamore PD, Williamson JR, Lehmann R (1997) The Pumilio protein binds RNA through a conserved domain that defines a new class of RNA-binding proteins. *Rna* 3(12):1421–1433
82. Wickens M, Bernstein DS, Kimble J, Parker R (2002) A PUF family portrait: 3'UTR regulation as a way of life. *Trends in genetics : TIG* 18(3):150–157
83. Spassov DS, Jurecic R (2003) The PUF family of RNA-binding proteins: does evolutionarily conserved structure equal conserved function? *IUBMB life* 55(7):359–366. doi:10.1080/15216540310001603093
84. Vessey JP, Vaccani A, Xie Y, Dahm R, Karra D, Kiebler MA, Macchi P (2006) Dendritic localization of the translational repressor pumilio 2 and its contribution to dendritic stress granules. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 26(24):6496–6508. doi:10.1523/JNEUROSCI.0649-06.2006
85. Murata Y, Wharton RP (1995) Binding of pumilio to maternal hunchback mRNA is required for posterior patterning in Drosophila embryos. *Cell* 80(5):747–756
86. Sonoda J, Wharton RP (2001) Drosophila brain tumor is a translational repressor. *Genes & development* 15(6):762–773. doi:10.1101/gad.870801
87. Chagnovich D, Lehmann R (2001) Poly(A)-independent regulation of maternal hunchback translation in the Drosophila embryo. *Proc Natl Acad Sci U S A* 98(20):11359–11364. doi:10.1073/pnas.201284398
88. Gerber AP, Luschnig S, Krasnow MA, Brown PO, Herschlag D (2006) Genome-wide identification of mRNAs associated with the translational regulator PUMILIO in Drosophila melanogaster. *Proc Natl Acad Sci U S A* 103(12):4487–4492. doi:10.1073/pnas.0509260103
89. Ye B, Petritsch C, Clark IE, Gavis ER, Jan LY, Jan YN (2004) Nanos and pumilio are essential for dendrite morphogenesis in Drosophila peripheral neurons. *Current biology : CB* 14(4):314–321. doi:10.1016/j.cub.2004.01.052
90. Menon KP, Sanyal S, Habara Y, Sanchez R, Wharton RP, Ramaswami M, Zinn K (2004) The translational repressor pumilio regulates presynaptic morphology and controls postsynaptic accumulation of translation factor eIF-4E. *Neuron* 44(4):663–676. doi:10.1016/j.neuron.2004.10.028
91. Menon KP, Andrews S, Murthy M, Gavis ER, Zinn K (2009) The translational repressors Nanos and pumilio have divergent effects on presynaptic terminal growth and postsynaptic glutamate receptor subunit composition. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 29(17):5558–5572. doi:10.1523/JNEUROSCI.0520-09.2009
92. Dubnau J, Chiang AS, Grady L, Barditch J, Gossweiler S, McNeil J, Smith P, Buldoc F, Scott R, Certa U, Broger C, Tully T (2003) The staufer/pumilio pathway is involved in Drosophila long-term memory. *Current biology : CB* 13(4):286–296
93. Siemen H, Colas D, Heller HC, Brustle O, Pera RA (2011) Pumilio-2 function in the mouse nervous system. *PloS one* 6(10):e25932. doi:10.1371/journal.pone.0025932
94. Sun W, Wagnon JL, Mahaffey CL, Briese M, Ule J, Frankel WN (2013) Aberrant sodium channel activity in the complex seizure disorder of Celf4 mutant mice. *The Journal of physiology* 591(Pt 1):241–255. doi:10.1113/jphysiol.2012.240168
95. Chatelier A, Zhao J, Bois P, Chahine M (2010) Biophysical characterisation of the persistent sodium current of the Nav1.6 neuronal sodium channel: a single-channel analysis. *Pflugers Archiv: European journal of physiology* 460(1):77–86. doi:10.1007/s00424-010-0801-9
96. Wagnon JL, Mahaffey CL, Sun W, Yang Y, Chao HT, Frankel WN (2011) Etiology of a genetically complex seizure disorder in Celf4 mutant mice. *Genes brain and behavior* 10(7):765–777. doi:10.1111/j.1601-183x.2011.00717.x
97. Vacher H, Trimmer JS (2012) Trafficking mechanisms underlying neuronal voltage-gated ion channel localization at the axon initial segment. *Epilepsia* 53(Suppl 9):21–31. doi:10.1111/epi.12032
98. Shao D, Okuse K, Djamgoz MB (2009) Protein-protein interactions involving voltage-gated sodium channels: post-translational regulation, intracellular trafficking and functional expression. *The*

- international journal of biochemistry & cell biology 41(7):1471–1481. doi:[10.1016/j.biocel.2009.01.016](https://doi.org/10.1016/j.biocel.2009.01.016)
99. Ekberg JA, Boase NA, Rychkov G, Manning J, Poronnik P, Kumar S (2014) Nedd4-2 (NEDD4L) controls intracellular Na⁺-mediated activity of voltage-gated sodium channels in primary cortical neurons. *The Biochemical journal* 457(1):27–31. doi:[10.1042/BJ20131275](https://doi.org/10.1042/BJ20131275)
100. Laedermann CJ, Cachemaille M, Kirschmann G, Pertin M, Gosselin RD, Chang I, Albesa M, Towne C, Schneider BL, Kellenberger S, Abriel H, Decosterd I (2013) Dysregulation of voltage-gated sodium channels by ubiquitin ligase NEDD4-2 in neuropathic pain. *The Journal of clinical investigation* 123(7):3002–3013. doi:[10.1172/JCI68996](https://doi.org/10.1172/JCI68996)
101. Sun L, Gilligan J, Staber C, Schutte RJ, Nguyen V, O'Dowd DK, Reenan R (2012) A knock-in model of human epilepsy in *Drosophila* reveals a novel cellular mechanism associated with heat-induced seizure. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32(41):14145–14155. doi:[10.1523/JNEUROSCI.2932-12.2012](https://doi.org/10.1523/JNEUROSCI.2932-12.2012)
102. Brigo F, Ausserer H, Tezzon F, Nardone R (2013) When one plus one makes three: the quest for rational antiepileptic polytherapy with supraadditive anticonvulsant efficacy. *Epilepsy & behavior : E&B* 27(3):439–442. doi:[10.1016/j.yebeh.2013.03.010](https://doi.org/10.1016/j.yebeh.2013.03.010)