

Technical University of Denmark



Oxidative stability of krill oil (*Euphausia superba*)

Lu, Henna Fung Sieng; Jacobsen, Charlotte; Bruheim, I.; Griinari, M.; Oterhals, A.; Haugsjerd, B. O.; Vogt, G.

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Lu, H. F. S., Jacobsen, C., Bruheim, I., Griinari, M., Oterhals, A., Haugsjerd, B. O., & Vogt, G. (2013). Oxidative stability of krill oil (*Euphausia superba*). Abstract from 11th Euro Fed Lipid Congress, Antalya, Turkey.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Oxidative stability of krill oil (*Euphasia superba*)

F. S. H. Lu¹, C. Jacobsen¹, I. Bruheim², M. Griinari³,
A. Oterhals⁴, B. O. Haugsjerd⁵, G. Vogt⁶

¹Division of Industrial Food Research, Lipids and Oxidation Group, National Food Institute, Technical University of Denmark, Søtofts Plads, Building 221, 2800 Kgs, Lyngby, Denmark

²Olympic Seafood AS, P.O. Box 234, N-6099 Fosnavaag, Norway

³Clanet Ltd., Lotankatu 1, 02680 Espoo, Finland

⁴Nofima, Division Fisheries, Industry and Market, Marine Biotechnology Group, Kjerreidviken 16, 5141 Fyllingsdalen, Norway

⁵Nofima, Division Aquaculture, BioLab, Kjerreidviken 16, 5141 Fyllingsdalen, Norway

⁶Nofima, Division Food, Food and Health Group, Osloveien 1, 1430 Ås, Norway

Krill oil has been reported in many studies to have high oxidative stability when evaluated by peroxide value (PV) and anisidine value (AV). However, recent studies have shown that other compounds than primary and secondary oxidation products are formed when krill oil is exposed to oxidative conditions. These compounds include Strecker degradation compounds and pyrroles. Some of these compounds may have antioxidative effect. Commercial scale processing of krill prior to extraction may affect the oxidative stability of krill oil. Therefore, the main objective of this study was to compare lipid oxidation in krill oil produced in a commercial process and krill oil carefully extracted from frozen krill in the laboratory. Krill oil was incubated at different temperatures (20, 30 and 40 °C) for 1, 2, 3, 4 and 6 weeks, under conditions of constant stirring while being exposed to air. The oxidative stability of both oils was assessed by simple chemical analyses such as PV, AV, spectrophotometric determination at 400 nm, and also advanced determination methods such as a) measurement of secondary volatile oxidation products and Strecker degradation products by DHS-GC/MS, b) measurement of lipid classes especially phosphatidylcholine and phosphatidylethanolamine, c) measurement of antioxidants level such as tocopherols, rosemary, ascorbyl palmitate and astaxanthin upon storage. In addition, the effect of residues or impurities present in krill oil contributing to its oxidative stability were investigated by d) the determination of amino acids composition and transition metals such as Cu²⁺ and Fe²⁺.