

Microencapsulated sodium selenite supplementation in dairy cows: effects on selenium status

E. Grilli^{1†}, A. Gallo², M. Fustini¹, P. Fantinati³ and A. Piva¹

¹Dipartimento di Scienze Mediche Veterinarie, Università di Bologna, Bologna 40064, Italy; ²ISAN, Università Cattolica Sacro Cuore, Piacenza 29122, Italy;

³Vetagro SpA, Reggio Emilia 42124, Italy

(Received 5 October 2012; Accepted 22 July 2013; First published online 10 September 2013)

The objective of this study was to compare the efficiency of transfer of selenium (Se) to plasma and milk from inorganic sodium selenite, either free or microencapsulated, and from selenized yeast in dairy cows. The study consisted of an in situ-nylon bags incubation, and in an in vivo experiment to compare the Se status of cows supplemented with either sodium selenite, microencapsulated sodium selenite, or Se yeast. Thirty dairy cows, divided in five groups, were fed the following diets: the control group (CTR) received a total mixed ration supplemented with sodium selenite in order to have 0.3 mg/kg DM of total Se; 0.3M and 0.5M groups received the same control diet supplemented with lipid microencapsulated sodium selenite to provide 0.3 and 0.5 mg/kg DM of total Se, respectively; 0.3Y and 0.5Y groups received selenized yeast to provide 0.3 and 0.5 mg/kg of total Se, respectively. Cows were fed the supplements for 56 days during which milk, blood, and fecal samples were collected weekly to conduct analysis of Se and glutathione peroxidase (GSH-px) activity. Se concentration in the nylon bags was assessed to 72%, 64%, and 40% of the initial value (time 0) after 4, 8, and 24 h of incubation, respectively. In vivo, cows supplemented with 0.3 mg/kg of microencapsulated Se had higher milk Se concentration compared to CTR. The increment was more pronounced at the highest inclusion rate (0.5 mg/kg, 0.5M group). GSH-px activity was not significantly affected by treatments. The results indicate that lipid microencapsulation has the potential to protect nutrients from complete rumen reduction and that Se from microencapsulated selenite is incorporated in milk more efficiently than the free form. Microencapsulated sodium selenite was shown to be comparable to Se-yeast in terms of availability and incorporation in milk when fed at 0.3 mg/kg DM, whereas the inclusion in the diet at 0.5 mg/kg DM resulted in higher plasma and milk concentrations than selenized yeast.

Keywords: dairy cow, microencapsulation, selenium

Implications

Inorganic selenium (Se) supplements, that is selenate and selenite, have a reduced intestinal availability in ruminants, whereas organic Se, supplemented in the form of selenized yeast, has been demonstrated to be more efficiently stored in tissues and biological fluids like milk and blood than sodium selenite. In this study we evaluated lipid microencapsulated sodium selenite as an alternative source of Se in comparison with both sodium selenite and selenized yeast by assessing Se status in dairy cows. The results indicate that microencapsulation has the potential to improve the passage of non-reduced Se to the intestine and to increase Se concentration in milk.

Introduction

Selenium (Se) is an essential trace element crucial for the homeostasis of the oxidative status of animals. It plays an

important role as a cofactor of glutathione peroxidase (GSH-px) and of other enzymes of the antioxidant system (Rotruck *et al.*, 1973; Underwood and Suttle, 1999); Se contributes also to regulate the synthesis of thyroid hormones (Berry *et al.*, 1991), and improves the immune system function with particular reference to the cell-mediated immunity (Grasso *et al.*, 1990); in cows, it also reduces the prevalence and severity of mastitis (Smith *et al.*, 1984). Severe Se deficiencies affect fertility and reproduction, growth, and sanitary status (Weiss *et al.*, 1990; Underwood and Suttle, 1999).

In Europe, Se is authorized at a maximum of 0.5 mg/kg DM (Council of the European Communities, 1970), whereas in the US, the Federal and Drug Administration authorized Se supplementation in feeds at a maximum of 0.3 mg/kg DM (Food and Drug Administration, 1997).

Dietary sources of Se can be organic, such as Se-methionine or cysteine, or inorganic, such as sodium selenite or sodium selenate. Inorganic Se is one of the most widespread forms of Se supplementation in feeds, but the ability of ruminants to use

[†] E-mail: ester.grilli@unibo.it

inorganic Se, as well as other nutrients, is strongly affected by the highly reducing environment of the rumen. In fact, the rumen reduces selenate to selenite, which is, in turn, converted into low molecular weight insoluble forms, most likely non-absorbable (Wright and Bell, 1966; Weiss, 2005). Inorganic Se absorption coefficient in dairy cows is estimated to be 50% (Weiss, 2005), whereas organic Se, deriving from forages or grains in which it is incorporated in Se-methionine and Se-cysteine as a substitute of sulfur, is estimated to be 65%.

For this reason, the use of organic supplements, in which Se is incorporated in yeasts during fermentation, has recently become popular.

Lipid microencapsulation is an effective tool to prevent complete absorption of nutrients and additives in the duodenum and jejunum of monogastric animals (Piva *et al.*, 2007), as well as to protect nutrients from ruminal degradation (Gallo *et al.*, 2010).

Given the importance of increasing absorbable Se flow at intestinal level, the aim of this study was to compare the efficiency of transfer of Se to plasma and milk from inorganic sodium selenite, either free or microencapsulated, and from selenized yeast in dairy cows.

Material and methods

In situ nylon bag incubation of microencapsulated sodium selenite

Feed additive. Sodium selenite (SmartSel[®], Vetagro Spa, Reggio Emilia, Italy) was microencapsulated through a spray-chilling with hydrogenated fat embedding matrix method. The particle size of the final product was distributed as follows: 4.2% > 2000 µm, 1500 < 27.4% < 2000 µm, 1000 < 39.9% < 1500 µm, 500 < 26.4% < 1000 µm, and 2.1% < 500 µm.

Animals and diets. Two Holstein cows were fitted with rumen fistula and fed a total mixed ration (TMR) containing 40% grass hay, 20% corn silage, and 40% concentrates according to National Research Council (NRC) requirements (2001). Animals were fed twice per day.

In situ nylon bag incubation. The incubation was performed according to De Boer *et al.* (1987). Briefly, nylon bags (10 × 15 cm; pore size 45–60 µm) were filled with 3 g of lipid microencapsulated sodium selenite (providing 10 g/kg of Se according to company specifications), sealed, and incubated ventrally in the rumen for 0, 4, 8, and 24 h. Each sample incubation was repeated twice.

Nylon bags were incubated in the rumen in a polyester mesh bag (25 × 40 cm; 3 mm pore size) and were removed from the rumen at the same time to allow simultaneous washing. Once removed, samples were mechanically washed with running water at room temperature. Bags at time 0 were not incubated but just mechanically washed in order to estimate the initial amount of Se in the product. Washed samples were then freeze-dried for 60 h and the weight of residues was determined on the lyophilized sample.

Se analysis and calculations. The determination of Se was performed through inductively coupled plasma-mass spectrometry ICP-MS (Agilent 7500ce, CA, USA) according to Moschini *et al.* (2010).

In vivo study

Animals and diets. The experiment was conducted between 24 October 2010 and 29 December 2010, at the Centro di Ricerca per la Zootecnia e l'Ambiente (CERZOO; San Bonico, Piacenza, Italy), which is a Good Laboratory Practice-certified facility, and operates according to the procedure of animal protection and welfare (directive No 86/609/EEC).

Thirty Holstein lactating cows were selected from the herd and randomly divided in five groups with similar average days in milking (161.6 ± 113.8) and number of lactations (3.0 ± 1.7). At the end of 10 days of pre-experimental adaptation, animals were randomly assigned to one of the dietary treatments which consisted of a basal TMR (0.18 mg/kg DM Se) differing only for Se supplementation source and concentration: the control group (CTR) diet was supplemented with 0.12 mg/kg sodium selenite in order to have 0.3 mg/kg DM of total Se (CTR), whereas treated groups diets were supplemented with microencapsulated sodium selenite (0.3M and 0.5M; SmartSel[®], Vetagro SpA, RE, Italy) or selenized yeast (0.3Y and 0.5Y; ALKOSEL[®] R397, Lallemand Animal Nutrition, Blagnac, France) in order to have a final concentration of 0.3 and 0.5 mg/kg DM of total Se. Animals were fed for 56 days a TMR containing 54% of corn silage, 11.5% and 5% of corn meal and flaked corn, respectively, 13% of a mixture of soybean cotton and gluten meal, and vitamin and mineral premix; 12.3% of dehydrated alfalfa and 4.2% of rye grass hay. The TMR was formulated according to NRC requirements (National Research Council (NRC), 2001). Ingredients were mixed in a horizontal mixer without the supplemental Se and administered to cows. Then, the different Se sources (i.e. sodium selenite, microencapsulated Se from sodium selenite, or Se enriched yeasts) were premixed with 1 kg of corn meal as a carrier and subsequently added directly to the TMR. Weighed experimental diets were individually offered twice daily and refusals were removed before the fresh TMR was offered. DM intake was calculated by subtracting refusals to the amount of TMR offered.

Cows were milked twice daily at 0500 and 1500 h and the herringbone milking parlor was equipped to record individual cow milk yield (Afimilk, S.A.E. AFIKIM, Kibbutz Afikim, Israel).

Feed sampling and analyses. TMR and feedstuffs samples were collected weekly throughout the study (from day 10 to day 56) and kept frozen (–20°C). At the end of the study, samples from each week were pooled in one sample and analyzed for the nutritional characteristics according the following methods: DM was determined gravimetrically drying the sample at 103°C to a constant weight (International Standards Organization (ISO), 6496:1999), CP, aNDF, and ADF were determined according to International Standards Organization (2006, 2009) method, and International Standards Organization (2008), respectively.

Starch was determined by polarimetric method according to International Standards Organization (1997).

Feedstuffs included in the TMR were also analyzed for Se content according to Calamari *et al.* (2010). The basal Se content was taken as a reference to calculate the supplemental Se to be added to the experimental diets.

Blood, milk, and feces sampling and analyses. Individual blood samples were collected at the beginning of the study (day 0), and weekly thereafter by jugular venipuncture using 10 cc heparinized (lithium heparin) disposable syringes (Becton Dickinson, Franklin Lakes, NJ, USA). Blood was centrifuged for 10 min at 3000 r.p.m.; plasma was transferred with Pasteur pipette in a plastic test tube and frozen before analysis. Plasma Se and plasma GSH-px were analyzed according to Calamari *et al.* (2010).

Individual milk samples were collected at the beginning of the study (day 0) and weekly thereafter. The two milk samples collected daily were pooled proportionally to the yield of each lactation and split into two aliquots. One was immediately analyzed for protein, fat, and lactose by MilkoScan (FT6000 Foss Electric, Hillerod, Denmark), whereas the other was frozen for Se analysis that was subsequently performed according to Calamari *et al.* (2010).

Fecal samples were collected at the beginning of the study (day 0), at day 14, day 28, day 42, and at the end of the study (day 56), frozen, and analyzed for Se content.

Table 1 Diet composition and nutrient analysis

Nutrient	% DM
CP	14.6
Ether extract	3.72
Starch	27.2
aNDF	38.6
ADF	24.1
ADL	3.43
Se (mg/kg DM)	0.18
Net energy of lactation (Mcal/kg DM) ^a	1.70

^aCalculated according to NRC (2001).

Statistical analysis. Data measured over time were subjected to ANOVA using the repeated measures in the mixed procedure of SAS (SAS Institute Inc., Cary, NC, USA; release 9.1, 2002–2003) in a completely randomized design. The statistical model included the fixed effect of treatment, time of measurement and (treatment × time) interaction. The random variable was the cow within treatment. Each variable analyzed was subjected to three covariance structures: autoregressive order AR (1), compound symmetry, and spatial power. Using the Akaike information criterion and the Schwarz Bayesian criterion, the compound symmetry was the covariance structure that fitted the model best for all considered variables.

Orthogonal contrasts were used to compare the supplementation effect (CTR v. 0.3M and 0.5M; CTR v. 0.3Y and 0.5Y) and a linear dose effect of Se supplementation was evaluated on microencapsulated Se (0.3M and 0.5M) and selenized yeast (0.3Y and 0.5Y). Differences were considered significant at $P < 0.05$.

Results and discussion

In situ nylon bag incubation of microencapsulated sodium selenite

The *in situ* nylon bag experiment was performed to assess the amount of Se in the lipid matrix throughout 24 h of incubation in the rumen. Se from sodium selenite in the microencapsulated product at time 0 was 8.36 g/kg and gradually decreased over time.

In particular, Se concentration in the nylon bags gradually decreased to 72%, 64%, and 40% of the initial value (time 0) after 4, 8, and 24 h of incubation, respectively. The limitation of nylon bag technique (i.e. the lack of a non-encapsulated control, due to selenite small particle size and high water solubility), and the inability of inductively coupled mass spectrometry to detect the form of Se, makes it impossible to make conclusions about rumen-protection of selenite. Nevertheless, assuming that all of the selenite entrapped in the matrix remained unmodified throughout the incubation and considering an average permanence-transit time in the rumen of 8 h, we would have an extent of rumen protection $\geq 64\%$.

Table 2 Effect of Se source supplementation in plasma, milk and fecal output of Se and plasma GSH-px concentration in dairy cows

	Treatments ²						Contrasts ¹			
							Treatment effect		Linear dose effect	
	CTR	0.3M	0.3Y	0.5M	0.5Y	s.e.	CTR v. 0.3M and 0.5M	CTR v. 0.3Y and 0.5Y	0.3M and 0.5M	0.3Y and 0.5Y
Plasma Se (ng/g)	82.75	88.24	83.41	99.48	90.51	2.99	***	ns	**	*
Plasma GSH-px (U/l)	46.78	50.54	49.43	48.10	49.66	2.51	ns	ns	ns	ns
Milk Se (ng/g)	19.05	26.32	23.27	41.49	26.70	2.60	***	*	**	ns
Feces Se (ng/g)	404	431	540	539	666	34	**	***	**	***

CTR = control group; GSH-px = glutathione peroxidase; ns = non-significant.

¹Contrasts significance of effects of treatment (Se source) and level of inclusion (0.3 and 0.5 mg/kg).

²Treatments: CTR, sodium selenite providing Se at 0.3 mg/kg DM; 0.3M and 0.5M, microencapsulated sodium selenite providing Se at 0.3 mg/kg and 0.5 mg/kg DM, respectively; 0.3Y and 0.5Y, selenized yeast providing Se at 0.3 mg/kg and 0.5 mg/kg DM, respectively.

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

In vivo study

Feed analyses and Se content. Nutritional values and the mean Se concentration of the basal diet are given in Table 1. Net energy of lactation of the TMR was calculated according to NRC (2001) considering the inclusion rates of ingredients in the diet.

Feed intake, milk production and Se concentration in blood, milk, and feces. Animals were in a good health status throughout the experiment and the average DM intake (24.2 ± 2.4 kg/head per day) was similar among experimental groups. Milk yield (average of 34 kg/head per day) and milk parameters were unaffected by the source or the inclusion level of Se (data not shown).

Plasma Se was generally not affected by the source of Se when fed at the lowest dose (0.3 mg/kg) even if there was a numerical increase in 0.3M group compared with CTR and 0.3Y (+6.6% and +5.8%, respectively; Table 2). Serum or plasma Se concentration are highly correlated to Se intake and for this reason they are considered a quick and effective tool to estimate Se blood status; serum or plasma Se should respond more quickly to changes in dietary intake than whole blood Se, which is, on the other side, more indicative of the historical nutritional status of the cow (Waldner *et al.*, 1998). Se concentrations indicative of a good Se status should be above 0.18 $\mu\text{g/ml}$ in whole blood or 0.08 $\mu\text{g/ml}$ in plasma, and supplemental Se at 0.3 mg/kg diet is usually enough to obtain those levels in lactating cows (Weiss *et al.*, 1990; Smith *et al.*, 1997; NRC, 2001). Cows entering this study had Se plasma concentration of 0.078 ± 0.0064 $\mu\text{g/g}$, and plasma Se increased over time, regardless of the treatment (day effect, $P < 0.05$; Table 2; Figure 1b). When Se intakes were the highest, as in 0.5 mg/kg fed groups, plasma Se responded better to microencapsulated supplementation than to selenized yeast. In fact, 0.5M group had 9.9% higher plasma Se than 0.5Y ($P < 0.05$; Table 2); the increment in plasma levels at steady state since the beginning of the study was almost double in 0.5M animals than in 0.5Y ones (29.5% and 17.8%, respectively).

Milk Se concentration was affected by both Se source and dose: in particular, cows fed microencapsulated Se at 0.3 mg/kg had 38% higher milk Se than control animals, whereas yeast-supplemented cows did not differ from control or 0.3M groups (Table 2); when Se was fed at 0.5 mg/kg, 0.5M cows had 55% higher Se concentration than 0.5Y ones ($P < 0.01$; Table 2; Figure 1a). If the ratio between milk and plasma Se concentration is indicative of the transfer of Se from blood to milk, as suggested by Calamari *et al.* (2010), microencapsulated selenite at 0.5 mg/kg was more efficient than the selenized yeast as the ratio was almost 38% higher ($P < 0.01$; Table 3). Within the 0.3 mg/kg dose there were no differences among groups (Table 3) although a slightly higher numerical trend was observed for 0.3M group.

When reporting Se excretion in milk to the ingested amount (mass of Se in milk over mass of ingested Se), the ratio averaged 8.6%, 10.9%, and 10.3% for CTR, 0.3M and 0.3Y group, respectively ($P < 0.001$; Table 3) and 10.5% and




Figure 1 Behavior of total Se in milk (a), plasma (b), and feces (c) in dairy cows that received diets containing either sodium selenite providing Se at 0.3 mg/kg DM (CTR) or microencapsulated sodium selenite providing Se at 0.3 mg/kg and 0.5 mg/kg DM (0.3M and 0.5M, respectively) and selenized yeast providing Se at 0.3 mg/kg and 0.5 mg/kg DM (0.3Y and 0.5Y, respectively). Values are least squares means, and vertical bars represent s.e. ($n = 6$).

7.1% for 0.5M, and 0.5Y, respectively ($P < 0.001$; Table 3). In this study, CTR group cows had an higher milk Se/ingested Se ratio compared with the 3.2% reported by Calamari *et al.* (2010). On the other hand, average Se secretion in milk of cows supplemented with selenized yeast is expected to be 10–18% depending on lactation stage, milk yield, protein yield, and Se intake (Heard *et al.*, 2007; Juniper *et al.*, 2008;

Table 3 Effect of Se source supplementation on Se milk excretion parameters

	Treatments ¹					s.e.	P-value
	CTR	0.3M	0.3Y	0.5M	0.5Y		
Se ingested (mg/day)	6.95	7.22	12.25	7.77	12.10	2.67	
Se in milk/Se ingested (%)	8.61 ^a	10.92 ^b	10.29 ^b	10.52 ^b	7.06 ^a	1.61	***
Se in milk/Se in plasma	0.23 ^a	0.29 ^a	0.24 ^a	0.40 ^b	0.29 ^b	0.06	**

Different superscripts in the same row significantly differ.

¹Treatments: CTR, sodium selenite providing Se at 0.3 mg/kg DM; 0.3M and 0.5M, microencapsulated sodium selenite providing Se at 0.3 mg/kg and 0.5 mg/kg DM, respectively; 0.3Y and 0.5Y, selenized yeast providing Se at 0.3 mg/kg and 0.5 mg/kg DM, respectively.

** $P < 0.01$, *** $P < 0.001$.

Calamari *et al.*, 2010; Walker *et al.*, 2010), whereas, in this study, 0.5Y cows had a surprisingly low milk Se concentration. Cows entering the study had a marginally adequate Se intake, with 60% of total Se (0.18 mg/kg out of 0.3 mg/kg) provided with the TMR. Therefore, the added amount, either from sodium selenite or Se yeast might have had a relatively minor impact on the total transfer. When Se was included at 0.5 mg/kg, instead, the Se supply from the TMR was relatively minor (36% of the total) and most of Se was provided with the additives, resulting in more appreciable differences for both milk and plasma concentrations.

Ortman and Pehrson (1999) observed a peak of 31 ng/g of Se in milk of cows supplemented with 0.3 mg/kg of Se from yeast and this peak was rapidly reached after one week since the beginning of the supplementation. In our experiment, we observed a peak at 23–27 ng/g after 1 week of supplementation with Se-yeast, with minor fluctuations over time (Figure 1a), without differences between low dose and high dose of Se-yeast. Fecal Se was 34% and 25% higher in 0.3Y compared with CTR and 0.3M, respectively, and 24% higher in 0.5Y than in 0.5M ($P < 0.01$, Table 2; Figure 1c). These data would suggest a poor Se retention in animals receiving the highest concentration of the yeast which is probably due to the source of Se itself.

GSH-px activity was not affected by the treatments (Table 2), though a numerical increase due to both microencapsulated Se and Se-yeasts was found (5–8% depending on the treatment). These results are aligned with those reviewed by Weiss (2005) and Ivancic and Weiss (2000), who reported a lack of increase of plasma GSH-px activity over a 42 days supplementation study with cows that had an adequate Se status at the beginning of the experiment. It seems in fact that the increase in GSH-px activity often described in literature might be more likely related to the amount of Se fed rather than to the source. In our study all of the animals rapidly reached a good Se status, as predicted by plasma concentration at 7 days, regardless of the treatment, and GSH-px activity remained fairly stable for the first 2 weeks of the experiment. The difference in GSH-px activity between inorganic and organic sources seems in fact to be the largest when cows are fed the lowest doses and are in a sub-clinical Se deficiency status (Knowles *et al.*, 1999; Weiss, 2005), whereas above the minimally adequate Se intake there is little correlation between plasma Se and GSH-px activity (Ullrey, 1987).

Conclusion

To our knowledge, this is the first study where microencapsulated sodium selenite was fed to cows in comparison with either free selenite or Se-yeast. Overall, the data provided with this study lead to the conclusion that lipid microencapsulation has the potential to protect nutrients from rumen degradability and theoretically increase the amount of rumen by-pass. In this study, the effectiveness of lipid microencapsulation is further substantiated by the results obtained *in vivo*. In fact, in cows fed with 0.3 mg/kg of Se as microencapsulated sodium selenite, Se was transferred to milk more efficiently than in cows fed the non-encapsulated form. Within the same Se dose, the microencapsulated form was comparable to Se-yeasts at 0.3 mg/kg, but when Se was included in the diet at 0.5 mg/kg, microencapsulated Se performed better than the Se-yeast.

Acknowledgments

The authors acknowledge the financial support of Vetagro for the study.

References

- Berry MJ, Banu L and Larsen PR 1991. Type I iodothyronine deiodinase is a selenocysteine-containing enzyme. *Nature* 349, 438–440.
- Calamari L, Petrera F and Bertin G 2010. Effects of either sodium selenite or Se yeast (Sc CNCM I-3060) supplementation on selenium status and milk characteristics in dairy cows. *Livestock Science* 128, 154–165.
- Council of the European Communities 1970. Council Directive 70/524/EEC concerning additives in feeding-stuffs. *Official Journal* 50, 1–17.
- De Boer G, Murphy JJ and Kennelly JJ 1987. Mobile nylon bag for estimating intestinal availability of rumen undegradable protein. *Journal of Dairy Science* 70, 977–982.
- Food and Drug Administration (FDA) 1997. Food additives permitted in feed and drinking water of animals; selenium. *Federal Register* 62, 44892–44894.
- Gallo A, Fusconi G, Fiorentini L, Grilli E, Fantinati P and Masoero F 2010. Blood methionine and lysine concentration in lactating dairy cows supplemented with commercial rumen-protected methionine and lysine products. In *Proceeding of the third International Symposium on Energy and Protein Metabolism and Nutrition (ISEP)*, 6–10 September 2010, Parma, Italy, p. 349.
- Grasso PJ, Scholz RW, Erskine RJ and Eberhart RJ 1990. Phagocytosis, bactericidal activity, and oxidative metabolism of mammary neutrophils from dairy cows fed selenium-adequate and selenium deficient diets. *American Journal of Veterinary Research* 51, 269.
- Heard JW, Stockdale CR, Walker GP, Leddin CM, Dunshea FR, McIntosh GH, Shields PM, McKenna A, Young GP and Doyle PT 2007. Increasing selenium

- concentration in milk: effects of amount of selenium from yeast and cereal grain supplements. *Journal of Dairy Science* 90, 4117–4127.
- International Standards Organization (ISO 10520:1997E) 1997. Native starch – determination of starch content – Ewers polarimetric method. ISO, Geneva, Switzerland.
- International Standards Organization (ISO 6496:1999) 1999. Animal feeding stuffs – determination of moisture and other volatile matter content. ISO, Geneva, Switzerland.
- International Standards Organization (ISO 16472:2006) 2006. Animal feeding stuffs – determination of amylase-treated neutral detergent fibre content (aNDF). ISO, Geneva, Switzerland.
- International Standards Organization (ISO 13906:2008) 2008. Animal feeding stuffs – determination of acid detergent fibre (ADF) and acid detergent lignin (ADL) contents. ISO, Geneva, Switzerland.
- International Standards Organization (ISO 5983:2009) 2009. Animal feeding stuffs – determination of nitrogen content and calculation of crude protein content. ISO, Geneva, Switzerland.
- Ivancic J and Weiss WP 2000. Effect of dietary sulfur and selenium concentrations on selenium balance of lactating holstein cows. *Journal of Dairy Science* 84, 225–232.
- Juniper DT, Phipps RH, Ramos-Morales E and Bertin G 2008. Effect of dietary supplementation with selenium-enriched yeast or sodium selenite on selenium tissue distribution and meat quality in beef cattle. *Journal of Animal Science* 86, 3100–3109.
- Knowles SO, Grace ND, Wurms K and Lee J 1999. Significance of amount and form of dietary selenium on blood, milk, and casein selenium concentrations in grazing cows. *Journal of Dairy Science* 82, 429–437.
- Moschini M, Battaglia M, Beone GM, Piva G and Masoero F 2010. Iodine and selenium carry over in milk and cheese in dairy cows: effect of diet supplementation and milk yield. *Animal* 4, 147–155.
- National Research Council (NRC) 2001. Nutrient requirements of dairy cattle, 7th revised edition. The National Academy Press, Washington, DC.
- Ortman K and Pehrson B 1999. Effect of selenate as a feed supplement to dairy cows in comparison to selenite and selenium yeast. *Journal of Animal Science* 77, 3365–3370.
- Piva A, Pizzamiglio V, Morlacchini M, Tedeschi M and Piva G 2007. Lipid microencapsulation allows slow release of organic acids and natural identical flavors along the swine intestine. *Journal of Animal Science* 85, 486–493.
- Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG and Hoekstra WG 1973. Selenium: biochemical role as a component of glutathione peroxidase. *Science* 179, 588–590.
- Smith KL, Hogan JS and Weiss WP 1997. Dietary vitamin E and selenium affect mastitis and milk quality. *Journal of Animal Science* 75, 1659–1665.
- Smith KL, Harrison JH, Hancock DD, Todhunter DA and Conrad HR 1984. Effect of vitamin E and selenium supplementation on incidence of clinical mastitis and duration of clinical symptoms. *Journal of Dairy Science* 67, 1293–1300.
- Underwood EJ and Suttle NF 1999. In the mineral nutrition of livestock, 3rd edition. CABI Publishing, Wallingford, UK.
- Ullrey DE 1987. Biochemical and physiological indicators of selenium status in animals. *Journal of Animal Science* 65, 1712–1726.
- Waldner C, Campbell J, Jim GK, Guichon PT and Booker C 1998. Comparison of 3 methods of selenium assessment in cattle. *Canadian Veterinary Journal* 39, 225–231.
- Walker GP, Dunshea FR, Heard JW, Stockdale CR and Doyle PT 2010. Output of selenium in milk, urine, and feces is proportional to selenium intake in dairy cows fed a total mixed ration supplemented with selenium yeast. *Journal of Dairy Science* 93, 4644–4650.
- Weiss WP 2005. Selenium sources for dairy cattle. In *Proceeding of the Tri-State Dairy Nutrition Conference*, Fort Wayne, IN, pp. 61–71.
- Weiss WP, Todhunter DA, Hogan JS and Smith KL 1990. Effect of duration of supplementation of selenium and vitamin E on periparturient dairy cows. *Journal of Dairy Science* 73, 3187–3194.
- Wright PL and Bell MC 1966. Comparative metabolism of selenium and tellurium in sheep and swine. *American Journal of Physiology* 211, 6–10.